Oxytocin reduces subjective fear in naturalistic social contexts via enhancing top-down middle

cingulate amygdala regulation and brain-wide multi-level fear representations

Authors

Kun Fu[1,2], Shuyue Xu[3,4], Zheng Zhang[1,2], Dan Liu[1,2], Ting Xu[5,6], Yuan Zhang[1,2], Feng Zhou[5,6],

Xiaodong Zhang[1,2], Chunmei Lan[1,2], Junjie Wang[1,2], Lan Wang[1,2], Jingxian He[1,2], Keith M

Kendrick[1,2], Bharat Biswal[2,7], Dezhong Yao[2], Zhen Liang[3,4]*, Weihua Zhao[1,2]*, Benjamin

Becker[1,2,8,9,10]*

Affiliations

[1] The Center of Psychosomatic Medicine, Sichuan Provincial Center for Mental Health, Sichuan

Provincial People's Hospital, University of Electronic Science and Technology of China, Chengdu,

China

[2] MOE Key Laboratory for Neuroinformation, School of Life Science and Technology, University of

Electronic Science and Technology of China, Chengdu, China

[3] School of Biomedical Engineering, Medical School, Shenzhen University, Shenzhen, China

[4] Guangdong Provincial Key Laboratory of Biomedical Measurements and Ultrasound Imaging,

Shenzhen, China

[5] Faculty of Psychology, Southwest University, Chongqing, China

[6] Key Laboratory of Cognition and Personality, Ministry of Education, Chongqing, China

[7] Department of Biomedical Engineering, New Jersey Institute of Technology, Newark, USA

[8] State Key Laboratory of Brain and Cognitive Sciences, The University of Hong Kong, Hong Kong,

China

[9] Department of Psychology, The University of Hong Kong, Hong Kong, China

[10] Lead contact

*Corresponding authors

bbecker@hku.hk (B. Becker, lead contact)

janezliang@szu.edu.cn (Z. Liang)

zarazhao@uestc.edu.cn (W. Zhao)

Abstract

Accumulating evidence from animal and human studies suggests a fear-regulating potential of the neuropeptide oxytocin (OT), yet the clinical translation into novel interventions for pathological fear requires a behavioral and neurofunctional characterization under close-to-real life conditions. Here, we combined a naturalistic fMRI-design inducing high and immersive fear experience in social and non-social contexts with a preregistered between-subjects randomized double-blind placebo-controlled intranasal OT trial (24 IU, n = 67 healthy men). OT selectively reduced subjective fear in social contexts but not in non-social contexts. Specifically, OT enhanced left middle cingulate cortex (lMCC) activation and its functional connectivity with the contralateral amygdala, with both neural indices significantly and inversely associated with subjective fear following OT. On the network level, OT enhanced communication between the dorsal attention network (DAN) with the fronto-parietal (FPN) and the default-mode network (DMN) as well as on the more fine-grained level brain-wide communication. Utilizing an independent activity-connectivity neuromarker for fear in naturalistic contexts (CAFE) confirmed that OT attenuated brain-wide fear expressions. Findings indicate an ecologically valid and social-specific fear-reducing potential of OT, highlighting its potential value as a treatment option for disorders characterized by excessive fear in social situations.

Keywords: Oxytocin, Subjective Fear, Naturalistic, fMRI, Social Anxiety

Introduction

Fear is an evolutionarily conserved and highly adaptive defensive avoidance response, fundamental to surviving and functioning in everyday life [1-4]. The experience of excessive and overwhelming fear represents a hallmark symptom of anxiety and trauma-related disorders [5-7]. These highly prevalent psychiatric disorders have become a leading cause of disability and are accompanied by personal suffering as well as an enormous socio-economic burden. The conventional pharmacological interventions targeting the classical neurotransmitter (GABA-ergic, serotonergic) systems [7-9] are characterized by moderate response rates and potential negative side effects ranging from dizziness towards addiction. While several promising candidate compounds have shown effects in animal models, no new pharmacological treatment for fear or anxiety has entered the clinical practice during the last decades (for recent non-pharmacological developments see e.g. [10,11]).

Based on promising findings in animal models and preclinical human studies neuropeptides, including oxytocin (OT), vasopressin or angiotensin II, have been suggested as promising new strategies to regulate fear and anxiety [12-16]. Most previous research in this field has focused on OT, a neuropeptide produced in the hypothalamus, and animal models [15,16] as well as preclinical human proof-of-concept studies demonstrated promising effects of OT on regulating negative emotional processing in the domains of fear and anxiety [16-20]. Some of the effects of OT appear to be specific to social contexts and stimuli, yet in the context of contradictory empirical evidence, debates on the social-specific effects of OT continue [21,22].

OT neurons from the hypothalamus project to a wide range of brain regions involved in fear, anxiety as well as social processes, including the amygdala, hippocampus, midbrain and frontal lobe [5,23], and previous studies indicate that intranasal OT can regulate these processes via influencing activity in the amygdala, dorsal anterior cingulate gyrus and medial prefrontal lobe [7,18-20,24-26]. However, despite the initially promising findings from these studies, the fear and anxiety-reducing effects of OT and the underlying neural circuits have been determined under laboratory experimental contexts that strongly differ from the dynamic contexts during which fear arises in everyday life [27]. Specifically, these studies combined the intranasal administration of OT with sparsely presented static experimental stimuli (e.g. pictures of fearful faces or scenes) and primarily focused on treatment-induced changes in brain regional activity during fMRI.

This approach stands in contrast to contemporary neuroscience models proposing that fear emerges in interaction between subjective appraisal and dynamic changes in the environment [28,29] and that the corresponding mental states are mediated by an intricate interplay among distributed yet interacting brain systems [30-32]. To account for the complexity of subjective affective experiences under ecologically valid conditions and to capture the underlying network-level communication in the brain, recent overarching conceptualizations have proposed the

combination of movie stimuli as highly immersive and dynamic stimuli with network-level

analyses [33,34]. Initial empirical studies confirmed that these stimuli can induce highly immersive

affective experiences in complex social situations [35,36] as well as the essential role that network-

level communication plays in establishing affective and social experiences under naturalistic

contexts [27,34,37,38].

Mental disorders, including fear-related disorders, have been reconceptualized as network-

level disorders that are characterized by dysregulated regional activity as well as network level

connectivity (putatively reflecting communication between brain regions) [32,39,40]. Together with a

growing number of studies underscoring the potential of OT to regulate neurofunctional network-

level interactions in fear-related networks across species [41-44], which may be partly explained by

the physiological properties of the oxytocin-signaling system, including an axonal release,

widespread receptor distribution and considerably longer half-life compared to classical

neurotransmitters [45], these findings indicate the necessity to account for network-level effects to

fully determine a regulatory effect of OT on fear experience and its therapeutic potential.

Against this background, the present pre-registered between-subjects randomized double-

blind placebo-controlled pharmaco-fMRI study (n = 67 healthy male participants) capitalized on

recent progress in naturalistic fMRI and network-level analyses to determine whether a single

dosage of intranasal OT (1) reduces the subjective experience of fear under highly immersive and

ecologically valid conditions, (2) modulates concomitant regional activity in brain systems

identified in previous studies, (3) modulates brain communication of the identified regions and

on the whole-brain network level, and (4) effects vary as a function of social context. Based on

previous studies [7,18-20] we hypothesized that intranasal OT would decrease subjective fear in

naturalistic contexts with more pronounced effects in the social domain while concomitantly

enhancing neural activity in regulatory brain regions and inhibiting activity in fear-processing

brain regions. On the network level, we expected that OT would enhance communication

between emotion-regulation brain regions and the amygdala as well as communication between

large-scale networks with pronounced effects in the social fear context.

Results

Demographics and potential confounders.

124   The OT (n = 33) and PLC (n = 34) groups were comparable with respect to socio-demographics,

125   mood and mental health indices arguing against nonspecific treatment effects (**Table 1**, all *ps* >

126   0.10).

Table 1. Demographics and potential confounders in the two treatment groups.

	Time	OT(M ± SD)	PLC(M ± SD)	t	p
Age		21.30 ± 2.05	20.76 ± 1.83	1.13	0.26
BDI II	Before	4.70 ± 3.83	5.38 ± 5.40	-0.60	0.55
BAI	Before	2.36 ± 2.56	2.53 ± 3.75	-0.21	0.83
LSAS	Before	31.36 ± 20.35	26.15 ± 19.47	1.07	0.29
SAI	Before	34.79 ± 6.63	33.15 ± 9.37	0.83	0.41
TAI	Before	37.82 ± 7.30	37.97 ± 9.99	-0.07	0.94
ERQ-Reappraisal	Before	31.00 ± 3.98	30.74 ± 4.15	0.27	0.79
ERQ-Suppression	Before	15.85 ± 4.72	14.82 ± 4.41	0.92	0.36
PANAS-P	Before[1]	28.85 ± 6.05	28.47 ± 7.17	0.23	0.82
	2 - 1	-6.12 ± 6.71	-4.06 ± 5.18	-1.41	0.16
	Treatment peak[2]	22.73 ± 5.95	24.41 ± 6.90	-1.07	0.29
	3 - 2	-2.85 ± 5.50	-1.62 ± 5.39	-0.93	0.36
	After experiment[3]	19.88 ± 6.71	22.79 ± 7.71	-1.65	0.10
	3 - 1	-8.97 ± 7.12	-5.68 ± 7.53	-1.84	0.07
PANAS-N	Before[1]	15.09 ± 5.17	14.09 ± 4.82	0.82	0.41
	2 - 1	-3.24 ± 4.70	-3.06 ± 4.06	-0.17	0.87
	Treatment peak[2]	11.85 ± 3.17	11.03 ± 2.32	1.21	0.23
	3 - 2	1.33 ± 2.97	1.65 ± 2.62	-0.46	0.65
	After experiment[3]	13.18 ± 4.15	12.68 ± 3.86	0.52	0.61
	3 - 1	-1.91 ± 6.07	-1.41 ± 5.51	-0.35	0.73

Values presented here as mean ± standard deviation (M ± SD). *BDI II* Beck Depression Inventory II, *BAI* Beck Anxiety Index, *LSAS* Liebowitz Social Anxiety Scale, *STAI* Spielberger State-Trait Anxiety Inventory, *ERQ* Emotion Regulation Questionnaire, *PANAS* Positive and Negative Affect Schedule, *OT* oxytocin, *PLC* placebo.

OT reduced subjective fear experience in social fear naturalistic contexts.

The ANOVA revealed a significant main effect of emotion ($F_{(1,65)} = 1434$, $p < 0.0001$, $\eta_p^2 = 0.957$, **Fig. 1a**) such that participants rated subjective fear considerably higher during fear trials (FS and FNS) than neutral ones, and a significant main effect of social ($F_{(1,65)} = 76.81$, $p < 0.0001$, $\eta_p^2 = 0.542$, **Fig. 1a**) reflecting lower fear during social trials (FS and NS) compared with non-social trials. With respect to the effects of OT we found a significant interaction effect between treatment and emotion ($F_{(3,65)} = 4.064$, $p = 0.048$, $\eta_p^2 = 0.059$, **Fig. 1a**), with post hoc tests indicating that OT significantly decreased fear experience in response to fearful stimuli in naturalistic social contexts but not in non-social contexts ($t_{(1,65)} = -2.252$, $p = 0.028$, Cohen's $d = -0.55$ under FS while $t_{(1,65)} = -1.475$, $p = 0.145$, Cohen's $d = -0.36$ under FNS, **Fig. 1b**). Finally, the interaction between emotion and social was significant ($F_{(3,65)} = 53.15$, $p < 0.0001$, $\eta_p^2 = 0.45$, **Fig. 1a**) reflecting that participants reported generally lower fear experience in social fear contexts. The main effect of treatment and other interactions were not significant. Together the results partly support that OT can reduce subjective fear in naturalistic social fear situation.

Fig. 1 The effects of OT on behavioral fear rating and cingulate activity. a OT reduced subjective fear rating in social fear naturalistic situation, **b** OT effects on fear rating under four conditions separately, **c** OT but not PLC could strengthen the activation in the left middle cingulate cortex (lMCC) under the interaction of valence and social. Particularly, **d** OT specifically increased the activation in lMCC under FS but not other conditions. **e** The activation in lMCC was significantly negatively correlated with subjective fear experience in OT group but not PLC group. OT oxytocin, PLC placebo, FS fear social, FNS fear non-social, NS neutral social, NNS neutral non-social. **** $p_{\text{FDR-corrected}} < 0.0001$, ** $p_{\text{FDR-corrected}} < 0.01$, * $p_{\text{FDR-corrected}} < 0.05$, ns not significant.

OT enhanced middle cingulate activation in the social fear contexts.

We initially examined brain regions participating in fear processing under naturalistic contexts with different social features compared with neutral control conditions, a series of corresponding voxel-wise one-sample t-tests conducted in the PLC group confirmed previous studies, suggesting that visual and temporal regions were strongly engaged during exposure to dynamic fear movie clips irrespective of social features (**Fig. S3**). Examining treatment effects revealed a significant interaction between treatment × emotion × social located in the left middle cingulate cortex (peak MNI, x, y, z = -7, -3, 30, $t_{(1,65)}$ = 3.67, k = 33, $p_{\text{FWE-peak(svc)}}$ = 0.019, **Fig. 1c**), with post-hoc t-tests on extracted activity measures from independently defined (atlas-based) lMCC indicating that compared to PLC, OT selectively enhanced activation in this region for FS ($t_{(1,65)}$ = 2.69, p = 0.009, **Fig. 1d**) but not other conditions (all ps > 0.05, **Fig. 1d**). Examining associations between lMCC activity (extracted parameter estimates from the atlas region) and subjective fear experience revealed a significant negative association in the OT (r = -0.387, p = 0.026, **Fig. 1e**) but not in PLC group (r = 0.015, p = 0.934, **Fig. 1e**). A Fisher r-z transformation indicated that

these correlation coefficients were significantly different between the OT and PLC groups (Z = -

1.65, one-tailed p = 0.0495).

OT enhanced MCC-amygdala functional communication consistently in the social

fear contexts.

Examining OT effects on the functional communication of the lMCC revealed a significant

interaction effect between treatment × emotion × social located in the right amygdala (rAmy)

(peak MNI, x, y, z = 22, -7, 13, $t_{(1,65)}$ = 3.81, k = 19, $p_{FWE-peak(svc)}$ = 0.021, **Fig. 2a**), with post-hoc t-

tests indicating that OT relative to PLC selectively enhanced lMCC-rAmy coupling in social fear

contexts ($t_{(1,65)}$ = 2.69, p = 0.009, **Fig. 2b**) but not other contexts (all ps > 0.05, FDR-corrected,

Fig. 2b), confirming social fear-specific effects of OT on the neural level. In addition, we found

that lMCC-rAmy functional coupling was significantly negatively associated with subjective fear

experience in the OT (r = -0.426, p = 0.013, **Fig. 2c**) but not PLC group (r = -0.026, p = 0.883, **Fig.**

2c, Fisher r-to-z transformation revealed a significant difference between the treatment groups,

Z = -1.68, one-tailed p = 0.0465).

Fig. 2 OT strengthened lMCC-rAmy communication specifically in social fear contexts. a Interaction effect between treatment × emotion × social on lMCC connectivity with the right amygdala **b** OT specifically increased functional connectivity between lMCC and rAmy in the FS contexts but not other conditions and **c** The functional connectivity between lMCC and rAmy was significantly negatively correlated with subjective fear experience in OT group but not PLC group. OT oxytocin, PLC placebo, FS fear social, FNS fear non-social, NS neutral social, NNS neutral non-social. ** $p_{FDR-corrected}$ < 0.01, * $p_{FDR-corrected}$ < 0.05, # $p_{un-corrected}$ < 0.05.

OT enhanced FPN-DAN and DMN-DAN coupling during social fear contexts.

Examining the effects of OT on the large-scale network level using separate two-sample t-tests on

the matrices of z-values for the two main contrasts of interest (FS − NS, FNS − NNS) and

following mantel test revealed that OT significantly affected network functional connectivity

differently under two contexts ($rho_{mantel\ test}$ = -0.1333, $p > 0.05$). Specifically, compared to PLC (all

ps > 0.05, **Fig. 3f**), OT significantly enhanced the functional communication between FPN and

DAN ($t_{(1,65)}$ = 2.75, p = 0.0077, Cohen's d = 0.672, **Fig. 3e**) as well as between DMN and DAN

($t_{(1,65)}$ = 2.02, p = 0.048, Cohen's d = 0.493, **Fig. 3e**) under the social fear but not the non-social

fear contexts.

Fig. 3 OT enhanced functional connectivity on the level of large-scale networks. **a b c d** Flow of z-value matrix calculation for the degree of response functional connectivity under specific contrasts. (1) The Fisher r-z matrices of each condition for each participant were subtracted to yield a new matrix of z-value corresponding to the contrast (e.g. $z_{FS − NS} = z_{FS} − z_{NS}$). (2) New matrices of z-value corresponding to a specific contrast were generated for each participant in each group, which were used as features in a two-sample t-test to derive the between-group differences under the specified contrast. **e** OT significantly strengthened the functional connectivity between FPN and DAN as well as DMN and DAN in the fear social-related contrast **f** but not in the fear non-social-related contrast. OT oxytocin, PLC placebo, FS fear social, FNS fear non-social, NS neutral social, NNS neutral non-social, VN visual network, SMN somatomotor network, DAN dorsal attention network, VAN ventral attention network, LN limbic network, FPN frontoparietal network, DMN default mode network, SCT subcortical network, B/C brainstem and cerebellum. ##$p_{un-corrected} < 0.01$, # $p_{un-corrected} < 0.05$.

OT modulated whole-brain functional connectivity in the social fear context.

We further conducted exploratory analyses to examine whether OT changes brain-wide

communications. Results showed that while under PLC only few positive connections increased

activity during the FS versus NS contrast (**Fig. 4a**), OT induced a more widespread connectivity

between several brain regions (**Fig. 4b**). Under the FNS versus NNS contrast, the PLC group

exhibited wide-spread connectivity (**Fig. 4c**), with no obvious changes following OT (**Fig. 4d**).

Further Mantel tests and Welch's t-test were conducted on the t-value matrices (FS − NS and

FNS − NNS) between both groups showed significant modulating effects of OT under both

contrasts (rho$_{mantel\ test}$ = 0.0335, $t_{(1,65)}$ = 9.05, $p < 0.0001$ under FS − NS, rho$_{mantel\ test}$ = 0.208,

$t_{(1,65)}$ = -2.51, $p = 0.012$ under FNS − NNS), particularly under the fear social contexts.

Fig. 4 Comparison between treatment groups for whole-brain functional connectivity during social and non-social fear separately. Under fear social contexts, **a** PLC group showed a sparse connectivity between brain regions **b** the OT group exhibited stronger functional interactions at the whole-brain level. In contrast, under the fear non-social context **c** PLC and **d** OT groups showed similar whole-brain functional connectivity patterns. OT oxytocin, PLC placebo, FS fear social, FNS fear non-social, NS neutral social, NNS neutral non-social, VN visual network, SMN somatomotor network, DAN dorsal attention network, VAN ventral attention network, LN limbic network, FPN frontoparietal network, DMN default mode network, SCT subcortical network, B/C brainstem and cerebellum.

OT attenuates the response of a synergistic brain connectivity- and activity-based

signature for subjective fear (CAFE)

To comprehensively determine and independently validate the social-specific effects of OT, we

utilized a recently developed neuromarker that accurately tracks subjective fear in naturalistic

contexts by synergistically capitalizing on activity and connectivity[27]. The CAFE succesfully

determined fear versus no-fear brain responses in the entire group with high accuracy (FS vs. NS:

accuracy = 97±2.1%, area under the curve (AUC) = 1.00, P < 0.0001; FNS vs. NNS: accuracy =

100±0%, AUC = 1.00, P < 0.0001; two-alternative forced-choice test, **Fig. 5a**) validating its high

potential to track subjective fear in dynamic contexts. In line with the univariate activation

analysis and functional connectivity analysis showing that neural distinctions between treatment

were only evident in the social fear contexts(FS—NS contrast), the whole-brain MVPA results revealed that the significant classification of PLC and OT treatments was demonstrated a 66% accuracy in the FS—NS contrast (mean$\pm$SD of accuracy = 66$\pm$5.8%, p = 0.02 , area under the curve (AUC) = 0.62, **Fig. 5b left pannel**), but not in the FNS—NS contrast (mean$\pm$SD of accuracy = 63$\pm$5.9%, p = 0.06, AUC = 0.60, **Fig. 5b right pannel**).

Fig. 5 CAFE classification results. a CAFE successfully classify fear with neutral in entire group with high accuracy. b Treatment classification results of CAFE response in social and non-social fear contexts. OT oxytocin, PLC placebo, FS fear social, FNS fear non-social, NS neutral social, NNS neutral non-social.

Discussion

The present study determined the acute behavioral and neural effects of a single intranasal OT administration on subjective fear experience in a naturalistic context that resembles dynamic fear processing in social and non-social contexts in a close-to-real life setting. On the behavioral level, OT selectively decreased the experience of subjective fear in social fear contexts, supporting a fear-regulating potential of OT under dynamic and ecologically valid social conditions. In the meanwhile, neural effects of OT were also primarily observed in social fear contexts, such that OT enhanced regional activation in the left MCC and its communication with the contralateral amygdala specifically during fear in social contexts with both neural indices exhibiting a negative association with subjective fear following OT but not PLC. Exploratory network-level analyses further revealed that OT significantly enhanced the functional connectivity between the DAN and the FPN/DMN, respectively, and brain-wide communication selectively in social fear contexts. OT additionally inhibited the CAFE response in social fear contexts but not in non-social fear

contexts, which independently validates that OT regulates the whole-brain expression of fear

under ecologically valid conditions and highlights the potential of multivariate neuroaffective

signatures to facilitate treatment evaluation utilizing precision-pharmaco-neuroimaging[13,46].

Together, the findings indicate that a single dose of intranasal OT has the potential to decrease

fear in close-to-real life settings by enhancing lMCC activation and its communication with the

amygdala while concomitantly affecting network-level communication of the DAN with brain

networks involved in regulatory control (FPN [47,48]) and social processing (DMN [49,50]) as well as

brain-wide functional communication and whole-brain fear expressions.

Several previous studies in rodents and humans suggested a fear- and anxiety-reducing

potential of OT, however, results remained inconsistent. Studies have described a promising fear-

and anxiety-reducing neurofunctional profile of OT across species, yet the neural effects on

amygdala-related circuits were often observed in the absence of changes in subjective fear

experience or under experimental laboratory settings with limited ecological validity [15,17-20,51].

In the context of increasing attempts to enhance the ecological validity of neuroimaging

research using dynamic movie stimuli [27,28,52], we here combined naturalistic fMRI employing fear-

inducing movie clips with a pre-registered randomized double-blind placebo-controlled OT-

administration trial. Across pilot experiments and treatment studies, the procedure induced a

highly immersive and comparably strong experience of fear under close-to-real life conditions.

Importantly, OT decreased subjective fear experience in the more naturalistic context with

moderate effect size in the social context. Although the pronounced fear-reducing effects of OT in

the social context may partly reflect the generally more moderate fear experience reported by

our participants in this context, pronounced effects of OT in social contexts have been

conceptualized in a number of overarching theories and have been reported in several previous

studies [21,22,53] including studies reporting decreased subjective anxiety [54]. The latter hypothesis

on the social-specific effects of OT is also mirrored in the brain functional findings, such that the

neurofunctional effects were selectively observed during fear experiences in social contexts.

With respect to regional activation, OT increased the activation of the left middle cingulate

cortex (lMCC). The MCC represents a highly integrative hub that has been involved in both, fear

evaluation [5,55-57] as well as the cognitive control of negative affective states including fear [58,59].

Concomitantly, OT enhanced functional connectivity from the lMCC to the contralateral amygdala

(rAmy), a pathway that has been previously involved in implementing top-down inhibitory control

over excessive fear-related amygdala engagement [55,57,60]. Effects of OT on this circuitry have been

previously reported in the context of reduced subjective anxiety in individuals with PTSD [61] and in

response to static fearful faces in patients with generalized social anxiety disorders [25]. Exploring

brain-behavior associations further revealed that OT established a negative association between

higher activity and connectivity in this circuit and subjective fear, underscoring the behavioral

relevance of the OT-induced neurofunctional changes. Together, our findings suggest that OT

exerts its fear-regulating effects under naturalistic social fear contexts via enhancing MCC

engagement and control over the amygdala.

On the large-scale network level, OT significantly enhanced the functional connectivity of

the DAN with the FPN as well as the DMN in the social fear contexts. The DAN is centered around

the intraparietal sulcus and the frontal eye fields and plays a pivotal role in top-down or goal-

directed control of attention [62-64], while the FPN supports goal-directed actions and cognitive

control [47,65,66] and the DMN supports self-referential [67,68] and social processing [49,50], with more

recent findings additionally suggesting a role in emotional experience in dynamic naturalistic

contexts [69,70]. While studies have demonstrated the effects of OT on the interaction between

these large-scale networks during the task-free state [71,72], the present findings underscore the

behavioral relevance of these effects of OT in terms of regulating emotional processes in social

contexts. Together with previous studies suggesting an association between dysfunctional

interaction between these large-scale networks and emotional dysregulations in anxiety and

fear-related disorders [32,68], the present findings support a potential therapeutic potential of OT

for these disorders [32,73,74].

Network-level analyses that explored the effects of OT on the fine-grained level of brain-

wide connectivity patterns suggested that OT may broadly facilitate brain-wide exchange of

information within and across networks particularly between the FPN and DMN in social fear

contexts. The results align with recent findings suggesting that emotional processes such as fear

and anxiety are processed in distributed networks and their interactions [27,34,75] as well as the

distributed network-level properties and effects of the OT signaling system [41,76,77]. Findings on the

whole-brain effects were additionally validated using a precision pharmaco-neuroimaging

approach which capitalizes on MVPA-based decoding. To this end we employed an independently

developed fMRI-based marker that accurately tracks subjective fear in naturalistic contexts

(CAFE[27]). The marker precisely tracked subjective fear under ecologically valid contexts in both

groups, yet showed an attenuated reactivity in the OT group, in particular in the social fear

context. Independent whole-brain-level MVPA analysis indicated that OT has the potential to

diminish distributed fear pattern. These findings underscore that OT may exert its effects on

human emotion processing under complex naturalistic conditions via regulating extensive brain

wide interactions. Together with icurrent reconceptualizations proposing that emotional

processes are represented in brain-wide distributions[75,78] and that mental disorders are not

related to dysfunctions of a single brain region or pathway but rather dysregulated brain-wide

communication [32,68,79], these findings further underscore the therapeutic potential of the OT

signaling system.

Findings have to be considered in the context of limitations of the present design, including

(1) to avoid variations in the effects of OT related to sex [80,81], we only focused on male

participants and future studies need to test generalization to females, (2) the study employed a

proof-of-concept design in healthy subjects and future studies need to validate the effects of OT

in populations with pathological fear and anxiety, (3) despite extensive pre-study testing the

videos of the social and non-social category differed in terms of the level of fear induced which

partly limits conclusions regarding social-specific effects of OT, (4) the videos included

homogenous brief episodes of fear and future studies need to examine effects of OT on fear

during longer time scales and longer episodes with dynamic fear [27,28].

Taken together, this study demonstrated that OT can reduce subjective fear experience

under naturalistic social fear conditions via enhancing lMCC engagement and its regulatory

control over the amygdala as well as enhancing communication between large-scale networks

and brain-wide communication. These findings consistently indicate a high translational

therapeutic potential of OT to regulate fear in dynamic and close-to-real life environments.

Effects were pronounced or restricted to the social contexts, supporting the role that OT may play

in social contexts, and rendering OT as a promising treatment for social context-specific fear

experiences, e.g. in social anxiety disorders. Findings indicate that OT may be used as a novel

therapeutic strategy to regulate excessive subjective fear symptoms in mental disorders [7,17,24] or

may help to prevent the development of these disorders during early stages or promote long-

term recovery [72] via improving the quality of life, and reduce the social and economic burden

related to fear-related disorders.

Methods

Participants

Sixty-nine right-handed male participants with normal or corrected-normal vision were recruited

from the University of Electronic Science and Technology of China. The sample size was

determined using G*Power v.3.1 [82,83] indicating that 62 participants were required to detect a

significant effect on the behavioral level ($f = 0.25$, $\alpha = 0.05$, $\beta = 0.80$, $2 \times 2 \times 2$ mixed ANOVA

interaction effect). The present study focused on male subjects to reduce variance related to sex

differences in the effects of OT [80,81] and adhered to validated exclusion criteria (details see

supplementary methods). The final sample size was n = 67 (mean $\pm$ SD, age = 21.03 $\pm$ 1.95

years) due to two data from two participants being excluded (withdraw due to personal reasons,

technical MRI issues). All participants provided written informed consent, protocols were pre-

registered on Clinical Trials.gov (clinicaltrials.gov/ NCT05892939), approved by the ethics

committee at the University of Electronic Science and Technology of China (UESTC-

1061423041725893) and in line with the latest Declaration of Helsinki.

Using a double-blind randomized, placebo-controlled, between-subject pharmacological

fMRI design, participants were randomly assigned to receive either a single intranasal dose of

oxytocin (OT, 24IU-Sichuan Defeng Pharmaceutical Co. Ltd, China) or a placebo (PLC, same

ingredients i.e. sodium chloride and glycerin but without the peptide). The spray bottles were

identical and dispensed by an independent researcher based on a computer-generated

randomization sequence to ensure double-blinding. The investigators involved in data acquisition

and analyses were blinded for group allocation. Following recent recommendations [84-86], the fMRI

assessment began 45 minutes after treatment administration.

To control for pre-treatment between-group differences, participants completed a series of

mood and mental health questionnaires (as detailed in **Table 1**). Participants were asked to guess

which treatment they had received after the experiment (with a non-significant treatment guess

$\chi^2 = 0.39$, $p = 0.53$, confirming successful double-blinding). Additionally, participants were

required to complete a surprise recognition task after the experiment that involved presenting

stills of the video clips shown during fMRI and new clips to test the attentive processing of the

stimuli. All participants achieved a re-recognition accuracy of over 75% and no significant

difference in performance was observed between the two treatment groups ($t_{(1,65)} = -0.693$, $p =$

0.491).

Naturalistic fear induction paradigm.

The task began 45 minutes after the treatment intervention (see **Fig. 6a**). During the naturalistic

fear induction paradigm, participants were instructed to watch a series of video clips attentively

and rate their level of fear experience after each clip. The clips lasted 25 seconds and were

followed by a 1-3 second jittered fixation cross to separate the stimuli from the rating period.

During the 5-second period following each stimulus, participants were asked to report their level

of subjective fear on a 9-point Likert scale, with 1 indicating no fear and 9 indicating very strong

fear. This was followed by a jittered inter-trial interval of 9-11 seconds, during which a fixation

cross was presented (see **Fig. 6b**). A total of 32 short video clips were presented over 2 runs, with

each run consisting of 16 randomly presented clips, half of which were neutral and half of which

were fear-inducing clips.

Given ongoing discussions about the social-specific effects of OT [87,88], we carefully pre-

selected a balanced set of stimuli that included social as well as non-social fearful or neutral

stimuli, respectively. This led to a set of four stimulus categories (fear social clips-FS, fear non-

social clips-FNS, neutral social clips-NS, neutral non-social clips-NNS, see **Fig. 6c**). For detailed

characterization of the video clips, see Supplementary Methods. The video clips used for the

naturalistic fear induction paradigm were selected from a larger database based on stimuli

evaluation ratings in an independent sample (n = 20; 10 females; age = 22.65 ± 2.01). Key

exclusion criteria for the video clips were reported in supplementary methods. Based on the

results of the behavioral ratings (see **Fig. S2**), 8 complimentary video clips were selected for each

of the four categories. The task was programmed in Python 3.7 using the PsychoPy package

(version 2022.2.4).

Fig. 6 Experimental timeline, paradigm and schematic depiction of stimuli. a The timeline of the whole experiment. **b**
Naturalistic fear induction and rating paradigm. **c** Example stimuli. In each run, 4 types of video clips were presented (FS, FNS,
NS, NNS) according to a randomized order. Abbreviations: OT oxytocin, PLC placebo, fMRI functional magnetic resonance
imaging, FS fear social, FNS fear non-social, NS neutral social, NNS neutral non-social. License: Images in 1a and 1b were
obtained from Flaticon.com under the free license with attribution.

Slected clips could reduce robust subjective fear feeling .

To initially ensure that the selected video clips were effective in inducing the intended emotions,

repeated measures ANOVA (sex was used as the independent variable, while emotion and social

were used as repeated variables) were conducted on five rating metrics (see supplementary

methods). Analyses were conducted using GraphPad Prism 9.0 (GraphPad Software, Inc., La Jolla,

CA).

Effects of OT on subjective fear experience.

To determine the effects of OT on subjective fear in naturalistic contexts and whether they vary

as a function of social context, we computed a repeated measures ANOVA with the between-

subject factor treatment (OT vs PLC) and the within-subject factors emotion (fear vs neutral) and

social (social vs non-social) and the dependent variable fear ratings during fMRI. Analyses were

379 conducted using GraphPad Prism 9.0 (GraphPad Software, Inc., La Jolla, CA), and corresponding

380 effect sizes were computed using JASP 0.18.3.0 (JASP Team, 2019; jasp-stats.org). All ANOVAs and

381 t-tests used false discovery rate (FDR) [89] recommended by Prism for multiple comparison

382 correction.

MRI data acquisition and preprocessing.

MRI data were collected on a 3.0-T GE Discovery MR750 system (General Electric Medical System,

Milwaukee, WI, USA) and preprocessed using FMRIPREP 21.0.0[90] (RRID: SCR_016216), a Nipype

1.6.1 based tool that integrates preprocessing routines from different software packages and

SPM 12 (Statistical Parametric Mapping; http://www.fil.ion.ucl.ac.uk/spm/; Wellcome Trust

Centre for Neuroimaging, detailed in supplementary methods).

Next, a voxel-wise general linear model (GLM) was conducted for each participant.

Specifically, a first-level model was designed that included separate regressors for the four

conditions and the rating period. In line with a previous study [28], we removed variance associated

with the mean, linear and quadratic trends, the average signal within anatomically-derived CSF

mask, the effects of motion estimated during the head-motion correction using an expanded set

of 24 motion parameters (six realignment parameters, their squares, their derivatives, and their

squared derivatives) and motion spikes (FMRIPREP default: FD > 0.5mm or standardized

DVARS > 1.5)[91].

Effects of OT on fear-related brain activity.

To examine the effects of OT on fear-related neural activity in naturalistic contexts, a GLM-based

model was established in SPM 12. The model incorporated five regressors: FS, FNS, NS, NNS, and

the rating period. We included the 24 head motion parameters as covariates of no interest. Key

first-level contrasts of interest were contrasts that modeled fear in social and non-social contexts

separately (i.e., [FS — NS] or [FNS — NNS]) as well as their interaction (i.e., [FS — NS]

— [FNS — NNS]). In line with the behavioral analyses, the effects of OT were determined using an

ANOVA with the factors emotion and social context, implemented in a partitioned error ANOVA

approach that subjected the first level interaction contrast to a voxel-wise two-sample t-test with

the factor treatment group on the second level. Based on our hypothesis and previous studies

reporting that fear is strongly associated with amygdala, cingulate cortex (particularly middle

cingulate cortex, MCC), insula, and ventromedial prefrontal cortex (vmPFC) and that oxytocin

mediates its anxiolytic effects via these regions [14,22,27,55,92,93], our analyses focused on atlas-

derived bilateral masks for the amygdala, MCC, insula and vmPFC using separate small volume

correction (SVC) and family-wise error (FWE) correction [94] applied at peak level.

To further disentangle complex interaction effects involving treatment and for display

purpose, parameter estimates were extracted using independent left MCC mask[95] using Marsbar

(http://marsbar.sourceforge.net/). The extracted Blood Oxygenation Level Dependent (BOLD)

signals were then subjected to appropriate post-hoc tests comparing group differences in each of

the separate conditions.

Additionally, to determine associations between the behavioral (subjective fear) and neural

effects of OT (both activation and functional connectivity between specific brain regions) we

conducted correlation analysis using JASP.

Effects of OT on functional communication of the regions identified.

We further examined whether the regions showing altered regional activation following OT also

changed their network-level communication via a general Psychophysiological interaction (gPPI)

analysis. gPPI examines functional connectivity between brain regions that is contingent on a

psychological context. We used the Generalized PPI toolbox [96] as implemented in SPM 12, to

conduct gPPI. This method includes additional task regressors to reduce the likelihood that the

functional connectivity estimates are driven simply by co-activation and has been widely used in

similar studies [97,98]. Based on our results, OT enhanced activation in the left middle cingulate

cortex (lMCC) under the three-way interaction (treatment × emotion × social), we therefore

computed the functional communication of this region (using an 8mm radius which centered in

peak Montreal Neurological Institute (MNI), x, y, z = -7, -3, 30) with other voxels in the whole

brain under the two-way interaction of emotion and social, and a two-sample t-test was

conducted to confirm the treatment effect of OT. Based on previous studies reporting effects of

OT on the functional communication of the amygdala [80,81], separate atlas-derived bilateral

masks[95] for the amygdala, MCC, insula and vmPFC were used for SVC analyses with FWE

correction [94] applied at peak level.

Marsbar was used to extract the lMCC - rAMY functional coupling using independent rAMY

mask[95] to further disentangle complex interaction effects involving treatment. The extracted

functional connectivity signals were then subjected to compare the treatment effect under four

conditions by using two-sample t-tests separately.

Effects of OT on large-scale brain network functional connectivity.

Given that several studies reported the effects of OT on large-scale network organization of the

brain [71,99], we examined OT effects on the whole-brain network level. To this end, time-series of

each region of interest (ROI) was extracted according to the template including 463 ROIs across

the whole brain (see supplementary methods), resulting in BOLD time series of the size of $r \times t$ (r

was the number of ROIs and t was the BOLD time series length) for each subject per run. A matrix

of size $r \times r$ was obtained by calculating Pearson's correlation between the average BOLD time

series of each pair of ROIs. Additionally, Fisher's z-transform was applied to improve the

normality of the correlation coefficients. The brain regions can be divided into nine networks

based on their functional divisions, and the neural activities of each network's regions were

averaged to calculate network functional connectivity.

During each run, four types of video clips were presented in a pseudo-randomized order. The

onset time of each video clip was shifted by 3 TRs to account for hemodynamic lag. To extract

neural activity time series for a specific condition, we used the time points and durations

specified for the appearance of different trials of the same type. This resulted in four $r \times d$

matrices (where r is the number of ROIs and d is the BOLD time series length of each condition).

Additionally, we obtained the $r \times r$ matrix between ROIs and the $n \times n$ matrix (where n is the

number of networks) between brain networks for different conditions.

After performing Fisher-Z transformation on the network functional connectivity matrices

for different conditions, matrices of Z-values corresponding to specific conditions

($Z_{FS}\backslash Z_{FNS}\backslash Z_{NS}\backslash Z_{FNS}$) are obtained. It is important to note that r values are not additive or

subtractive, but Z scores are summative. Matrix of Z-values corresponding to particular contrasts

(e.g. Z_{FS-NS}) can be computed by subtract the matrices of Z-values corresponding to different

conditions (e.g. $Z_{FS}-Z_{NS}$)[100,101]. Therefore, each participant in both groups would have their own

subtracted Z-values matrix under specific contrasts ($Z_{FS-NS}\backslash Z_{FNS-NNS}$). We then conducted two-

sample t-tests on each pair of network-level functional connectivity to determine the treatment

effect.

Effects of OT on whole-brain functional connectivity.

To further investigate the impact of OT on a finer spatial scale, we analyzed whole-brain

functional connectivity using paired t-tests for key contrasts (FS − NNS and FNS − NNS)

separately for the two groups. We applied FDR correction ($p_{FDR-corrected} < 0.05$) to identify the

most relevant functional connections for each contrast. A Mantel test [102,103] was performed on

the FDR-corrected functional connectivity matrices of the two groups, under specific contrasts, to

assess their similarity. Furthermore, to investigate treatment-specific effects on functional

connectivity strength at the whole-brain level, we transformed the upper triangular data of the t-

value matrices of the two groups under specific contrast into two $(r \times (r-1)) / 2$ by 1 feature

vectors, and performed Welch's t-test between them.

Effects of OT on synergistic brain connectivity- and activity-based signature for

subjective fear (CAFE).

We applied a recently developed whole-brain multivariate neuromarker for fear in naturalistic

contexts (CAFE[27]) which synergistically capitalizes on brain connectivity- and activity-based

signature to accurately track subjective fear in dynamic naturalistic contexts[27] to independently

validate effects of OT on the whole-brain expression of fear. First, we calculated the mean beta value

vector(one for each region; n=463 in total, see supplemental methods in detail) and functional

connectivity matrix (463×463) for each subject under the four conditions (FS\FNS\NS\NNS) and

applied dot-product the integrated one-dimensional vector (n = 107416 features in total) with the

CAFE weights to calculate the CAFE response for each subject under each of the four conditions.

Second, in line with the other analyses we build the contrasts of interest (FS-NS; FNS-NNS) and

calculated CAFE response of each contrast. To quantify the classification performance of the model

(classification performance between PLC and OT under same contrast condition), Receiver

Operating Characteristic (ROC) analysis was conducted.

Acknowledgement

This work was supported by the China MOST2030 Brain Project (Grant No. 2022ZD0208500-BB), the National Natural Science Foundation of China (Grant No. 82271583-BB), the National Key Research and Development Program of China (Grant No. 2018YFA0701400-BB), the Natural Science Foundation of Sichuan Province (Grant number 2022NSFSC1375-WZ) and a start-up grant from The University of Hong Kong. Disclaimer: Any opinions, findings, conclusions or recommendations expressed in this publication do not reflect the views of the Government of the Hong Kong Special Administrative Region or the Innovation and Technology Commission.

Author contributions

KF and BB[1] designed the study. KF, ZZ and DL conducted the experiment and collected the data. KF, SX, WZ, ZL performed the data analysis with the help of TX, YZ, XZ, CL, JW, LW, JH and BB[2]. KF and BB[1] wrote the manuscript. SX, FZ, KK, TX, YZ, DY, ZL and WZ critically revised the manuscript draft.

Note: BB[1], Benjamin Becker; BB[2], Bharat Biswal.

Competing interests

The authors declare no competing interests.

References

Osakada, T. *et al.* A dedicated hypothalamic oxytocin circuit controls aversive social learning. *Nature*, doi:10.1038/s41586-023-06958-w (2024).
Akinrinade, I. *et al.* Evolutionarily conserved role of oxytocin in social fear contagion in zebrafish. *Science* **379**, 1232-1237, doi:10.1126/science.abq5158 (2023).
Faul, L. *et al.* Proximal threats promote enhanced acquisition and persistence of reactive fear-learning circuits. *Proceedings of the National Academy of Sciences* **117**, 16678-16689, doi:10.1073/pnas.2004258117 (2020).
Becker, B. & Zhou, F.   (2020).
Zhou, F. *et al.* A distributed fMRI-based signature for the subjective experience of fear. *Nature Communications* **12**, 6643, doi:10.1038/s41467-021-26977-3 (2021).
Bramson, B., Meijer, S., van Nuland, A., Toni, I. & Roelofs, K. Anxious individuals shift emotion control from lateral frontal pole to dorsolateral prefrontal cortex. *Nature Communications* **14**, 4880, doi:10.1038/s41467-023-40666-3 (2023).
Kreuder, A.-K. *et al.* Common and dissociable effects of oxytocin and lorazepam on the neurocircuitry of fear. *Proceedings of the National Academy of Sciences* **117**, 11781-11787, doi:10.1073/pnas.1920147117 (2020).
Szuhany, K. L. & Simon, N. M. Anxiety Disorders: A Review. *JAMA* **328**, 2431-2445, doi:10.1001/jama.2022.22744 (2022).
Slee, A. *et al.* Pharmacological treatments for generalised anxiety disorder: a systematic review and network meta-analysis. *The Lancet* **393**, 768-777, doi:https://doi.org/10.1016/S0140-6736(18)31793-8 (2019).

10 Taschereau-Dumouchel, V. *et al.* Towards an unconscious neural reinforcement intervention
for common fears. *Proceedings of the National Academy of Sciences* **115**, 3470-3475,
doi:10.1073/pnas.1721572115 (2018).
11 Zhao, Z. *et al.* Real-Time Functional Connectivity-Informed Neurofeedback of Amygdala-
Frontal Pathways Reduces Anxiety. *Psychotherapy and Psychosomatics* **88**, 5-15,
doi:10.1159/000496057 (2019).
12 Paul, J. M. *et al.* Limbic Neuropeptidergic Modulators of Emotion and Their Therapeutic
Potential for Anxiety and Post-Traumatic Stress Disorder. *The Journal of Neuroscience* **41**, 901,
doi:10.1523/JNEUROSCI.1647-20.2020 (2021).
13 Zhang, R. *et al.* Angiotensin II Regulates the Neural Expression of Subjective Fear in Humans:
A Precision Pharmaco-Neuroimaging Approach. *Biological Psychiatry: Cognitive Neuroscience*
and Neuroimaging **8**, 262-270, doi:https://doi.org/10.1016/j.bpsc.2022.09.008 (2023).
14 Zhou, F. *et al.* Human Extinction Learning Is Accelerated by an Angiotensin Antagonist via
Ventromedial Prefrontal Cortex and Its Connections With Basolateral Amygdala. *Biol Psychiat*
86, 910-920, doi:https://doi.org/10.1016/j.biopsych.2019.07.007 (2019).
15 Knobloch, H. S. *et al.* Evoked Axonal Oxytocin Release in the Central Amygdala Attenuates
Fear Response. *Neuron* **73**, 553-566, doi:https://doi.org/10.1016/j.neuron.2011.11.030
(2012).
16 Neumann, I. D. & Slattery, D. A. Oxytocin in General Anxiety and Social Fear: A Translational
Approach. *Biol Psychiat* **79**, 213-221, doi:https://doi.org/10.1016/j.biopsych.2015.06.004
(2016).
17 Eckstein, M. *et al.* Oxytocin Facilitates the Extinction of Conditioned Fear in Humans. *Biol*
Psychiat **78**, 194-202, doi:https://doi.org/10.1016/j.biopsych.2014.10.015 (2015).
18 Kou, J. *et al.* Anxiolytic Effects of Chronic Intranasal Oxytocin on Neural Responses to Threat
Are Dose-Frequency Dependent. *Psychotherapy and Psychosomatics* **91**, 253-264,
doi:10.1159/000521348 (2022).
19 Xin, F. *et al.* Oxytocin Differentially Modulates Amygdala Responses during Top-Down and
Bottom-Up Aversive Anticipation. *Adv Sci* **7**, 2001077,
doi:https://doi.org/10.1002/advs.202001077 (2020).
20 Liu, C. *et al.* Oxytocinergic Modulation of Threat-Specific Amygdala Sensitization in Humans Is
Critically Mediated by Serotonergic Mechanisms. *Biological Psychiatry: Cognitive*
Neuroscience and Neuroimaging **6**, 1081-1089,
doi:https://doi.org/10.1016/j.bpsc.2021.04.009 (2021).
21 Shamay-Tsoory, S. G. & Abu-Akel, A. The Social Salience Hypothesis of Oxytocin. *Biol Psychiat*
79, 194-202, doi:https://doi.org/10.1016/j.biopsych.2015.07.020 (2016).
22 Yao, S. *et al.* Oxytocin Facilitates Approach Behavior to Positive Social Stimuli via Decreasing
Anterior Insula Activity. *Int J Neuropsychoph* **21**, 918-925, doi:10.1093/ijnp/pyy068 (2018).
23 Oliver, J. R., Alexandra, C. P., Brian, C. & Christian, G. The translational neural circuitry of
anxiety. *Journal of Neurology, Neurosurgery & Psychiatry* **90**, 1353,
doi:10.1136/jnnp-2019-321400 (2019).
24 Scheele, D. *et al.* Trauma Disclosure Moderates the Effects of Oxytocin on Intrusions and
Neural Responses to Fear. *Psychotherapy and Psychosomatics* **88**, 61-63,
doi:10.1159/000496056 (2019).
25 Gorka, S. M. *et al.* Oxytocin Modulation of Amygdala Functional Connectivity to Fearful Faces
in Generalized Social Anxiety Disorder. *Neuropsychopharmacology* **40**, 278-286,
doi:10.1038/npp.2014.168 (2015).
26 Eckstein, M. *et al.* Oxytocin Facilitates Pavlovian Fear Learning in Males.
Neuropsychopharmacology **41**, 932-939, doi:10.1038/npp.2015.245 (2016).
27 Feng, Z. *et al.* Capturing dynamic fear experiences in naturalistic contexts: An ecologically
valid fMRI signature integrating brain activation and connectivity. *bioRxiv*,
2023.2008.2018.553808, doi:10.1101/2023.08.18.553808 (2024).
28 Chang, L. J. *et al.* Endogenous variation in ventromedial prefrontal cortex state dynamics
during naturalistic viewing reflects affective experience. *Science Advances* **7**, eabf7129,
doi:10.1126/sciadv.abf7129.
29 Meer, J. N. v. d., Breakspear, M., Chang, L. J., Sonkusare, S. & Cocchi, L. Movie viewing elicits
rich and reliable brain state dynamics. *Nature Communications* **11**, 5004,
doi:10.1038/s41467-020-18717-w (2020).

30 Adolphs, R. How should neuroscience study emotions? by distinguishing emotion states,
concepts, and experiences. *Soc Cogn Affect Neur* **12**, 24-31, doi:10.1093/scan/nsw153 (2017).
31 Pessoa, L. A Network Model of the Emotional Brain. *Trends in Cognitive Sciences* **21**, 357-371,
doi:10.1016/j.tics.2017.03.002 (2017).
32 Wen, Z., Seo, J., Pace-Schott, E. F. & Milad, M. R. Abnormal dynamic functional connectivity
during fear extinction learning in PTSD and anxiety disorders. *Molecular Psychiatry* **27**, 2216-
2224, doi:10.1038/s41380-022-01462-5 (2022).
33 Eickhoff, S. B., Milham, M. & Vanderwal, T. Towards clinical applications of movie fMRI.
NeuroImage **217**, 116860, doi:https://doi.org/10.1016/j.neuroimage.2020.116860 (2020).
34 Saarimäki, H. *et al.* Classification of emotion categories based on functional connectivity
patterns of the human brain. *NeuroImage* **247**, 118800,
doi:https://doi.org/10.1016/j.neuroimage.2021.118800 (2022).
35 de Vries, I. E. J. & Wurm, M. F. Predictive neural representations of naturalistic dynamic input.
Nature Communications **14**, 3858, doi:10.1038/s41467-023-39355-y (2023).
36 Redcay, E. & Moraczewski, D. Social cognition in context: A naturalistic imaging approach.
NeuroImage **216**, 116392, doi:https://doi.org/10.1016/j.neuroimage.2019.116392 (2020).
37 Lee, J.-J. *et al.* A neuroimaging biomarker for sustained experimental and clinical pain. *Nature
Medicine* **27**, 174-182, doi:10.1038/s41591-020-1142-7 (2021).
38 Hayoung, S., Bo-yong, P., Hyunjin, P. & Won Mok, S. Cognitive and Neural State Dynamics of
Narrative Comprehension. *The Journal of Neuroscience* **41**, 8972,
doi:10.1523/JNEUROSCI.0037-21.2021 (2021).
39 Etkin, A. *et al.* Using fMRI connectivity to define a treatment-resistant form of post-traumatic
stress disorder. *Science Translational Medicine* **11**, eaal3236,
doi:10.1126/scitranslmed.aal3236 (2019).
40 Xu, X. *et al.* Intrinsic connectivity of the prefrontal cortex and striato-limbic system
respectively differentiate major depressive from generalized anxiety disorder.
Neuropsychopharmacology **46**, 791-798, doi:10.1038/s41386-020-00868-5 (2021).
41 Jiang, X. *et al.* Intrinsic, dynamic and effective connectivity among large-scale brain networks
modulated by oxytocin. *NeuroImage* **227**, 117668,
doi:https://doi.org/10.1016/j.neuroimage.2020.117668 (2021).
42 Zhao, Z. *et al.* Oxytocin differentially modulates specific dorsal and ventral striatal functional
connections with frontal and cerebellar regions. *NeuroImage* **184**, 781-789,
doi:https://doi.org/10.1016/j.neuroimage.2018.09.067 (2019).
43 Liu, N. *et al.* Oxytocin modulates fMRI responses to facial expression in macaques.
Proceedings of the National Academy of Sciences **112**, E3123-E3130,
doi:10.1073/pnas.1508097112 (2015).
44 Wahis, J. *et al.* Astrocytes mediate the effect of oxytocin in the central amygdala on neuronal
activity and affective states in rodents. *Nat Neurosci* **24**, 529-541, doi:10.1038/s41593-021-
00800-0 (2021).
45 Bethlehem, R. A. I., van Honk, J., Auyeung, B. & Baron-Cohen, S. Oxytocin, brain physiology,
and functional connectivity: A review of intranasal oxytocin fMRI studies.
Psychoneuroendocrino **38**, 962-974, doi:https://doi.org/10.1016/j.psyneuen.2012.10.011
(2013).
46 Paloyelis, Y. Using Multivariate Neurofunctional Signatures to Dissect the Specific Mental
Processes Modulated by Novel Pharmacological Targets. *Biological Psychiatry: Cognitive
Neuroscience and Neuroimaging* **8**, 233-234, doi:https://doi.org/10.1016/j.bpsc.2023.01.002
(2023).
47 Cole, M. W. *et al.* Multi-task connectivity reveals flexible hubs for adaptive task control. *Nat
Neurosci* **16**, 1348-1355, doi:10.1038/nn.3470 (2013).
48 Uddin, L. Q. Cognitive and behavioural flexibility: neural mechanisms and clinical
considerations. *Nature Reviews Neuroscience* **22**, 167-179, doi:10.1038/s41583-021-00428-w
(2021).
49 Simony, E. *et al.* Dynamic reconfiguration of the default mode network during narrative
comprehension. *Nature Communications* **7**, 12141, doi:10.1038/ncomms12141 (2016).
50 Axelrod, V., Rees, G. & Bar, M. The default network and the combination of cognitive
processes that mediate self-generated thought. *Nature Human Behaviour* **1**, 896-910,
doi:10.1038/s41562-017-0244-9 (2017).

51   Peter, K. *et al.* Oxytocin Modulates Neural Circuitry for Social Cognition and Fear in Humans.
The Journal of Neuroscience **25**, 11489, doi:10.1523/JNEUROSCI.3984-05.2005 (2005).
52   Sonkusare, S., Breakspear, M. & Guo, C. Naturalistic Stimuli in Neuroscience: Critically
Acclaimed. *Trends in Cognitive Sciences* **23**, 699-714,
doi:https://doi.org/10.1016/j.tics.2019.05.004 (2019).
53   Menon, R. & Neumann, I. D. Detection, processing and reinforcement of social cues:
regulation by the oxytocin system. *Nature Reviews Neuroscience* **24**, 761-777,
doi:10.1038/s41583-023-00759-w (2023).
54   Xu, X., Li, J., Chen, Z., Kendrick, K. M. & Becker, B. Oxytocin reduces top-down control of
attention by increasing bottom-up attention allocation to social but not non-social stimuli – A
randomized controlled trial. *Psychoneuroendocrino* **108**, 62-69,
doi:https://doi.org/10.1016/j.psyneuen.2019.06.004 (2019).
55   Qi, S. *et al.* How cognitive and reactive fear circuits optimize escape decisions in humans.
Proceedings of the National Academy of Sciences **115**, 3186-3191,
doi:10.1073/pnas.1712314115 (2018).
56   Mobbs, D., Trimmer, P. C., Blumstein, D. T. & Dayan, P. Foraging for foundations in decision
neuroscience: insights from ethology. *Nature Reviews Neuroscience* **19**, 419-427,
doi:10.1038/s41583-018-0010-7 (2018).
57   Tashjian, S. M., Zbozinek, T. D. & Mobbs, D. A Decision Architecture for Safety Computations.
Trends in Cognitive Sciences **25**, 342-354, doi:10.1016/j.tics.2021.01.013 (2021).
58   van Heukelum, S. *et al.* Where is Cingulate Cortex? A Cross-Species View. *Trends in
Neurosciences* **43**, 285-299, doi:10.1016/j.tins.2020.03.007 (2020).
59   Loh, K. K. *et al.* Cognitive control of orofacial motor and vocal responses in the ventrolateral
and dorsomedial human frontal cortex. *Proceedings of the National Academy of Sciences* **117**,
4994-5005, doi:10.1073/pnas.1916459117 (2020).
60   Shackman, A. J. *et al.* The integration of negative affect, pain and cognitive control in the
cingulate cortex. *Nature Reviews Neuroscience* **12**, 154-167, doi:10.1038/nrn2994 (2011).
61   Koch, S. B. J. *et al.* Intranasal Oxytocin Normalizes Amygdala Functional Connectivity in
Posttraumatic Stress Disorder. *Neuropsychopharmacology* **41**, 2041-2051,
doi:10.1038/npp.2016.1 (2016).
62   Patel, G. H. *et al.* Functional evolution of new and expanded attention networks in humans.
Proceedings of the National Academy of Sciences **112**, 9454-9459,
doi:10.1073/pnas.1420395112 (2015).
63   Womelsdorf, T. & Everling, S. Long-Range Attention Networks: Circuit Motifs Underlying
Endogenously Controlled Stimulus Selection. *Trends in Neurosciences* **38**, 682-700,
doi:10.1016/j.tins.2015.08.009 (2015).
64   Rosenberg, M. D., Finn, E. S., Scheinost, D., Constable, R. T. & Chun, M. M. Characterizing
Attention with Predictive Network Models. *Trends in Cognitive Sciences* **21**, 290-302,
doi:10.1016/j.tics.2017.01.011 (2017).
65   Dixon, M. L. *et al.* Heterogeneity within the frontoparietal control network and its
relationship to the default and dorsal attention networks. *Proceedings of the National
Academy of Sciences* **115**, E1598-E1607, doi:10.1073/pnas.1715766115 (2018).
66   Keller, A. S. *et al.* Hierarchical functional system development supports executive function.
Trends in Cognitive Sciences **27**, 160-174, doi:10.1016/j.tics.2022.11.005 (2023).
67   Satpute, A. B. & Lindquist, K. A. The Default Mode Network's Role in Discrete
Emotion. *Trends in Cognitive Sciences* **23**, 851-864, doi:10.1016/j.tics.2019.07.003 (2019).
68   Sylvester, C. M. *et al.* Functional network dysfunction in anxiety and anxiety disorders. *Trends
in Neurosciences* **35**, 527-535, doi:10.1016/j.tins.2012.04.012 (2012).
69   Xu, S. *et al.* Functional connectivity profiles of the default mode and visual networks reflect
temporal accumulative effects of sustained naturalistic emotional experience. *NeuroImage*
269, 119941, doi:https://doi.org/10.1016/j.neuroimage.2023.119941 (2023).
70   Brandman, T., Malach, R. & Simony, E. The surprising role of the default mode network in
naturalistic perception. *Communications Biology* **4**, 79, doi:10.1038/s42003-020-01602-z
(2021).
71   Xin, F. *et al.* Oxytocin Modulates the Intrinsic Dynamics Between Attention-Related Large-
Scale Networks. *Cerebral Cortex* **31**, 1848-1860, doi:10.1093/cercor/bhy295 (2021).

72   Davies, C. *et al.* Connectome dysfunction in patients at clinical high risk for psychosis and
modulation by oxytocin. *Molecular Psychiatry*, doi:10.1038/s41380-024-02406-x (2024).
73   Chai, Y. *et al.* Functional connectomics in depression: insights into therapies. *Trends in*
Cognitive Sciences **27**, 814-832, doi:10.1016/j.tics.2023.05.006 (2023).
74   Yan, C.-G. *et al.* Reduced default mode network functional connectivity in patients with
recurrent major depressive disorder. *Proceedings of the National Academy of Sciences* **116**,
9078-9083, doi:10.1073/pnas.1900390116 (2019).
75   Liu, X. *et al.* A neural signature for the subjective experience of threat anticipation under
uncertainty. *Nature Communications* **15**, 1544, doi:10.1038/s41467-024-45433-6 (2024).
76   Zhang, H., Chen, K., Bao, J. & Wu, H. Oxytocin enhances the triangular association among
behavior, resting-state, and task-state functional connectivity. *Human Brain Mapping* **44**,
6074-6089, doi:https://doi.org/10.1002/hbm.26498 (2023).
77   Quintana, D. S. *et al.* Oxytocin pathway gene networks in the human brain. *Nature*
Communications **10**, 668, doi:10.1038/s41467-019-08503-8 (2019).
78   Gan, X. *et al.* A neurofunctional signature of subjective disgust generalizes to oral distaste and
socio-moral contexts. *Nature Human Behaviour* **8**, 1383-1402, doi:10.1038/s41562-024-
01868-x (2024).
79   Deco, G. & Kringelbach, Morten L. Great Expectations: Using Whole-Brain Computational
Connectomics for Understanding Neuropsychiatric Disorders. *Neuron* **84**, 892-905,
doi:10.1016/j.neuron.2014.08.034 (2014).
80   Gao, S. *et al.* Oxytocin, the peptide that bonds the sexes also divides them. *Proceedings of the*
National Academy of Sciences **113**, 7650-7654, doi:10.1073/pnas.1602620113 (2016).
81   Ma, X. *et al.* Sex- and context-dependent effects of oxytocin on social sharing. *NeuroImage*
183, 62-72, doi:https://doi.org/10.1016/j.neuroimage.2018.08.004 (2018).
82   Faul, F., Erdfelder, E., Buchner, A. & Lang, A.-G. Statistical power analyses using G*Power 3.1:
Tests for correlation and regression analyses. *Behavior Research Methods* **41**, 1149-1160,
doi:10.3758/BRM.41.4.1149 (2009).
83   Liu, Y. *et al.* Oxytocin modulates social value representations in the amygdala. *Nat Neurosci*
22, 633-641, doi:10.1038/s41593-019-0351-1 (2019).
84   Watanabe, T. *et al.* Mitigation of Sociocommunicational Deficits of Autism Through Oxytocin-
Induced Recovery of Medial Prefrontal Activity A Randomized Trial. *Jama Psychiat* **71**, 166-
175, doi:10.1001/jamapsychiatry.2013.3181 (2014).
85   Spengler, F. B. *et al.* Kinetics and Dose Dependency of Intranasal Oxytocin Effects on
Amygdala Reactivity. *Biol Psychiat* **82**, 885-894, doi:10.1016/j.biopsych.2017.04.015 (2017).
86   Lan, C. M. *et al.* Oxytocinergic Modulation of Stress-Associated Amygdala-Hippocampus
Pathways in Humans Is Mediated by Serotonergic Mechanisms. *Int J Neuropsychoph* **25**, 807-
817, doi:10.1093/ijnp/pyac037 (2022).
87   Bartz, J. A., Zaki, J., Bolger, N. & Ochsner, K. N. Social effects of oxytocin in humans: context
and person matter. *Trends in Cognitive Sciences* **15**, 301-309, doi:10.1016/j.tics.2011.05.002
(2011).
88   Andari, E. *et al.* Promoting social behavior with oxytocin in high-functioning autism spectrum
disorders. *Proceedings of the National Academy of Sciences* **107**, 4389-4394,
doi:10.1073/pnas.0910249107 (2010).
89   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*
(Methodological) **57**, 289-300, doi:10.1111/j.2517-6161.1995.tb02031.x (1995).
90   Esteban, O. *et al.* fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nature*
Methods **16**, 111-116, doi:10.1038/s41592-018-0235-4 (2019).
91   Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L. & Petersen, S. E. Spurious but
systematic correlations in functional connectivity MRI networks arise from subject motion.
NeuroImage **59**, 2142-2154, doi:https://doi.org/10.1016/j.neuroimage.2011.10.018 (2012).
92   Motzkin, J. C., Philippi, C. L., Wolf, R. C., Baskaya, M. K. & Koenigs, M. Ventromedial Prefrontal
Cortex Is Critical for the Regulation of Amygdala Activity in Humans. *Biol Psychiat* **77**, 276-
284, doi:https://doi.org/10.1016/j.biopsych.2014.02.014 (2015).
93   Mihov, Y. *et al.* Mirroring Fear in the Absence of a Functional Amygdala. *Biol Psychiat* **73**, e9-
e11, doi:https://doi.org/10.1016/j.biopsych.2012.10.029 (2013).

94 Yekutieli, D. & Benjamini, Y. Resampling-based false discovery rate controlling multiple test
procedures for correlated test statistics. *Journal of Statistical Planning and Inference* **82**, 171-
196, doi:https://doi.org/10.1016/S0378-3758(99)00041-5 (1999).
95 Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic
Functional Connectivity MRI. *Cerebral Cortex* **28**, 3095-3114, doi:10.1093/cercor/bhx179
(2018).
96 McLaren, D. G., Ries, M. L., Xu, G. & Johnson, S. C. A generalized form of context-dependent
psychophysiological interactions (gPPI): A comparison to standard approaches. *NeuroImage*
61, 1277-1286, doi:https://doi.org/10.1016/j.neuroimage.2012.03.068 (2012).
97 Fung, B. J., Qi, S., Hassabis, D., Daw, N. & Mobbs, D. Slow escape decisions are swayed by trait
anxiety. *Nature Human Behaviour* **3**, 702-708, doi:10.1038/s41562-019-0595-5 (2019).
98 Xu, T. *et al.* Angiotensin blockade enhances motivational reward learning via enhancing
striatal prediction error signaling and frontostriatal communication. *Molecular Psychiatry* **28**,
1692-1702, doi:10.1038/s41380-023-02001-6 (2023).
99 Brodmann, K., Gruber, O. & Goya-Maldonado, R. Intranasal Oxytocin Selectively Modulates
Large-Scale Brain Networks in Humans. *Brain Connectivity* **7**, 454-463,
doi:10.1089/brain.2017.0528 (2017).
100 Ruslan, M., Irina, K., Alexander, K., Denis, C. & Maxim, K. Comparison of whole-brain task-
modulated functional connectivity methods for fMRI task connectomics. *bioRxiv*,
2024.2001.2022.576622, doi:10.1101/2024.01.22.576622 (2024).
101 Schmälzle, R. *et al.* Brain connectivity dynamics during social interaction reflect social
network structure. *Proceedings of the National Academy of Sciences* **114**, 5153-5158,
doi:10.1073/pnas.1616130114 (2017).
102 Foster, Brett L., Rangarajan, V., Shirer, William R. & Parvizi, J. Intrinsic and Task-Dependent
Coupling of Neuronal Population Activity in Human Parietal Cortex. *Neuron* **86**, 578-590,
doi:10.1016/j.neuron.2015.03.018 (2015).
103 Giordano, B. L. *et al.* The representational dynamics of perceived voice emotions evolve from
categories to dimensions. *Nature Human Behaviour* **5**, 1203-1213, doi:10.1038/s41562-021-
01073-0 (2021).
789