Contribution of Beta-Lactamases and Efflux Pumps to Multidrug Resistance in *Pseudomonas aeruginosa* Isolates from ICU Patients in Northeast Brazil

Marília S. Maia [1, 2]#, Lavouisier F.B. Nogueira [1, 2]#, Marco A.F Clementino [1, 2]#, Jose Q.S. Filho [1, 3], Ila F.N. Lima [4], José K. Sousa [1, 5]*, Deiziane V.S. Costa [6], Jorge L.N. Rodrigues [4], Luciana V.C. Fragoso [4], Alexandre Havt [1, 2], and Aldo A.M. Lima [1, 2, 3,]*

[1] Institute of Biomedicine, School of Medicine, Federal University of Ceara, Fortaleza-Ceara 60430-270, Brazil.
[2] Postgraduate Medical Microbiology Program, Department of Medicine, Federal University of Ceara, Fortaleza-Ceara 60430-270, Brazil.
[3] Postgraduate Medical Sciences Program, Department of Medicine, Federal University of Ceara, Fortaleza-Ceara 60430-270, Brazil.
[4] Hospital Universitário Walter Cantídio, Federal University of Ceara, Fortaleza-Ceara 60430-270, Brazil.
[5] Division of Pediatric Gastroenterology, Hepatology, & Nutrition, University of Virginia, Charlottesville, VA 22903, United States
[6] Division of Infectious Diseases and International Health, University of Virginia, Charlottesville, VA 22903, United States

* Correspondence: alima@ufc.br; ztx7ex@virginia.edu.
# authors contributed equally to this work
e-Mails: mmaia800@gmail.com, lavouisier@gmail.com, mclementino@ufc.br, jquirinof@gmail.com, ila.lima@ebserh.gov.br, kleybson.jks@gmail.com, deiziane2009@gmail.com, jorge1207@gmail.com, 1uciana.vladia@gmail.com, ahavt@ufc.br, alima@ufc.br

Abstract

Pseudomonas aeruginosa is an opportunistic pathogen with high clinical relevance in intensive care units (ICU) due to its elevated resistance to various antimicrobials, which lead to high morbidity and mortality in patients in critical situations. In this study, we aimed to detect variants of genes encoding β-lactamases and efflux pumps in *P. aeruginosa* isolates resistant to β-lactams, fluoroquinolones and aminoglycosides. All genes belonging to the subfamilies were included in this study: bla_{SHV}, bla_{TEM}, bla_{NDM}, bla_{KPC}, bla_{GES}, bla_{CTX-M}. In addition, we investigated the most relevant variants of the bla_{OXA} subfamily and genes belonging to the efflux pumps of the Mex family. We tested 54 isolates of *P. aeruginosa* with a high prevalence of resistance to the antimicrobials piperacillin/tazobactam, ceftazidime, cefepime, imipenem and meropenem. Resistance genes related to carbapenems and spectrum β-lactamases extended were found, which included bla_{KPC} genes (81.49%), bla_{CTXM-2} (72.22%) and

bla_{CTXM-1} (66.66%). In relation to the presence of Mex family efflux pumps genes,

100% of positivity were detected. These findings suggest that *P. aeruginosa* isolates

exhibit an arsenal of genes encoding β-lactamases able to induce phenotypic patterns of

resistance to several antimicrobials commonly used as first-line treatment.

Key words: *P. aeruginosa*; Antimicrobial Resistance; β-lactamases genes; efflx pumps

genes; Intensive Care Unit.

Author Summary

Since the introduction of the use of antimicrobials, resistance to antimicrobials has been

growing and becoming a global public health problem, as it leads to ineffective

treatment and an increased risk of mortality. *P. aeruginosa* is included in the World

Health Organization (WHO) critical list of bacteria that have a higher rate of resistance

to antimicrobials, requiring constant epidemiological investigation of the strains,

especially in hospital environments, to correctly approach them. In this work, we used a

methodology that detects 740 variants of different classes of β-lactamases to evaluate

the genotype of the study strains against the phenotype found. We evidenced a high

prevalence of strains carrying genes related to carbapenems and extended-spectrum β-

lactamases, demonstrating a correlation with the phenotypes. Furthermore, we found a

100% positivity rate among the efflux pumps tested belonging to the MEX family.

Introduction

Antimicrobial-resistant Gram-negative bacteria (GNB) are of particular concern

worldwide due to their high morbidity and mortality, particularly in intensive care units

(ICUs) [1]. The emergence and spread of microbial resistance are global problems that

has profound consequences for public health and the economy [2].

Due to the selective pressure exerted by the frequent use of antimicrobials and

the long stay of patients with compromised immune systems, infections related to

multidrug-resistant (MDR) microorganisms are prevalent in ICU facilities [3,4].

Healthcare-associated infections (HAIs) caused by MDR are considered a threat to

public health, as they limit or even make available therapeutic options unfeasible, which

increases the severity and, consequently, the morbidity and mortality rates of those

affected, increasing patient length of stay and hospital costs [1].

In this scenario, *Pseudomonas aeruginosa*, an important opportunistic Gram-

negative pathogen that causes a variety of infectious diseases, has acquired resistance to

several antimicrobials in recent times [5,6]. According to the latest antimicrobial

resistance surveillance report from the European Center for Disease Prevention and

Control (ECDC), who evaluated *P. aeruginosa* isolates from 2013 to 2016, the average

of *P. aeruginosa* isolates with combined resistance (resistance to three or more

antimicrobial groups, including carbapenems) was 12.9% [7]. Furthermore, a study

carried out in Singapore observed a percentage of 11.5% of health-related infections

caused by *P. aeruginosa*, in which 23% of these isolates were resistant to carbapenems

[8]. *P. aeruginosa* is included in the "critical" category of the World Health

Organization (WHO) priority list of bacterial pathogens for which research and

development of new antimicrobials are urgently needed [1].

The main mechanisms involved in antimicrobial resistance in *P. aeruginosa*

strains are enzymatic mechanisms, where β-lactamases stand out, and efflux pumps.

[5,9]. *P. aeruginosa* isolates contain a variety of β-lactamases, such as the

carbapenemases KPC, GES, IMP, VIM, NDM and SPM, and are distributed throughout

the world [10,11,12].

Efflux pumps are related to resistance to several antimicrobials, particularly

those belonging to the AcrAB-TolC and Mex pump families belonging to the

nodulation-division resistance superfamily (RND). Efflux pumps not only mediate

intrinsic and acquired microbial resistance, but are also involved in other functions,

including bacterial stress response and pathogenicity [13,14,15].

Based on this, we sought to detect variants of genes encoding β-lactamases and

efflux pumps, belonging to the RND superfamily, in *P. aeruginosa* isolates resistant to

β-lactams, fluoroquinolones and aminoglycosides isolated from patients admitted to an

intensive care ward a tertiary care health unit in the city of Fortaleza-CE, with the

purpose of tracking the resistance profile of the strains isolated.

Material and methods

Obtaining bacterial isolates and identification

For the following study, 259 samples were collected from patients admitted to

the clinical ICU of a tertiary care health unit in the city of Fortaleza-CE in the period

from April 23, 2019 to May 29, 2021, who presented an infectious condition (defined

by the attending ward physicians) involving different anatomical sites. Those

microorganisms that were identified as Gram-negative bacteria, resistant to one or more

groups of antimicrobials were included: β-lactams, fluoroquinolones, aminoglycosides,

macrolides, and glycopeptides. These samples were made available for study upon

approval by the research ethics committee of the health unit in question (n° 3.143.258),

in which the data were anonymized before accessing them, with the need for consent

being waived by the ethics committee in question.

With the aim of diagnosing potential etiological agents, blood culture samples

and other noble liquids were inoculated in specific bottles and incubated in the

equipment BacT/Alert® 3D (BioMérieux, Marcy l'Etoile, France). The other biological

materials were sown in specific culture media using a qualitative or quantitative sowing technique, in order to obtain isolated colonies. Plates were incubated at 37±2°C for 18-24 hours.

The bacterial isolates were identified and tested for their susceptibility to antimicrobials using the automated method VITEK® 2 Compact (BioMérieux, Marcy l'Etoile, France), according to the manufacturer's recommendations. Minimum inhibitory concentrations were interpreted according to the Clinical and Laboratory Standards Institute [16] and with the Brazilian committee on antimicrobial susceptibility testing [17,18]. For quality control of sensitivity tests, strains from the American Type Culture Collection (ATCC) were used. Specimens that had a resistance profile that fit the research objectives were included in the study.

Selection of analyzed strains

A total of 258 isolates were analyzed, among which those identified as *P. aeruginosa* were included in this study. Strains of *P. aeruginosa* resistant to one or more groups of β-lactam antimicrobials, fluoroquinolones and aminoglycosides were then used to investigate their genetic resistance profile. MDR was defined as multidrug resistant to ≥1 agent in ≥3 classes of antimicrobials and extending drug resistance (XDR) as drug susceptibility to ≤2 classes of antimicrobials agents.

Extraction of bacterial DNA

The isolates maintained in trypticase soy broth were incubated overnight in a bacteriological oven at 37°C, at 200 rpm, for subsequent extraction of bacterial DNA. To extract genetic material, the Wizard Genomic DNA Purification extraction and purification kit was used (Promega, Madison, USA), according to the manufacturer's recommendations.

After extraction, all samples were quantified by spectrophotometry using the

NanoDropTM 2000 (Thermo Fisher Scientific, Waltham, Massachusetts, USA) and

stored in a -80°C freezer until used in the experiments.

Detection of resistance-related genes by molecular biology

The selected strains of *P. aeruginosa* were analyzed by real-time PCR (qPCR)

using QuantStudio 3® (Applied Biosystems, Massachusetts, USA), using specific

primers, for the detection of 13 genes encoding efflux pumps (**Table 1**) and 14 genes

encoding β-lactamases described by Nogueira and collaborators [19]. Reactions were

standardized using positive controls and negative controls (milliQ water) to determine

the most efficient qPCR conditions.

Table 1 Description of genes related to encoding efflux pumps.

Efflux pumps	Gene	Function	Primers	Primer sequence (5'-3')	Product size (bp)	At (°C)
MexAB-OprM	mexA	membrane fusion protein	MexA F	GACAAGTGGCTGGTTACCGA	96	63
			MexA R	CACGGTCTTCACCTCGACAC		
	mexB	Inner membrane transporter	MexB F	GTCGATTCCGTTCTCGGTGA	158	65
			MexB R	ACTCCACGATGAGAATGGCG		
MexCD-OprJ	mexC	membrane fusion protein	MexC F	CGTGCAATAGGAAGGATCGG	104	60
			MexC R	TCCACCGGCAACACCATTT		
	mexD	Inner membrane transporter	MexD F	TACACCCTGATCCCGTCCAT	170	64
			MexD R	ATGATCCGCTCGACGTTCTC		
MexEF-OprN	mexE	membrane fusion	MexE F	CTGAGCTTCACCCGGATCAC	140	64

					Bp	At
		protein	MexE R	GCGTCGAAGTAGGCGTAGAC		
	mexF	Inner membrane transporter	MexF F	TCATCAAGGTCAGCGACACC	169	64
			MexF R	CACTCGTAGGTCATGCCGTT		
MexGHI-OpmD	mexG	membrane protein	MexG F	TGCAGCGCTTCATCGATAACT	120	64
			MexG R	GGCTGGCCTGATAGTCGAAC		
	mexH	membrane fusion protein	MexH F	TCATCAAGGTCAGCGACACC	124	65
			MexH R	CACTCGTAGGTCATGCCGTT		
MexVW-OprM	mexV	membrane fusion protein	MexV F	TACTGTTCCTTTCCGGCGAC	170	61
			MexV R	TTCGCTTTTCGAGATGGCCT		
	mexW	membrane protein	MexW F	CCTCGGTCTACATCGGCATC	181	65
			MexW R	CCGAGGGTCTTCACCACTTC		
MexXY-OprM	mexX	membrane protein	MexX F	TGCTGTTCCAGATCGACCCT	127	60
			MexX R	TCCTTGATCAGGTCGGCGTA		
	mexY	membrane fusion protein	MexY F	GTCAACCAAATGACCGCCAC	134	65
			MexY R	ATGTTGTAGCTCACGCCCTC		

At= Annealing temperature; Bp= Base pairs; Primer F = Forward; Primer R = Reverse

Data Analysis

Fisher's test was used to compare the proportion of occurrence of a variable

between groups. The confidence interval was 95% and the test was considered

statistically significant when $P < 0.05$, using the IBM Statistical Package for the Social

Sciences Version 21 program (SPSS Statistics Software, Nova York, EUA).

Results

Prevalence of *P. aeruginosa* in the study

In this study, 258 Gram-negative bacteria were isolated. Among these isolates,

(20.9%, 54/258) were identified as *P. aeruginosa*, which were isolated from different

anatomical sites of hospitalized patients. Most of the *P. aeruginosa* strains were isolated

from tracheal aspirate (74.1%, 40/54), followed by blood samples (11.11%, 6/54),

bronchoalveolar lavage (7.40 %, 4/54), abdomen cavity (1.85%, 1/54), tissue fragment

(1.85%, 1/54), sacral pressure ulcer (1.85%, 1/54) and tendon (1.85%, 1/54).

Antimicrobial resistance phenotype tested in *P. aeruginosa* isolates

Fifteen antimicrobials from different classes were tested, in which *P.*

aeruginosa isolates obtained a resistance rate of 100% to cefoxitin (32/32),

ceftriaxone (25/25), cefuroxime axetil (32/32) and tigecycline (32/32), confirming the

intrinsic resistance of this pathogen to these antimicrobials. In addition, most of the *P.*

aeruginosa isolates were also resistant to piperacillin/tazobactam (74.4%, 32/43),

ceftazidime (63.4%, 33/52) and cefepime (63.4%, 33/52), imipenem (61.2%, 30/ 49),

meropenem (56.2%, 27/48), ciprofloxacin (54.9%, 28/51), gentamicin (41.5%, 22/53)

and amikacin (28.3%, 15/53) (**Figure 1**). Interestingly, none of the isolates

demonstrate a phenotype of resistance to colistin, ceftazidime/avibactam and

ceftolozone/tazobactam.

Figure 1. Prevalence of *P. aeruginosa* isolates resistant to β-lactam antimicrobials

(A) and aminoglycosides, quinolones and glycylcycline (B) evaluated, demonstrating

a phenotypic profile of resistance to multiple antimicrobials in more than 50% of the

isolates, with the exception of colistin, ceftazidime/ avibactam, ceftolozone/

tazobactam, gentamicin and amikacin.

The tested isolates of *P. aeruginosa* obtained a percentage of (61.11%, 33/54)

for MDR microorganisms, followed by a percentage of (55.5%, 30/54) for XDR

microorganisms and a percentage of (46.2%, 25/54) for carbapenem-resistant

microorganisms.

Detection of resistance genes in *P. aeruginosa* isolates by qPCR

For the detection of genes encoding β-lactamases, the highest percentage

found was the bla_{KPC} gene (81.49%, 44/54), followed by bla_{CTXM-2} with (72.22%,

39/54), bla_{CTXM-1} (66.66%, 36/54), bla_{CTXM-5} (59.25%, 32/54), bla_{SHV} (40.74%,

22/54), bla_{TEM} and $bla_{OXA-23-like}$ (38.89%, 21/54), bla_{NDM} (29.6%, 16/54), bla_{GES} and

$bla_{OXA-51-like}$ (24%, 13/54), $bla_{OXA-24/40-like}$ (20.4%, 11/54), bla_{CTXM-3} (11.11%, 6/54),

bla_{CTXM-4} (9.26%, 5/54) and there was no detection of the $bla_{OXA-48-like}$ gene in the

isolates (**Figure 2).**

Figure 2. Prevalence of β-lactam resistance genes among *P. aeruginosa* isolates,

grouped according to the Ambler classification, (A) in which there is a high detection

rate for class A ESBL (bla_{CTX-M} do clade 2, 1 and 5 respectively; bla_{SHV}; bla_{TEM}), (B)

marked presence of the bla_{KPC} carbapenemase, (C) in addition to the identification of

$bla_{OXA-23-like}$, $bla_{OXA-51-like}$ and $bla_{OXA-24/40-like}$ commonly associated with *Acinetobacter*

baumannii strains.

Association of the presence of resistance genes with the resistance phenotype

The association between the prevalence of the 14 genes encoding β-lactamases

and the β-lactam resistance phenotype identified in *P. aeruginosa* isolates was

evaluated. The distribution of the percentage of genes detected in the isolates tested

for the antimicrobials cefepime and ceftazidime showed a higher positivity rate for

the genes bla_{KPC} (82.7%, 43/52), bla_{CTXM-2} (75%, 39/52), bla_{CTXM-1} (69.2%, 36/52),

bla_{CTXM-5} (61.5%, 32/52) and bla_{SHV} (42.3%, 22/52). The presence of the bla_{TEM} (P=

0.017; OR 5.66; 95% CI 1.38-23.21) and $blaOXA$-23-like (P=0.041; OR 3.98; 95% CI

1.08 – 14.58) genes were associated with phenotypic resistance to the antimicrobials

cefepime and ceftazidime (**Table 2)**.

Table 2 Percentage of positivity of genes encoding β-lactamases in *P. aeruginosa*

isolates in the groups resistant and susceptible to the antimicrobials cefepime and

ceftazidime.

Coding genes β-lactamases	Total N=52(%)	Resistant N=33(%)	Susceptible N=19(%)	P valor	OR	95% CI
bla_{KPC}	43 (82.7)	27 (81.8)	16 (84.2)	1.0000	0.84	0.18 – 3.85
bla_{CTXM-2}	39 (75)	25 (75.8)	14 (73.7)	1.0000	1.11	0.30 – 4.07
bla_{CTXM-1}	36 (69.2)	24 (72.7)	12 (63.2)	0.5405	1.55	0.46 – 5.20
bla_{CTXM-5}	32 (61.5)	19 (57.6)	13 (68.4)	0.5579	0.62	0.19 – 2.05
bla_{SHV}	22 (42.3)	13 (39.4)	9 (47.37)	0.3969	0.58	0.18 – 1.82
$bla_{OXA-23-like}$	21 (40.4)	17 (51.5)	4 (21)	**0.041***	3.98	1.08 – 14.58
bla_{TEM}	20 (38.5)	17 (51.5)	3 (15.8)	**0.017***	5.66	1.38-23.21
bla_{NDM}	16 (30.7)	10 (30.3)	6 (31.6)	1.0000	0.94	0.27 – 3.19
bla_{GES}	13 (25)	8 (24.2)	5 (26.3)	1.0000	0.89	0.24 – 3.27

$bla_{OXA-51-like}$	13 (25)	11 (33.3)	2 (10.5)	0.0987	4.25	0.82 – 21.79
$bla_{OXA-24/40-like}$	10 (19.2)	8 (24.2)	2 (10.5)	0.2925	2.72	0.51 – 14.42
bla_{CTXM-3}	6 (11.5)	3 (9)	3 (15.8)	0.6563	0.53	0.09 – 2.95
bla_{CTXM-4}	5 (9.3)	4 (12.1)	1 (5.3)	0.6410	2.48	0.25 – 24.02
$bla_{OXA-48-like}$	0	0	0	0	0	0

OR= *Odds ratio*; CI= Confidence interval.

The most frequently identified genes in the isolates tested for the antimicrobial

piperacillin in combination with the β-lactamase inhibitor tazobactam were bla_{KPC},

bla_{CTXM-2}, bla_{CTXM-1}, bla_{CTXM-5}, bla_{SHV} and $bla_{OXA-23-like}$ with percentages of 81.4%

(35/43), 74.4% (32/43), 67.4% (29/43), 65.1% (28/43), 41.8% (18/43) and 41.8%

(18/43), respectively. There was no statistical significance between the genes studied

and the phenotype.

The percentage of positivity of the genes encoding β-lactamases detected in

the isolates tested for the antimicrobial imipenem indicated a higher positivity rate for

the genes bla_{KPC}, bla_{CTXM-2}, bla_{CTXM-1}, bla_{CTXM-5} and bla_{SHV} with percentages of

79.6% (39/49), 71.4% (35/49), 67.3% (33/49), 57.1% (28/49) and 38.7% (19/49),

respectively. The presence of the bla_{TEM} gene achieved statistical significance with

phenotypic resistance to the antimicrobial imipenem (P=0.018; OR 5.33; 95% CI

1.28-22.20) (**Table 3**).

Table 3 Percentage of positivity of genes encoding β-lactamases in *P. aeruginosa*

isolates in the resistant and susceptible groups to the antimicrobial imipenem.

Coding genes β-lactamases	Total N=49(%)	Resistant N=30(%)	Susceptible N=19(%)	P valor	OR	95% CI
bla_{KPC}	39 (79.6)	24 (80)	15 (78.9)	1.0000	1.06	0.25 – 4.41
bla_{CTXM-2}	35 (71.4)	20 (66.6)	15 (78.9)	0.5185	0.53	0.13 – 2.03
bla_{CTXM-1}	33 (67.3)	19 (63.3)	14 (73.7)	0.5412	0.61	0.17 – 2.18
bla_{CTXM-5}	28 (57.1)	15 (50)	13 (68.4)	0.2467	0.46	0.13 – 1.53
bla_{SHV}	19 (38.7)	10 (33.3)	9 (47.3)	0.3774	0.55	0.17 – 1.80
$bla_{OXA-23-like}$	18 (36.7)	13 (43.3)	5 (26.3)	0.3621	2.14	0.61 – 7.48
bla_{TEM}	18 (36.7)	15 (50)	3 (15.7)	**0.018***	5.33	1.28 – 22.20
bla_{NDM}	15 (30.6)	7 (23.3)	8 (42.1)	1.0000	0.87	0.27 – 2.81
$bla_{OXA-51-like}$	12 (24.5)	10 (33.3)	2 (10.5)	0.0948	4.25	0.81 – 22.14
bla_{GES}	12 (24.5)	6 (20)	6 (31.6)	0.4975	0.54	0.14 – 2.02
$bla_{OXA-24/40-like}$	8 (16.3)	6 (20)	2 (10.5)	0.4583	2.12	0.38 – 11.83
bla_{CTXM-3}	5 (10.2)	3 (10)	2 (10.5)	1.0000	0.94	0.14 – 6.25
bla_{CTXM-4}	3 (6.1)	3 (10)	0	0.2730	4.96	0.24 – 101.7
$bla_{OXA-48-like}$	0	0	0	0	0	0

OR= *Odds ratio*; CI= Confidence interval.

The distribution of the percentage of positivity of the genes encoding β-

lactamases detected in the isolates tested for the antimicrobial meropenem showed a

higher rate for the genes bla_{KPC}, bla_{CTXM-2}, bla_{CTXM-1}, bla_{CTXM-5} and bla_{SHV} with

percentages of 79.6% (39/49), 71.4% (35/49), 67.3% (33/49), 57.1% (28/49) and 38.7%

(19/49), respectively. The presence of bla_{TEM} (P= 0.034; OR 4.57; 95% CI 1.21-17.23)

and $bla_{OXA-51-like}$ (P=0.043; OR 5 .58; 95% CI 1.06 – 29.20) genes achieved statistical

significance with phenotypic resistance to the antimicrobial meropenem (**Table 4**).

Table 4 Percentage of positivity of genes encoding β-lactamases in *P. aeruginosa*

isolates in the resistant and susceptible groups to the antimicrobial meropenem.

Coding genes β-lactamases	Total N=48(%)	Resistant N=27(%)	Susceptible N=21(%)	P valor	OR	95% CI
bla_{KPC}	38 (79.1)	21 (77.7)	17 (80.9)	1.0000	0.82	0.19 – 3.40
bla_{CTXM-2}	35 (72.9)	18 (66.6)	17 (80.9)	0.3377	0.47	0.12 – 1.81
bla_{CTXM-1}	35 (72.9)	19 (70.3)	16 (76.2)	0.7502	0.74	0.20 – 2.72
bla_{CTXM-5}	28 (58.3)	13 (48.1)	15 (71.4)	0.1437	0.37	0.11 – 1.24
bla_{SHV}	21 (43.7)	10 (37)	11 (52.4)	0.3820	0.53	0.16 – 1.70
$bla_{OXA-23-like}$	19 (39.5)	13 (48.1)	6 (28.6)	0.2369	2.32	0.69 – 7.79
bla_{TEM}	18 (37.5)	14 (51.8)	4 (19)	**0.034***	4.57	1.21 – 17.23
bla_{NDM}	16 (33.3)	7 (25.9)	9 (42.8)	0.2373	0.46	0.13 – 1.58
$bla_{OXA-51-like}$	12 (25)	10 (37)	2 (9.5)	**0.043***	5.58	1.06 – 29.20
bla_{GES}	13 (27)	6 (22.2)	7 (33.3)	0.5161	0.57	0.15 – 2.06
$bla_{OXA-24/40-like}$	8 (16.6)	6 (22.2)	2 (9.5)	0.4371	2.71	0.48 – 15.11
bla_{CTXM-3}	6 (12.5)	3 (11.1)	3 (14.3)	1.0000	0.75	0.13 – 4.16

*bla*CTXM-4	2 (4.2)	2 (7.4)	0	0.4973	4.21	0.19 – 92.73
*bla*OXA-48-like	0	0	0	0	0	0

OR= *Odds ratio*; CI= Confidence interval.

The genes that encode the efflux pumps of the MEX family were evaluated by

operon, in which the distribution of the percentage of positivity found in the tested

isolates was 100% for the genes MEX-AB, MEX-EF, MEX-GH, MEX-VW, MEX-XY

and followed by a percentage of 96.3% for the MEX-CD gene. No statistical tests were

performed regarding efflux pumps due to the high positivity of the genes.

Discussion

P. aeruginosa is an opportunistic pathogen with considerable clinical importance

in intensive care units, mainly due to its high resistance to several antimicrobials,

making effective treatment impossible, leading to high morbidity and mortality in these

patients in critical situations [20].

This study found a prevalence of 20.9% of colonization by this microorganism,

among Gram-negative isolates resistant to at least one antimicrobial, which represents a

similar percentage with other studies, as demonstrated in a report issued by ANVISA

[21], where *P. aeruginosa* was the third most isolated microorganism among Gram-

negative bacteria reported in bloodstream infections, with a percentage of 16.2%. In the

United States, Sader and collaborators, analyzing samples of Gram-negative bacteria

isolated from ICU and non-ICU patients, collected in 70 hospital centers, from 2018 to

2020, found a prevalence of 23.5% of *P. aeruginosa* isolated in patients admitted to

ICU [22]. Dunphy and collaborators carried out a study in outpatient and hospital

environments in regions of Virginia, United States, over a period of one year (2019-

2020), collecting several samples, in which they identified a prevalence of 60% of *P.*

aeruginosa isolates from different anatomical sites, analyzing 971 isolates collected

from 590 patients [23], in which these studies demonstrate a variability in the detection

rate of this pathogen.

In this study, the largest number of *P. aeruginosa* isolates was obtained through

bronchial lavage secretions with a percentage of 74.1%, similar results were reported in

other studies in which *P. aeruginosa* was the most isolated pathogen in lung secretions

(bronchial lavage and tracheal aspirate) [23,24]. Respiratory infections caused by this

pathogen are quite common, especially in hospital environments, and may be related to

the handling of invasive procedures on patients, such as tracheal intubation, aspirations

and the need for mechanical pulmonary ventilation [11,20].

In order to identify the phenotypic profile of antimicrobial resistance of *P.*

aeruginosa strains isolated in the hospital environment, antimicrobials from different

classes were tested, in which the highest resistance rate of *P. aeruginosa* isolates was

100% for cephalosporins of second and third generation cefoxitin, ceftriaxone and

cefuroxime axetil. Several studies corroborate this result demonstrating that the only

third generation cephalosporin with action on strains of *P. aeruginosa* is ceftazidime

[25,26].

In agreement with the findings of this study, research carried out in the United

States by Sader and collaborators [22] on *P. aeruginosa* strains collected over a two-

year period and a study carried out in Rio de Janeiro over a 20-year period [27], found

similar results for the antimicrobials piperacillin with tazobactam, ceftazidime,

cefepime and meropenem in strains of *P. aeruginosa*. These studies indicated a growing

increase in the rate of resistance to antimicrobials over the years.

Regarding third-generation cephalosporins in combination with β-lactamases

inhibitors (ceftazidime with avibactam and ceftolozone with tazobactam), in our study

there was no detection of the resistance phenotype. This result is in line with what was

expected since ceftolozone with tazobactam is an antipseudomonal antimicrobial that

acts on multidrug-resistant strains of *P. aeruginosa* [28,29]. The antimicrobial

ceftazidime with avibactam acts on multidrug-resistant *P. aeruginosa* strains that are

ESBL, AmpC and KPC positive, as this inhibitor removes the serine residue that is the

active portion of the enzyme, however, these antimicrobials do not act on metallo β-

lactamases positive strains [30], indicating the action of β-lactamases inhibitors in light

of the high positivity of the presence of genes encoding KPC in our study.

The susceptibility rates obtained in our study are in agreement with studies that

have reported good activity of ceftazidime associated with avibactam and ceftolozone

associated with tazobactam. López-Calleja and collaborators [31] carried out a study in

a university hospital located in Spain, in which they analyzed the susceptibility profile

of 12 MDR strains of *P. aeruginosa* and 117 XDR strains of *P. aeruginosa* against

ceftolozone associated with tazobactam, and detected a susceptibility rate of 92.2% in

these isolates. In the same context, research carried out by the Global Antimicrobial

Testing Leadership and Surveillance Program, collected 2.521 clinical isolates of *P.*

aeruginosa from 41 medical centers in 10 countries in Latin America, from 2017 to

2019, in which they found a rate of 86.9% of isolates susceptible to ceftazidime

associated with avibactam [32]. Therefore, these studies corroborate our results,

showing that third-generation cephalosporins in combination with β-lactamases

inhibitors are a good therapeutic option.

In order to identify the presence and prevalence of genes encoding β-lactamases

in the university hospital, we analyzed 14 genes, where the most prevalent gene was

bla_{KPC} with 81.49% positivity in the isolates. In a study carried out by Hu and

collaborators [33], researching strains of carbapenem-resistant *P. aeruginosa* isolated

from the intestine, from 2014 to 2019, the presence of the bla_{KPC-2} gene was identified in

21.1% of the isolates. Likewise, a prospective and observational cohort study, carried

out from 2018 to 2019, analyzed 972 patients admitted with confirmed infection by

carbapenem-resistant *P. aeruginosa* in 44 hospitals, distributed across three continents,

identifying the presence of the bla_{KPC-2} gene in 49% of isolates, demonstrating the

circulation of this gene that confers resistance to carbapenems in various parts of the

world [34]. Carbapenems have the widest spectrum of action for the treatment of

infections caused by multidrug-resistant bacteria, which is why the emergence and

spread of carbapenemases has become a major public health problem. Our study

detected a rate well above the studies cited, as we used a bla_{KPC} primer that detects 19

variants, expanding the detection of this β-lactamase, in contrast to just one variant that

the studies above investigated.

To understand the association of the β-lactam resistance phenotype with genes

related to β-lactamases, we correlated the phenotype and genotype and found a

statistical association between resistance to ceftazidime and cefepime with the presence

of the bla_{TEM} and $bla_{OXA-23-like}$ genes. This can be explained by the fact that the bla_{TEM}

gene has several variants that are ESBLs, capable of hydrolyzing third and fourth

generation cephalosporins [35]. In this case, our study used primers that have the ability

to detect 167 variants of the bla_{TEM} gene, of which many of these variants are ESBLs.

Regarding the $bla_{OXA-23-like}$ gene, which encodes an enzyme capable of hydrolyzing all

cephalosporins and carbapenems [36,37], was the first-class D carbapenemase to be

discovered in a strain of *Acinetobacter baumannii*, in 1985, coincidentally the same

year that imipenem was approved for clinical use [38]. Genes belonging to the $bla_{OXA-23-like}$

group can be transmitted by plasmids between species, in which studies have

detected the role of plasmids and transposons in the dissemination of the $bla_{OXA-23\ gene}$,

which has been found in *Acinetobacter* spp., *P. aeruginosa*, as well as species belonging

to the *Enterobacteriaceae* [36,39,40,41].

The presence of the bla_{TEM} gene was also associated with resistance to

carbapenems, imipenem and meropenem; there is no previous information in the

literature that corroborates this result in *P. aeruginosa*. However, a study carried out by

Han and colleagues [42], analyzing strains of *A. baumannii* resistant to carbapenems in

a university hospital in China, found an association between the presence of the bla_{TEM}

and $bla_{OXA-23-like}$ genes with the carbapenem resistance phenotype. In research carried

out in an Iranian hospital, from 2015 to 2017, resistant strains of *P.aeruginosa* were

isolated from patients with nosocomal and non-nosocomial infections and the presence

of the $bla_{OXA-23-like}$ gene was identified in 11.19% of the isolates [43], evidencing the

circulation of the OXA group gene in species other than *Acinetobacter*.

Finally, the presence of the $bla_{OXA-51-like}$ gene was associated with resistance to

meropenem. $bla_{OXA-51-like}$ encodes a carbapenemase belonging to Ambler class D, which

are enzymes intrinsic to *A. baumannii* and are found naturally in the chromosome of this

species [36], however, in 2009, Lee and colleagues reported the presence of the bla_{OXA-51}

gene in a non-*Acinetobacter* species, in which they identified this gene with an

ISAba1 insertion, which was plasmid-encoded and the surrounding sequences suggested

that its origin was from *A. baumannii*. In this study, the enzyme encoded by bla_{OXA-51}

had the capacity to increase the minimum inhibitory concentration (MIC) of meropenem

by up to 256 times, thus conferring resistance to it [44]. However, there is only one

study in the literature that reports the presence of the bla_{OXA-51} gene in *P. aeruginosa*

strains, which were isolated from outpatients in Maranhão, Brazil [45]. These studies

demonstrate the spread of the bla_{OXA-51} gene beyond strains of *A. baumannii*, where this

gene was considered a marker for these species.

Regarding the presence of Mex family efflux pumps in *P. aeruginosa* isolates,

our study showed a positivity of 96.3% for the gene that encodes the MexCD pump and

a positivity of 100% for the rest of the Mex pumps tested. There was no statistical

association of efflux pumps in relation to the resistance phenotype due to the high

positivity in the isolates, but studies have shown that efflux pumps in *P. aeruginosa* are

among one of the main resistance mechanisms that lead to the emergence of MDR

strains and XDR. This primary resistance mechanism is responsible for reducing the

intracellular concentration of the drug to a subinhibitory concentration at which other

resistance mechanisms for specific classes can evolve and be selected, increasing

antimicrobial resistance [46,47].

The present study had some limitations, such as: susceptible strains of *P.*

aeruginosa were not included in the study for a more detailed epidemiological analysis,

no research was carried out into the existence of clones of *P. aeruginosa* circulating in

the ICU, no sequencing was carried out positive controls for a more precise validation

of all variants that our method aims to detect and the isolates were not tested for all

antimicrobials in a homogeneous way due to clinical implications and procedures

adopted by the hospital analyzed in the study. However, our work innovated in the use

of primers that could detect a greater number of genes encoding β-lactamases, thus

expanding their diagnosis.

The *P. aeruginosa* strains isolated in this study showed a high rate of phenotypic

resistance to several antimicrobials, in which these strains have a repertoire of genes

encoding β-lactamases capable of inducing these phenotypic patterns of resistance to

antimicrobials commonly used for these infections, making treatment difficult or even

impossible for the patient.

Author Contributions:

Bibliographic review and data collection, Marília S. Maia, Lavouisier F.B. Nogueira, Marco A.F Clementino and Aldo A.M. Lima; Methodology, Lavouisier F.B. Nogueira, Marília S. Maia, Marco A.F Clementino, Alexandre Havt, Ila F.N. Lima, Jorge L.N. Rodrigues, Luciana V. C. Fragoso and Aldo A. M. Lima; Bioinformatics, Lavouisier F.B. Nogueira and Marco A.F Clementino; *In vitro* validation of primers and sample testing, Marília S. Maia, Lavouisier F.B. Nogueira,; Data analysis, Jose Q.S. Filho; Deiziane V.S. Costa, José K. Sousa, and Aldo A.M. Lima.

Written review and editing, Marília S. Maia, Lavouisier F.B. Nogueira, and Aldo A.M. Lima; Supervision, Alexandre Havt and Aldo A.M. Lima; Project administration, Aldo A.M. Lima; Resource acquisition, Aldo A.M. Lima.

Funding: This research was funded by CNPq (http://www.cnpq.br), grant numbers 402607/2018-0 and 408549/2022-0 and FUNCAP (https://www.funcap.ce.gov.br/), grant number: OFÍCIO Nº 102/2021 – DINOV.

Data Availability Statement: The data and reports relating to this study will be available via direct request via email to the corresponding author.

Conflicts of Interest: The authors declare that there are no financial or personal conflicts of interest that may have influenced the work.

References

1.	Who. Media Center. WHO publishes list of bacteria for which new antibiotics are urgently needed. World Health Organization: 27 Fev., 2017.

2.	Fleece ME, Pholwat S, Mathers AJ, Houpt ER. Molecular diagnosis of antimicrobial resistance in Escherichia coli. Expert Review of Molecular Diagnostics. 2018;18: 207–217. doi:10.1080/14737159.2018.1439381

3.   Amaral MVF, Roza LEM, Tonin JJ, Coronel JL, Zambon JDS, Vidaletti TB, et

al. Prevalência de microorganismos multirresistentes em um hospital universitário -

análise de uma série histórica. Anais do II Congresso Brasileiro de Medicina Hospitalar.

São Paulo: Editora Edgard Blücher; 2014. Available: http://dx.doi.org/10.5151/medpro-

ii-cbmh-067

4.   Carvalho, M.R., Moreira, I.C., Neta, F.L., Guimaraes, M.D., Viana, V.G.,

Oliveira, F.W. Incidência de bactérias multirresistentes em uma unidade de terapia

intensiva. Revista Interdisciplinar. 2015;8: 75-85.

5.   Botelho J, Grosso F, Peixe L. WITHDRAWN: Antibiotic resistance in

Pseudomonas aeruginosa – mechanisms, epidemiology and evolution. Drug Resistance

Updates. 2019. doi:10.1016/j.drup.2019.07.001

6.   Jurado-Martín I, Sainz-Mejías M, McClean S. *Pseudomonas aeruginosa*: An

Audacious Pathogen with an Adaptable Arsenal of Virulence Factors. International

Journal of Molecular Sciences. 2021;22: 3128. doi:10.3390/ijms22063128

7.   European Center for Disease Prevention and Control (ECDC), Surveillance of

antimicrobial resistance in Europe, Annual report of the European Antimicrobial

Resistance Surveillance Network (EARS-Net), 2016.

8.   Cai Y, Venkatachalam I, Tee NW, et al. Prevalence of Healthcare-Associated

Infections and Antimicrobial Use Among Adult Inpatients in Singapore Acute-Care

Hospitals: Results From the First National Point Prevalence Survey. Clin Infect Dis.

2017;64(suppl_2):S61-S67. doi:10.1093/cid/cix103

9.   Rossi E, La Rosa R, Bartell JA, et al. *Pseudomonas aeruginosa* adaptation and

evolution in patients with cystic fibrosis. Nat Rev Microbiol. 2021;19(5):331-342.

doi:10.1038/s41579-020-00477-5

10. Escandón-Vargas K, Reyes S, Gutiérrez S, Villegas MV. The epidemiology of

carbapenemases in Latin America and the Caribbean. Expert Rev Anti Infect Ther.

2017;15(3):277-297. doi:10.1080/14787210.2017.1268918.

11. Tenover FC, Nicolau DP, Gill CM. Carbapenemase-producing *Pseudomonas*

aeruginosa - an emerging challenge. Emerg Microbes Infect. 2022;11(1):811-814.

doi:10.1080/22221751.2022.2048972

12. Walters MS, Grass JE, Bulens SN, et al. Carbapenem-Resistant *Pseudomonas*

aeruginosa at US Emerging Infections Program Sites, 2015. Emerg Infect Dis.

2019;25(7):1281-1288. doi:10.3201/eid2507.181200

13. Li XZ, Plésiat P, Nikaido H. The challenge of efflux-mediated antibiotic

resistance in Gram-negative bacteria. Clin Microbiol Rev. 2015;28(2):337-418.

doi:10.1128/CMR.00117-14

14. Lv L, Wan M, Wang C, et al. Emergence of a Plasmid-Encoded Resistance-

Nodulation-Division Efflux Pump Conferring Resistance to Multiple Drugs, Including

Tigecycline, in *Klebsiella pneumoniae*. mBio. 2020;11(2):e02930-19.

doi:10.1128/mBio.02930-19.

15. Ni RT, Onishi M, Mizusawa M, et al. The role of RND-type efflux pumps in

multidrug-resistant mutants of *Klebsiella pneumoniae*. Sci Rep. 2020;10(1):10876.

doi:10.1038/s41598-020-67820-x

16. CLSI, Clinical and Laboratory Standards Institute, M100, 30th ed., January,

2020.

17. BrCAST, Brazilian Committee on Antimicrobial Susceptibility Testing, March,

2020.

18. BrCAST, Brazilian Committee on Antimicrobial Susceptibility Testing, May,

2021.

19. Nogueira LFB, Clementino MAF, Maia MS, et al. Molecular insights into β-lactams resistance in *Klebsiella pneumoniae* clinical isolates with a focus on Multidrug Resistance and virulence. medRxiv 2023.12.30.23300671; doi:10.1101/2023.12.30.23300671.

20. Urzedo JE, de Paula Menezes R, Porto JP, et al. High mortality by nosocomial infections caused by carbapenem-resistant *P. aeruginosa* in a referral hospital in Brazil: facing the perfect storm. J Med Microbiol. 2020;69(12):1388-1397. doi:10.1099/jmm.0.001273

21. Agência Nacional de Vigilância Sanitária (ANVISA). rede nacional de monitoramento da resistência microbiana em serviços de saúde, Boletim Segurança do Paciente e Qualidade em Serviços de Saúde nº 28 - Avaliação Nacional dos indicadores de IRAS e RM, Brasília: ANVISA, 2021.

22. Sader HS, Mendes RE, Streit JM, Carvalhaes CG, Castanheira M. Antimicrobial susceptibility of Gram-negative bacteria from intensive care unit and non-intensive care unit patients from United States hospitals (2018-2020). Diagn Microbiol Infect Dis. 2022;102(1):115557. doi:10.1016/j.diagmicrobio.2021.115557

23. Dunphy LJ, Kolling GL, Jenior ML, et al. Multidimensional Clinical Surveillance of *Pseudomonas aeruginosa* Reveals Complex Relationships between Isolate Source, Morphology, and Antimicrobial Resistance. mSphere. 2021;6(4):e0039321. doi:10.1128/mSphere.00393-21

24. de Souza Nunes LH, Bernardi Lora JF, Fanhani Cracco LA, et al. *Pseudomonas aeruginosa* in tracheal aspirate: Colonization, infection, and recurrence. Clin Respir J. 2023;17(5):439-446. doi:10.1111/crj.13612

25. Babich T, Naucler P, Valik JK, et al. Ceftazidime, Carbapenems, or Piperacillin-tazobactam as Single Definitive Therapy for *Pseudomonas aeruginosa* Bloodstream

Infection: A Multisite Retrospective Study. Clin Infect Dis. 2020;70(11):2270-2280. doi:10.1093/cid/ciz668

26. Freifeld AG, Bow EJ, Sepkowitz KA, et al. Clinical practice guideline for the use of antimicrobial agents in neutropenic patients with cancer: 2010 update by the infectious diseases society of america. Clin Infect Dis. 2011;52(4):e56-e93. doi:10.1093/cid/cir073

27. de Oliveira Santos IC, Pereira de Andrade NF, da Conceição Neto OC, et al. Epidemiology and antibiotic resistance trends in clinical isolates of *Pseudomonas aeruginosa* from Rio de janeiro - Brazil: Importance of mutational mechanisms over the years (1995-2015). Infect Genet Evol. 2019;73:411-415. doi:10.1016/j.meegid.2019.05.015

28. Haidar G, Philips NJ, Shields RK, et al. Ceftolozane-Tazobactam for the Treatment of Multidrug-Resistant *Pseudomonas aeruginosa* Infections: Clinical Effectiveness and Evolution of Resistance. Clin Infect Dis. 2017;65(1):110-120. doi:10.1093/cid/cix182

29. Munita JM, Aitken SL, Miller WR, et al. Multicenter Evaluation of Ceftolozane/Tazobactam for Serious Infections Caused by Carbapenem-Resistant Pseudomonas aeruginosa. Clin Infect Dis. 2017;65(1):158-161. doi:10.1093/cid/cix014

30. Drawz SM, Papp-Wallace KM, Bonomo RA. New β-lactamase inhibitors: a therapeutic renaissance in an MDR world. Antimicrob Agents Chemother. 2014;58(4):1835-1846. doi:10.1128/AAC.00826-13

31. López-Calleja AI, Morilla Morales E, Nuñez Medina R, et al. Antimicrobial activity of ceftolozane-tazobactam against multidrug-resistant and extensively drug-resistant *Pseudomonas aeruginosa* clinical isolates from a Spanish hospital. Rev Esp Quimioter. 2019;32(1):68-72

32.   Karlowsky JA, Kazmierczak KM, Valente MLNF, et al. In vitro activity of

ceftazidime-avibactam against Enterobacterales and *Pseudomonas aeruginosa* isolates

collected in Latin America as part of the ATLAS global surveillance program, 2017-

2019. Braz J Infect Dis. 2021;25(6):101647. doi:10.1016/j.bjid.2021.101647

33.   Hu Y, Qing Y, Chen J, et al. Prevalence, Risk Factors, and Molecular

Epidemiology of Intestinal Carbapenem-Resistant *Pseudomonas aeruginosa*. Microbiol

Spectr. 2021;9(3):e0134421. doi:10.1128/Spectrum.01344-21

34.   Reyes J, Komarow L, Chen L, et al. Global epidemiology and clinical outcomes

of carbapenem-resistant *Pseudomonas aeruginosa* and associated carbapenemases

(POP): a prospective cohort study. Lancet Microbe. 2023;4(3):e159-e170.

doi:10.1016/S2666-5247(22)00329-9

35.   Castanheira M, Simner PJ, Bradford PA. Extended-spectrum β-lactamases: an

update on their characteristics, epidemiology and detection. JAC Antimicrob Resist.

2021;3(3):dlab092. doi:10.1093/jacamr/dlab092

36.   Evans BA, Amyes SG. OXA β-lactamases. Clin Microbiol Rev. 2014;27(2):241-

263. doi:10.1128/CMR.00117-13.

37.   Kaitany KC, Klinger NV, June CM, et al. Structures of the class D

Carbapenemases OXA-23 and OXA-146: mechanistic basis of activity against

carbapenems, extended-spectrum cephalosporins, and aztreonam. Antimicrob Agents

Chemother. 2013;57(10):4848-4855. doi:10.1128/AAC.00762-13

38.   Lyon JA. Imipenem/cilastatin: the first carbapenem antibiotic. Drug Intell Clin

Pharm. 1985;19(12):895-899.

39.   Dias VC, Diniz CG, Peter AC, et al. Epidemiological characteristics and

antimicrobial susceptibility among carbapenem-resistant non-fermenting bacteria in

Brazil. J Infect Dev Ctries. 2016;10(6):544-553. doi:10.3855/jidc.6640

40.   Liu LL, Ji SJ, Ruan Z, et al. Dissemination of blaOXA-23 in *Acinetobacter* spp.

in China: main roles of conjugative plasmid pAZJ221 and transposon Tn2009.

Antimicrob Agents Chemother. 2015;59(4):1998-2005. doi:10.1128/AAC.04574-14

41.   Zong G, Zhong C, Fu J, et al. The carbapenem resistance gene bla_{OXA-23} is

disseminated by a conjugative plasmid containing the novel transposon Tn6681 in

Acinetobacter johnsonii M19. Antimicrob Resist Infect Control. 2020;9(1):182.

doi:10.1186/s13756-020-00832-4

42.   Han L, Lei J, Xu J, Han S. blaOXA-23-like and blaTEM rather than blaOXA-

51-like contributed to a high level of carbapenem resistance in *Acinetobacter baumannii*

strains from a teaching hospital in Xi'an, China. Medicine (Baltimore).

2017;96(48):e8965. doi:10.1097/MD.0000000000008965

43.   Rouhi S, Ramazanzadeh R. Prevalence of $bla_{Oxacillinase-23}$and $bla_{Oxacillinase-24/40}$-type

Carbapenemases in *Pseudomonas aeruginosa* Species Isolated From Patients With

Nosocomial and Non-nosocomial Infections in the West of Iran. Iran J Pathol.

2018;13(3):348-356.

44.   Lee YT, Turton JF, Chen TL, et al. First identification of blaOXA-51-like in

non-baumannii *Acinetobacter* spp. J Chemother. 2009;21(5):514-520.

doi:10.1179/joc.2009.21.5.514

45.   Nitz F, de Melo BO, da Silva LCN, et al. Molecular Detection of Drug-

Resistance Genes of bla_{OXA-23}-bla_{OXA-51} and *mcr-1* in Clinical Isolates of *Pseudomonas*

aeruginosa. Microorganisms. 2021;9(4):786. doi:10.3390/microorganisms9040786

46.   Lorusso AB, Carrara JA, Barroso CDN, Tuon FF, Faoro H. Role of Efflux

Pumps on Antimicrobial Resistance in *Pseudomonas aeruginosa*. Int J Mol Sci.

2022;23(24):15779. doi:10.3390/ijms232415779

47.   Oliveira WK, Ferrarini M, Morello LG, Faoro H. Resistome analysis of

bloodstream infection bacterial genomes reveals a specific set of proteins involved in

antibiotic resistance and drug efflux. NAR Genom Bioinform. 2020;2(3):lqaa055.

doi:10.1093/nargab/lqaa055.

(C)

Percentage (%)

(B)

(A)

(B)

(A)