# Estimating Rates of Change to Interpret Quantitative Wastewater Surveillance of Disease Trends

David A. Holcomb[a,b], Ariel Christensen[a,c], Kelly Hoffman[b], Allison Lee[b], A. Denene Blackwood[d], Thomas Clerkin[d], Javier Gallard-Góngora[d], Angela Harris[e], Nadine Kotlarz[f], Helena Mitasova[g,h], Stacie Reckling[c,g], Francis L. de los Reyes III[e], Jill R. Stewart[b], Virginia T. Guidry[c], Rachel T. Noble[b,d], Marc L. Serre[b], Tanya P. Garcia[i,†], Lawrence S. Engel[a,†,*]

[a] Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

[b] Department of Environmental Sciences and Engineering, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

[c] Occupational & Environmental Epidemiology Branch, Division of Public Health, North Carolina Department of Health and Human Services, Raleigh, NC, USA

[d] Institute of Marine Sciences, Department of Earth, Marine and Environmental Sciences, University of North Carolina at Chapel Hill, Morehead City, NC, USA

[e] Department of Civil, Construction and Environmental Engineering, North Carolina State University, Raleigh, NC, USA

[f] Department of Biological Sciences, North Carolina State University, Raleigh, NC, USA

[g] Center for Geospatial Analytics, North Carolina State University, Raleigh, NC, USA

[h] Department of Marine, Earth and Atmospheric Sciences, North Carolina State University, Raleigh, NC, USA

[i] Department of Biostatistics, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

[†] TG and LE are co-senior authors

[*] Department of Epidemiology, University of North Carolina, CB #7435, Chapel Hill, North Carolina 27599-7435, USA; Larry.Engel@unc.edu; 919-962-2756

The authors declare they have no conflicts of interest related to this work to disclose.

**Abstract**

*Background:* Wastewater monitoring data can be used to estimate disease trends to inform public health responses. One commonly estimated metric is the rate of change in pathogen quantity, which typically correlates with clinical surveillance in retrospective analyses. However, the accuracy of rate of change estimation approaches has not previously been evaluated.

*Objectives:* We assessed the performance of approaches for estimating rates of change in wastewater pathogen loads by generating synthetic wastewater time series data for which rates of change were known. Each approach was also evaluated on real-world data.

*Methods:* Smooth trends and their first derivatives were jointly sampled from Gaussian processes (GP) and independent errors were added to generate synthetic viral load measurements; the range hyperparameter and error variance were varied to produce nine simulation scenarios representing different potential disease patterns. The directions and magnitudes of the rate of change estimates from four estimation approaches (two established and two developed in this work) were compared to the GP first derivative to evaluate classification and quantitative accuracy. Each approach was also implemented for public SARS-CoV-2 wastewater monitoring data collected January 2021 – May 2024 at 25 sites in North Carolina, USA.

*Results:* All four approaches inconsistently identified the correct direction of the trend given by the sign of the GP first derivative. Across all nine simulated disease patterns, between a quarter and a half of all estimates indicated the wrong trend direction, regardless of estimation approach. The proportion of trends classified as plateaus (statistically indistinguishable from zero) for the North Carolina SARS-CoV-2 data varied considerably by estimation method but not by site.

*Discussion:* Our results suggest that wastewater measurements alone might not provide sufficient data to reliably track disease trends in real-time. Instead, wastewater viral loads could be combined with additional public health surveillance data to improve predictions of other outcomes.

## Introduction

The use of wastewater surveillance to monitor infectious disease expanded dramatically during the global coronavirus disease 2019 (COVID-19) pandemic, with thousands of monitoring sites active across dozens of countries by early 2023.[1] Wastewater monitoring offers attractive features for augmenting surveillance of a wide range of pathogens and other population health-relevant targets, such as toxic metals and endogenous biomarkers, and may be particularly well-suited as an early warning system for outbreaks of novel pathogens and variants.[2–7] However, assessing disease *trends* using wastewater surveillance faces an inherent challenge of interpretation: unlike traditional population metrics derived from counts of infected, symptomatic, or hospitalized individuals, the quantity of pathogen markers (e.g., gene targets) measured in wastewater cannot be used directly as a proxy for community disease burden. The loads of pathogen markers present in wastewater are broadly proportional to the number of infected individuals shedding the pathogen in their feces—demonstrated for SARS-CoV-2 by widely reported positive associations between wastewater viral loads and reported COVID-19 cases—but numerous biological, environmental, and site-specific factors can differentially impact measurements of wastewater pathogen loads at any given place and time.[8–12]

A common strategy for interpreting wastewater pathogen loads is to estimate traditional disease metrics like incidence rate or effective reproduction number.[12–17] Such metrics are typically estimated by exploiting statistical associations between wastewater pathogen loads and reported cases, hospitalizations, etc. at a given site or by constructing mechanistic models of fecal shedding to estimate the number of community infections required to produce the pathogen loads measured in the community's wastewater. These strategies require additional assumptions and data to implement, such as geographically and temporally aligned population surveillance data or pathogen-specific fecal shedding distributions. Such data, however, are often unavailable for novel pathogens and are subject to change unpredictably over the course of an outbreak or pandemic.[17–19]

An alternative strategy for interpreting wastewater pathogen loads is to assess trends over time

by comparing the loads measured at different time points within the same location. Because assessing

trends within-site helps to control for site-specific factors that influence pathogen load measurements, an

increasing trend in wastewater measurements should correspond to an increase in infections in the

community. Useful information about the direction and speed at which community infection trends are

changing may therefore be inferred solely on the basis of wastewater measurements by estimating the

slope of the wastewater trend at specific times, where the sign and magnitude of the slope provide the

direction and rate of change, respectively.[20]

During the COVID-19 pandemic, the United States Centers for Disease Control and Prevention

(CDC) described a simple regression-based approach for estimating the rate of change in SARS-CoV-2

wastewater trends over small subsets of wastewater viral load data.[21,22] A refinement of this approach

was suggested that uses reported daily COVID-19 case counts to impute wastewater viral loads on

unmonitored days before applying linear regression to estimate rates of change.[20] Both approaches

produce estimated slopes (the rate of change) and associated standard errors that can be used for trend

classification: a positive and statistically significant slope means the trend is increasing, a negative and

significant slope means the trend is decreasing, and a slope that is not statistically significant (regardless

of the sign) indicates the trend is not meaningfully changing and is classified as a plateau.[18] Rate of

change estimates from both approaches have been compared with population-based metrics (e.g.,

reported cases) but, to the best of our knowledge, the estimation performance and trend classification

accuracy of either approach has not yet been evaluated.[18,20]

We developed a simulation-based approach to evaluate rate of change estimates using synthetic

time series data for which the underlying smooth trends and their rates of change were known exactly.

We sampled from Gaussian processes (GP) to jointly simulate smooth wastewater viral load trends and

their first derivatives.[23–25] Independent random errors were introduced to the simulated trends to

generate synthetic measurements of wastewater viral loads, varying the smoothness of the trends and the

magnitude of the errors to represent a range of potential infectious disease patterns. We evaluated four

rate of change estimation approaches: the linear regression approach described by CDC, the multivariate

imputation approach proposed by Al-Faliti et al. (2022), a modified univariate imputation approach

requiring only wastewater measurements, and a continuous smoothing approach using generalized

additive models (GAM) with numerical approximation to estimate the smooth trend and its first

derivative.[20,21,26,27] These candidate approaches were applied to the synthetic wastewater data and

evaluated by comparing their rate of change estimates to the simulated GP derivatives. Finally, all four

approaches were applied to public wastewater monitoring data from 25 North Carolina sewersheds to

assess the impact of estimation method on the interpretation of trends in a real-world context.

**Methods**

*Rate of Change Estimation Approaches*

All analyses were performed in R version 4.2.2.[28,29] R packages used are denoted by italics.

Rolling Regressions by Sampling Event

CDC National Wastewater Surveillance System (NWSS) recommends analyzing trends in

measured wastewater viral loads by fitting simple linear regression models to a minimum of the three

most-recent wastewater samples for a given location. These models use log-transformed viral load ($\log_{10}$

gene copies/day) as the response variable and date as the predictor variable.[20–22] When fit to three

observations of weekly wastewater samples or five observations of twice-weekly samples, the regression

coefficient corresponds to the slope of the trend—the average daily change in viral load—over the

preceding ~15 days. The estimated rate of change can also be expressed as percent daily change (PDC),

enabling more direct comparison with trends in other metrics.[21] We estimated the rate of change on each

day of sample collection by fitting rolling linear models to wastewater viral loads measured on the

estimation day and the preceding four sampling events (five observations total).[21,30] For event $i$ of $N$

sampling events, let $\boldsymbol{y}_i = [y_{i-4}, \ldots, y_i]^T$ denote the $\log_{10}$-transformed wastewater viral loads and

$\boldsymbol{z}_i = [z_{i-4}, \ldots, z_i]^T$ denote the sampling dates. The rate of change estimate is given by $\beta$ in

$$\boldsymbol{y}_i = \beta_0 + \boldsymbol{z}_i \beta + \boldsymbol{\epsilon}_i, \tag{1}$$

where we assume independent, normally distributed residuals $\boldsymbol{\epsilon}_i = [\epsilon_{i-4}, \ldots, \epsilon_i]^T$.

Rolling Regressions on Imputed Daily Observations

Multivariate Imputation

While clinical and syndromic surveillance of infectious disease outcomes (e.g., incident cases,

hospitalizations, and deaths) are generally reported at daily resolution, wastewater surveillance programs

typically sample less frequently, often once or twice a week.[22,31] Rate of change estimates based on

small subsets of wastewater observations are subject to substantial uncertainty and temporal

variability.[18,32] To address the temporal sparsity of wastewater data, Al-Faliti et al. used daily reported

cases to impute wastewater viral loads on unsampled days.[20] Daily rates of change were estimated using

rolling linear models applied over 21- or 28-day subsets of the imputed daily wastewater viral loads.

Following the approach of Al-Faliti et al., we constructed five complete daily viral load datasets using

the *mice* package to implement multivariate imputation using chained equations (MICE) with random

forest models.[33] We used $\log_{10}$ wastewater viral loads and a 7-day moving average of daily cases as

inputs to MICE. We modified the original approach slightly by log-transforming the averaged cases for

computational stability and specifying a consistent 20 iterations of the MICE algorithm for each dataset

we imputed.[34] From the five resulting complete daily datasets, we selected the realization demonstrating

the highest Spearman rank correlation between 7-day average cases and the imputed daily wastewater

viral loads for downstream analyses, as specified by the method developers. For the selected

multivariate-imputed daily dataset, we estimated the rate of change on each original sampling day by

applying the rolling linear model approach described previously to the 21 daily observations ending on

the estimation day.

Univariate Imputation

The multivariate imputation approach relied on daily reported case data but a key motivation for

estimating rates of change in wastewater surveillance data is to enable identification and interpretation

of infectious disease trends using only wastewater surveillance data (i.e., when reported case data are

unavailable or inadequate). We therefore also implemented a univariate time series imputation approach

that used only the measured wastewater viral loads to impute viral loads on unmonitored days.

Univariate imputation was conducted by Kalman smoothing on structural time series models using the

*imputeTS* package.[35] Kalman smoothing has previously been shown to flexibly estimate smooth trends

in wastewater viral loads on both synthetic and various real-world wastewater surveillance data.[32]

However, as a discrete-time autoregressive approach that assumes equal-sized time steps, Kalman

smoothing does not provide a direct way to estimate the rate of change and corresponding uncertainty of

the modeled trend. As such, we used Kalman smoothing trend estimates to impute wastewater viral

loads on unsampled days, then estimated the rate of change on each original sampling day by applying

21-day window rolling linear models to the imputed dataset.

<u>First Derivatives of Smooth Functions of Time</u>

Substantial fluctuations over short timescales in both measured wastewater viral loads and

reported infections have motivated the use of a variety of smoothing approaches to better characterize

infectious disease trends from noisy surveillance data.[20,26,36] Many common smoothing techniques,

including simple moving averages and locally weighted scatterplot smoothing (LOESS), use the values

of neighboring observations within a user-defined window to estimate smoothed values.[17,18,20,37] Such

techniques are entirely data-dependent and do not have simple mathematical representations like those

from the methods previously presented. Common time series approaches like Kalman smoothing assume

equally spaced time steps and can only provide smooth estimates at discrete time points.[38,39] By contrast,

approaches that estimate continuous, smooth functions of time from the observed data can be evaluated

at any arbitrary time point to obtain the corresponding estimate of the smooth trend. This feature

provides a straightforward means of estimating the rate of change in the smooth trend at any moment

during the monitoring period using finite differences to numerically approximate the first derivative.[38]

We used generalized additive models to estimate smoothed wastewater viral loads as continuous

functions of time (see Supplemental Material, Smoothing with Generalized Additive Models). GAMs

are a flexible extension of the generalized linear model that have previously been shown to provide

accurate estimates of wastewater SARS-CoV-2 viral loads.[26,27,40] We used the *mgcv* package to estimate

GAMs by restricted maximum likelihood (REML) using $\log_{10}$ wastewater viral load as the response and

study date as a smooth predictor term. We specified a cubic regression spline basis and the lesser of 100

or half the number of observations as the maximum basis dimension.[27,41] Daily first derivatives and their

corresponding pointwise 95% confidence intervals (CI) were estimated from GAM fits using the *gratia*

package.[42]

*Simulating Differentiable Time Series*

<u>Simulating Smooth Trends with Known Rates of Change</u>

We simulated wastewater trends and corresponding rates of change by sampling from Gaussian

processes with squared exponential kernel covariance functions. A GP represents a distribution over all

the possible smooth functions of a continuous domain (e.g., time) and is defined by its covariance

function $k(z_i, z_j)$ that relates any pair of time points $z_i$, $z_j$ on that domain.[23,38,43] The time-derivative of a

GP is also a GP with a covariance kernel function $k'(z_i, z_j)$ equal to the derivative of the original

covariance function with respect to times $z_i$ and $z_j$.[24,25] This feature enables us to simulate both a smooth

trend and its instantaneous rate of change at any finite set of time points by jointly sampling from the GP

and its derivative, which follow a multivariate normal distribution (see Supplemental Material, Gaussian

Process Derivatives).[25,44]

The squared exponential kernel covariance function is given by

$$k(z_i, z_j | \alpha, \rho) = \alpha^2 \exp\left(-\frac{1}{2}\left(\frac{z_i - z_j}{\rho}\right)^2\right), \tag{2}$$

where $\alpha$ is the marginal standard deviation, a scale hyperparameter that controls the magnitude of the

covariance.[25] The rate at which correlation decays with increasing distance between times $z_i$ and $z_j$ is

controlled by the range hyperparameter $\rho$. Smaller values of $\rho$ indicate that correlation decays more

quickly, so that each observation provides less information about observations at other time points. The

result is a more rapidly changing, wiggly function. Conversely, larger values of $\rho$ mean each

observation offers greater information about its temporal neighbors, producing a more slowly changing,

or smoother, trend. We implemented the squared exponential kernel and its derivatives (see

Supplemental Material, Squared Exponential Kernel Function) as **R** functions and jointly sampled trend

observations and derivatives using the *mvnfast* package.[44,45]

<u>Generating Synthetic Wastewater Measurements</u>

Synthetic wastewater viral load time series data (in $\log_{10}$ copies/day) were generated by jointly

sampling a smooth trend and its first derivative from a GP at 1000 sequential integer locations to

represent a 1000-day monitoring period with daily trend realizations. Independent random errors

$\epsilon_t^{ww} \sim N(0, \sigma^2)$ with standard deviation $\sigma$ were independently sampled on each day $t$ and added to the

corresponding trend value $x_t$ to simulate the daily wastewater $\log_{10}$ viral load measurement

$y_t^{ww} = x_t + \epsilon_t^{ww}$. We down-sampled the synthetic wastewater measurements by selecting every third,

then fourth, observation in an alternating pattern to represent a typical twice-weekly wastewater

sampling frequency, denoted $y_i^{ww,obs} = y_{t[i]}^{ww}$ for the $i^{th}$ of $N$ simulated sampling events.[46] Although our

primary aim was estimating rates of change using wastewater measurements alone, implementing the

multivariate imputation approach required simulating daily reported case counts that shared an

underlying trend with the simulated wastewater viral loads. We sampled daily cases from a Poisson

distribution with the daily log-incidence rate given by the sum of the GP trend, independent Gaussian

error, and a constant mean log-incidence rate determined mechanistically for an assumed sewershed

population of 200,000 (see Supplemental Material, Simulating Reported Case Counts).[12]

*Evaluating Performance of Rate of Change Estimation Approaches*

<u>Simulation Scenarios</u>

We evaluated the rate of change estimation performance for three values of $\rho$ so as to have

varying smoothness of trends, and three values of $\sigma$ so as to control the magnitude of variation of

synthetic observations around the trend. In total, we had nine simulation scenarios. Range

hyperparameter values of $\rho = 90$, $\rho = 30$, and $\rho = 15$ days were selected to produce more smooth

(slowly changing), moderately smooth, and less smooth (wiggly) trends, respectively.[38] The GP

marginal standard deviation $\alpha = 1$ was used for all simulations, such that the uncorrelated variance $\sigma^2$

was a quarter of the autocorrelated trend variance $\alpha^2$ for the less-noisy condition $\sigma = 0.5$;

approximately half the trend variance for the moderate condition $\sigma = 0.75$; and equal to the trend

variance for $\sigma = 1$. We generated 1000 synthetic datasets under each scenario and estimated the rate of

change on each designated "sampling event" day (i.e., the third and seventh day of each seven-day

period) by each of the four approaches (rolling linear models, multivariate imputation, univariate

imputation, and generalized additive models).

246           Implementing an estimation approach over the entire synthetic dataset corresponds to a

retrospective analysis in which previously collected data are analyzed to characterize past trends.

However, active wastewater monitoring programs are primarily concerned with identifying changes in

infection trends in near real-time, updating estimates as new data become available.[22] For the rolling

linear model approach these analyses are identical, as only the five most recent observations are

analyzed for each estimation day. By default, the imputation and GAM approaches make use of the

entire set of observations, but in real-time analyses they would be limited to only the data collected up to

each estimation time point. Accordingly, we also implemented a modified local GAM approach

("rolling GAM") that refit the GAM to only the subset of data already observed by the day for which the

rate of change was being estimated. Locally restricted rolling imputations were not implemented:

multiple imputation was too computationally intensive to feasibly perform across all simulation

scenarios and iterations, while for univariate imputation, the maximum likelihood estimation

underpinning the Kalman smoothing too often failed to converge, halting the simulations. The global

imputation approaches provide upper bounds on the performance of these approaches by incorporating

future information into the imputations while only using a limited window of antecedent imputed

observations to estimate the rate of change on a given day.

<u>Performance Metrics</u>

For each simulated 1000-day surveillance period, we estimated the rate of change by each approach on all days with a corresponding synthetic wastewater viral load measurement after the first 90 days (a three-month baseline data collection period to accommodate the different minimum sample sizes required by each approach), for a total of 260 estimates per approach. The rate of change estimate $RC_{t[i]}^{k}$ for approach $k$ was compared with the sampled GP first derivative $x'_{t[i]}$ on day $t$ corresponding to viral load observation $i$ to assess pointwise estimation performance. The pointwise performance indicators were summarized across all estimates for a given simulated time series to calculate performance metrics for each approach. Performance metric distributions were further characterized as the median, 2.5%, and 97.5% quantiles of each metric across all 1000 simulations of each simulation scenario. Quantitative accuracy was assessed by the root mean square error, $RMSE^{k} = \sqrt{\frac{1}{N}\sum_{i=1}^{N}\left(RC_{t[i]}^{k} - x'_{t[i]}\right)^{2}}$.[17] The 95% CI coverage (the proportion of 95% CIs containing the true rate of change) and average width (distance between the upper and lower 95% CI bounds) served as indicators of quantitative precision.

Each estimate was also classified as increasing or decreasing according to the sign of the point estimate (positive or negative, respectively, as the point estimate was never exactly zero).[18] The rate of change estimate at observation $i$ was considered a true positive (TP) when both the point estimate and true rate of change were positive ($RC_{t[i]}^{k} > 0$ and $x'_{t[i]} > 0$); a true negative (TN) for $RC_{t[i]}^{k} < 0$ and $x'_{t[i]} < 0$; a false positive (FP) for $RC_{t[i]}^{k} > 0$ but $x'_{t[i]} < 0$; and a false negative (FN) for $RC_{t[i]}^{k} < 0$ but $x'_{t[i]} > 0$. We assessed binary classification performance by sensitivity, the proportion of true increasing trends correctly classified as increasing, and specificity, the proportion of true decreasing trends correctly classified as decreasing:

$$senstivity^k = \frac{TP^k}{TP^k + FN^k}$$

$$specificity^k = \frac{TN^k}{TN^k + FP^k}$$

(3)

284    CDC suggests a third class, plateau, corresponding to low rates of change that may not warrant a

285   response.[18,21] In practice, however, plateaus are classified on the basis of a statistical test and identify

286   trends with rates of change that the estimator cannot confidently differentiate from zero. Because the

287   simulated true trend and its rate of change are known exactly, there is no directly equivalent definition

288   available to classify true plateaus for evaluating multiclass performance. We instead incorporated the

289   concept of varying confidence in class predictions by considering the probability that the trend was

290   increasing. We estimated the probability of an increasing trend by computing the proportion of a rate of

291   change estimate's 95% CI that was greater than zero. A CI that covered only positive rates of change

292   was assigned a 100% probability of belonging to the increasing class, while a CI that included only

293   negative rates of change was considered to have a 0% probability of increasing. For a CI that included

294   zero, we computed the probability of an increasing trend by dividing the rate of change value at the CI's

295   upper bound by the width of the CI. We performed receiver operating characteristic (ROC) curve

296   analysis with the *yardstick* package to incorporate trend class probability into the binary classification

297   performance assessment.[47] Sensitivity and specificity were calculated using each observed class

298   probability as the threshold for classifying an increasing trend, which generated an ROC curve tracing

299   the sensitivity-specificity trade-off across probability thresholds. We used the area under the curve

300   (AUC) to assess classification performance when treating the rate of change estimation approach as a

301   probabilistic classifier.[48]

*Application: North Carolina SARS-CoV-2 Wastewater Viral Loads*

We obtained publicly available data on wastewater SARS-CoV-2 per-capita viral loads and

COVID-19 cases for 25 North Carolina sewersheds from the NC Department of Health and Human

Services (NCDHHS) COVID-19 Wastewater Monitoring Dashboard.[49] Detailed procedures for sample

collection, laboratory analysis, and data processing have been described previously.[46,50] Ten sewersheds

began reporting viral loads in January 2021, with nine sewersheds added in June 2021, five more in

October–November 2021, and a single addition in March 2022. We analyzed data collected through

24 May 2023, when COVID-19 case reporting ended statewide. The publicly available data listed a

count of 2 for any day with $1 - 4$ cases to protect privacy and no value (missing) for days with no new

cases, with COVID-19 incidence reported as daily new cases per 10,000 sewershed population. We

scaled by the reported sewershed population and rounded to the nearest integer to recover daily case

counts, substituting 0 for missing counts and a random integer from $1 - 4$ with equal probability for any

recovered counts of 2. Wastewater per-capita viral loads were $\log_{10}$-transformed for all analyses,

yielding units of $\log_{10}$ copies/person/day; data were provided with imputed values already substituted for

non-detects, as described previously.[46,50]

For each sewershed, we estimated the rate of change on the date of each wastewater sample

(after an initial 90-day baseline monitoring period) via the four estimation approaches and classified

each estimate as increasing, decreasing, or plateau.[21] We compared rate of change estimates to the first

derivative of the global trend estimated by the GAM fit to the full dataset for each sewershed ("global

GAM"). Agreement between the global GAM estimates and the four local estimation approaches

informed only by antecedent observations was assessed using the same metrics as for the simulation

study.

*Approval and Availability Statement*

No human participants were involved in this research. All analyses were performed on synthetic

or publicly available, aggregated data and did not require ethical approval. The code and data to perform

these analyses are freely available in a permanent online repository at

https://doi.org/10.17605/OSF.IO/BPGN4 (see Supplemental Material, Analysis Code). The original NC

sewershed monitoring data may be accessed at https://covid19.ncdhhs.gov/dashboard/data-behind-

dashboards.

**Results**

*Simulation Study*

<u>Scenarios</u>

Figure 1 presents examples of the synthetic wastewater viral load data we generated for each

simulation scenario. As expected, the $\rho = 90$ days, $\sigma = 0.5 \log_{10}$ copies/day scenario produced the

smoothest trend and least noisy observations, whereas the $\rho = 15$ days, $\sigma = 1 \log_{10}$ copies/day

scenario produced the most wiggly trend with the noisiest observations. The trend first derivatives,

corresponding to the rate of change, were influenced only by the value of $\rho$ (Figure 1b). The smoothest

scenarios ($\rho = 90$) produced rates of change with the smallest magnitudes, while the wigglier $\rho = 15$

scenarios produced much larger magnitude rates of change.

344
345 Figure 1. Illustrative realizations of (a) simulated wastewater viral load trend (black line) and synthetic
346 observations (blue points) and (b) first derivative of simulated wastewater viral load trend for nine
347 scenarios with varying specifications of the Gaussian process (GP) range parameter ρ and independent
348 random error standard deviation $\sigma$
349

350   Rate of Change Estimates

351           Under the moderate $\rho = 30$ days, $\sigma = 0.75 \log_{10}$ copies/day simulation scenario, the global

GAM approach generally produced smooth estimates that largely tracked both the simulated trend

(Figure 2a) and its rate of change (Figure 2b). By contrast, the four local estimation approaches yielded

more disjointed estimates that broadly oscillated around the true rate of change, with the rolling GAM

and univariate imputation point estimates appearing to track the truth more closely and the rolling linear

model swinging more dramatically between estimates. The rolling linear model estimates also exhibited

the highest uncertainty, with the widest 95% CIs on average (Table S1); univariate imputation estimates

typically had the narrowest CIs, which frequently did not include the true rate of change given by the GP

derivative. Both the global and rolling GAM estimates generally covered the true rate of change with

their 95% CIs. The rolling GAM estimates had greater uncertainty. This uncertainty resulted because

each estimate was made at the extreme of the range of the observed data without the benefit of future

observations to the right of the estimation point that were available to the global GAM (except for the

final observation, for which the approaches, as expected, converged to identical estimates).[40] Both

imputation approaches and the rolling GAM also appeared to lag somewhat during periods of more

rapid change in the trend, observable in Figure 2b as the right-ward shift in the estimated rate of change

relative to the GP derivative.

367
368 Figure 2. True (simulated) and estimated (a) viral load trend and (b) rate of change in viral load by each
369 of the candidate estimation approaches for one realization of the moderately smooth, moderately noisy
370 ($\rho = 30, \sigma = 0.75$) simulation scenario. The synthetic measurements used to fit all models are
371 displayed as green points.
372

 Estimation Performance

Across all nine scenarios, the global GAM—which utilized more data than an ongoing, real-time

wastewater monitoring program would have access to—consistently produced the most accurate

estimates, as indicated by lowest RMSE (Figure 3a). For the smoothest scenarios ($\rho = 90$ days), the

rolling GAM exhibited similarly high accuracy regardless of the magnitude of the noise parameter $\sigma$,

followed by the univariate imputation approach, multivariate imputation, and finally by the rolling linear

model approach, which was considerably less accurate and more impacted by increasing noise variance.

However, the differences in accuracy between approaches diminished for less-smooth trends as RMSE

increased, such that the RMSE distributions were similar across all four local estimation approaches

(median RMSE: $0.06 - 0.08$ $\Delta\log_{10}$ copies/day) for the least-smooth ($\rho = 15$) scenarios with low ($\sigma =$

$0.5$) and moderate ($\sigma = 0.75$) noise variance.

Although the rolling linear model estimates generally had the widest 95% CIs (Figure 2b), they

also most consistently included the true rate of change in about 95% of intervals, the target coverage

proportion (Figure 3b). The uncertainty of global GAM estimates, while much narrower than for the

rolling linear model, was overly conservative, with median coverage >95% for the smoother and less

noisy scenarios. Univariate imputation had coverage proportions appreciably <95% for all but the

smoothest scenarios. The multivariate imputation approach also generally had coverage proportions that

were <95%. The median 95% CI coverage of rolling and global GAM estimates remained relatively

high across all scenarios, but both approaches demonstrated large variability in interval coverage across

iterations of the more challenging (less smooth, noisier) scenarios. GAMs appear to be susceptible to

estimating inappropriately smooth trends under such conditions, as observed in the essentially flat trend

with narrow 95% CIs estimated by the global GAM for the $\rho = 30, \sigma = 1$ scenario in Figure S4.[26] Such

over-smoothing appears to occur more frequently when GAMs are fit to high-variance (relative to $\alpha$)

synthetic measurements generated from wiggly trends, which can produce an essentially uniform cloud

of observations that envelop and obscure the trend ($\rho = 30, \sigma = 1$ scenario in Figure 1a).

The relative performance of each approach at correctly classifying the trend as increasing or

decreasing was consistent across the nine simulation scenarios (Figure 3c). The global GAM

consistently demonstrated the highest median AUC, although AUCs were more variable across

iterations of the wigglier and noisier scenarios. Under the smoothest trend scenarios ($\rho = 90$), the

rolling GAM estimates had the highest median AUC among the local estimation approaches, but also

greater variability in AUC. Rolling linear models exhibited the lowest median AUC at just over 50%—

only slightly better than chance. Intriguingly, the AUC of rolling linear models improved as trend

smoothness decreased and the AUC of all other approaches degraded; for the wiggliest scenarios

($\rho = 15$), the rolling linear model median AUC was equal to or greater than any of the other local

models. The rolling GAM approach provided the least accurate trend classifications in the least smooth,

most noisy scenario, likely related to the propensity for GAMs to over-smooth under such conditions.[26]

By contrast, the univariate imputation approach performed fairly consistently across less-smooth

scenarios, regardless of the noise variance, potentially due to the overly precise estimates providing

more weight when the sign was correct (as was the case for the majority of estimates). However, no

local estimation approach consistently achieved median sensitivities, specificities, or AUCs above 80%

for any scenario (Table S1), indicating that none of these approaches would reliably classify pandemic

trends as increasing or decreasing in near real-time as part of an ongoing wastewater monitoring

program.

## (a) root mean square error

## (b) 95% CI coverage

## (c) area under the curve (binary classifier)

Figure 3. Median and 2.5$^{th}$ – 97.5$^{th}$ percentiles of performance metrics for candidate rate of change
estimation approaches across 1000 realizations of each of the nine simulation scenarios. Vertical dotted
lines indicate the target performance for each metric.

The 25 NC sewersheds that we analyzed served populations ranging from 3500 – 550,000 people

and were monitored over periods of 431 – 871 days, with SARS-CoV-2 viral loads reported for 121 –

245 wastewater samples at each site (Table S2). The lowest median per-capita viral loads were observed

in the Wilmington sewershed, at 3.7 million copies/person/day (interquartile range [IQR] 7.6 million

copies/person/day), with the highest viral loads observed at the Cary 3 site (median (IQR): 36.8 (44.3)

million copies/person/day). The measured wastewater viral loads and global GAM-estimated trend,

along with the corresponding rate of change estimates by each approach assessed in the simulation

study, are presented separately for each sewershed in Figures S7 – S31. Concordance between the global

GAM rate of change estimates and the local model estimates were similar between approaches except

for rolling linear model estimates, which exhibited somewhat higher and more variable RMSE across

the 25 sewersheds (Figure S6). As in the simulation study, the rolling linear model estimates had the

widest 95% CIs but also included the mean global GAM estimate at approximately the target proportion

of 95%. Likewise, the univariate imputation approach had the narrowest average 95% CIs, which failed

to cover the majority of GAM estimates.

Under the three-class system, in which rate of change estimates are classified as increasing or

decreasing only when their 95% confidence intervals exclude zero, the univariate imputation approach

also consistently identified the highest proportion of estimates as clearly increasing or decreasing and

the fewest as plateaus across all sites (Figure 4). The majority of estimates were classified as plateaus by

each of the other approaches, while only one site (Roanoke Rapids) was majority plateau (53%) by the

univariate imputation approach (Table S3). The proportion of estimates deemed plateaus was highest

(often >90%) by either the rolling linear model or the rolling GAM approaches, except in Jacksonville,

where the global GAM classified all but one estimate as plateau (Figure S20). Approximately three-

quarters of global GAM estimates and two-thirds of multivariate imputation estimates were identified as

plateaus. The relative frequency at which each estimation approach classified trends as plateaus matched

between the NC data and the simulated data under the moderate ($\rho = 30$) and least smooth ($\rho = 15$)

scenarios, although the global GAM produced fewer plateau calls than did the univariate imputation for

the smoothest ($\rho = 90$) simulation scenarios (Figure S5). All approaches identified both increasing and

decreasing trends at each site, but we observed appreciable variation between approaches in the ratio of

increasing to decreasing trend classifications.

Figure 4. Percentage of estimates classified as decreasing (purple), increasing (mauve), or plateau (orange) trends by each rate of change estimation approach for 25 North Carolina Sewersheds.

**Discussion**

Approaches for classifying viral load trends and estimating their rates of change in wastewater

monitoring programs have not, to our knowledge, previously been compared for accuracy and reliability.

Because trends and their slopes are not directly measured, we generated synthetic wastewater viral load

time series using Gaussian processes to simulate a range of potential patterns of disease trends with

known first derivatives. We implemented four rate of change estimation approaches—two previously

reported and two developed herein—representing routine trend assessments from a typical, twice-

weekly wastewater monitoring program that could reasonably be utilized by public health authorities

without specialized statistical expertise. When applied to the synthetic time series data, all of the

approaches showed only modest reliability in identifying the correct direction of the trend. The median

agreement between the signs of the estimated and true rates of change typically ranged between 50%

and 75%, corresponding to at least one out of every four trend assessments (i.e., once every two weeks)

providing misleading conclusions about the direction of the trend. This result raises concerns about the

ability to take appropriate action (e.g., issuing alerts in response to apparently increasing trends)

informed by such trend classification approaches.

The addition of plateau-class trends in a three-class system partially addresses this modest

reliability by requiring greater certainty before identifying trends as increasing or decreasing. As

currently defined in wastewater surveillance applications, plateau status is not an inherent property of

the trend but rather the product of a conventional decision criterion that the rate of change estimate was

not significantly different from zero at the 5% significance level.[51] Both the magnitude of the estimate

and the precision with which it was estimated affect whether statistical significance was achieved,

meaning that trends classified as plateau may have been changing too slowly to warrant attention or that

the rate of change estimate was too uncertain to confidently determine the direction of change. The

choice of estimation approach consistently impacted the proportion of estimates deemed plateau, though

these patterns appeared mostly related to the precision of the rate of change estimates, with the

inappropriately precise imputation-based approaches identifying a larger proportion of non-plateau

trends. The plateau concept implies a trade-off between making a potentially incorrect determination

(e.g., classifying as decreasing a trend that is truly increasing) against failing to make a determination

(i.e., classifying as plateau) when conditions are in fact meaningfully changing. However, by conflating

two distinct concerns—the magnitude of change and estimation uncertainty—the existing plateau

definition does not directly address key questions of what is a meaningful rate of change to warrant

further attention and how to balance the costs of incorrect action vs. inaction when meaningful changes

in the trend are underway. Such considerations may be informally addressed by selecting more or less

aggressive classification approaches or by varying the significance threshold. Greater transparency

would be afforded by explicitly specifying the threshold above which rates of change would be

considered meaningful, as when Keshaviah et al. specified the doubling of wastewater viral quantities as

a component of an algorithm to detect infection surges.[36] As of December 2023, CDC NWSS similarly

published fixed categories defined by the percent change in viral load over the previous 15 days to

classify trends, with changes $\geq 100\%$ considered "large" increases.[52,53]

Both quantitative and classification accuracy differed between approaches, but the differences

were generally observed only for smoother trends. As trends oscillated more rapidly with decreasing $\rho$,

method performance degraded such that all were essentially equally poor. This dependence on

characteristics of the trend itself suggests that even highly performing approaches may not be suitable in

all contexts and anticipated trend smoothness should be considered when selecting a rate of change

estimation approach to apply at a given monitoring site and time. While trend smoothness may be

directly characterized by estimating GP range hyperparameters from previously collected monitoring

data, fitting GPs is non-trivial—particularly estimating the weakly identified covariance function

hyperparameters—and may be generally infeasible outside academic settings.[43,54,55] Informal

smoothness assessments may prove sufficient, for example by visually comparing global GAM-

estimated trends to representative GP simulations across a range of $\rho$ values. Both strategies rely on the

assumption that future trends will be similar to those already observed, for which adequate data to

characterize trend smoothness may be unavailable, particularly after the emergence of a novel pathogen.

To aid interpretability, we simulated wastewater trends using a simple GP with zero global mean

and a squared exponential covariance function with constant marginal standard deviation $\alpha$ and a single

temporal range hyperparameter $\rho$ across the entire simulated monitoring period. These conditions do not

fully reflect the real-world complexity of infectious disease trends. However, the flexibility of GPs

readily allows extension of our simulation approach to represent smooth trends of far greater

complexity. For example, the GP may be specified with multiple additive Matérn covariance functions

(of which the squared exponential is a special case) to introduce fluctuations at multiple time scales, or

with alternative covariance structures that represent specific physical processes (though non-Matérn

functions may not be readily differentiable).[23,43,46,56,57]

The broad utility of wastewater surveillance, particularly in detecting emerging pathogen

lineages, has been widely demonstrated.[58–60] However, effectively leveraging wastewater measurements

alone to estimate the rate of change of disease trends in real-time (i.e., on the day of the most recent

measurement) has proved particularly challenging. While retrospective estimation of temporal trends

and their rates of change by global GAMs was reasonably accurate in our simulation study and credible

when applied to observed SARS-CoV-2 viral loads in NC sewersheds, real-time estimates are made at

the extreme range of the data where uncertainty is greatest (as illustrated by the rolling GAM's much

noisier estimates and wider CIs relative to the global GAM). Coupling the most recent viral load

measurement with recent estimated trend values provides a good indication of the neighborhood of

potential values the current trend may take, but is less informative about where the trend is heading.[61]

Rather than attempting to estimate instantaneous rates of change, relative metrics with more

retrospective features, such as the more recently developed CDC NWSS Wastewater Viral Activity

Level metric that compares current viral loads to a long-running, site-specific baseline value, may offer

a more reliable basis for understanding community disease burdens using only wastewater data.[62] As

wastewater surveillance data become increasingly relied on, any such wastewater-only metric must be

thoroughly evaluated before being used to identify public health-relevant differences in community

disease burden. We suggest that additional studies focus on using wastewater measurements in context

with other public health metrics, such as hospitalizations or emergency department visits, to enhance

predictions and updating current models to adapt to changing disease dynamics.[63]

**Acknowledgment**

We gratefully acknowledge the assistance of Mitham Al-Faliti and Jeseth Delgado Vega in

implementing the multivariate imputation approach. Megan Lott and Joe Brown provided insightful

commentary. This work was supported by the CDC National Wastewater Surveillance System through

the Epidemiology and Laboratory Capacity Cooperative Agreement with North Carolina Department of

Health and Human Services, with additional support from the National Institute for Occupational Health

and Safety (T42OH008673) and the NSF RAPID program (project #2029866). We also thank Steven

Berkowitz and all contributors to the NC Wastewater Monitoring Network, as well as its earlier

incarnation as the NC Wastewater Pathogen Research Network, which received crucial support from the

NC Policy Collaborative.

# References

(1)  Naughton, C. C.; Roman, F. A., Jr; Alvarado, A. G. F.; Tariqi, A. Q.; Deeming, M. A.; Kadonsky, K. F.; Bibby, K.; Bivins, A.; Medema, G.; Ahmed, W.; Katsivelis, P.; Allan, V.; Sinclair, R.; Rose, J. B. Show Us the Data: Global COVID-19 Wastewater Monitoring Efforts, Equity, and Gaps. *FEMS Microbes* **2023**, *4*, xtad003. https://doi.org/10.1093/femsmc/xtad003.

(2)  Bivins, A.; North, D.; Ahmad, A.; Ahmed, W.; Alm, E.; Been, F.; Bhattacharya, P.; Bijlsma, L.; Boehm, A. B.; Brown, J.; Buttiglieri, G.; Calabro, V.; Carducci, A.; Castiglioni, S.; Cetecioglu Gurol, Z.; Chakraborty, S.; Costa, F.; Curcio, S.; de los Reyes, F. L.; Delgado Vela, J.; Farkas, K.; Fernandez-Casi, X.; Gerba, C.; Gerrity, D.; Girones, R.; Gonzalez, R.; Haramoto, E.; Harris, A.; Holden, P. A.; Islam, Md. T.; Jones, D. L.; Kasprzyk-Hordern, B.; Kitajima, M.; Kotlarz, N.; Kumar, M.; Kuroda, K.; La Rosa, G.; Malpei, F.; Mautus, M.; McLellan, S. L.; Medema, G.; Meschke, J. S.; Mueller, J.; Newton, R. J.; Nilsson, D.; Noble, R. T.; van Nuijs, A.; Peccia, J.; Perkins, T. A.; Pickering, A. J.; Rose, J.; Sanchez, G.; Smith, A.; Stadler, L.; Stauber, C.; Thomas, K.; van der Voorn, T.; Wigginton, K.; Zhu, K.; Bibby, K. Wastewater-Based Epidemiology: Global Collaborative to Maximize Contributions in the Fight Against COVID-19. *Environmental Science & Technology* **2020**, acs.est.0c02388. https://doi.org/10.1021/acs.est.0c02388.

(3)  Levy, J. I.; Andersen, K. G.; Knight, R.; Karthikeyan, S. Wastewater Surveillance for Public Health. *Science* **2023**, *379* (6627), 26–27. https://doi.org/10.1126/science.ade2503.

(4)  Rao, G.; Capone, D.; Zhu, K.; Knoble, A.; Linden, Y.; Clark, R.; Lai, A.; Kim, J.; Huang, C.-H.; Bivins, A.; Brown, J. Simultaneous Detection and Quantification of Multiple Pathogen Targets in Wastewater. *PLOS Water* **2024**, *3* (2), e0000224. https://doi.org/10.1371/journal.pwat.0000224.

(5)  Valdivia-Carrera, C. A.; Ho-Palma, A. C.; Munguia-Mercado, A.; Gonzalez-Pizarro, K.; Ibacache-Quiroga, C.; Dinamarca, A.; Stehlík, M.; Rusiñol, M.; Girones, R.; Lopez-Urbina, M. T.; Basaldua Galarza, A.; Gonzales-Gustavson, E. Surveillance of SARS-CoV-2, Rotavirus, Norovirus Genogroup II, and Human Adenovirus in Wastewater as an Epidemiological Tool to Anticipate Outbreaks of COVID-19 and Acute Gastroenteritis in a City without a Wastewater Treatment Plant in the Peruvian Highlands. *Science of The Total Environment* **2023**, 167161. https://doi.org/10.1016/j.scitotenv.2023.167161.

(6)  Thai, P. K.; O'Brien, J. W.; Banks, A. P. W.; Jiang, G.; Gao, J.; Choi, P. M.; Yuan, Z.; Mueller, J. F. Evaluating the In-Sewer Stability of Three Potential Population Biomarkers for Application in Wastewater-Based Epidemiology. *Science of The Total Environment* **2019**, *671*, 248–253. https://doi.org/10.1016/j.scitotenv.2019.03.231.

(7)  Roy, S.; Tang, M.; Edwards, M. A. Lead Release to Potable Water during the Flint, Michigan Water Crisis as Revealed by Routine Biosolids Monitoring Data. *Water Research* **2019**, *160*, 475–483. https://doi.org/10.1016/j.watres.2019.05.091.

(8)  Graham, K. E.; Loeb, S. K.; Wolfe, M. K.; Catoe, D.; Sinnott-Armstrong, N.; Kim, S.; Yamahara, K. M.; Sassoubre, L. M.; Mendoza Grijalva, L. M.; Roldan-Hernandez, L.; Langenfeld, K.; Wigginton, K. R.; Boehm, A. B. SARS-CoV-2 RNA in Wastewater Settled Solids Is Associated with COVID-19 Cases in a Large Urban Sewershed. *Environ. Sci. Technol.* **2021**, *55* (1), 488–498. https://doi.org/10.1021/acs.est.0c06191.

(9)  Mantilla-Calderon, D.; Huang, K. (Kevin); Li, A.; Chibwe, K.; Yu, X.; Ye, Y.; Liu, L.; Ling, F. Meta-Analyses on SARS-CoV-2 Viral RNA Levels in Wastewater and Their Correlations to Epidemiological Indicators. *Environ. Sci.: Water Res. Technol.* **2022**, *8* (7), 1391–1407. https://doi.org/10.1039/D2EW00084A.

(10) Hoar, C.; Chauvin, F.; Clare, A.; McGibbon, H.; Castro, E.; Patinella, S.; Katehis, D.; Dennehy, J. J.; Trujillo, M.; Smyth, D. S.; Silverman, A. I. Monitoring SARS-CoV-2 in Wastewater during New York City's Second Wave of COVID-19: Sewershed-Level Trends and Relationships to

Publicly Available Clinical Testing Data. *Environ. Sci.: Water Res. Technol.* **2022**, *8* (5), 1021–
1035. https://doi.org/10.1039/D1EW00747E.

(11) Lott, M. E. J.; Norfolk, W. A.; Dailey, C. A.; Foley, A. M.; Melendez-Declet, C.; Robertson, M. J.;
Rathbun, S. L.; Lipp, E. K. Direct Wastewater Extraction as a Simple and Effective Method for
SARS-CoV-2 Surveillance and COVID-19 Community-Level Monitoring. *FEMS Microbes* **2023**,
*4*. https://doi.org/10.1093/femsmc/xtad004.

(12) Soller, J.; Jennings, W.; Schoen, M.; Boehm, A.; Wigginton, K.; Gonzalez, R.; Graham, K. E.;
McBride, G.; Kirby, A.; Mattioli, M. Modeling Infection from SARS-CoV-2 Wastewater
Concentrations: Promise, Limitations, and Future Directions. *Journal of Water and Health* **2022**.
https://doi.org/10.2166/wh.2022.094.

(13) Peccia, J.; Zulli, A.; Brackney, D. E.; Grubaugh, N. D.; Kaplan, E. H.; Casanovas-Massana, A.;
Ko, A. I.; Malik, A. A.; Wang, D.; Wang, M.; Warren, J. L.; Weinberger, D. M.; Arnold, W.;
Omer, S. B. Measurement of SARS-CoV-2 RNA in Wastewater Tracks Community Infection
Dynamics. *Nat Biotechnol* **2020**, *38* (10), 1164–1167. https://doi.org/10.1038/s41587-020-0684-z.

(14) Fitzgerald, S. F.; Rossi, G.; Low, A. S.; McAteer, S. P.; O'Keefe, B.; Findlay, D.; Cameron, G. J.;
Pollard, P.; Singleton, P. T. R.; Ponton, G.; Singer, A. C.; Farkas, K.; Jones, D.; Graham, D. W.;
Quintela-Baluja, M.; Tait-Burkard, C.; Gally, D. L.; Kao, R.; Corbishley, A. Site Specific
Relationships between COVID-19 Cases and SARS-CoV-2 Viral Load in Wastewater Treatment
Plant Influent. *Environ. Sci. Technol.* **2021**. https://doi.org/10.1021/acs.est.1c05029.

(15) Omori, R.; Miura, F.; Kitajima, M. Age-Dependent Association between SARS-CoV-2 Cases
Reported by Passive Surveillance and Viral Load in Wastewater. *Science of The Total*
*Environment* **2021**, *792*, 148442. https://doi.org/10.1016/j.scitotenv.2021.148442.

(16) McMahan, C. S.; Self, S.; Rennert, L.; Kalbaugh, C.; Kriebel, D.; Graves, D.; Colby, C.; Deaver, J.
A.; Popat, S. C.; Karanfil, T.; Freedman, D. L. COVID-19 Wastewater Epidemiology: A Model to
Estimate Infected Populations. *The Lancet Planetary Health* **2021**, *5* (12), e874–e881.
https://doi.org/10.1016/s2542-5196(21)00230-8.

(17) Huisman, J. S.; Scire, J.; Caduff, L.; Fernandez-Cassi, X.; Ganesanandamoorthy, P.; Kull, A.;
Scheidegger, A.; Stachler, E.; Boehm, A. B.; Hughes, B.; Knudson, A.; Topol, A.; Wigginton, K.
R.; Wolfe, M. K.; Kohn, T.; Ort, C.; Stadler, T.; Julian, T. R. Wastewater-Based Estimation of the
Effective Reproductive Number of SARS-CoV-2. *Environmental Health Perspectives* **2022**, *130*
(5), 057011. https://doi.org/10.1289/EHP10050.

(18) Kotlarz, N.; Holcomb, D. A.; Pasha, A. B. M. T.; Reckling, S.; Kays, J.; Lai, Y.-C.; Daly, S.;
Palani, S.; Bailey, E.; Guidry, V. T.; Christensen, A.; Berkowitz, S.; Hoppin, J. A.; Mitasova, H.;
Engel, L. S.; de los Reyes, F. L.; Harris, A. Timing and Trends for Municipal Wastewater, Lab-
Confirmed Case, and Syndromic Case Surveillance of COVID-19 in Raleigh, North Carolina.
*American Journal of Public Health* **2023**, *113* (1), 79–88.
https://doi.org/10.2105/AJPH.2022.307108.

(19) Chen, W.; Bibby, K. Making Waves: Establishing a Modeling Framework to Evaluate Novel
Targets for Wastewater-Based Surveillance. *Water Research* **2023**, *245*, 120573.
https://doi.org/10.1016/j.watres.2023.120573.

(20) Al-Faliti, M.; Kotlarz, N.; McCall, C.; Harris, A. R.; Smith, A. L.; Stadler, L. B.; de los Reyes, F.
L.; Delgado Vela, J. Comparing Rates of Change in SARS-CoV-2 Wastewater Load and Clinical
Cases in 19 Sewersheds Across Four Major Metropolitan Areas in the United States. *ACS EST*
*Water* **2022**. https://doi.org/10.1021/acsestwater.2c00106.

(21) CDC. *Wastewater Surveillance Data Reporting and Analytics: Trends*; Website; Centers for
Disease Control and Prevention, 2023.
https://web.archive.org/web/20230923022331/https://www.cdc.gov/nwss/reporting.html.

(22) Kirby, A. E.; Walters, M. S.; Jennings, W. C.; Fugitt, R.; LaCross, N.; Mattioli, M.; Marsh, Z. A.;
Roberts, V. A.; Mercante, J. W.; Yoder, J.; Hill, V. R. Using Wastewater Surveillance Data to

Support the COVID-19 Response — United States, 2020–2021. *MMWR Morb. Mortal. Wkly.*
*Rep.* **2021**, *70* (36), 1242–1244. https://doi.org/10.15585/mmwr.mm7036a2.
(23) Rasmussen, C. E.; Williams, C. K. I. *Gaussian Processes for Machine Learning*; The MIT Press,
2005. https://doi.org/10.7551/mitpress/3206.001.0001.
(24) Solak, E.; Murray-smith, R.; Leithead, W.; Leith, D.; Rasmussen, C. Derivative Observations in
Gaussian Process Models of Dynamic Systems. In *Advances in Neural Information Processing*
*Systems*; MIT Press, 2002; Vol. 15.
https://proceedings.neurips.cc/paper_files/paper/2002/file/5b8e4fd39d9786228649a8a8bec4e008-
Paper.pdf.
(25) McHutchon, A. J. *Nonlinear Modelling and Control Using Gaussian Processes*; Dissertation;
University of Cambridge, 2014. https://mlg.eng.cam.ac.uk/pub/pdf/Mch14.pdf.
(26) Arabzadeh, R.; Grünbacher, D. M.; Insam, H.; Kreuzinger, N.; Markt, R.; Rauch, W. Data Filtering
Methods for SARS-CoV-2 Wastewater Surveillance. *Water Science and Technology* **2021**, *84* (6),
1324–1339. https://doi.org/10.2166/wst.2021.343.
(27) Wood, S. N. *Generalized Additive Models: An Introduction with R*, 2nd ed.; Chapman and
Hall/CRC, 2017. https://doi.org/10.1201/9781315370279.
(28) R Core Team. *R: A Language and Environment for Statistical Computing*; Software version 4.2.2;
Vienna, Austria, 2022. https://www.R-project.org/.
(29) Wickham, H.; Averick, M.; Bryan, J.; Chang, W.; McGowan, L. D.; François, R.; Grolemund, G.;
Hayes, A.; Henry, L.; Hester, J.; Kuhn, M.; Pedersen, T. L.; Miller, E.; Bache, S. M.; Müller, K.;
Ooms, J.; Robinson, D.; Seidel, D. P.; Spinu, V.; Takahashi, K.; Vaughan, D.; Wilke, C.; Woo,
K.; Yutani, H. Welcome to the tidyverse. *Journal of Open Source Software* **2019**, *4* (43), 1686.
https://doi.org/10.21105/joss.01686.
(30) Vaughan, D.; Dancho, M. *Tibbletime: Time Aware Tibbles*; R package version 0.1.8.9000; 2023.
https://github.com/business-science/tibbletime.
(31) CDC. *Developing a Wastewater Surveillance Sampling Strategy*; Website; Centers for Disease
Control and Prevention, 2023.
https://web.archive.org/web/20231030161926/https://www.cdc.gov/nwss/sampling.html.
(32) Cluzel, N.; Courbariaux, M.; Wang, S.; Moulin, L.; Wurtzer, S.; Bertrand, I.; Laurent, K.; Monfort,
P.; Gantzer, C.; Guyader, S. L.; Boni, M.; Mouchel, J.-M.; Maréchal, V.; Nuel, G.; Maday, Y. A
Nationwide Indicator to Smooth and Normalize Heterogeneous SARS-CoV-2 RNA Data in
Wastewater. *Environment International* **2022**, *158*, 106998.
https://doi.org/10.1016/j.envint.2021.106998.
(33) van Buuren, S.; Groothuis-Oudshoorn, K. Mice: Multivariate Imputation by Chained Equations in
R. *Journal of Statistical Software* **2011**, *45* (3). https://doi.org/10.18637/jss.v045.i03.
(34) van Buuren, S. *Flexible Imputation of Missing Data, Second Edition*, 2nd ed.; Chapman and
Hall/CRC: New York, 2018. https://doi.org/10.1201/9780429492259.
(35) Moritz, S.; Bartz-Beielstein, T. imputeTS: Time Series Missing Value Imputation in R. *The R*
*Journal* **2017**, *9* (1), 207–218. https://doi.org/10.32614/RJ-2017-009.
(36) Keshaviah, A.; Huff, I.; Hu, X. C.; Guidry, V.; Christensen, A.; Berkowitz, S.; Reckling, S.; Noble,
R. T.; Clerkin, T.; Blackwood, D.; McLellan, S. L.; Roguet, A.; Musse, I. Separating Signal from
Noise in Wastewater Data: An Algorithm to Identify Community-Level COVID-19 Surges in
Real Time. *Proceedings of the National Academy of Sciences* **2023**, *120* (31), e2216021120.
https://doi.org/10.1073/pnas.2216021120.
(37) Rauch, W.; Schenk, H.; Insam, H.; Markt, R.; Kreuzinger, N. Data Modelling Recipes for SARS-
CoV-2 Wastewater-Based Epidemiology. *Environmental Research* **2022**, *214*, 113809.
https://doi.org/10.1016/j.envres.2022.113809.
(38) Simpson, G. L. Modelling Palaeoecological Time Series Using Generalised Additive Models.
*Frontiers in Ecology and Evolution* **2018**, *6*. https://doi.org/10.3389/fevo.2018.00149.

(39) Bertels, X.; Hanoteaux, S.; Janssens, R.; Maloux, H.; Verhaegen, B.; Delputte, P.; Boogaerts, T.;
van Nuijs, A. L. N.; Brogna, D.; Linard, C.; Marescaux, J.; Didy, C.; Pype, R.; Roosens, N. H. C.;
Van Hoorde, K.; Lesenfants, M.; Lahousse, L. Time Series Modelling for Wastewater-Based
Epidemiology of COVID-19: A Nationwide Study in 40 Wastewater Treatment Plants of
Belgium, February 2021 to June 2022. *Science of The Total Environment* **2023**, *899*, 165603.
https://doi.org/10.1016/j.scitotenv.2023.165603.
(40) Clark, N. J.; Wells, K. Dynamic Generalised Additive Models (DGAMs) for Forecasting Discrete
Ecological Time Series. *Methods in Ecology and Evolution* **2022**. https://doi.org/10.1111/2041-
210x.13974.
(41) Wood, S. N. Fast Stable Restricted Maximum Likelihood and Marginal Likelihood Estimation of
Semiparametric Generalized Linear Models. *Journal of the Royal Statistical Society: Series B
(Statistical Methodology)* **2011**, *73* (1), 3–36. https://doi.org/10.1111/j.1467-9868.2010.00749.x.
(42) Simpson, G. L. *Gratia: Graceful Ggplot-Based Graphics and Other Functions for GAMs Fitted
Using Mgcv*; R package version 0.8.1.34; 2023. https://gavinsimpson.github.io/gratia/.
(43) Betancourt, M. *Robust Gaussian Process Modeling*; Blog Post; 2020.
https://web.archive.org/web/20240320133727/https://betanalpha.github.io/assets/case_studies/gau
ssian_processes.html.
(44) Susmann, H. *Derivatives of a Gaussian Process*; Blog post; 2020.
https://web.archive.org/web/20240410015709/http://herbsusmann.com/2020/07/06/gaussian-
process-derivatives/.
(45) Fasiolo, M. *An Introduction to Mvnfast.*; R package version 0.2.7; 2016.
http://mfasiolo.github.io/mvnfast/.
(46) Hoffman, K.; Holcomb, D.; Reckling, S.; Clerkin, T.; Blackwood, D.; Beattie, R.; de los Reyes, F.;
Harris, A.; Mitasova, H.; Kotlarz, N.; Stewart, J.; Kazenelson, J.; Cahoon, L.; Frampton, A.;
Munir, M.; Lee, A.; Berkowitz, S.; Noble, R.; Guidry, V. T.; Engel, L.; Serre, M.; Christensen, A.
Using Detrending to Assess SARS-CoV-2 Wastewater Loads as a Leading Indicator of
Fluctuations in COVID-19 Cases at Fine Temporal Scales: Correlations across Twenty
Sewersheds in North Carolina. *PLOS Water* **2023**, *2* (10), e0000140.
https://doi.org/10.1371/journal.pwat.0000140.
(47) Kuhn, M.; Vaughan, D.; Hvitfeldt, E. *Yardstick: Tidy Characterizations of Model Performance*; R
package version 1.2.0; 2023. https://yardstick.tidymodels.org.
(48) Fawcett, T. An Introduction to ROC Analysis. *Pattern Recognition Letters* **2006**, *27* (8), 861–874.
https://doi.org/10.1016/j.patrec.2005.10.010.
(49) NCDHHS. *Wastewater Monitoring Dashboard*; Website; North Carolina Department of Health and
Human Services, 2023.
https://web.archive.org/web/20231206225713/https://covid19.ncdhhs.gov/dashboard/wastewater-
monitoring.
(50) Beattie, R. E.; Blackwood, A. D.; Clerkin, T.; Dinga, C.; Noble, R. T. Evaluating the Impact of
Sample Storage, Handling, and Technical Ability on the Decay and Recovery of SARS-CoV-2 in
Wastewater. *PLOS ONE* **2022**, *17* (6), 1–16. https://doi.org/10.1371/journal.pone.0270659.
(51) Greenland, S. Divergence versus Decision P-Values: A Distinction Worth Making in Theory and
Keeping in Practice: Or, How Divergence P-Values Measure Evidence Even When Decision P-
Values Do Not. *Scandinavian Journal of Statistics* **2023**, *50* (1), 54–88.
https://doi.org/10.1111/sjos.12625.
(52) CDC. *COVID Data Tracker: Wastewater Surveillance*; Website; Centers for Disease Control and
Prevention, 2023. https://web.archive.org/web/20231217025913/https://covid.cdc.gov/covid-data-
tracker/#wastewater-surveillance.
(53) CDC. *Using Wastewater Data to Inform Public Health Action*; Website; Centers for Disease
Control and Prevention, 2023.

https://web.archive.org/web/20231209003517/https://www.cdc.gov/respiratory-viruses/whats-
new/using-wastewater-data.html.

(54) Zhang, H. Inconsistent Estimation and Asymptotically Equal Interpolations in Model-Based
Geostatistics. *Journal of the American Statistical Association* **2004**, *99* (465), 250–261.
https://doi.org/10.1198/016214504000000241.

(55) Stan Development Team. 10.3 Fitting a Gaussian Process. In *Stan User's Guide*; 2022. https://mc-
stan.org/docs/stan-users-guide/fit-gp.html#fit-gp.section.

(56) Holcomb, D. A.; Messier, K. P.; Serre, M. L.; Rowny, J. G.; Stewart, J. R. Geostatistical Prediction
of Microbial Water Quality Throughout a Stream Network Using Meteorology, Land Cover, and
Spatiotemporal Autocorrelation. *Environmental Science & Technology* **2018**, *52* (14), 7775–7784.
https://doi.org/10.1021/acs.est.8b01178.

(57) Peterson, E. E.; Theobald, D. M.; Ver Hoef, J. M. Geostatistical Modelling on Stream Networks:
Developing Valid Covariance Matrices Based on Hydrologic Distance and Stream Flow.
*Freshwater Biology* **2007**, *52* (2), 267–279. https://doi.org/10.1111/j.1365-2427.2006.01686.x.

(58) Karthikeyan, S.; Levy, J. I.; De Hoff, P.; Humphrey, G.; Birmingham, A.; Jepsen, K.; Farmer, S.;
Tubb, H. M.; Valles, T.; Tribelhorn, C. E.; Tsai, R.; Aigner, S.; Sathe, S.; Moshiri, N.; Henson,
B.; Mark, A. M.; Hakim, A.; Baer, N. A.; Barber, T.; Belda-Ferre, P.; Chacón, M.; Cheung, W.;
Cresini, E. S.; Eisner, E. R.; Lastrella, A. L.; Lawrence, E. S.; Marotz, C. A.; Ngo, T. T.;
Ostrander, T.; Plascencia, A.; Salido, R. A.; Seaver, P.; Smoot, E. W.; McDonald, D.; Neuhard,
R. M.; Scioscia, A. L.; Satterlund, A. M.; Simmons, E. H.; Abelman, D. B.; Brenner, D.; Bruner,
J. C.; Buckley, A.; Ellison, M.; Gattas, J.; Gonias, S. L.; Hale, M.; Hawkins, F.; Ikeda, L.; Jhaveri,
H.; Johnson, T.; Kellen, V.; Kremer, B.; Matthews, G.; McLawhon, R. W.; Ouillet, P.; Park, D.;
Pradenas, A.; Reed, S.; Riggs, L.; Sanders, A.; Sollenberger, B.; Song, A.; White, B.; Winbush,
T.; Aceves, C. M.; Anderson, C.; Gangavarapu, K.; Hufbauer, E.; Kurzban, E.; Lee, J.; Matteson,
N. L.; Parker, E.; Perkins, S. A.; Ramesh, K. S.; Robles-Sikisaka, R.; Schwab, M. A.; Spencer, E.;
Wohl, S.; Nicholson, L.; Mchardy, I. H.; Dimmock, D. P.; Hobbs, C. A.; Bakhtar, O.; Harding,
A.; Mendoza, A.; Bolze, A.; Becker, D.; Cirulli, E. T.; Isaksson, M.; Schiabor Barrett, K. M.;
Washington, N. L.; Malone, J. D.; Schafer, A. M.; Gurfield, N.; Stous, S.; Fielding-Miller, R.;
Garfein, R. S.; Gaines, T.; Anderson, C.; Martin, N. K.; Schooley, R.; Austin, B.; MacCannell, D.
R.; Kingsmore, S. F.; Lee, W.; Shah, S.; McDonald, E.; Yu, A. T.; Zeller, M.; Fisch, K. M.;
Longhurst, C.; Maysent, P.; Pride, D.; Khosla, P. K.; Laurent, L. C.; Yeo, G. W.; Andersen, K.
G.; Knight, R. Wastewater Sequencing Reveals Early Cryptic SARS-CoV-2 Variant
Transmission. *Nature* **2022**. https://doi.org/10.1038/s41586-022-05049-6.

(59) Smyth, D. S.; Trujillo, M.; Gregory, D. A.; Cheung, K.; Gao, A.; Graham, M.; Guan, Y.;
Guldenpfennig, C.; Hoxie, I.; Kannoly, S.; Kubota, N.; Lyddon, T. D.; Markman, M.; Rushford,
C.; San, K. M.; Sompanya, G.; Spagnolo, F.; Suarez, R.; Teixeiro, E.; Daniels, M.; Johnson, M.
C.; Dennehy, J. J. Tracking Cryptic SARS-CoV-2 Lineages Detected in NYC Wastewater. *Nature
Communications* **2022**, *13* (1), 635. https://doi.org/10.1038/s41467-022-28246-3.

(60) Jahn, K.; Dreifuss, D.; Topolsky, I.; Kull, A.; Ganesanandamoorthy, P.; Fernandez-Cassi, X.;
Bänziger, C.; Devaux, A. J.; Stachler, E.; Caduff, L.; Cariti, F.; Corzón, A. T.; Fuhrmann, L.;
Chen, C.; Jablonski, K. P.; Nadeau, S.; Feldkamp, M.; Beisel, C.; Aquino, C.; Stadler, T.; Ort, C.;
Kohn, T.; Julian, T. R.; Beerenwinkel, N. Early Detection and Surveillance of SARS-CoV-2
Genomic Variants in Wastewater Using COJAC. *Nature Microbiology* **2022**.
https://doi.org/10.1038/s41564-022-01185-x.

(61) Simpson, G. L. *Extrapolating with B Splines and GAMs*; Blog post; 2020.
https://web.archive.org/web/20231003073128/https://fromthebottomoftheheap.net/2020/06/03/ext
rapolating-with-gams/.

(62) CDC. *About Wastewater Data: Data Methods*; Website; Centers for Disease Control and
Prevention, 2023.

https://web.archive.org/web/20231204153532/https://www.cdc.gov/nwss/about-data.html#data-
method.
(63) Li, X.; Liu, H.; Gao, L.; Sherchan, S. P.; Zhou, T.; Khan, S. J.; van Loosdrecht, M. C. M.; Wang,
Q. Wastewater-Based Epidemiology Predicts COVID-19-Induced Weekly New Hospital
Admissions in over 150 USA Counties. *Nature Communications* **2023**, *14* (1), 4548.
https://doi.org/10.1038/s41467-023-40305-x.
801