RSV-Related Healthcare Burden: A Prospective Observational Study in a Resource-Constrained

Setting

Senjuti Saha PhD[1#], Sudipta Saha SM[1,2], Naito Kanon MSc[1], Yogesh Hooda PhD[1], Mohammad Shahidul

Islam PhD[1], Shuborno Islam BSc[1], Zabed Bin Ahmed MSc[1], Md Jahangir Alam FCPS[3], Ataul Mustufa Anik

MSc[1], Probir K Sarkar FCPS[3], Mohammed Rizwanul Ahsan, DCH[4], Md. Ruhul Amin FCPS[1], Samir K Saha

PhD[1,5]

1. Child Health Research Foundation, Dhaka, Bangladesh

2. Department of Social and Behavioural Sciences, Harvard T. H. Chan School of Public Health, Boston,

USA

3. Department of Pediatric Respiratory Medicine (Pulmonology), Bangladesh Shishu Hospital and

Institute, Dhaka, Bangladesh

4. Department of Emergency, Observatory and Referral, Bangladesh Shishu Hospital and Institute,

Dhaka, Bangladesh

5. Department of Microbiology, Bangladesh Shishu Hospital and Institute, Dhaka, Bangladesh

Correspondence to

Senjuti Saha, PhD

Deputy Executive Director

Child Health Research Foundation

23/3 Khilji Road, Mohammadpur

Dhaka 1207, Bangladesh

Email: senjutisaha@chrfbd.org

Abstract

Background: Respiratory syncytial virus (RSV) is a leading cause of pediatric hospitalizations globally, impacting overstretched health systems. Comprehensive data is vital for informing RSV vaccination policies.

Methods: From January to December 2019, a prospective study was conducted at Bangladesh's largest pediatric hospital to evaluate RSV's burden on the health system. We analyzed hospitalization rates, lengths of stay, and outcomes for children under five using WHO criteria and qPCR testing. We also examined survival probability for children denied admission due to bed shortages, compared with those admitted using the Kaplan-Meier method, and estimated the effects of a maternal vaccine using Monte-Carlo simulations.

Findings: Out of 40,664 children admitted, 31,692 were under five. Of these, 19,940 were eligible for study inclusion with 7,191 meeting inclusion criteria; 6,149 (86%) had samples taken, with 1,261 (21%) testing positive for RSV. The hospital incidence rate was 465 per 10,000 admissions. The median age of RSV patients was 3 months, with a median stay of 5 days. RSV accounted for 8,274 bed days of the total 151,110 observed bed days. In-hospital mortality was 1.9%, increasing to 9.9% during a 90-day follow-up over telephone. Additionally, 9,169 children were denied admission during the study period; of these, 5,969 under-five children were approached, and outcomes of 2,850 admitted versus 3,928 refused were followed. The hazard ratio for death was 1.37 for refused versus admitted children, highest within neonates at 1.7. A 70% efficacious vaccine would equate to adding 50 hospital beds and averting 195 deaths.

Interpretation: RSV significantly contributes to pediatric hospitalizations in Bangladesh, exacerbating healthcare burdens and increasing mortality risks. Maternal vaccination could significantly reduce both direct and indirect RSV burdens, enhancing healthcare capacity and benefiting overall child health in resource-limited settings.

Funding: Bill and Melinda Gates Foundation

Introduction

Respiratory syncytial virus (RSV) is the leading cause of acute lower respiratory infection in young children. Annually, RSV infections are estimated to result in 33 million episodes, 3.6 million hospitalizations, and 26,300 deaths [1]. Low- and middle-income countries (LMICs) bear a disproportionate burden of the disease, where over 95% of RSV infections and more than 97% of RSV-attributable deaths occur [1].

Emerging interventions, such as maternal vaccines and long-acting monoclonal antibodies, promise to reduce RSV-related morbidity and mortality [2–4]. There is also growing evidence suggesting that these interventions might have broader benefits, like reducing respiratory illnesses caused by other pathogens, decreasing recurrent hospitalizations, and improving lung health [5–7]. Policymakers across countries are grappling with decisions about introduction, and which groups should be prioritized for these interventions. Making informed decisions necessitates up-to-date and locally relevant epidemiological data. However, there exist large data gaps from LMICs.

In Bangladesh, a densely populated lower-middle income country, with a birth cohort of almost 3 million[8] the healthcare system faces unique challenges due to the high burden of communicable and non-communicable diseases[9]. Data from the country's largest pediatric hospital indicate that approximately 20% of children needing hospitalization—whether for communicable or non-communicable diseases—have to be turned away due to bed shortages [10]. Timely and appropriate care is vital for child survival and well-being. In settings where healthcare resources are stretched thin, a high RSV burden can exacerbate capacity issues, potentially leading to adverse health outcomes for children who cannot be admitted for various reasons. Thus, it is imperative to consider both the direct and indirect effects of RSV infections on child health and the broader health system when making informed policy decisions.

Two prominent multi-country studies, Aetiology of Neonatal Infections in South Asia (ANISA) [11], and Pneumonia Etiology Research for Child Health (PERCH) [12], have highlighted RSV as a primary etiological

agent in sepsis and respiratory tract infections in Bangladesh and other countries in Asia and Africa.

While these studies underscore RSV's role in community-based child morbidity, gaps remain in

understanding the extent of RSV-associated hospitalizations and the broader impact on the healthcare

capacity in resource-constrained settings.

To address the current gaps in data and estimate the potential impact of RSV vaccines, we conducted a

prospective, observational study at Bangladesh's largest pediatric hospital. We aimed to quantify RSV-

associated hospitalizations, bed usage, and outcomes. We also assessed the wider health implications

for children denied admission due to bed shortages, providing crucial data to inform policy and

intervention strategies.

Methods

Ethical considerations

This study was approved by the ethical review board of the Bangladesh Shishu Hospital and Institute

(BSHI). We obtained written informed consent from guardians of children admitted in the hospital

from whom nasopharyngeal samples were collected. We obtained informed verbal consent from

caregivers when they were registered in the study after hospital admission refusal for telephone follow

up. We also obtained informed verbal consent from caregivers of admitted patients selected for

follow-up over telephone.

Study Design and Setting

From January to December 2019, we conducted a prospective observational surveillance at the BSHI,

previously known as Dhaka Shishu Hospital (DSH), located in Dhaka, Bangladesh. BSHI, with its 653-bed

capacity (2019), is Bangladesh's largest pediatric hospital. It offers primary to tertiary care for patients

up to 18 years old. Admission decisions for the in-patient department (IPD) are made by physicians in

the Emergency Room (ER).

This study had two main components: (i) active surveillance within the IPD to detect RSV infections;

and (ii) active surveillance of health outcomes of patients who had to be denied hospitalization due to

bed shortages in the IPD, despite needing hospitalization, and a control group of patients who were

admitted in the IPD.

RSV Surveillance in the IPD

Children aged 0–59 months admitted to the wards selected for the study with 414 beds were

evaluated by study physicians using the WHO RSV hospital-based surveillance case definition [13] (Text

S1, appendix page 3). The remaining 239 beds of the total 653 were not selected because they were in

wards that do not admit patients with possible infectious diseases (surgery, nephrology, oncology and

cardiology wards). Eligible children were enrolled in the study after obtaining written consent from

caregivers and obtaining a nasopharyngeal swab sample. Nasopharyngeal swabs were collected by

trained nurses and tested using established qPCR methods (Text S2, appendix page 3).

Refusal Surveillance in the ER

Upon arrival at BSHI, families with children potentially needing hospitalization are triaged to the

Emergency Room (ER). Here, a physician evaluates the child's condition to determine if immediate

hospitalization is necessary. Beds are available on a first come first serve basis. If the physician decides

that the child requires hospitalization, but no beds are available, families are referred or advised to

seek care elsewhere. During this study, when a child was refused admission due to bed shortages

despite requiring hospitalization, trained research assistants documented the ER physician's diagnosis

for the patients, and upon obtaining verbal consent from the caregiver, they collected basic details,

including the child's age, family address, and contact numbers.

Health Outcome Follow-Up

Among the <5 y children refused admission due to shortage of beds and whose contact information

was obtained, families were selected using a computer-based randomization algorithm, and contacted

via telephone approximately 2 weeks after hospital-refusal, and the child's health status ("alive" or

"deceased"), and date of death (if applicable) was recorded based on caregivers' response. If the child

was alive during the 2-week follow-up, a second follow-up call was made three months post refusal. In

parallel, from the cohort of children <5 years who were admitted during the same time, children were

randomly chosen for follow-up in the same manner. Additional information on in-hospital mortality

was obtained from hospital records.

Survival curves were computed using the Kaplan-Meier method. Date of death was collected during

follow-up, or from hospital records for admitted cases. Conservative censoring assumptions were

made – children refused admission and not followed-up at 14-days were considered censored at day 0,

and those not followed up at 90-days were considered censored at 14-days (last successful contact).

Admitted children's censoring time was dealt with similarly, except that hospital discharge (either alive

or through mortality) was additional information. Multivariate Cox Proportional-Hazards models were

used to estimate hazard ratios adjusted for diagnosis cluster (Table S1, appendix page 5), age in

months, and sex. We also obtained stratified estimates of hazard ratios.

Estimated impact of RSV maternal vaccine impact analysis (Monte Carlo Simulation)

A Monte Carlo simulation was conducted to estimate the direct and indirect effects of RSV vaccination

on hospital admissions and 90-day mortality. The simulation used empirical data on the total number

of children requiring admission in 2019 across all age groups and diagnoses, length of stay, and

estimated 90-day survival probabilities among refused and admitted cases based on follow-up. We

explored the impact on total deaths within 3 months of hospital admission or refusal when a fixed

number of beds were occupied under a baseline scenario reflecting the empirical data. Additionally,

scenarios with various reductions in the number of RSV cases were simulated. The median value across

1000 simulations and the 95% prediction intervals are reported. Further methodological details are

provided in Text S3, appendix page 4.

Role of funding source

The funder had no role in study design, data collection, data analysis, data interpretation, writing of

the report, or the decision to submit this manuscript. The corresponding authors had full access to all

the data in the study and had final responsibility for the decision to submit for publication.

Results

Cases Hospitalized with RSV:

In 2019, a total of 49,833 children sought care at BSHI and were advised admission, of whom 40,664

could be admitted based on the available beds (Figure 1, Table 1). 31,692 of them were <5, and of

them 19,940 were admitted in the screening wards. Of them, 7,191 met the inclusion criteria of the

study, and a sample was successfully collected from 6,149 (86%) of these children. Samples could not

be collected from the remaining 1,042 eligible cases due to various reasons including the guardian's

refusal to participate in the study, use of oxygen mask, and discharge before enrollment or sample

collection (Table S2, appendix page 5)). 92% (5658/6149) of all samples were collected within 72 hours

of hospitalization (Table S3, appendix page 6).

All 6,149 samples were tested for the presence of RSV using qPCR, and of them 1,261 (21%) samples

tested positive. Assuming 1,042 untested eligible cases among 7191 had the same positivity as those

tested, the overall proportion of all hospital admissions that were RSV positive was 4.65% or 465 RSV

cases per 10,000 <5 hospitalizations (95% CI: 443 to 489).

Figure 1. Flowchart of sample collection and patient follow-up. The figure illustrates the process of sample collection and subsequent patient follow-up for children under five admitted to BSHI or refused admission due to unavailability of bed. Reasons for failures of sample collection and details on those lost to follow-up are provided in Supplementary Appendix Tables S2, S4 - S7.

197   The highest number of patients were enrolled in the study between June and September (Figure 2A),

198   and these four months accounted for over 70% of the total RSV-positive cases (882/1261) (Figure 2B).

199   During July, August and September, more than 20% of the 414 beds were due to RSV positive cases

200   (Figure 2C).

202   **Table 1: Clinical attributes study participants enrolled in the study stratified by RSV status.**

	Overall	RSV Negative/ Not Tested	RSV Positive	p
Number of Patients	**19940**	**18679**	**1261**	
Sex - Female (%)	7582 (38.0)	7196 (38.5)	386 (30.6)	<0.001
Age in months (median [IQR])	5.00 [0.00, 16.00]	5.00 [0.00, 17.00]	3.00 [1.00, 8.00]	<0.001
< 2 wks	5364 (26.9)	5293 (28.3)	71 (5.6)	
2 wks - 3 mnths	4039 (20.3)	3413 (18.3)	626 (49.6)	
4 - 6 mnths	1648 (8.3)	1464 (7.8)	184 (14.6)	
7 mnths - 2 yrs	5693 (28.6)	5381 (28.8)	312 (24.7)	
2 - 5yrs	3196 (16.0)	3128 (16.7)	68 (5.4)	
In-hospital mortality (%)	1266 (6.3)	1242 (6.6)	24 (1.9)	<0.001
Length of Stay (median [IQR])	4.00 [2.00, 7.00]	4.00 [2.00, 7.00]	5.00 [4.00, 8.00]	<0.001
Diagnosis Cluster (%)				<0.001
Respiratory Manifestation	5026 (25.2)	4065 (21.8)	961 (76.2)	
Gastrointestinal Manifestation	2192 (11.0)	2178 (11.7)	14 (1.1)	
Perinatal Asphyxia	2178 (10.9)	2164 (11.6)	14 (1.1)	
Febrile Illness	1464 (7.3)	1445 (7.7)	19 (1.5)	
Systemic Infections	1421 (7.1)	1352 (7.2)	69 (5.5)	
Preterm low-birth weight	899 (4.5)	887 (4.7)	12 (1.0)	
Other	6758 (33.9)	6586 (35.3)	172 (13.6)	

204
Of the positive cases, 30.6% were females. The median age of PCR-confirmed RSV cases was 3 months

(IQR 1.3, 8.3 days) and 70% (881 of 1261) of the cases were within the first six months (Fig 3A, Table

1)). Notably, 17% of infections occurred in the first 4 weeks of life, highlighting the vulnerability of very

young babies to RSV. Median hospital stay of RSV cases was 5 days (IQR 4 – 8), and medial stay of all

RSV-negative cases and non-tested cases were 4 days (IQR 2 – 7). Throughout the year, RSV cases

accounted for 8,274 (5.5%) of the total 151,110 bed days observed.

Figure 2. RSV Infections and Bed Occupancy in BSHI in 2019. A. The total number of patients enrolled into the study each month, and the number of samples collected from these patients. B. Percentage of RSV positive cases detected each month. C. Percentage of occupied beds with RSV positive cases across different months. The dip in bed occupancy seen in mid-August is because of Eid-ul-Adha holidays.

In total, 24 cases (1.9%) of all positive cases died during their stay at the hospital (Table 1). Median age

of these cases was 130 days (IQR 77 - 175) and died between 0 - 26 days of hospitalization.

Figure 3. Age distribution of RSV cases and length of hospital stay. A. Age distribution of RSV cases (n
= 1,261) in weeks, where the green line represents RSV test positivity in different age groups, the solid
orange line indicates the median age, and the dotted orange line shows the mean age. B. Length of
hospital stay in days of RSV cases (n = 1,261), with the solid orange line representing the median length
of stay and the dotted orange line the mean.

Impact of Bed Shortages on Non-Admitted Patients

During the study period, of the total 49,833 children requiring hospitalization, the hospital was unable

to admit 9,169 (18.4%) patients requiring admission due to lack of beds (Figure 1). Of them, we were

able to approach and collect contact information of 5,969 families for follow-up; 3,200 patients were

not approached because they either came to the hospital at night when the study research assistants

were absent (n = 2,324) or were >5 years of age (n = 876). Of the families who provided verbal consent

and contact information for telephone follow-up, a random selection of 3,928 (43%) patients were

chosen for follow-up. Out of these, 3,435 (87%) could be successfully contacted and followed up at 2

weeks. The primary reasons for missing follow-up included switched-off mobile phones, incorrect

phone numbers, or no response to the call (Table S4, appendix page 6). During the first follow up, 2931

(85.3%) children were reported to be alive by their guardian, and 504 (14.7%) were reported to have

died. Of the 2,931 patients alive, 2631 (89.8%) could be subsequently contacted during the 3-month

follow up, and 2525 (96%) were reported to be alive, and an additional 106 (4%) had died. The survival

probability at the end of follow up was 0.805 [95% CI: 0.791, 0.819] for refused patients (Figure 4a).

Figure 4. Survival Analysis of Patients followed-up. A. Survival probabilities of patients who were
refused admission with those who were admitted. Cases were censored at discharge if there was no
follow-up or at the last successful follow-up. B. Survival curve for only RSV-positive cases: the 24 cases
that resulted in in-hospital deaths, alongside the additional 205 cases that were followed up, of whom
18 subsequently died.

2,850 admitted <5-year children were randomly selected for follow-up during the same time. Of them,

2,527 (89%) were successfully followed up at 2 weeks. Reasons for unsuccessful follow up included

those above, in addition to missing contact information in the hospital records (Supplementary table

6). During the first follow up, 2302 (91.1%) children were reported to be healthy, and 225 (8.9%) were

reported to have died (Table 2). Of the 2,302 patients alive 1,965 (85.3%) could be subsequently

contacted during the 3-month follow up, of which 1,883 (95.8%) were alive, and an additional 82

(4.2%) had died (Figure 4). The survival probability at the end of follow up was 0.874 [95% CI: 0.861,

0.887] among those admitted (Figure 4a).

Table 2: Clinical attribute and outcome of participants admitted or refused admissions who were followed-up. The table represents data of participants for whom data could be collected. Diagnoses was could not be collected for 18% of participants, 14-day mortality was missing for 12% and 90-day mortality was missing for 21%.

	Overall	Admitted	Refused	p-value
Number of Patients	**6772**	**2845**	**3927**	
Sex - Female (%)	2582 (38.1)	1071 (37.6)	1511 (38.5)	0.502
Age in months (median [IQR])	3.00 [0.00, 12.00]	4.00 [0.00, 15.00]	3.00 [0.00, 11.00]	0.006
< 2 wks	2017 (29.8)	804 (28.3)	1213 (30.9)	<0.001
2 wks - 3 mnths	1430 (21.1)	600 (21.1)	830 (21.1)	
4 - 6 mnths	622 (9.2)	226 (7.9)	396 (10.1)	
7 mnths - 2 yrs	1869 (27.6)	764 (26.9)	1105 (28.1)	
2 - 5yrs	834 (12.3)	451 (15.9)	383 (9.8)	
ER Diagnosis Cluster (%)				
Respiratory Manifestation	1463 (26.5)	351 (20.1)	1112 (29.4)	
Gastrointestinal Manifestation	475 (8.6)	182 (10.4)	293 (7.8)	
Perinatal Asphyxia	702 (12.7)	211 (12.1)	491 (13.0)	
Febrile Illness	207 (3.7)	101 (5.8)	106 (2.8)	<0.001
Systemic Infections	496 (9.0)	145 (8.3)	351 (9.3)	
Preterm low-birth weight	382 (6.9)	78 (4.5)	304 (8.0)	
Other	1799 (32.6)	677 (38.8)	1122 (29.7)	
14-day Mortality (%)	729 (12.2)	225 (8.9)	504 (14.7)	<0.001
90-day Mortality (%)	917 (17.2)	307 (14.0)	610 (19.4)	<0.001

Overall comparison between the two groups revealed a hazard ratio (HR) of 1.69 (CI 1.48 - 1.94) of death in children who were refused admission compared to children who were admitted (Table 3). The HR was 1.37 (CI 1.56 - 1.79) in models adjusted for age, diagnosis and sex. Models stratified by age groups showed that HR of death was significantly greater for refused cases among babies in their first month of life (neonates) (HR 1.93 [CI 1.62 - 2.30]) but not in other age groups. When stratified by

275 diagnosis, HR of death was significantly greater among those with perinatal asphyxia (HR 2.22 [1.68,

276 2.92]), systemic infections (HR 1.79 [1.13, 2.85]), and preterm low-birth weight 1.85 [1.32, 2.59].

278 **Table 3: Hazard ratio (HR) of refused vs admitted participants adjusted by age, sex, and diagnosis.**
279

Adjusted by	Strata	HR [95% CI]
		1.69 [1.48, 1.69]
Age, Sex, Diagnosis		**1.56 [1.36, 1.79]**
Sex, Diagnosis	< 1 mnth	**1.93 [1.62, 2.3]**
Sex, Diagnosis	2-3 mnths	1.19 [0.89, 1.58]
Sex, Diagnosis	4 mnths - 1 yr	1.47 [0.87, 2.48]
Sex, Diagnosis	1-5 yrs	1.05 [0.66, 1.68]
Age, Diagnosis	Male	**1.59 [1.34, 1.9]**
Age, Diagnosis	Female	**1.49 [1.2, 1.84]**
Age, Sex	Respiratory Manifestation	0.96 [0.7, 1.31]
Age, Sex	Perinatal Asphyxia	**2.22 [1.68, 2.92]**
Age, Sex	Systemic Infections	**1.79 [1.13, 2.85]**
Age, Sex	Preterm low-birth weight	**1.85 [1.32, 2.59]**
Age, Sex	Febrile Illness	0.79 [0.14, 4.37]
Age, Sex	Gastrointestinal Manifestation	1.37 [0.51, 3.71]
Age, Sex	Other	**1.32 [1.02, 1.71]**

Post-hospitalization RSV-associated mortality

While the study was not designed to specifically follow-up RSV positive cases, of the 2,527 admitted

children randomly selected and successfully followed up, there were 205 RSV positive cases. Of these,

18 children died within the 90-day follow-up period. Their median age was 107.5 days (IQR 44.75 - 157)

Considering both in-hospital mortality of all RSV admissions and mortality during follow-up of a subset

of cases revealed that survival decreased substantially over time, with a 90-day survival probability of

0.901 [95% CI: 0.865, 0.938] at the end of the follow-up period (Figure 4b).

Potential Impact of Maternal RSV Vaccination

In a Monte Carlo simulation of hospital admissions and refusals for the baseline scenario, based on the

empirical data and a 650-bed capacity, on average, 49,968 children required admissions (95%

prediction intervals (PI): 49,500, 50,379), and 9,221 (95% PI: 8,869, 9,760) cases were refused

admission over the course of the year. This was in concordance with the true burden of total cases

requiring admission and refusals – 49,833 and 9,169 respectively. In the vaccine scenario with 70%

reduction in RSV cases requiring admission, 48,895 (95% PI: 48,514, 49,315) cases required admission,

and 8,103 cases were refused (7,728, 8.695) – a 16.6% refusal rate. Even though total cases requiring

admissions in this scenario remained as high as 97.9% (48895/49968) of the baseline scenario, the

number of refused cases was 87.8% (8103/9221), due to the freeing up of bed spaces. Figure 5

demonstrates the effects of a 70% reduction in RSV admissions through vaccination, showing its impact

on deaths and hospital bed utilization. The plot compares mortality with increase in available beds,

demonstrating how reducing RSV cases is akin to increasing bed capacity. Overall, there were 7,002

deaths (6829, 7167) in the baseline scenario, and 6807 deaths (6629, 6930) in the vaccine scenario, a

difference of 195 deaths. A reduction in RSV cases was found to decrease both hospital RSV incidence

and the overall number of RSV cases seeking admission at the hospital. The estimated reduction in

deaths with the 70% efficacious vaccine would be equivalent of adding roughly 50 beds to the hospital

although, there is considerable overlap of the variability in the estimates from the different scenarios.

313
314 **Figure 5. Impact of RSV Vaccination on Hospital Utilization and 90-day Mortality Rates of all children**
315 **who require admission at BSHI.** Effects of a 70% reduction in RSV admissions through vaccination,
316 showing its impact on deaths and hospital bed utilization among children under five. The plot
317 compares mortality against the effective increase in available beds, demonstrating how reducing RSV
318 cases is akin to increasing bed capacity.
319

Discussion

Informed decision-making on interventions to prevent RSV infections in young infants requires

contemporary and locally relevant epidemiological data. Our study demonstrates the significant impact

of RSV on healthcare resources in Bangladesh, a country characterized by a high burden of infectious

diseases and a substantial demand for pediatric care amidst resource limitations.

Our study recorded an overall RSV positivity rate of 21% among the 6,149 patients admitted to the

largest pediatric hospital of Bangladesh with respiratory illnesses, corresponding to an RSV hospital

incidence of 465 per 10,000 <5 admissions. The median age of confirmed RSV cases was 3 months.

These findings are concordant with those of the PERCH study that reported that in Bangladesh 31.2%

of children hospitalized with respiratory tract infections tested positive for RSV [12].

The in-hospital mortality rate for RSV-positive cases was 1.6%, which escalated to 9.9% during the 90-

day follow-up of discharged patients. The median age of recorded RSV-associated deaths was 4.2

months. These data are comparable to a Zambian study that found RSV in 9% of community deaths

and 5% of in-hospital deaths [14]. In Child Health and Mortality Prevention Surveillance (CHAMPS) study,

RSV was found in 5.5% of all deaths, with variability across countries: 9.7% in Mali, 10.7% in Ethiopia,

and a lower 2% in Bangladesh [15]. Half of the deaths in the CHAMPS study occurred in infants younger

than six months.

These findings underscore the critical need for maternal immunization strategies, such as vaccines like

ABRYSVO™, which has demonstrated 81.8% efficacy in preventing RSV infections within the first three

months of life [16]. Our estimates suggest that maternal immunization could potentially reduce the

hospital incidence from 465 to 140 cases per 10,000 <5 admissions.

Beyond the direct impact of RSV infections, our findings also demonstrate the considerable burden

infections impose on the healthcare system. For a minimum of three months during the study year,

RSV cases occupied more than 20% of the beds under observation, with a median hospital stay for RSV-

positive cases at 5 days—longer than the overall median length of stay of 4 days for all admitted

patients. In 2019, RSV cases accounted for 8,274 of the total 151,110 bed days observed. As the largest

pediatric tertiary hospital in Bangladesh, BSHI, even with its total of 653 beds, is insufficient.

Consequently, clinicians often must refuse admission due to bed unavailability [10]. We followed over

5,000 admitted and refused children up to 90 days to determine the difference in health outcomes and

the wider effects of high-burden infections like RSV on the healthcare system. The cumulative

mortality proportion for children denied admission was 19.4%, significantly higher than the 14% among

those admitted. This discrepancy in mortality risk is quantified by a hazard ratio (HR) of 1.69 across all

children under five.

The HR for death was 1.37 among children refused admission compared to those admitted, adjusted

for age, diagnosis, and sex. The highest mortality risk was observed in neonates requiring

hospitalization within their first month of life, with a HR of 1.93. This disparity highlights the essential

role of healthcare access in determining child survival outcomes. Monte Carlo simulation using a 70%

efficacious vaccine is estimated to be equivalent to adding approximately 50 beds to the hospital, and

averting 195 deaths in only one hospital, each year.

The issue of bed shortages, a prevalent concern across many LMICs, is pressing. In Bangladesh, the rate

of 0.8 beds per 1,000 population is markedly lower than in developed nations such as 2.9 in the USA

and 2.5 in the UK [17]. With limited infrastructure, preventative strategies become even more vital. Over

the past decades, the introduction of multiple vaccines in Bangladesh and improved health systems

have led to a decrease in under-5, yet the rate 33 death in 1,000 live birth remain high, underscoring

the need for further disease prevention measures [18]. Given the considerable proportion of hospital

beds occupied by RSV cases during peak periods, introducing an RSV vaccine could significantly impact

bed availability for critically ill infants. Other interventions like administration of monoclonal antibodies

in neonates and infants can also be considered. However, the cost and logistical hurdles associated

with monoclonal antibody treatments, such as nirsemivab, highlight the need for affordable, scalable,

and equitable solutions, particularly in LMICs like Bangladesh where 69% of health expenditures are

out-of-pocket [19].

While our 2019 study centered on RSV, a significant dengue outbreak during the same period

compounded the strain on hospital resources [20]. In addition, other infectious etiologies of respiratory

infections remain to be explored, as evidenced by the high number of cases meeting the inclusion

criteria for RSV October to December, despite the absence of RSV detection.

The findings of this study must be viewed considering certain limitations. Primary among them is the

necessity to conduct the study in a single hospital due to the high volume of cases and the extensive

follow-up required for thousands of patients. While the authors' lived experiences suggest that similar

circumstances are likely in other resource-limited settings, a multi-site research would be helpful in

developing a more detailed understanding of the RSV burden among children throughout Bangladesh

and potentially the wider region. Furthermore, the study's duration was constrained to one year,

shortened by the COVID-19 pandemic in the following year. A longitudinal study spanning multiple

years could provide more information on the patterns of RSV seasonality, but previous reports from

Bangladesh suggest an unpredictable pattern of RSV seasonality [21]. And the overall high burden of RSV

in the study hospital aligns with previous studies [11,12]. This study also did not investigate concurrent

infections alongside RSV, nor was it set up to confirm deaths directly attributable to RSV. Lastly,

potential biases due to human factors may have influenced the clinicians' admissions decisions during

times of resource scarcity, even though such decisions should theoretically be unaffected by bed

availability constraints.

The broader context and global health implications of our study are significant. The challenges

highlighted related to shortage of hospital beds in places with high burden of disease are common in

many LMICs and the global health community can learn from these findings particularly regarding

resource allocation and estimate potential impact of preventive strategies like vaccination. This study

also contributes to the growing body of evidence supporting the development and distribution of RSV

vaccines or other interventions, which could significantly impact child health globally, specifically in

low-resource countries where burden is the highest.

Declaration of Interests

The authors declare no conflict of interest.

Author contributions

S.S. contributed to conceptualization, funding acquisition, study design, implementation, monitoring,

data analysis, drafting the initial manuscript, and preparation of figures. S.Sa. was involved in data

analysis, preparation of figures, and manuscript editing. N.K. contributed to data analysis and editing.

Y.H. participated in data analysis, preparation of figures, and manuscript editing. M.S.I. was responsible

for analysis and coordination of the study, and also contributed to editing. S.I. carried out specimen

extraction and PCR testing. Z.B.A. coordinated and trained doctors and nurses, assisted in the

implementation of the study, in addition to contributing to manuscript editing. M.J.A. was involved in

monitoring and management activities, as well as editing. A.M.A. performed data cleaning. P.K.S.

contributed to management and patient data analysis. M.R.A. was involved in implementation and

management. M.R.A. contributed to editing. S.K.S. contributed to conceptualization, funding

acquisition, study design, implementation, monitoring, and manuscript editing. All authors reviewed

the manuscript and agreed on the contents.

Declaration of Generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used ChatGPT to check grammar, spellings, and

improve readability of the manuscript. After using this tool/service, the authors reviewed and edited

the content as needed and take full responsibility for the content of the publication.

References

1 Li Y, Wang X, Blau DM, *et al.* Global, regional, and national disease burden estimates of acute lower

respiratory infections due to respiratory syncytial virus in children younger than 5 years in 2019: a

systematic analysis. *Lancet* 2022; **399**: 2047–64.

2 Baral R, Higgins D, Regan K, Pecenka C. Impact and cost-effectiveness of potential interventions

against infant respiratory syncytial virus (RSV) in 131 low-income and middle-income countries using

a static cohort model. *BMJ Open* 2021; **11**: e046563.

3 Muller William J., Madhi Shabir A., Seoane Nuñez Beatriz, *et al.* Nirsevimab for Prevention of RSV in

Term and Late-Preterm Infants. *New England Journal of Medicine* 2023; **388**: 1533–4.

4 Commissioner O of the. FDA Approves First Vaccine for Pregnant Individuals to Prevent RSV in

Infants. FDA. 2023; published online Aug 22. https://www.fda.gov/news-events/press-

announcements/fda-approves-first-vaccine-pregnant-individuals-prevent-rsv-infants (accessed April

13, 2024).

5   Zar HJ, Nduru P, Stadler JAM, *et al.* Early-life respiratory syncytial virus lower respiratory tract
infection in a South African birth cohort: epidemiology and effect on lung health. *The Lancet Global*
Health 2020; **8**: e1316–25.

6   Abreo A, Wu P, Donovan BM, *et al.* Infant Respiratory Syncytial Virus Bronchiolitis and Subsequent
Risk of Pneumonia, Otitis Media, and Antibiotic Utilization. *Clin Infect Dis* 2020; **71**: 211–4.

7   Feikin DR, Karron RA, Saha SK, *et al.* The full value of immunisation against respiratory syncytial virus
for infants younger than 1 year: effects beyond prevention of acute respiratory illness. *The Lancet*
Infectious Diseases 2023; **0**. DOI:10.1016/S1473-3099(23)00568-6.

8   Bangladesh | Gavi, the Vaccine Alliance. 2018; published online Nov 29.
https://www.gavi.org/programmes-impact/country-hub/south-east-asia/bangladesh (accessed April
13, 2024).

9   Islam SMS, Uddin R, Das S, *et al.* The burden of diseases and risk factors in Bangladesh, 1990–2019: a
systematic analysis for the Global Burden of Disease Study 2019. *The Lancet Global Health* 2023; **11**:
e1931–42.

10 Saha S, Santosham M, Hussain M, Black RE, Saha SK. Rotavirus Vaccine will Improve Child Survival by
More than Just Preventing Diarrhea: Evidence from Bangladesh. *Am J Trop Med Hyg* 2018; **98**: 360–
3.

11 Saha SK, Schrag SJ, El Arifeen S, *et al.* Causes and incidence of community-acquired serious
infections among young children in south Asia (ANISA): an observational cohort study. *The Lancet*
2018; **392**: 145–59.

12 O'Brien KL, Baggett HC, Brooks WA, *et al.* Causes of severe pneumonia requiring hospital admission
in children without HIV infection from Africa and Asia: the PERCH multi-country case-control study.
The Lancet 2019; **394**: 757–79.

13 Case definitions. https://www.who.int/teams/global-influenza-programme/global-respiratory-
syncytial-virus-surveillance/case-definitions (accessed Jan 6, 2024).

14 Gill CJ, Mwananyanda L, MacLeod WB, *et al.* Infant deaths from respiratory syncytial virus in Lusaka,
Zambia from the ZPRIME study: a 3-year, systematic, post-mortem surveillance project. *The Lancet*
Global Health 2022; **10**: e269–77.

15 Blau DM, Baillie VL, Els T, *et al.* Deaths Attributed to Respiratory Syncytial Virus in Young Children in
High-Mortality Rate Settings: Report from Child Health and Mortality Prevention Surveillance
(CHAMPS). *Clin Infect Dis* 2021; **73**: S218–28.

16 Fleming-Dutra KE, Jones JM, Roper LE, *et al.* Use of the Pfizer Respiratory Syncytial Virus Vaccine
During Pregnancy for the Prevention of Respiratory Syncytial Virus-Associated Lower Respiratory
Tract Disease in Infants: Recommendations of the Advisory Committee on Immunization Practices -
United States, 2023. *MMWR Morb Mortal Wkly Rep* 2023; **72**: 1115–22.

17 World Bank Open Data. World Bank Open Data. https://data.worldbank.org (accessed April 21,
2024).

18 SDG Tracker. https://sdg.gov.bd/page/indicator-wise/1/30/3/0#1 (accessed April 21, 2024).

19 2023-05-31-17-02-bed461250ff1b439a0338ab3ddfea682.pdf.
https://heu.portal.gov.bd/sites/default/files/files/heu.portal.gov.bd/page/364480c2_26f9_4bbb_bf
71_8b4ac613d96a/2023-05-31-17-02-bed461250ff1b439a0338ab3ddfea682.pdf (accessed April 14,
2024).

20 Yesmin S, Sarmin S, Ahammad AM, Rafi MdA, Hasan MJ. Epidemiological Investigation of the 2019
Dengue Outbreak in Dhaka, Bangladesh. *J Trop Med* 2023; **2023**: 8898453.

21 Hossain ME, Rahman MZ, Islam MdM, *et al.* Pre COVID-19 molecular epidemiology of respiratory
syncytial virus (RSV) among children in Bangladesh. *Heliyon* 2022; **8**: e11043.

489