The Effectiveness of Using a Clinical Support Tool in Managing Adolescents with Non-Traumatic Knee Pain (MAP-Knee): protocol for a cluster-randomised superiority trial with delayed intervention and embedded realist evaluation

Authors: Henrik Riel[1,2,3], Simon Kristoffer Johansen[1,3], Erika Maria Andersen[1], Malene Kjær Bruun[3], Niels Henrik Bruun[4], Chris Djurtoft [3], Simon Doessing [5,6,7], Tina Heyckendorff-Diebold[8], Per Hölmich[9], Martin Bach Jensen [3], Søren Kaalund[10], Niels-Christian Kaldau[9], Tommy Frisgaard Oehlenschlaeger[5,6], Charlotte Overgaard[11], Ole Rahbek[12], Dorthe Brøndum Rasmussen[13,14], Susanne Olesen Schaarup[14], Thomas Sørensen[15], Louise Lund Holm Thomsen[11], Michael Skovdal Rathleff[1,3,17]

Affiliations:
1. Department of Health Science and Technology, Aalborg University, Aalborg, Denmark.
2. Department of Physiotherapy, University College of Northern Denmark, Aalborg, Denmark.
3. Center for General Practice at Aalborg University, Department of Clinical Medicine, Aalborg University, Aalborg, Denmark.
4. Research Data and Biostatistics, Aalborg University Hospital, Aalborg, Denmark
5. Institute of Sports Medicine Copenhagen, Department of Orthopaedic Surgery, University Hospital Bispebjerg-Frederiksberg, Copenhagen, Denmark.
6. Center for Healthy Aging, Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark.
7. Section for Sports Traumatology, Department of Orthopedic Surgery, University Hospital Bispebjerg-Frederiksberg Hospital, Copenhagen, Denmark.
8. Department of Orthopaedic Surgery, Lillebælt Hospital, Vejle, Denmark.
9. Sports Orthopedic Research Center Copenhagen (SORC-C), Department of Orthopedic Surgery, Copenhagen University Hospital Amager-Hvidovre, Copenhagen, Denmark.
10. Sports Medicine Center, North Denmark Regional Hospital, Frederikshavn, Denmark.

11. Unit of Health Promotion, Department of Public Health, University of Southern Denmark,

Esbjerg, Denmark.

12. Department of Orthopedics, Aalborg University Hospital, Aalborg, Denmark.

13. Department of Physiotherapy and Occupational Therapy, Næstved-Slagelse-Ringsted

Hospital, Naestved, Denmark.

14. Department of Elective Orthopaedics, Naestved Hospital of Zealand, Naestved, Denmark.

15. Department of Physio- and Occupational Therapy, Lillebaelt Hospital - University Hospital

of Southern Denmark, Vejle, Denmark.

16. Department of Physical Therapy and Occupational Therapy, Aalborg University Hospital,

Aalborg, Denmark.

Corresponding author

Henrik Riel, PhD, Department of Health Science and Technology, Faculty of Medicine, Aalborg

University, Selma Lagerløfs Vej 249, 9260 Gistrup. E-mail: hriel@dcm.aau.dk. Telephone:

+4530201570

Abstract

Background

Knee pain affects one in three adolescents, which makes it one of the most common pain sites. Guideline recommendations about the clinical selection of patients likely to benefit from interventions are unclear, which leads to treatment heterogeneity and the potential of wasted resources among adolescents with a good prognosis. In contrast, adolescents with a poorer prognosis may not receive sufficient care. A newly developed clinical decision-support tool (The MAP-Knee Tool) intends to support clinicians in engaging with patients and adjusting the clinicians' evidence-based practices to accommodate patient preferences and treatment needs via a shared decision-making process. The aims of this trial are 1) to investigate the effectiveness of using a clinical decision-support tool (The MAP-Knee Tool) compared with usual care in adolescents with non-traumatic knee pain in reducing pain measured by KOOS-Child Pain after 12 weeks and 2) to investigate how the intervention worked, for whom, why and under which circumstances applying realist evaluation methodology.

Methods

This trial is a cluster-randomised superiority trial with a delayed intervention and a realist evaluation. Six hospital departments start with a usual care period of 4 months before randomly crossing over to using the intervention (The MAP-Knee Tool) after 4, 6, or 8 months, respectively. We will recruit 290 adolescents suffering from non-traumatic knee pain diagnoses who are followed for one year, with the change in KOOS-Child Pain after 12 weeks considered the primary endpoint. Secondary outcomes include 1) Global Rating of Change, 2) EQ-5D-Youth, 3) Anterior Knee Pain Youth, 4) the International Physical Activity Questionnaire short version, and 5) sports participation. The realist evaluation will utilise a prospective, qualitative approach for collecting data needed to develop and test a program theory and identify context-mechanism-outcome configurations essential for understanding how outcomes are achieved within specific contexts.

Discussion

This trial focuses on how the initial clinical encounter can be improved to meet the support and management needs of adolescents with chronic knee pain seeking treatment for knee pain in secondary care and investigates how the intervention worked, for whom, why and under which circumstances.

Trial registration

Clinicaltrials.gov (NCT05791513). Prospectively registered on March 30th, 2023.

Background

From the age of 5–9 years to the age of 10–14, there is an almost fourfold increase in the years lived with disability as a result of musculoskeletal conditions (1). In Danish primary care, there is an eightfold increase in contacts in our healthcare sector because of knee symptoms between 5–9 and 10–19 years of age and approximately half of these adolescents are referred to the secondary care sector (2,3). This makes knee pain one of the most common pain sites experienced by youths and adolescents and can affect their school attendance, participation in hobbies and social activities, and mental health (4–6). The most common non-traumatic knee pain diagnoses are Patellofemoral Pain and Osgood-Schlatter Disease; however, patellar tendinopathy, Sinding-Larsen Johansson, and iliotibial pain syndrome are frequent differential diagnoses.(7–9)

Knee pain affects one in three adolescents, and around 60 % will seek medical care due to knee pain (10). Data from prospective cohorts show that 40-50% will still experience knee pain and symptoms after 2-5 years (11–13). A cohort study from our group documented that 70% of adolescents with Patellofemoral Pain reduced or stopped participating in sports during a two-year period, and they had significantly worse EQ-5D Index and VAS scores than those without knee pain (12). This underlines the need for effective treatments to prevent long-term disability and physical inactivity.

There are several management strategies for adolescents suffering from knee pain. These range from minimal support, including patient education and leaflets regarding self-management, to specialised supervised exercise therapy (14–17). Because of insufficient evidence, guideline recommendations about the clinical selection of patients likely to benefit from referral to additional interventions are unclear, which leads to treatment heterogeneity (3,11). Therefore, resources may be wasted among adolescents with a good prognosis, while adolescents with a poorer prognosis may not receive sufficient care. Patient decision aids can support clinicians in engaging with patients and adjusting their evidence-based practices to accommodate patient preferences and treatment needs via a shared decision-making process. Such aids may be especially relevant in conditions with several treatment options available and thus may improve care for adolescents suffering from non-traumatic knee pain.(18–20)

We recently developed a clinical decision-support tool intending to support shared decision-making between adolescents, medical doctors, and parents. The development included systematic literature

searches, an Argumentative Delphi process, and end-user involvement (21), ultimately leading to

the MAP-Knee Tool (22), which clinicians may use during consultation with adolescents with non-

traumatic knee pain. The tool is intended to support the entire consultation, from diagnosing the

condition and explaining the condition to providing information regarding possible management

options. The feasibility of using the tool in a clinical setting has been established (Bruun et al. *in*

preparation), but whether the MAP-Knee Tool is associated with improved long-term outcomes

among adolescents with non-traumatic knee pain compared to usual care is unknown.

The overall aims of this trial are 1) to investigate the effectiveness of using our novel clinical

decision-support tool compared with usual care in adolescents with non-traumatic knee pain in

reducing pain measured by KOOS-Child Pain after 12 weeks and 2) to investigate how the

intervention worked, for whom, why and under which circumstances using a qualitative realist

evaluation (23). We hypothesise that adolescents with non-traumatic knee pain who have been

consulting a clinician who uses the MAP-Knee Tool will be superior in self-reported recovery and

physical function after 12 weeks compared to adolescents who have been consulting a clinician not

using the MAP-Knee Tool.

Methods

Design and setting

The trial is designed as a cluster-randomised superiority trial with a delayed intervention and a

realist evaluation component (23). All hospitals start with a control period of 4 months before

randomly crossing over to using the intervention after 4, 6, or 8 months. The trial is conducted

across six hospital departments in Denmark (Aalborg University Hospital, Amager-Hvidovre

Hospital, Bispebjerg Hospital, North Denmark Regional Hospital Frederikshavn, Næstved Hospital,

and Vejle Hospital). Future reporting of the trial will follow CONSORT guidelines for reporting

non-pharmacologic treatments (24). Reporting of this protocol follows the SPIRIT statement (25).

The trial was planned in accordance with the PREPARE Trial guide (26). Before the inclusion of

the first participant, the trial was registered on clinicaltrials.gov (NCT05791513), and the trial was

approved by the Ethics Committee of the North Denmark Region (N-20220043).

The design of the realist evaluation was anchored in the realist evaluation circle to facilitate theory

generation, which included formulating an initial theory of how the intervention brings about

change, generating hypotheses about essential mechanisms and processes, collection and analysis of

empirical data and program theory testing (23,27). The realist evaluation will involve several data

collections conducted before, during and after the stepped wedge trial to support theory gleaning,

refinement and consolidation (28). Data will be collected via document analysis, realist interviews

with researchers who developed the MAP-Knee Tool, clinicians who will be implementers of the

MAP-Knee Tool and adolescents with non-traumatic knee pain who received treatments with the

MAP-Knee Tool. The data will be analysed prospectively using the linked coding approach by

Jackson and Kolla (29) to identify context-mechanism-outcome (CMO) strings within the coded

texts and Gilmore et al.'s (30) memo-coding technique to extract CMO configurations from our data

and inform the refinement and consolidation of the intervention program theory (23,31). The

program theory will synthesise insights from the document and realist interviews with insights from

the quantitative data collected at the 12-week follow-up (27,32) and evaluated by clinicians using

Mukumbang et al.'s (33) approach for including program theory segments during realist interviews

to obtain feedback and consolidate program theories and context-mechanism-outcome

configurations.

Adolescents with non-traumatic knee pain will be recruited when they arrive at one of the six

hospitals due to a referral from their general practitioner. In contrast, clinicians and adolescents with

non-traumatic knee pain for the realist evaluation will be included prospectively based on

information power before, during and after the initiation of the MAP-Knee trial (34). Thus,

recruitment is based on the natural flow of patients with non-traumatic knee pain seen at the

hospital and the availability of clinicians who have used the MAP-Knee Tool in their treatment of

adolescents with non-traumatic knee pain. Adolescents will only attend the baseline examination at

the hospital, whereas a link to questionnaires is emailed to them for the 12-week and 52-week

follow-ups using REDCap (Vanderbilt University, Nashville, TN, USA). They may have additional

appointments at the hospital during the trial, depending on the treatment plan, but they will not

answer questionnaires during these clinical encounters.

Roles and responsibilities

The project manager is an experienced physiotherapist who coordinates the trial. When potential

participants arrive at the hospital, they are informed about the trial by either a project nurse or

project secretary. This person collects written informed consent from participants and their

parents/legal guardians if the participant is under the age of 18 years and collects data using

questionnaires. After this, the participant is seen by a clinician (orthopaedic surgeon,

physiotherapist, rheumatologist, general practitioner). The type of clinician varies from one hospital

to another due to differences in how each hospital is organised. The realist analysis will be led by a

researcher with a background in Information Science and extensive experience in planning and

conducting qualitative studies with youths with chronic pain and healthcare providers, supported by

experts in conducting realist evaluations in healthcare settings. He is supervised by experts in

conducting realist evaluations (CO and LLH).

Eligibility criteria

The inclusion criteria for the adolescents are: 1) having been referred to a hospital due to non-

traumatic knee pain and 2) being aged between 10 and 19 years. Exclusion criteria: 1) knee pain

with a traumatic origin or 2) lack of ability to cooperate. Any orthopaedic or similar hospital

department in Denmark to which adolescents suffering from non-traumatic knee pain may be

referred was eligible for participation. Any medical doctor or physiotherapist employed at either of

the study sites who regularly sees adolescents with non-traumatic knee pain is eligible to participate

in the trial. The inclusion criteria for the researchers participating in the realist evaluation were: 1)

having participated in the conceptualisation, development or testing of the MAP-Knee Tool, or 2)

any of the relevant sub-studies, or 3) planning of the MAP-Knee Trial. Finally, the inclusion criteria

for clinicians included in the realist evaluation were 1) being a healthcare provider (e.g.,

physiotherapist, orthopaedic surgeon, rheumatologist) who is 2) employed at one of the study sites,

and 3) either had or was likely to use the MAP-Knee Tool when treating adolescents with non-

specific knee pain.

Interventions

Before crossover occurs at the hospital, the adolescents will be diagnosed and treated as usual at the

treating clinician's discretion. Based on our previous research on the usual care pathway, this will

be heterogeneous and include advice to wait and see, imaging (most often MRI and x-ray), or a

rehabilitation plan for treatment within the municipality setting (3,11).

After crossing over to using the MAP-Knee Tool, the treating clinician will use the MAP-Knee Tool

together with the adolescent. The tool was designed to support the entire consultation, from

diagnosing the condition (Patellofemoral Pain, Osgood-Schlatter, Sinding-Larsen-Johansson,

growth pain, patellar tendinopathy, or iliotibial band syndrome) to deciding on future management.

The MAP-Knee Tool includes four separate components: 1) a tool for diagnosing the most common

types of non-traumatic knee pain (SMILE) (35), 2) credible explanations of the aetiology and

pathogenesis specific to the diagnosis based on multiple methods with an iterative design, 3) a

presentation of prognostic factors based on an individual participant data meta-analysis(36), and 4)

an option grid that presents the users of the tool with pros and cons of commonly used management

options, derived from a systematic literature search of systematic and narrative reviews within non-

traumatic adolescent knee pain. An overarching focus of all components was to support shared

decision-making and base decisions on all three pillars of evidence-based medicine: patient values,

clinical expertise, and relevant research.(37) Therefore, the tool should not provide clinicians using

the MAP-Knee Tool with definitive answers simply based on the available evidence. After the

initial prototype of the tool, including all four components, had been made, we performed end-user

testing using think-aloud sessions with adolescents suffering from non-traumatic knee pain,

adolescents with no history of knee pain, and medical doctors.(22)

Before the start of crossover, there will be a 2-week transition period for training the clinicians in

using the support tool in their clinical practice. The clinicians will participate in a 1-hour training

session with the project manager, where they will be educated on using the MAP-Knee Tool. After

this session, they will be asked to use the tool for one week in clinical practice, and a follow-up

session will be held where they can share their experiences with using the tool and ask the project

manager questions. Then, the clinicians can use the MAP-Knee Tool for another week before the

patients seen with the tool are included in the trial. The patients seen during the 2-week transition

period will not be included in the trial. To improve long-term adherence to using the tool, there will

be a booster session similar to the follow-up after the first week in the transition period after four

weeks.

The research activities related to the realist evaluation will run in parallel with the activities of the

MAP-Knee Trial while moving from the initial theory-gleaning phase and formulating preliminary

program theories to phase 2, which entailed the data collection during the crossover of hospital

department 1 and 2 (23). The realist evaluation will utilise a prospective, qualitative approach for

collecting the qualitative data needed to develop and test a program theory and identify context-

mechanism-outcome configurations (23), delineating the micro, meso, and macro-level barriers and

facilitators for archiving positive health outcomes from interventions with the MAP-Knee Tool

(38).

Documentation analysis

All available documents (protocols, logic models, pre-preprints, peer-reviewed studies) will be

collected, categorised, and analysed through a realist thematic analysis approach by Wiltshire and

Ronkainen (39) using NVivo 14 coding software to identify thematic patterns across texts and using

this insight to glean the underlying causal mechanisms which researchers envisioned would

facilitate behavioural change and tool acceptance during the conceptualisation and design of the

MAP-Knee Tool.

Realist interviews

We will conduct realist interviews with relevant stakeholders using the principles for planning and

conducting realist interviews by Manzanos (28) and Mukumbang et al. (31) to maintain theoretical

awareness during initial theory gleaning and subsequent theory testing interviews. Theory-gleaning

interviews with the researchers behind the MAP-Knee Tool will be conducted before the first

departments cross over, using an interview guide with open and probing questions informed by the

document analysis. Additionally, one realist interview will be conducted with a clinician

(physiotherapist, orthopaedic surgeon, rheumatologist) during the time of the crossover to identify

the clinicians' expectations, thoughts, needs, and contextual challenges related to using the MAP-

Knee tool when treating adolescents with non-specific knee pain. The theory-gleaning interviews

will inform the initial program theory (IPT). The initial theory will be tested and expanded during

the intervention with the MAP-Knee tool. The testing will be conducted through (n=3) realist

interviews with clinicians (physiotherapists, orthopaedic surgeons, rheumatologists) per department,

and (n=6-8) adolescents with knee pain, who received treatment with the MAP-Knee Tool across

the timespan of the intervention. By exploring how clinicians and adolescents with non-traumatic

knee pain experienced the MAP-Knee Tool facilitated the treatment of non-traumatic knee pain, we

will test, revise and expand the initial program theory and identify the CMO configurations present

within the use situation.

To ensure the integrity of the contents of the program theory and the identified CMO

configurations, the program theory will be tested and validated through realist interviews with

selected clinicians from each department using the program theory to test and consolidate findings

(33). The realist interviews will test, revise, and finalise the program theory to ensure no significant

points are missed during the data collection and analysis. This will ensure the program theory's

relevance, scalability, and transferability and validate the identified CMO configurations.

Randomisation

Hospitals are randomly allocated to start the interventional period after 4, 6, or 8 months,

respectively, using a random list generator (www.random.org). The project manager generated the

list and will inform the hospitals when they should cross over as late as possible but no later than

one month before the crossover.

Variables

Descriptives

During baseline, we will collect the following data: age, height, weight, location (unilateral or

bilateral) and duration of the knee pain, pain frequency, pain intensity during the week prior to

baseline measured on a 0 to 10 Numerical Rating Scale (0=no pain, 10= worst pain imaginable),

and pain medication use. At each follow-up, adolescents will be asked which treatment they have

received and which clinicians they may have consulted.

Primary outcome

The primary outcome will be change in KOOS-Child pain subscale (40) from baseline to the 12-

week follow-up. KOOS-Child consists of 39 items divided into five subscales (Pain, Symptoms,

Activities of Daily Living, Sport/Recreation, and Quality of Life) ranging from 0 to 100, with 100

being optimum, and we use a Danish translation of the questionnaire. This questionnaire has been

found to have good psychometric properties and is recommended for children and adolescents

suffering from knee disorders.(40) The questionnaire will be completed at baseline and the 12-week

and 52-week follow-ups.

Secondary outcomes

Secondary outcomes include: 1) Global Rating of Change (GROC), 2) EQ-5D-Youth, 3) Anterior

Knee Pain Youth (AKP-Youth), 4) the International Physical Activity Questionnaire short version

(IPAQ), and 5) sports participation.

We will use the GROC to measure the participants' self-reported recovery on a 7-point Likert scale

ranging from "much improved" to "much worse". Participants are categorised as improved if they

rate themselves as "much improved" or "improved" (category 6-7) and categorised as not improved

if they rate themselves from "slightly improved" to "much worse" (category 1-5). Health-related

quality of life will be estimated by the EQ-5D-Youth questionnaire, an adapted version of the EQ-

5D used in an adult population. Yet, it still consists of the same five subscales, which are mobility,

self-care, usual activities, pain and discomfort, and anxiety and depression.(41) We will use our

newly developed questionnaire (AKP-Youth) as a secondary measure of condition severity. The

AKP-Youth contains 23 items that are divided into four overarching domains of impact: symptoms,

limitations in physical activity, limitations in social activities, and emotional impact of pain.

Changes in physical activity level will be measured by the IPAQ. A Danish translation of the

original questionnaire will be used. The IPAQ is the most used questionnaire for measuring physical

activity and consists of 9 items that provide information on the time spent performing vigorous and

moderate activities, the time spent walking, and the time spent sedentary during the past week. The

IPAQ estimates the total weekly physical activity measured in MET minutes per week and the total

minutes spent sitting.(42,43) Change in sports participation will be explored using a questionnaire

in which participants are asked whether they have participated in leisure sports activities, type of

sports activities and weekly frequency. All questionnaires will be completed at baseline and at the

12-week and 52-week follow-ups. The GROC is only being answered during the 12-week and 52-

week follow-ups.

Adverse events

The participants will be able to report any adverse events to the primary investigator when they

occur by phone, SMS, or e-mail. Adverse events will be graded 1 to 5 according to the Common

Terminology Criteria for Adverse Events v4.03 (44). A medical doctor will assess and grade the

adverse event. Participants will not be withdrawn from the trial due to an adverse event. If a

participant experiences an adverse event and requests withdrawal from the study, data until the

adverse event occurred will be included in the analyses. The project manager will report any

incidents to the sponsor as quickly as possible and no later than 15 days after the participant reports

the event. Sponsor will report any severe adverse events (grade 3-5) to the Ethics Committee of

North Denmark Region no later than seven days after being informed.

Sample size

Trial

The trial is powered to detect a between-group difference of at least 10 points on the KOOS-Child questionnaire pain subscale scale (0 to 100) (40). Based on a standard deviation of 22 points, type I error α =0.05 and power = 0.90, a conservative estimate of four steps and an intraclass correlation coefficient = 0.01, we will need at least 240 participants (20 per step per cluster). We will increase the sample size to 290 participants across the six clusters to account for a larger outcome variance.

Realist Evaluation

Participants for the realist evaluation will be selected progressively based on an ongoing assessment of participants' knowledge power (34) and how their experiences and knowledge inform the theory-gleaning, testing and consolidation phases of the realist evaluation (45). We anticipate that 15-20 clinician interviews and 4-8 interviews with adolescents with non-traumatic knee pain treated with the MAP-Knee Tool will be sufficient to inform the development, testing and consolidation of a program theory and ensure data completeness throughout the project.

Statistical analyses

All statistical analyses will be performed according to a pre-established analysis plan by a statistician. This plan is written with the statistician and will be published on the Aalborg University website before the inclusion of the last participant. Stata ver. 18 will be used as statistical software. We will use Q-Q plots and histograms to assess data distribution. The primary intention-to-treat analysis will investigate the between-group difference in KOOS-Child pain at the 12-week follow-up using a linear mixed effects model with the participant as random effect. The baseline value, time (12 and 52 weeks), group allocation (using or not using MAP-Knee Tool) and term for the interaction between time and group will be treated as fixed-effect variables. We will apply the same model for the other continuous outcomes. The relative risk (RR) will be calculated for the dichotomised GROC to determine the probability of being improved after 12 and 52 weeks. The number needed to treat will be calculated as 1/risk difference.

Qualitative data analysis

The data collected from the realist interviews with researchers, clinicians, stakeholders, and
adolescents with non-traumatic knee pain will be collected, stored, and analysed prospectively
using Gilmore et al.'s (30) memo-coding technique for extracting CMO configurations from
qualitative data. The memo-coding technique was selected because it focuses on extracting CMO
configurations and incorporating them into a program theory through a five-step process, which
included data preparation, CMO configuration extraction, program theory development, refinement,
and synthesis across multiple data collections. The data analysis will be conducted using NVivo 14
to sort and organise the data. The linked coding approach by Jackson and Kolla (29) will be used to
identify CMO strings within the raw data before using the memo-coding technique to derive CMO
configurations from the data. By adopting the memo-coding technique for organising and
synthesising our data, we obtain a foundation for bridging the gap between identifying 'empirical
themes', exploring 'co-occurring phenomena' within the data and extracting the 'structures and
mechanisms' which generate the actual phenomena present within the use situation and informing
our program theory (30,33).

The analysis will be conducted in steps by two researchers (SKJ; EMA) and involve the uptake and
multiple datatypes (document, researcher, clinician, clinical staff, adolescents) to inform the
ongoing development of the program theory (23,27,32). The data preparation and transcription will
be conducted using Whisper 1.1 (Open AI, San Francisco, CA, USA) AI transcription software and
ExpressScribe software v. 7.01 (NCH Software, Canberra, Australia) to check and resolve
transcription errors within the data before analysis. The text files from the document-, researcher-,
clinician and validation interviews will be uploaded and analysed sequentially using NVivo 14
analysis using the following steps.

The initial transcription of interview data will be undertaken using the Whisper 1.1 tool. The AI-
transcribed files will be checked manually by a researcher using ExpressScribe 7.05 transcription
software to ensure transcript integrity. Before the analysis, the preliminary CMOs from the initial
program theories (IPT) gleaned during the document analysis were entered as nodes within NVivo
14. During the data coding, potential CMOs will be identified via the linked coding approach and
added to an existing node (IPT) or a new node linked to an existing node (30). All nodes with
encoded data will be reviewed and subjected to interpretation using linked memos to summarise the
contents, extract CMO configurations and identify their internal relationships. From this, the

extracted CMO configurations will be organised hierarchically, using the child nodes to evaluate, support, refute, revise, and expand the IPTs through a memo writeup process. The revised IPTs and CMO configurations from each analysis will be integrated within a logic model (33) to inform the formulation of a program theory as the analysis progresses, to identify the change mechanisms acting as barriers and facilitators for supporting patient education, stratification and shared decision-making when using the MAP-Knee Tool in complex treatment settings.

Validation interviews

The contents of the program theory and identified CMO configurations will be tested and subjected to evaluation through realist interviews with selected clinicians from each department who had used the MAP-Knee Tool when treating adolescents with non-traumatic knee pain. The validation interviews will combine principles from realist interviewing using selected parts of the program theory to inform discussions (31) to solicit participants' feedback and suggestions for optimising the program theory and CMO configurations to reflect their experiences by using the MAP-Knee Tool, and the individual and contextual barriers and facilitators for supporting patient education, stratification, and shared decision-making.

Data monitoring and quality assurance

All data will be stored electronically and handled according to the General Data Protection Regulation. Data safety may be overseen unannounced by the Danish Data Protection Agency. Participant data will be stored in REDCap, whereas data processor agreements, collaboration agreements between the project group and hospitals, and protocols will be stored on a secure server at Aalborg University. Data collection instruments have been developed in REDCap to prevent data entry errors so that required data must be included or an error will be displayed. Validation of each field has been chosen (e.g., if the data format does not appear to be a date in the field 'Date', an error is displayed). Data are checked once per week by the project manager to ensure no missing data, and participants and/or their parents are contacted if they do not respond to the questionnaires sent to them for the 12-week and 52-week follow-ups. All data will be kept for ten years after the trial is completed per the European Code of Conduct for Research Integrity.

Discussion

Moving from "what we prescribe" to "how it is delivered"

Previous trials to support adolescents with chronic knee pain have focused on what clinicians
should deliver (e.g., exercise, orthoses, taping or load management (14,16,17,46,47)). This trial
focuses on how the initial clinical encounter and clinicians' decision-making can be improved to
meet the support and management needs of adolescents with chronic knee pain seeking treatment
for knee pain in secondary care. Previous studies have highlighted how adolescents' formation of
self-management strategies does not occur within a vacuum (48) and how the lack of validation
from HCPs, peers and parents, uncertainty about the severity of the condition, excessive use of
diagnostic imaging and 'watchful waiting' without explanations may negatively influence
adolescents' formation of strategies for mastering their knee pain (49–52). Contrarily, patients
highlight how validation from clinicians, a name, and an explanation for why their knee pain
emerged enabled adolescents to adjust their pain beliefs and commence the work related to
accepting and exploring how to self-manage their knee pain in everyday situations (52). Studies
from other domains have described how informed parents can help facilitate adolescents' transition
into self-management by providing comfort, management instructions, problem-solving assistance
and engaging with external actors (teachers, trainers, etc.) (53,54), adolescents with knee pain and
parents described how they needed different types of information from HCPs to undertake this task
(51). Whereas shared decision-making has been highlighted as a viable method for merging
evidence-based treatments and patients' desired outcomes when prescribing treatments in clinical
settings (55), the question of how patient education is delivered to enable adolescents and parents to
engage in an informed negotiation of roles and management tasks is unexplored (56). Thus, we
anticipate that the inclusion of the MAP-Knee Tool can help shift the clinicians' focus from simply
prescribing treatments to how treatments and patient education can be delivered to make it
'actionable' for adolescents with knee pain by visualising the clinicians' decision-making process,
similar to what Star (57) describes within their boundary object—ensuring that clinicians
understand how the MAP-Knee Tool is not a treatment in itself but rather a tool for merging their
existing clinical expertise with patient-centred care principles via a systematised approach.
462

A tool that can be refined when new evidence emerges
The MAP-Knee Tool should not be viewed as a static tool but as a tool that can be updated when
new evidence emerges. This will ensure that clinicians have information regarding best practice
readily available to them. Therefore, it may be preferable to convert the tool into an online version
in the future. The current physical form of the tool is not seen as crucial as it is merely a delivery

vehicle of the content, which is the fundamental aspect of the MAP-Knee Tool. Yet, using a leaflet

can extend the clinical encounter, and the use of written materials has been found to aid patients

with chronic pain conditions in reducing pain catastrophising.(58)

Limitations of the trial design

An important potential limitation of the trial is the nature of the design in which recruitment relies

on the natural flow of patients referred to the hospitals. This does not allow adjusting the

recruitment strategy if the expected recruitment rate is unmet. Based on our past experiences when

recruiting adolescents with non-traumatic knee pain, the recruitment rate required to achieve the

sample size was feasible. However, due to a lower-than-expected recruitment rate during the first

two months of recruitment, we had to postpone the first crossover from after two months to after

four months of using usual care. To balance the periods of all hospitals using usual care with all

hospitals using The MAP-Knee Tool, we also extended the period after the last crossover from two

to four months. Due to the learning curve when starting to use the tool, it may be challenging for

clinicians to become confident in using it if the recruitment rate is not somewhat consistent. This

could potentially limit the implementation. Another challenge that could hamper recruitment is the

fact that the ethics committee did not approve only obtaining informed consent from a single

parent/legal guardian. Therefore, we must obtain retrospective informed consent from the other

parent in cases where only one parent attends the clinical examination at the hospital. If the other

parent cannot be contacted or does not want to sign the informed consent form, the participant will

need to be excluded.

Moving from clinical trials to understanding "how and why things work"

While multiple studies have explored implementing decision-making tools to support the cultivation

of patient-centred and collaborative care approaches in primary and secondary care settings (59–

62), these studies commonly focus on measuring the effects of interventions. This is problematic,

as studies from implementation science highlight how care settings are complex, dynamic, and

adaptive systems (63) and how introducing a tool for, e.g. optimising HCP workflows, may ripple

into other domains and influence the tasks, roles, division of labour and community and cultural

aspects within cares setting (64,65). Furthermore, introducing tools may also have unforeseen

impacts on workflows, collaborations, and communications within the organisations, resulting in

implementers having to use extra resources to find workarounds when using the tool (66). Thus, the

historical and traditional focus on exploring the effect of interventions has resulted in a dearth of

knowledge on how future treatment and supporting tools should be modelled to support their

meaningful integration into complex treatment settings. This is also true in Danish secondary care.

By adopting a realist perspective on the testing and evaluating the MAP-Knee Tool, an optic is

obtained that enables us to transcend the scope of the clinical trial and explore how, why, and which

circumstances result in the outcomes of the MAP-Knee Trial (23). Furthermore, by adapting the

view of interventions as programs as described by Pawson and Tilley (23), the qualitative and

quantitative data collected during the intervention with the MAP-Knee Tool can be synthesised to

identify the causal powers influencing the implementation of the tool on a micro, meso, and macro

level. By identifying the different context mechanisms activated by the MAP-Knee Tool

intervention, visualised within a program theory, the study has the potential to extract novel, general

insights on why some interventions fail and for whom, while others succeed, of high relevance for

future studies in secondary care.

References

1. Murray CJ, Richards MA, Newton JN, Fenton KA, Anderson HR, Atkinson C, et al. UK health performance: findings of the Global Burden of Disease Study 2010. The Lancet. 2013;381(9871):997–1020.

2. Bjerrum L, Ertmann RK, Jarbøl DE, Jensen MB, Kristensen JK, Maagaard R. Almen medicin. 1st ed. Hunskår S, editor. Munksgaard; 2014.

3. Skovdal Rathleff M, Rams Rathleff C, Lykkegaard Olesen J, Roos EM, Rasmussen S, Andreucci A, et al. Care-seeking behaviour of adolescents with patellofemoral pain: a retrospective cohort study. F1000Research 2022 11:161 [Internet]. 2022 Feb 9 [cited 2022 Sep 15];11:161. Available from: https://f1000research.com/articles/11-161

4. Brattberg G. Do pain problems in young school children persist into early adulthood? A 13-year follow-up. European Journal of Pain. 2004 Jun 1;8(3):187–99.

5. Incledon E, O'Connor M, Giallo R, Chalkiadis GA, Palermo TM. Child and Family Antecedents of Pain During the Transition to Adolescence: A Longitudinal Population-Based Study. J Pain. 2016 Nov 1;17(11):1174–82.

6. Fuss S, Pagé MG, Katz J. Persistent pain in a community-based sample of children and adolescents: Sex differences in psychological constructs. Pain Research & Management : The Journal of the Canadian Pain Society. 2011;16(5):303.

7. Patel DR, Villalobos A. Evaluation and management of knee pain in young athletes: overuse injuries of the knee. Transl Pediatr. 2017 Jul 1;6(3):19098–198.

8. Barber Foss KD, Myer GD, Chen SS, Hewett TE. Expected Prevalence From the Differential Diagnosis of Anterior Knee Pain in Adolescent Female Athletes During Preparticipation Screening. J Athl Train. 2012 Sep 1;47(5):519–24.

9. Stracciolini A, Casciano R, Levey Friedman H, Stein CJ, Meehan WP, Micheli LJ. Pediatric Sports Injuries. https://doi.org/101177/0363546514522393. 2014 Feb 24;42(4):965–72.

10. Rathleff MS, Roos EM, Olesen JL, Rasmussen S. High prevalence of daily and multi-site pain – a cross-sectional population-based study among 3000 Danish adolescents. BMC Pediatr. 2013 Nov 19;13(1):191.

11. Rathleff MS, Skuldbøl SK, Rasch MNB, Roos EM, Rasmussen S, Olesen JL. Care-seeking behaviour of adolescents with knee pain: a population-based study among 504 adolescents. BMC Musculoskelet Disord. 2013;14:225.

12. Rathleff MS, Rathleff CR, Olesen JL, Rasmussen S, Roos EM, Crossley K, et al. Is Knee Pain During Adolescence a Self-limiting Condition?: Prognosis of Patellofemoral Pain and Other Types of Knee Pain. Am J Sports Med. 2016 May 1;44(5):1165–71.

13. Rathleff MS, Holden S, Straszek CL, Olesen JL, Jensen MB, Roos EM. Five-year prognosis and impact of adolescent knee pain: a prospective population-based cohort study of 504 adolescents in Denmark. BMJ Open [Internet]. 2019 May 1 [cited 2022 Mar 25];9(5):24113. Available from: /pmc/articles/PMC6549701/

14. Rathleff MS, Roos EM, Olesen JL, Rasmussen S. Exercise during school hours when added to patient education improves outcome for 2 years in adolescent patellofemoral pain: a cluster randomised trial. Br J Sports Med. 2014;49(6):406–12.

15. Rathleff MS, Rathleff CR, Holden S, Thorborg K, Olesen JL. Exercise therapy, patient education, and patellar taping in the treatment of adolescents with patellofemoral pain: a prospective pilot study with 6 months follow-up. Pilot Feasibility Stud [Internet]. 2018 Apr 25 [cited 2022 Jun 1];4(1). Available from: /pmc/articles/PMC5899375/

16. Rathleff MS, Graven-Nielsen T, Hölmich P, Winiarski L, Krommes K, Holden S, et al. Activity Modification and Load Management of Adolescents With Patellofemoral Pain: A

560  Prospective Intervention Study Including 151 Adolescents. American Journal of Sports
561  Medicine. 2019 Jun 1;47(7):1629–37.

562 17. Rathleff MS, Winiarski L, Krommes K, Graven-Nielsen T, Hölmich P, Olesen JL, et al.
563  Activity Modification and Knee Strengthening for Osgood-Schlatter Disease: A Prospective
564  Cohort Study. Orthop J Sports Med. 2020 Apr 1;8(4).

565 18. Stacey D, Légaré F, Lewis K, Barry MJ, Bennett CL, Eden KB, et al. Decision aids for
566  people facing health treatment or screening decisions. Vol. 2017, Cochrane Database of
567  Systematic Reviews. John Wiley and Sons, Inc. and the Cochrane Library; 2017.

568 19. van der Horst DEM, Garvelink MM, Bos WJW, Stiggelbout AM, Pieterse AH. For which
569  decisions is Shared Decision Making considered appropriate? – A systematic review. Patient
570  Educ Couns. 2023 Jan 1;106:3–16.

571 20. Zadro JR, Traeger AC, Décary S, O'Keeffe M. Problem with patient decision aids. Vol. 26,
572  BMJ Evidence-Based Medicine. [London]: BMJ Publishing Group,; 2021. p. 180–3.

573 21. Djurtoft C, Bruun MK, Riel H, Hoegh MS, Darlow B, Rathleff MS. How do we explain
574  painful non-traumatic knee conditions to adolescents? A multiple-method study to develop
575  credible explanations. European Journal of Pain. 2023;00:1.

576 22. Riel H, Bruun MK, Djurtoft C, Jensen MB, Kaalund S, Leeuwen G van, et al. Development
577  of a clinical decision-support tool for Management of Adolescent knee Pain (The MAP-Knee
578  Tool). medRxiv. 2023 Jan 12;2023.01.11.23284426.

579 23. Pawson R, Tilley N. An Introduction to Scientific Realist Evaluation. In: Evaluation for the
580  21st Century: A Handbook [Internet]. 2455 Teller Road, Thousand Oaks California 91320
581  United States: SAGE Publications, Inc.; 1997 [cited 2022 May 19]. p. 405–18. Available
582  from: https://methods.sagepub.com/book/evaluation-for-the-21st-century/n29.xml

583 24. Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of
584  interventions: template for intervention description and replication (TIDieR) checklist and
585  guide. BMJ. 2014 Mar 7;348(mar07 3):g1687–g1687.

586 25. Chan AW, Tetzlaff JM, Gøtzsche PC, Altman DG, Mann H, Berlin J a, et al. SPIRIT 2013
587  explanation and elaboration: guidance for protocols of clinical trials. BMJ. 2013;346:e7586.

588 26. Bandholm T, Christensen R, Thorborg K, Treweek S, Henriksen M. Preparing for what the
589  reporting checklists will not tell you: the PREPARE Trial guide for planning clinical research
590  to avoid research waste. Br J Sports Med. 2017 Sep 7;bjsports-2017-097527.

591 27. F Kazi MA, to Mansoor F Kazi CA. Realist Evaluation for Practice. The British Journal of
592  Social Work. 2003 Sep 1;33(6):803–18.

593 28. Manzano A. The craft of interviewing in realist evaluation. Evaluation. 2016 Jul
594  1;22(3):342–60.

595 29. Jackson SF, Kolla G. A New Realistic Evaluation Analysis Method.
596  https://doi.org/101177/1098214012440030 [Internet]. 2012 Mar 16 [cited 2024 Mar
597  21];33(3):339–49. Available from:
598  https://journals.sagepub.com/doi/10.1177/1098214012440030

599 30. Gilmore B, McAuliffe E, Power J, Vallières F. Data Analysis and Synthesis Within a Realist
600  Evaluation: Toward More Transparent Methodological Approaches. Int J Qual Methods
601  [Internet]. 2019 Jul 1 [cited 2024 Mar 21];18. Available from:
602  https://journals.sagepub.com/doi/10.1177/1609406919859754

603 31. Mukumbang FC, Marchal B, Van Belle S, van Wyk B. Using the realist interview approach
604  to maintain theoretical awareness in realist studies. Qualitative Research. 2019 Oct
605  23;20(4):485–515.

32. Ebenso B, Manzano A, Uzochukwu B, Etiaba E, Huss R, Ensor T, et al. Dealing with context
in logic model development: Reflections from a realist evaluation of a community health
worker programme in Nigeria. Eval Program Plann. 2019 Apr 1;73:97.
33. Mukumbang FC, Marchal B, Van Belle S, van Wyk B. Using the realist interview approach
to maintain theoretical awareness in realist studies. Qualitative Research [Internet]. 2019 Oct
23;20(4):485–515. Available from: https://doi.org/10.1177/1468794119881985
34. Malterud K, Siersma VD, Guassora AD. Sample Size in Qualitative Interview Studies. Qual
Health Res [Internet]. 2016 Nov 10 [cited 2018 Apr 30];26(13):1753–60. Available from:
http://journals.sagepub.com/doi/10.1177/1049732315617444
35. Guldhammer C, Holden S, Sørensen ME, Olesen JL, Jensen MB, Rathleff MS. Development
and validation of the Sorting non-trauMatIc adoLescent knEe pain (SMILE) tool – a
development and initial validation study. Pediatric Rheumatology. 2021 Dec 6;19(1):110.
36. Holden S, Kasza J, Winters M, van Middelkoop M, Rathleff MS. Prognostic factors for
adolescent knee pain: an individual participant data meta-analysis of 1281 patients. Pain.
2021 Jun;162(6):1597–607.
37. Sackett DL, Rosenberg WMC, Gray JAM, Haynes RB, Richardson WS. Evidence based
medicine: what it is and what it isn't. 1996. Clin Orthop Relat Res. 2007 Jan
13;455(7023):3–5.
38. Tracing genres through organizations : a sociocultural approach to information design.
Cambridge, Mass.: MIT Press; 2003. xi, 246 s. (Acting with technology).
39. Wiltshire G, Ronkainen N. A realist approach to thematic analysis: making sense of
qualitative data through experiential, inferential and dispositional themes. J Crit Realism
[Internet]. 2021 Mar 15;20(2):159–80. Available from:
https://doi.org/10.1080/14767430.2021.1894909
40. Örtqvist M, Iversen MD, Janarv PM, Broström EW, Roos EM. Psychometric properties of
the Knee injury and Osteoarthritis Outcome Score for Children (KOOS-Child) in children
with knee disorders. Br J Sports Med. 2014 Oct 1;48(19):1437–46.
41. Wille N, Badia X, Bonsel G, Burström K, Cavrini G, Devlin N, et al. Development of the
EQ-5D-Y: a child-friendly version of the EQ-5D. Quality of Life Research. 2010
Aug;19(6):875.
42. van Poppel MNM, Chinapaw MJM, Mokkink LB, van Mechelen W, Terwee CB. Physical
Activity Questionnaires for Adults. Sports Medicine. 2010 Jul;40(7):565–600.
43. Craig CL, Marshall AL, Sjöström M, Bauman AE, Booth ML, Ainsworth BE, et al.
International physical activity questionnaire: 12-country reliability and validity. Med Sci
Sports Exerc. 2003 Aug;35(8):1381–95.
44. Common Terminology Criteria for Adverse Events (v4.03). U.S.Department of Health and
Human Services; 2010.
45. Manzano A. The craft of interviewing in realist evaluation. Evaluation [Internet]. 2016 Jul 1
[cited 2024 Feb 5];22(3):342–60. Available from:
https://journals.sagepub.com/doi/10.1177/1356389016638615
46. Willy RW, Hoglund LT, Barton CJ, Bolgla LA, Scalzitti DA, Logerstedt DS, et al.
Patellofemoral pain clinical practice guidelines linked to the international classification of
functioning, disability and health from the academy of orthopaedic physical therapy of the
American physical therapy association. Journal of Orthopaedic and Sports Physical Therapy.
2019 Sep 1;49(9):CPG1–95.
47. Collins NJ, Barton CJ, Van Middelkoop M, Callaghan MJ, Rathleff MS, Vicenzino BT, et al.
2018 Consensus statement on exercise therapy and physical interventions (orthoses, taping
and manual therapy) to treat patellofemoral pain: recommendations from the 5th

International Patellofemoral Pain Research Retreat, Gold Coast, Australia, 2017. Br J Sports Med. 2018 Sep 1;52(18):1170–8.

48. Modi AC, Pai AL, Hommel KA, Hood KK, Cortina S, Hilliard ME, et al. Pediatric Self-management: A Framework for Research, Practice, and Policy. Pediatrics [Internet]. 2012 Feb [cited 2024 Mar 21];129(2):e473. Available from: /pmc/articles/PMC9923567/

49. Glaviano NR, Holden S, Bazett-Jones DM, Singe SM, Rathleff MS. Living well (or not) with patellofemoral pain: A qualitative study. Physical Therapy in Sport. 2022 Jul 1;56:1–7.

50. Andreucci A, Rathleff MS, Reuther FØ, Hussein M, Rahimzai S, Linnemann TD, et al. "I had already tried that before going to the doctor" – exploring adolescents' with knee pain perspectives on 'wait and see' as a management strategy in primary care; a study with brief semi-structured qualitative interviews. Scand J Pain [Internet]. 2023 Apr 1 [cited 2024 Mar 21];23(2):341–52. Available from: https://www-degruyter-com.zorac.aub.aau.dk/document/doi/10.1515/sjpain-2022-0038/html

51. Johansen SK, Kanstrup AM, Haseli K, Stenmo VH, Thomsen JL, Rathleff MS. Exploring User Visions for Modeling mHealth Apps Toward Supporting Patient-Parent-Clinician Collaboration and Shared Decision-making When Treating Adolescent Knee Pain in General Practice: Workshop Study. JMIR Hum Factors [Internet]. 2023 [cited 2024 Mar 21];10(4). Available from: /pmc/articles/PMC10182461/

52. Johansen SK, Holden S, Pourbordbari N, Jensen MB, Thomsen JL, Rathleff MS. PAINSTORIES – Exploring the Temporal Developments in the Challenges, Barriers, and Self-Management Needs of Adolescents with Longstanding Knee Pain: A Qualitative, Retrospective Interview Study with Young Adults Experiencing Knee Pain Since Adolescence. J Pain. 2022 Apr 1;23(4):577–94.

53. Leonard BJ, Garwick A, Adwan JZ. Adolescents' Perceptions of Parental Roles and Involvement in Diabetes Management. J Pediatr Nurs. 2005 Dec 1;20(6):405–14.

54. Cha YJ, Saxena A, Wou A, Lee J, Newman MW, Park SY. Transitioning Toward Independence: Enhancing Collaborative Self-Management of Children with Type 1 Diabetes. Conference on Human Factors in Computing Systems - Proceedings [Internet]. 2022 Apr 29 [cited 2024 Mar 21]; Available from: https://dl.acm.org/doi/10.1145/3491102.3502055

55. Elwyn G, Frosch D, Thomson R, Joseph-Williams N, Lloyd A, Kinnersley P, et al. Shared Decision Making: A Model for Clinical Practice. J Gen Intern Med [Internet]. 2012 Oct [cited 2024 Mar 21];27(10):1361. Available from: /pmc/articles/PMC3445676/

56. Sandman L. The concept of negotiation in shared decision making. Health Care Analysis [Internet]. 2009 Sep 7 [cited 2024 Mar 21];17(3):236–43. Available from: https://link-springer-com.zorac.aub.aau.dk/article/10.1007/s10728-008-0103-y

57. Star SL. This is Not a Boundary Object: Reflections on the Origin of a Concept. Sci Technol Human Values [Internet]. 2010;35(5):601–17. Available from: http://www.jstor.org/stable/25746386

58. Gallagher L, McAuley J, Moseley GL. A randomized-controlled trial of using a book of metaphors to reconceptualize pain and decrease catastrophizing in people with chronic pain. Clinical Journal of Pain. 2013 Jan;29(1):20–5.

59. Alsulamy N, Lee A, Thokala P, Alessa T. What Influences the Implementation of Shared Decision Making: An Umbrella Review. Patient Educ Couns. 2020 Dec 1;103(12):2400–7.

60. Boland L, Graham ID, Légaré F, Lewis K, Jull J, Shephard A, et al. Barriers and facilitators of pediatric shared decision-making: a systematic review. Implement Sci [Internet]. 2019 Jan 18 [cited 2024 Mar 25];14(1). Available from: /pmc/articles/PMC6339273/

61.   Légaré F, Ratté S, Gravel K, Graham ID. Barriers and facilitators to implementing shared
decision-making in clinical practice: Update of a systematic review of health professionals'
perceptions. Patient Educ Couns. 2008 Dec 1;73(3):526–35.

62.   Grenfell J, Soundy A. People's Experience of Shared Decision Making in Musculoskeletal
Physiotherapy: A Systematic Review and Thematic Synthesis. Behavioral Sciences
[Internet]. 2022 Jan 1 [cited 2024 Mar 25];12(1). Available from:
/pmc/articles/PMC8773142/

63.   Ellis B. Complexity in practice: understanding primary care as a complexadaptive system. J
Innov Health Inform [Internet]. 2010 [cited 2024 Mar 21];18(2):135. Available from:
https://openurl.ebsco.com/contentitem/doi:10.14236%2Fjhi.v18i2.763?sid=ebsco:plink:crawl
er&id=ebsco:doi:10.14236%2Fjhi.v18i2.763

64.   Bardram J, Doryab A. Activity analysis: Applying activity theory to analyze complex work in
hospitals. Proceedings of the ACM Conference on Computer Supported Cooperative Work,
CSCW [Internet]. 2011 [cited 2024 Mar 21];455–64. Available from:
https://dl.acm.org/doi/10.1145/1958824.1958895

65.   Engestrom Y. Activity theory as a framework for analyzing and redesigning work.
Ergonomics [Internet]. 2000 Jul 1 [cited 2024 Mar 21];43(7):960–74. Available from:
https://www-tandfonline-com.zorac.aub.aau.dk/doi/abs/10.1080/001401300409143

66.   Spinuzzi Clay. Tracing genres through organizations. Vol. 1. Mit Press; 2003.
719