Title

Effect of a needs-based model of care on the characteristics of healthcare services in

England: the i-THRIVE National Implementation Programme

Short title

Effect of needs-based care on healthcare services

Authors

R Sippy[1], L Efstathopoulou[2,3], E Simes[2,3], M Davis[2,3], S Howell[2,3], B Morris[2], O Owrid[2,3], N

Stoll[2,3], P Fonagy[2,3], A Moore[1,2,3]*

[1]Department of Psychiatry, University of Cambridge, Cambridge, United Kingdom

[2]Anna Freud National Centre for Children and Families, London, United Kingdom

[3]Department of Department of Clinical, Educational and Health Psychology, University

College London, London, United Kingdom

* corresponding author, am2708@medschl.cam.ac.uk, Clifford Allbutt Building, Hills Road,

Cambridge, CB2 0XY

Word count

3,584

Abstract

Aims: Developing integrated mental health services that focus on the needs of children and

young people is a key policy goal in England. The THRIVE Framework and its associated

implementation programme, i-THRIVE, are now used in areas covering over 65% of

England's children. This study explores the experiences of staff involved with the i-THRIVE

programme, assesses its effectiveness, and examines how local system working

relationships influence the programme's success.

Methods: The i-THRIVE programme was evaluated using a quasi-experimental study

among twenty participating sites (ten implementation and ten comparison sites).

Measurements included surveys of staff and leaders at each site and assessment of the

"THRIVE-like" features of each site. Additional site-level characteristics were collected from

health system reports. The effect of i-THRIVE was evaluated using a four-group propensity-

score weighted difference-in-differences model; the moderating effect of local system

working relationships was evaluated with a difference-in-difference-in-differences model.

Results: Staff at implementation sites were more likely to report using THRIVE in their own

practice and exhibited better knowledge of THRIVE principles than comparison sites. The

mean improvement among i-THRIVE sites was 16.7, and 8.8 among comparison sites. The

results show that strong working relationships in the local system significantly enhances the

effectiveness of the i-THRIVE programme. Sites with highly effective working relationships

showed a notable improvement in "THRIVE-like" features, with an average increase of 16.41

points (95% confidence interval: 1.69–31.13, p-value: 0.031) compared to control sites. In

contrast, sites with ineffective working relationships did not benefit from the i-THRIVE

programme (-2.76, 95% confidence interval: -18.25–12.73, p-value: 0.708). This influence of

working relationship effectiveness was consistent across various levels of THRIVE features.

Conclusions: The findings underscore the importance of working relationship effectiveness

in the successful adoption and implementation of health policies like i-THRIVE.

Background

Mental healthcare has numerous well-supported models. Globally, systems are challenged

to provide appropriate services efficiently, necessitating reform of current care models

(Hodgins et al., 2024). However, the continued presence of inadequate care is mainly due to

difficulties in applying these changes (Shortell et al., 1993; Grol and Grimshaw, 2003), as

systemic transformation is a complex process (Best et al., 2012). The transformative "Future

in Mind" report (Department of Health, 2015) suggested deviating from a generic, tiered

service model to a more responsive model for the specific mental health needs of local

young populations. It called for service providers and funders to overhaul children and young

people's mental health (CYPMH) services into comprehensive systems, offering a range of

services from prevention to risk management. The "Future in Mind" report focuses on

reforming a flawed system of service provision, characterised by disunity, inefficiency, and

limited access to services. Since 2016, Child and Adolescent Mental Health Services

(CAMHS) within England's National Health Service (NHS) have seen significant changes,

advancing more comprehensive approaches to care (Rocks et al., 2020).

The THRIVE Framework provides a set of principles to guide system reformation,

summarising the needs of CYP into five groups: Getting Advice, Getting Risk Support,

Getting Help, Getting More Help, and Thriving (Wolpert et al.). Support is provided to CYP in

these groups using a set of guiding principles. The THRIVE principles of care encompass

characteristics of support at three levels: macro, meso, and micro. At the macro level,

characteristics include interagency function and cooperation. This means a CAMHS system

following the "THRIVE-like" approach would involve supporting bodies such as educational

and social services in its policy-making and service delivery (Moore et al., 2023). At the

meso level, they include a needs-based perspective focusing on CYP and their support

services. A site adhering to meso-level THRIVE principles would be expected to have a

network of community providers (Moore et al., 2023). At the micro level, these characteristics

involve the interactions between CYP, their families, and healthcare professionals; we would

expect to see shared decision-making, with everyone understanding the needs of the child

and interventions being used (Moore et al., 2023).

The National i-THRIVE Programme (NIP) assists CAMHS sites in adopting the THRIVE

principles, which prioritise patient needs and cohesive service provision through

collaborative care networks (Moore et al., 2023). Over 65% of children and young people

(CYP) live in areas where i-THRIVE has been adopted (i-THRIVE Team). i-THRIVE was

created following implementation science guidelines to facilitate the adoption of THRIVE

principles by CAMHS (Moore et al., 2016). The NIP is explained in a published study

protocol (Moore et al., 2023). Briefly, i-THRIVE translates the complex aspects of a

'THRIVE-like' system into practical structures and tools for CAMHS to use in their

transformation. The NIP guides CAMHS staff and leaders in developing local models based

on THRIVE principles and creating detailed plans for implementation over four phases, using

six components (Moore et al., 2023). The implementation strategies in the NIP are drawn

from the Quality Implementation Framework (Meyers et al., 2012) and the Normalisation

Processing Theory (May, 2006). The embodiment of THRIVE principles depends not only on

efforts by the CAMHS site but also on broader community involvement (Wolpert et al.),

requiring effective working relationships among local systems. According to the

implementation strategy clusters created by the Expert Recommendations for Implementing

Change study (Waltz et al., 2015), the implementation strategies used in the NIP include use

of evaluative and iterative strategies, providing interactive assistance, adaptation and

tailoring to context, development of stakeholder interrelationships, training and education of

stakeholders, engagement of consumers, utilisation of financial strategies, and changes to

infrastructure.

Evaluation of implementation strategies is needed to understand the impacts of the

implementation and determine if it should be adopted for use in real-world scenarios. The

evaluation could include several outcome metrics, such as acceptability, feasibility, fidelity or

cost (Smith and Hasan, 2020). Fidelity is a particularly important metric, as it helps to

determine if observed effects can be attributed to the intervention of interest; poor

implementation may explain cases with no observed effects (Sanders et al., 2022). In this

work, we evaluate the impact of the NIP implementation model on fidelity to the THRIVE

principles of care; a healthcare service unit that embodies the THRIVE principles of care

would exhibit high fidelity to THRIVE at macro, meso, and micro levels.

Although in use since 2016, the NIP as an implementation model has not been assessed.

We focus on the average effect of NIP on the THRIVE fidelity of implementation sites. For

this estimate, we need to know the outcome at each site if NIP had not been implemented

(the potential outcome or unobserved counterfactual) (Stuart, 2010). The difference in

differences (DiD) approach compares changes in outcomes between implementation and

control sites before and after the intervention (Stuart et al., 2014). A key assumption for DiD

is that the average outcomes for both groups would have similar trends over time (Xu, 2017),

which may not always be plausible. Participation in NIP is voluntary, meaning sites that

choose to participate may differ from those that do not. The four-group propensity score-

weighted DiD method overcomes these issues by adjusting the implementation and control

groups while considering time factors. This adjustment ensures comparability among the

groups and reduces bias in estimating the desired effect. It's particularly useful when group

composition changes during the study, such as when health practitioners move between

Clinical Commissioning Groups (CCGs).

In this study, we evaluate the alignment of frontline staff with THRIVE principles, and assess

the effect of the NIP on THRIVE fidelity. We hypothesize that i-THRIVE had a positive effect

on THRIVE fidelity. Additionally, we examine how the effectiveness of local system working

relationships moderates the impact of the NIP on THRIVE fidelity.

Methods

<u>Study Setting and Design</u>

The study protocol is detailed elsewhere (Moore et al., 2023). In brief, we selected twenty

CAMHS sites in England: ten of these sites have been using i-THRIVE since 2016

(NIP/implementation sites), and ten others were using different approaches for

transformation (comparison sites). Details on the implementation strategies and delivery of

NIP are in Supplemental Material S1. This study reporting conforms to the Template for

Intervention Description and Replication (TIDieR) statement (Hoffmann et al., 2014); the

TIDieR checklist is provided in Supplemental Material S2.

The NIP sites were Bexley, Cambridgeshire and Peterborough, Camden, Hertfordshire,

Luton, Manchester, Stockport, Tower Hamlets, Waltham Forest, and Warrington. The

comparison sites included Bradford, Ipswich and East Suffolk, Lewisham, Norfolk,

Northampton, Portsmouth, Southampton, Stoke-on-Trent, Sunderland, and South

Worcestershire. Sites will be pseudonymised as Site A—T. Additional site details are in

Supplemental Material S3. Implementation of THRIVE started in April 2016; data from April

2015 to March 2016 were used as the pre-implementation period. Data from April 2018 to

March 2019 were used as the post-implementation period. To examine staff alignment with

and understanding of THRIVE principles, a survey was administered. To assess site

transformation or implementation, site leaders were also surveyed. To assess whether

THRIVE principles were embraced by sites, fidelity scores were assigned. To understand

whether the NIP impacted site adoption of THRIVE principles, fidelity scores from pre- and

post-implementation were compared between implementation and comparison sites, while

adjusting for site-level characteristics (auxiliary data).

<u>Measurements and Data</u>

Surveys

Two surveys were designed based on the RE-AIM Adoption Framework (Glasgow et al.,

2019), a guide for evaluation of programmes. Both surveys were primarily quantitative in

nature, but several questions requested open-ended responses. Copies of the surveys are in

Supplemental Materials S4—S6. The staff survey assessed the staff awareness of THRIVE

and use of THRIVE principles. Site leaders distributed the survey to professionals involved in

providing and commissioning CYPMH services at both i-THRIVE and comparison sites. The

survey was administered in August 2019, October 2019, and January 2020.

A survey for transformation leads was also conducted among i-THRIVE and comparison

sites. This survey, completed by programme managers, gathered information about site

transformation activities and the support they received. It was distributed from January to

May 2020. Additional details on the surveys are in Supplemental Material S3.

Principal Outcome: Fidelity

The primary focus is the degree to which sites follow the THRIVE principles. The THRIVE

principles of care encompass macro-, meso-, and micro-level features. We assessed these

features in participating sites using the i-THRIVE Assessment Tool (Moore et al., 2023),

wherein a score is assigned to each of 75 items describing THRIVE principles (1: low, 4:

high). Purposive sampling was used to recruit three staff members at macro (senior

leadership), meso (service management), and micro (frontline staff) levels for in-depth

interviews. Interview transcripts were scored by evaluators: sites received an overall fidelity

(300 possible) and level-specific scores (macro: 84 possible, meso: 104 possible, micro: 112

possible). Higher scores indicate better adherence to THRIVE principles. Fidelity to i-

THRIVE was measured at each site before and after the intervention by at least two

independent evaluators, with scores averaged for each site and period. When there was not

enough information to assign a score, we estimated the missing scores using the mean

difference between pre- and post-intervention scores for the site, based on the score for the

available period. Additional details on fidelity ratings are in Supplemental Materials S3.

Auxiliary Data

To calculate propensity score weights, it is important to identify site characteristics that might

cause selection bias or have a confounding effect (Stuart et al., 2014). These characteristics

were measured in 2016 and 2019, unless stated otherwise. They included population density

(total population per square kilometre) for each CCG (Office for National Statistics, 2016;

Office for National Statistics, 2017), annual funding support (£100,000 increments) per CCG

(NHS England, 2017), Indices of Multiple Deprivation (IMD) ranks (Department for

Communities and Local Government, 2015; Department for Communities and Local

Government, 2019), the initial number of CCGs per site (NHS England, 2019), and

compliance in 2017 with the CCG Improvement and Assessment Framework Mental Health

Transformation Milestones (NHS England, 2017). For sites comprising multiple CCGs, we

summed the annual funding and averaged the population density, effectiveness of working

relationships in the local system, and transformation compliance across CCGs. Site-level

IMD ranks were calculated using the method recommended by the Office of National

Statistics (Noble et al., 2019).

<u>Statistical Analysis</u>

We analysed the staff survey results using the chi-squared test and Cramer's V as a

measure of association between i-THRIVE/comparison sites and staff responses

("yes"/"no"). We used a binomial distributed generalised linear model with a log link for site-

specific results, excluding "I don't know" responses. To determine if any site had an unusual

probability of a "yes" response, we compared the "yes" probability for each site to the

average among comparison sites.

The responses from the survey of the implementation leads are in Supplemental Material

S1. Their experience of the delivery of i-THRIVE was summarised as proportions.

To assess inter-rater reliability for the fidelity scores, we used Krippendorf's alpha (see

Supplemental Material S3 for more details).

Voluntary participation in health policy implementations like the NIP can lead to selection

bias, as the sites that choose to participate may differ from those that do not. To correct for

this, we applied propensity score weighting (Rosenbaum and Rubin, 1985) to equalise the

distribution of characteristics between the implementation and comparison sites. We used a

four-group weighting method (pre-implementation i-THRIVE, post-implementation i-THRIVE,

pre-implementation comparison, and post-implementation comparison) to align

characteristics across all groups with those of the pre-implementation i-THRIVE sites (Stuart

et al., 2014). These propensity scores were calculated using a multinomial model with five

site characteristics (population density, annual funding, IMD rank, the number of CCGs per

site, and transformation compliance). Characteristic balance was checked using the

standardised difference in means (Stuart, 2010). The impact of NIP on fidelity was estimated

221 using maximum-likelihood repeated measures linear regression with an auto-regressive

222 correlation structure, weighted with the calculated propensity scores. To account for

223 remaining characteristic imbalances, we included population density, IMD, and

224 transformation compliance in the final model. The results from this model represent the four-

225 group weighted DiD effect estimate.

226 To assess the reliability of our fidelity results, we conducted a series of sensitivity analyses.

227 We employed different methods to estimate the effect of the NIP, including the standard

228 (unweighted) DiD and alternative model specifications. We examined the impact of non-

229 compliant control sites by repeating the analysis, excluding these sites. More details can be

230 found in Supplemental Material S7.

231 To examine variations in the effect of i-THRIVE, we investigated a possible effect moderator,

232 specifically the quality of local system working relationships (NHS England, 2021) on i-

233 THRIVE implementation. Further details are available in Supplemental Material S9.

234 All data visualisation, cleaning, and propensity score modelling was performed in R version

235 4.2.3 (The R Foundation for Statistical Computing, Vienna, Austria) and RStudio version

236 2023.03.0 (RStudio, Inc., Boston, MA, USA), with R packages cowplot, extrafont, ggpattern,

237 ggspatial, gridExtra, haven, irr, nnet, RColorBrewer, readxl, reshape2, sf, tableone, and

238 tidyverse (Venables and Ripley, 2002; Wickham, 2007; Neuwirth, 2014; Auguie, 2017;

239 Wickham and Miller, 2017; Pebesma, 2018; Gamer et al., 2019; Wickham et al., 2019;

240 Wilke, 2020; FC et al., 2022; Yoshida and Bartel, 2022; Chang, 2023; Dunnington, 2023;

241 Pebesma and Bivand, 2023; Wickham and Bryan, 2023). The simple DiD, four-group

242 weighted DiD, and effect modification analyses were performed in SAS® version 3.81 (SAS

243 Institute Inc., Cary, NC, USA) in SAS® Studio.

245 **Results**

246 <u>Surveys</u>

247 The staff survey had 689 responses across 19 sites (no responses from Luton). Detailed

248 survey results are in Supplemental Material S8.

While the THRIVE Framework was widely known, more implementation respondents

recognised it (83.9% vs. 70.5%, p<0.0001). A higher proportion of implementation

respondents reported using THRIVE principles in their daily practice (58.5% vs. 49.0%,

p=0.03). Implementation respondents were more likely to score perfectly on a test of the

THRIVE Framework (34.1% vs. 22.9%, p=0.001).

The transformation leads survey included eight managers from seven implementation sites

and eight managers from seven comparison sites. Notably, managers from four comparison

sites (Sites E, J, K, and S) reported using THRIVE as their service transformation model.

This prompted a more detailed examination of staff survey results concerning THRIVE

implementation at comparison sites.

The site-level analysis of survey results can be found in Supplemental Material S8. Among

those who provided a yes/no response at comparison sites, J and K reported a higher

likelihood of implementing THRIVE. The odds of respondents reporting site implementation

of THRIVE were 4.43 in J (95% CI: 2.32–8.47) compared to other comparison sites (76.5%

vs. 59.5%), and 4.43 in K (95% CI: 1.33–14.80) compared to other comparison sites (76.5%

vs. 59.5%). Among respondents at comparison sites, there was no difference in personal

use of THRIVE principles compared to other comparison sites. Regarding knowledge of the

THRIVE Framework, respondents from J had a higher probability of achieving a perfect

score on the quiz: the odds of scoring perfectly were 3.73 (95% CI: 2.25–6.19), compared to

other comparison sites (37.7% vs. 22.9%).

THRIVE Fidelity

During the pre-implementation period, i-THRIVE sites had an average fidelity score of 149.0

(range: 132.0—180.2) and comparison sites had an average score of 133.4 (range: 113.0—

158.2). Following implementation, i-THRIVE sites had an average score of 166.6 (range:

145.5—195.0), while comparison sites had an average of 142.2 (range: 132.0 —175.0). The

mean difference between pre- and post-implementation among i-THRIVE sites was 16.7;

among comparison sites the mean difference was 8.8. Two sites had incomplete fidelity

score information: the macro-level components for Site T during the post-implementation

period and the meso-level components for Site F during the pre-implementation period

(scores were assigned as outlined in the methods section). Detailed fidelity scores by level

and site are illustrated in Figure 1, and a map showing the changes in scores by site

throughout the study is presented in Figure 2.

Figure 1: Fidelity Scores by Site

Total fidelity scores during the pre- and post-implementation periods are represented by bar

height, with patterned overlay to indicate component levels (macro, meso, micro).

Implementation sites are in panel A (blue) while comparison sites are in panel B (red).

Figure 2: Change in Fidelity Scores Over Study Period

The difference in total fidelity scores during the pre- and post-implementation periods are

represented by colour, with an increased score in green and a decreased score in red, on a

background map of Clinical Commissioning Groups in panel A. Study sites have a bold

outline (comparison sites in red, implementation sites in blue). Inset maps for North

West/Midlands, London, and South East are in panels B–D, respectively.

Site characteristics, adjusted using four-group propensity-score weighting, are in Table S6,

Supplemental Material S8. In our analysis of the weighted standardised differences, we

identified some remaining imbalances, specifically in population density for the pre-

implementation control group, IMD rank for both control groups, and transformation

compliance for both control groups. These covariates were included in our effect estimate

model, ensuring a more accurate assessment of NIP impact.

The NIP effect estimates are presented in Table 1. Our analysis reveals that the overall

fidelity scores were moderately influenced by the NIP. Specifically, i-THRIVE sites showed

an average improvement of 7.05 points (95% CI: -4.47–18.57). The most notable

improvements were at the macro level, where i-THRIVE sites increased by an average of

2.92 points (95% CI: -1.09–6.92), followed by the meso level with an average increase of

2.76 points (95% CI: -1.98–7.51), and the micro level with an average increase of 1.39

points (95% CI: -3.94–6.72). None of these improvements were statistically significant.

When comparing the four-group weighted DiD with the standard DiD analyses

(Supplemental Material S7), we found comparable effects on overall and macro-level fidelity.

There were shifts in the impacts on meso- and micro-level fidelity. This suggests that lower-

level fidelity was more sensitive to the disparities between i-THRIVE and comparison

groups, which were corrected through the four-group propensity-score weighting approach.

Alternative modelling approaches produced results similar to our analysis. The exclusion of

non-compliant comparison sites (i.e. Site J; see Supplemental Material S7) did not alter the

results.

Among the study sites, five implementation sites and five comparison sites were found to

have highly effective working relationships among their local systems (Supplemental

Material S9). We observed a moderating effect of local systems working relationship

effectiveness on the impact of the NIP for overall and macro-level fidelity. i-THRIVE was

found to be more effective at sites with highly effective working relationships. The detailed

results are in Table 2. i-THRIVE sites with highly effective working relationships showed

increased fidelity scores compared to comparison sites with highly effective working

relationships. The most significant impact was on overall fidelity scores (16.41, 95% CI:

1.69–31.13), followed by macro-level scores (6.95, 95% CI: 2.15–11.75). The moderating

influence of highly-effective working relationships on meso-level and micro-level fidelity was

modest (5.52, 95% CI: -0.66–11.71 and 3.95 points, 95% CI: -3.24–11.15, respectively).

Notably, there was no discernible impact of the NIP on sites with ineffective working

relationships across any fidelity level.

Discussion

A major obstacle in implementing any complex intervention is ensuring it becomes a routine

practice across the entire organisation. This concept is often described as making it 'the way

we do things around here' (Haines et al., 2004; Proctor et al., 2011). Both THRIVE and i-

THRIVE are frequently mentioned in child mental health policy documents, by NHS

England's regional transformation boards, and in media related to child mental health. It is

encouraging for policymakers to note that over 70% of staff at comparison sites were aware

of THRIVE, and nearly 23% exhibited perfect knowledge of THRIVE principles. At the site

level, respondents from most i-THRIVE sites had an increased odds of reporting site

implementation of THRIVE, compared to the average among comparison sites, though two

comparison sites (Sites J and K) also had an increased odds of reporting site

implementation of THRIVE. When asked about personal use of THRIVE principles,

respondents from many i-THRIVE sites had an increased odds of reporting personal use,

compared to the average among comparison sites, and with Site J also exhibiting an

increased odds of reporting personal use. This motivated a sensitivity analysis in which Site

J was excluded from the fidelity analysis; the results were not affected. Overall, staff at i-

THRIVE sites were significantly more likely to be familiar with, understand, and apply

THRIVE principles in their daily work. This indicates that the NIP aids in embedding THRIVE

principles within an organisation.

Previous research has found staff development and training are key to the successful

implementation of a new intervention (Rahman et al., 2012; Resnick et al., 2018). Training

activities and attendance at both i-THRIVE Academy and Community of Practice events

were reported among implementation sites, and staff from implementation sites exhibited

knowledge of THRIVE principles. Continued engagement of staff will likely be important for

the integration of THRIVE, and formal assessments or competency checks may be useful to

monitor THRIVE delivery (Sanders et al., 2022). In addition to educational activities, several

of the implementation strategies used by the NIP, including site diagnostics activities,

coaching and support, and involving key partners were found to have positive results at least

75% of the time in a recent review (Ashcraft et al., 2024)

Implementation research generates critical evidence for selecting effective strategies for

health system transformation (Ashcraft et al., 2024). Like the NIP, most implementations

include multiple strategies (Ashcraft et al., 2024), meaning it is difficult to attribute a

successful implementation to any single strategy. There are few examples of implementation

research for mental health, many of which focus on implementation effectiveness (Wells et

al., 2013; Chung et al., 2014; Kilbourne et al., 2014; Waxmonsky et al., 2014; Chung et al.,

2015; Sinnema et al., 2015; Morton et al., 2020; Ruud et al., 2021; Bartels et al., 2022;

Toropova et al., 2022) with a minority examining the implementation process more broadly

(Williams et al., 2017; Bauer et al., 2019; Leone et al., 2022). Among studies of the

implementation process or implementation fidelity, a facilitator-implemented collaborative

care model resulted in some improvements to team function; adoption of collaborative care

model processes varied widely by site (Bauer et al., 2019). Another study examined

mediating factors for clinician adoption of evidence-based practices following an

organisation-supported implementation strategy, reporting high fidelity to the strategy at both

clinician and organisation levels (Williams et al., 2017). Although implementation studies

would benefit from enrolment of many participating sites to reduce the impact of between-

site variation, the enrolment of a high number of sites presents many logistical challenges.

Integrated care models, such as THRIVE, are used in Europe, Australia, and North America,

and have been tailored for use in services for CYP (Hodgins et al., 2024). A review of

integrated care models for youth discusses common features, including multidisciplinary staff

members able to partner with external organisations and managers committed to integration,

joint planning, and stakeholder partnership (Hodgins et al., 2024). The THRIVE Framework

targets a local community, and relies on the involvement of multiple agencies to transform

how these agencies provide mental healthcare for CYP (Wolpert et al.). Many of the

implementation sites hosted events to build stakeholder relationships. In this study, we found

that multi-agency cooperation was critical to the implementation of the NIP itself: the strength

of working relationships in the local system moderated the effect of the NIP on THRIVE

fidelity among sites. For future users of the NIP, an evaluation of local working relationships

or pre-implementation efforts to engage with community stakeholders and strengthen these

relationships would be worthwhile.

On average, the NIP had a modest impact on THRIVE fidelity among sites, without reaching

the level of a statistically-significant change. With a small number of sites included, the study

may have been under-powered. When implementation studies find a null effect, it is difficult

to know if the effect is truly null or if the implementation itself was incorrect (Sanders et al.,

2022), or if implementation success depends on site characteristics (Augustsson et al.,

2015). Site-level changes in THRIVE fidelity had high variability; even when an

implementation is supported in the same way for all sites, variation can occur (Augustsson et

al., 2015). To better understand the impact of NIP, we examined the moderating effect of an

important site characteristic: the strength of local working relationships at the site, finding

that the NIP improved THRIVE fidelity (overall and at macro level) at implementation sites

with strong local working relationships, compared to comparison sites with strong local

working relationships. A similar result was found in a study of an organisational-level

intervention for occupational health, where the implementation worked the best among units

with strong collaboration (Augustsson et al., 2015). To explain these results, the authors

suggest that the intervention was a better fit for those units or that the units were more

capable in adapting the intervention. Preliminary work can be done to prepare sites for an

implementation; for those sites interested in using the NIP, efforts to strengthen local

working relationships prior to implementation should be considered.

The future direction of this research will involve comprehensive evaluations of the service

and clinical impacts of the NIP. These evaluations will encompass a range of critical factors,

including the accessibility and efficiency of services, clinical outcomes, patient experiences,

and the specific impacts on various sub-groups within the patient population. The latter will

particularly focus on racial or ethnic minorities and distinct diagnostic categories, ensuring a

broad and inclusive understanding of the NIP's effectiveness.

Limitations

There are several limitations to consider in this work. For the survey of site staff, the

response rate was low (28.5%) compared to staff surveys in other implementation studies

(46.8—83.1%) (Leone et al., 2022; Toropova et al., 2022). We would expect this to cause

some bias in our results if the response rate was differential by implementation/comparison

site groups, but there was no difference in response rate by these groups. The low response

rate may indicate that the survey results are not generalisable. Future research should

consider incentives to encourage survey completion.

When introducing new health policies, understanding their impact and identifying the

contexts in which they are most effective is crucial. Estimating the mean effect presents

several methodological challenges, especially when data from pre-implementation periods

are scarce or when only a few sites are involved. Unlike randomised controlled trials, the

adoption of health policies is not random. This means that the characteristics of the

implementation and control sites are likely to vary, leading to potential selection bias. Yet, it

is a useful methodological approach for evaluating complex interventions selected and

implanted by policymakers. It is often unclear which characteristics influence a site's

decision to adopt a health policy. Even with numerous characteristics measured, careful

consideration of each characteristic's role is necessary. Confounding is another potential

bias, where some characteristics may influence both the decision to implement and the site's

capacity to do so effectively.

To adjust for these biases, various methods are available, but these methods can

themselves introduce bias. Sensitivity analyses are essential in gaining a deeper

understanding of the data. The four-group propensity-score weighting DiD method is

designed to mitigate potential selection biases and confounding factors (Stuart et al., 2014).

However, a key assumption of DiD, the parallel trends assumption, is not verifiable.

Violations of the parallel trends assumption can lead to issues with time-varying confounding

(Xu, 2017), complicating the interpretation of results. Thus, while our study provides valuable

insights into the effectiveness of the NIP, these issues must be considered when interpreting

the findings.

The variable transformation approaches used by comparison sites could be considered a

limitation, as this could complicate the interpretation of results. Strictly speaking, a

comparison group where all sites were transforming their CAMHS into the THRIVE

Framework would allow us to test the implementation strategies of the NIP specifically (i.e.

all sites seeking to fit the THRIVE Framework but the implementation group testing the NIP

implementation strategies). Four comparison sites reported using THRIVE as their

transformation model, but this sample size is too small for a full analysis. The comparison

group simply represents routine implementation, a common approach for implementation

research control groups (Smith and Hasan, 2020).

Conclusions

This study's investigation into the effectiveness of the NIP in England offers significant

insights into the implementation of complex health interventions, particularly in the field of

CAMHS. The findings underscore the importance of effective working relationships among

local systems in the successful adoption and implementation of health policies like i-

THRIVE. Specifically, the study demonstrates that sites with highly effective working

relationships exhibit substantial improvements in adhering to THRIVE principles, as

evidenced by the increase in fidelity scores.

The broad awareness of the THRIVE framework among staff, even in comparison sites,

highlights the programme's permeation in the field of CYPMH. However, implementation

strategies are critical in embedding these principles more deeply within organisations. This

distinction is crucial for policymakers and healthcare leaders aiming to foster more effective,

integrated mental health services tailored to the needs of CYP.

Methodologically, the study navigates the challenges of evaluating health policy

implementations in non-randomised settings. The use of four-group propensity-score

weighted DiD analysis is an effective approach to address potential biases, such as

selection bias and confounding factors. However, it also highlights the inherent complexities

and limitations in evaluating policy impact in real-world settings. The study's sensitivity

analyses further strengthen the validity of its findings.

In conclusion, the NIP presents a promising model for CAMHS. Its emphasis on effective

working relationships and the tailored approach to implementation are key factors in its

success. The insights from this study contribute valuable knowledge to the ongoing efforts to

improve mental health services and can guide future policies and programmes aimed at

enhancing the well-being of CYP.

Financial support

This study is independent research funded by the National Institute for Health and Care

Research (NIHR) Applied Research Collaboration North Thames (grant number CRJM,

46AZ74-/RCF/CLAHRC/UCL004, 52AZ95/AA2/UCL5, 52AZ95/AA2/UCL6, UCLP) and

UCLPartners, both awarded to AM. The views expressed in this publication are those of the

author(s) and not necessarily those of the NIHR, the Department of Health and Social Care

or UCLPartners. All research at the Department of Psychiatry in the University of Cambridge

is supported by the NIHR Cambridge Biomedical Research Centre (BRC-1215-20014) and

NIHR Applied Research Centre. The views expressed are those of the author(s) and not

necessarily those of the NIHR or the Department of Health and Social Care.

Conflicts of Interest

AM was the lead for i-THRIVE between January 2015 and January 2017 during which she

led the development of the i-THRIVE implementation approach and established the

community of practice. She was an employee of one i-THRIVE implementation site during

the evaluation.

LE was an employee of one i-THRIVE implementation site during the evaluation.

PF is currently on the board of the i-THRIVE Implementation Programme. Anna Freud

Centre hosted the development and on-going implementation of the THRIVE Framework in

collaboration with the Tavistock & Portman NHS Foundation Trust. He is Programme

Director of Mental Health at UCLPartners which provides programme management of the i-

THRIVE Implementation Programme.

Ethical Standards

The NHS/University Joint Research Office reviewed this study and classified it as a service

evaluation, meaning it does not need ethical approval or a review by the Research Ethics

Committee (IRAS application number: 250439). For the qualitative parts of the study,

participants included implementation leads, managers, and front-line staff, who were

interviewed about their service transformation projects. At the start of each interview, verbal

informed consent was obtained, which included explicit permission for the interview to be

recorded, analysed, and used in the service evaluation. This consent was recorded, and

transcripts of each interview were made to document it.

The data for the quantitative evaluation is retrospective and was routinely collected from

service records. It was anonymised at the source by the business intelligence staff at each

trust, with all identifying information removed. After review by local information governance

leads, this anonymised data was provided to the evaluation team. As the data is

retrospective and de-identified, consent for its use in the evaluation is neither legally nor

ethically required. The requirement for ethical approval was therefore waived by the joint

research office.

Availability of Data and Materials

All datasets and code are accessible on GitHub (Sippy, 2023).

Acknowledgements

The authors are indebted to the clinicians, managers and academics who have created

THRIVE and their generous help and thoughtful advice throughout the evaluation to assess

its effectiveness, in particular Sophie Dunn. We would also like to thank Rudolf Cardinal for

help with the research.

References

Ashcraft LE, Goodrich DE, Hero J, Phares A, Bachrach RL, Quinn DA, Qureshi N,
Ernecoff NC, Lederer LG, Scheunemann LP, et al (2024) A systematic review of
experimentally tested implementation strategies across health and human service
settings: evidence from 2010-2022. Implementation Science **19**: 43

Auguie B (2017) gridExtra: Miscellaneous functions for "Grid" graphics.

Augustsson H, Von Thiele Schwarz U, Stenfors-Hayes T, Hasson H (2015) Investigating
Variations in Implementation Fidelity of an Organizational-Level Occupational Health
Intervention. International Journal of Behavioral Medicine **22**: 345–355

Bartels SJ, Aschbrenner KA, Pratt SI, Zubkoff L, Jue K, Williams G, Godfrey MM,
Cohen MJ, Banerjee S, Xie H, et al (2022) Virtual Learning Collaborative Compared
to Technical Assistance as a Strategy for Implementing Health Promotion in Routine
Mental Health Settings: A Hybrid Type 3 Cluster Randomized Trial. Administration
and Policy in Mental Health **49**: 1031–1046

Bauer MS, Miller CJ, Kim B, Lew R, Stolzmann K, Sullivan J, Riendeau R, Pitcock J,
Williamson A, Connolly S, et al (2019) Effectiveness of Implementing a
Collaborative Chronic Care Model for Clinician Teams on Patient Outcomes and
Health Status in Mental Health: A Randomized Clinical Trial. JAMA network open **2**:
e190230

Best A, Greenhalgh T, Lewis S, Saul JE, Carroll S, Bitz J (2012) Large-system
transformation in health care: a realist review. The Milbank Quarterly **90**: 421–456

Chang W (2023) extrafont: Tools for Using Fonts.

Chung B, Ngo VK, Ong MK, Pulido E, Jones F, Gilmore J, Stoker-Mtume N, Johnson
M, Tang L, Wells KB, et al (2015) Participation in Training for Depression Care
Quality Improvement: A Randomized Trial of Community Engagement or Technical
Support. Psychiatric Services (Washington, DC) **66**: 831–839

Chung B, Ong M, Ettner SL, Jones F, Gilmore J, McCreary M, Sherbourne C, Ngo V,
Koegel P, Tang L, et al (2014) 12-month outcomes of community engagement
versus technical assistance to implement depression collaborative care: a partnered,
cluster, randomized, comparative effectiveness trial. Annals of Internal Medicine **161**:
S23-34

Department for Communities and Local Government (2015) English Indices of
Deprivation 2015 File 5 Scores for the Indices of Deprivation.

Department for Communities and Local Government (2019) English Indices of
Deprivation 2019 File 5 Scores for the Indices of Deprivation.

Department of Health (2015) Future in mind: Promoting, protecting and improving our
children and young people's mental health and wellbeing.

Dunnington D (2023) ggspatial: Spatial data framework for ggplot2.

FC M, Davis TL, ggplot2 authors (2022) ggpattern: 'ggplot2 Pattern Geoms.

Gamer M, Lemon J, ¡puspendra.pusp22@gmail.com¿ IFPS (2019) irr: Various
coefficients of interrater reliability and agreement.

Glasgow RE, Harden SM, Gaglio B, Rabin B, Smith ML, Porter GC, Ory MG,
Estabrooks PA (2019) RE-AIM Planning and Evaluation Framework: Adapting to
New Science and Practice With a 20-Year Review. Frontiers in Public Health **7**: 64

Grol R, Grimshaw J (2003) From best evidence to best practice: Effective implementation
of change in patients' care. Lancet **362**: 1225–1230

Haines A, Kuruvilla S, Borchert M (2004) Bridging the implementation gap between
knowledge and action for health. Bulletin of the World Health Organization **82**: 724–
731; discussion 732

Hodgins M, McHugh C, Eapen V, Georgiou G, Curtis J, Lingam R (2024) Creation of the
Youth Integration Project Framework: A Narrative Synthesis of the Youth Mental
Health Integrated Care Literature. International Journal of Integrated Care **24**: 5

Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, Altman DG,
Barbour V, Macdonald H, Johnston M, et al (2014) Better reporting of
interventions: template for intervention description and replication (TIDieR) checklist
and guide. British Medical Journal Clinical Research **348**: g1687

i-THRIVE Team National i-THRIVE Community of Practice. Anna Freud National Centre for
Children and Families

Kilbourne AM, Almirall D, Goodrich DE, Lai Z, Abraham KM, Nord KM, Bowersox NW
(2014) Enhancing outreach for persons with serious mental illness: 12-month results
from a cluster randomized trial of an adaptive implementation strategy.
Implementation science: IS **9**: 163

Leone SS, Smeets O, Lokman S, Boon B, van der Poel A, Van Doesum T, Shields-
Zeeman L, Kramer J, Smit F (2022) Comparing a ses-sensitive and an all-ses
implementation strategy to improve participation rates of patients with a lower
socioeconomic background in a web-based intervention for depressive complaints: a
cluster randomised trial in primary care. BMC primary care **23**: 205

May C (2006) A rational model for assessing and evaluating complex interventions in health
care. BMC health services research **6**: 86

Meyers DC, Durlak JA, Wandersman A (2012) The quality implementation framework: a
synthesis of critical steps in the implementation process. American Journal of
Community Psychology **50**: 462–480

Moore A, Jenkins P, Harris R, Fonagy P, Wolpert M (2016) The crisis in CAMHS: Can i-
Thrive provide a solution? British Association for Counselling and Psychotherapy
Children, Young People and Families 4–10

Moore A, Lindley Baron-Cohen K, Simes E, Chen S, Fonagy P (2023) A protocol for a
multi-site cohort study to evaluate child and adolescent mental health service
transformation in England using the i-THRIVE model. PLoS ONE **18**: e0265782

Morton I, Hurley B, Castillo EG, Tang L, Gilmore J, Jones F, Watkins K, Chung B,
Wells K (2020) Outcomes of two quality improvement implementation interventions
for depression services in adults with substance use problems. The American
Journal of Drug and Alcohol Abuse **46**: 251–261

Neuwirth E (2014) RColorBrewer: ColorBrewer Palettes.

NHS England (2017) Mental Health Five Year Forward View Dashboard.

NHS England (2019) CCG Allocations 2019/20 to 2023/24 - Technical Guidance.

NHS England (2021) CCG Improvement and Assessment Framework (CCG IAF) Data
Extract 2017/18.

Noble S, McLennan D, Noble M, Plunkett E, Gutacker N, Silk M, Wright G (2019) The
English Indices of Deprivation 2019.

Office for National Statistics (2016) CCG (July 2015) Ultra Generalised Clipped
Boundaries in England.

Office for National Statistics (2017) CCG (April 2018) Ultra Generalised Clipped
Boundaries in England.

Pebesma E (2018) Simple features for R: Standardized support for spatial vector data. The
R Journal **10**: 439–446

Pebesma E, Bivand R (2023) Spatial data science: With applications in R. doi:
10.1201/9780429459016

Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A, Griffey R,
Hensley M (2011) Outcomes for implementation research: conceptual distinctions,
measurement challenges, and research agenda. Administration and Policy in Mental
Health **38**: 65–76

Rahman AN, Applebaum RA, Schnelle JF, Simmons SF (2012) Translating research into
practice in nursing homes: can we close the gap? The Gerontologist **52**: 597–606

Resnick B, Kolanowski A, Van Haitsma K, Galik E, Boltz M, Ellis J, Behrens L,
Flanagan NM, Eshraghi KJ, Zhu S (2018) Testing the evidence integration triangle
for implementation of interventions to manage behavioral and psychological
symptoms associated with dementia: Protocol for a pragmatic trial. Research in
Nursing & Health **41**: 228–242

Rocks S, Fazel M, Tsiachristas A (2020) Impact of transforming mental health services for
young people in England on patient access, resource use and health: a quasi-
experimental study. British Medical Journal Open **10**: e034067

Rosenbaum PR, Rubin DB (1985) Constructing a Control Group Using Multivariate
Matched Sampling Methods That Incorporate the Propensity Score. The American
Statistician **39**: 33

Ruud T, Drake RE, Šaltytė Benth J, Drivenes K, Hartveit M, Heiervang K, Høifødt TS,
Haaland VØ, Joa I, Johannessen JO, et al (2021) The Effect of Intensive
Implementation Support on Fidelity for Four Evidence-Based Psychosis Treatments:
A Cluster Randomized Trial. Administration and Policy in Mental Health **48**: 909–920

Sanders G, Griffiths C, Flint S, Christensen A, Gately P (2022) Implementation fidelity of
an Integrated Healthy Lifestyle Service: a process evaluation. Perspectives in Public
Health **142**: 278–286

Shortell SM, Gillies RR, Anderson DA, Mitchell JB, Morgan KL (1993) Creating
organized delivery systems: The barriers and facilitators. Hospital and Health
Services Administration **38**: 447–466

Sinnema H, Majo MC, Volker D, Hoogendoorn A, Terluin B, Wensing M, van Balkom A
(2015) Effectiveness of a tailored implementation programme to improve recognition,
diagnosis and treatment of anxiety and depression in general practice: a cluster
randomised controlled trial. Implementation science: IS **10**: 33

Sippy R (2023) rsippy/THRIVE-fidelity: submission. doi: 10.5281/ZENODO.10402515

Smith JD, Hasan M (2020) Quantitative approaches for the evaluation of implementation
research studies. Psychiatry Research **283**: 112521

Stuart EA (2010) Matching Methods for Causal Inference: A Review and a Look Forward.
Stat Sci. doi: 10.1214/09-STS313

Stuart EA, Huskamp HA, Duckworth K, Simmons J, Song Z, Chernew ME, Barry CL
(2014) Using propensity scores in difference-in-differences models to estimate the
effects of a policy change. Health Services and Outcomes Research Methodology
14: 166–182

Toropova A, Björklund C, Bergström G, Elinder LS, Stigmar K, Wåhlin C, Jensen I,
Kwak L (2022) Effectiveness of a multifaceted implementation strategy for improving
adherence to the guideline for prevention of mental ill-health among school personnel
in Sweden: a cluster randomized trial. Implementation science: IS **17**: 23

Venables WN, Ripley BD (2002) Modern Applied Statistics with S, 4th Edition. Springer,
New York

Waltz TJ, Powell BJ, Matthieu MM, Damschroder LJ, Chinman MJ, Smith JL, Proctor
EK, Kirchner JE (2015) Use of concept mapping to characterize relationships
among implementation strategies and assess their feasibility and importance: results
from the Expert Recommendations for Implementing Change (ERIC) study.
Implementation Science **10**: 109

Waxmonsky J, Kilbourne AM, Goodrich DE, Nord KM, Lai Z, Laird C, Clogston J, Kim
HM, Miller C, Bauer MS (2014) Enhanced fidelity to treatment for bipolar disorder:

results from a randomized controlled implementation trial. Psychiatric Services
(Washington, DC) **65**: 81–90

Wells KB, Jones L, Chung B, Dixon EL, Tang L, Gilmore J, Sherbourne C, Ngo VK,
Ong MK, Stockdale S, et al (2013) Community-partnered cluster-randomized
comparative effectiveness trial of community engagement and planning or resources
for services to address depression disparities. Journal of General Internal Medicine
28: 1268–1278

Wickham H (2007) Reshaping Data with the reshape Package. Journal of Statistical
Software **21**: 1–20

Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G,
Hayes A, Henry L, Hester J, et al (2019) Welcome to the Tidyverse. Journal of
Open Source Software **4**: 1686

Wickham H, Bryan J (2023) readxl: Read Excel Files.

Wickham H, Miller E (2017) haven: Import and Export "SPSS", "Stata" and "SAS" Files.

Wilke CO (2020) cowplot: Streamlined plot theme and plot annotations for "ggplot2."

Williams NJ, Glisson C, Hemmelgarn A, Green P (2017) Mechanisms of Change in the
ARC Organizational Strategy: Increasing Mental Health Clinicians' EBP Adoption
Through Improved Organizational Culture and Capacity. Administration and Policy in
Mental Health **44**: 269–283

Wolpert M, Harris DR, Hodges DS, Fuggle DP, James DR, McKenna DC, Jones DM,
Fonagy P, Fleming DI, Munk DS, et al THRIVE Framework for system change.

Xu Y (2017) Generalized Synthetic Control Method: Causal Inference with Interactive Fixed
Effects Models. Political Analysis **25**: 57–76

Yoshida K, Bartel A (2022) tableone: Create "table 1" to describe baseline characteristics
with or without propensity score weights.

700
701

Table 1: Estimates of Association Between the National i-THRIVE Programme and THRIVE Fidelity			
Outcome	Estimated Score Change	95% CI	p-value
Overall Fidelity	7.05	-4.47–18.57	0.212
Macro-level Fidelity	2.92	-1.09–6.92	0.141
Meso-level Fidelity	2.76	-1.98–7.51	0.234
Micro-level Fidelity	1.39	-3.94–6.72	0.588
CI = confidence interval			

Table 2: Effect Modification by Working Relationship Quality on the Impact of the National i-THRIVE Programme				
Outcome	Working Relationship Quality	Estimated Score Change	95% CI	p-value
Overall Fidelity	Highly effective	16.41	1.69–31.13	0.031
	Ineffective	-2.76	-18.25–12.73	0.708
Macro-level Fidelity	Highly effective	6.95	2.15–11.75	0.007
	Ineffective	-1.24	-6.29–3.80	0.605
Meso-level Fidelity	Highly effective	5.52	-0.66–11.71	0.076
	Ineffective	-0.75	-7.26–5.76	0.808
Micro-level Fidelity	Highly effective	3.96	-3.24–11.15	0.257
	Ineffective	-0.78	-8.35–6.79	0.828
CI = confidence interval				