The Relationship Between Triglyceride to High-Density Lipoprotein Cholesterol Ratio and Cardiovascular High-Risk: A Cross-Sectional Investigation

Pengcheng Li,[1,2][†] Jirui Cai,[2][†] Shuaifang Yuan,[2] Yapeng Li,[1] Huiting Shi,[1] Cui Liang,[1] Bing He,[2] Qiaotao Xie,[2] Baocang Lei,[2] Jing Bai,[2] Nan Wang,[2] Dongliang Liu,[2] Qichao Wang,[2] Jianwei Xiong,[2] Jin Wang,[2] Haoran Wang,[2]

[1]The First Affiliated Hospital of Zhengzhou University, Zhengzhou, China

[2]Luohe Central Hospital, Luohe Medical College, Luohe, China

Corresponding to: Haoran Wang, Luohe Central Hospital, Luohe Medical College, Luohe, China

Alternate corresponding author: Jin Wang, Luohe Central Hospital, Luohe Medical College, Luohe, China

[†]Pengcheng Li and Jirui Cai contributed equally to this work.

Abstract

Aims

The triglyceride to high-density lipoprotein cholesterol (TG/HDL-C) ratio reflects the balance between atherogenic and anti-atherogenic lipid fractions. The study aims to investigate the utility of the TG/HDL-C ratio in identifying high-risk groups for cardiovascular disease (CVD) within the general population.

Methods

The current study was a branch of the ChinaHEART cohort in middle China that involved a total of 6,593 community-dwelling adults. We examine the association between TG/HDL-C and CVD high-risk in a cross-sectional investigation.

Results

Results from restricted cubic spline and ROC analyses revealed a significant relationship between higher TG/HDL-C levels and increased odds of cardiovascular high-risk, with the ratio exhibiting good discriminatory power (AUC=0.819 in the fully adjusted model). Logistic

regression further supported this association, indicating a 1.21-fold rise in

the odds of high-risk for each unit increase in TG/HDL-C.

Conclusions

The TG/HDL ratio demonstrates potential as a valuable clinical tool

for CVD prevention. By incorporating the TG/HDL ratio into risk

assessment models, healthcare professionals can enhance risk

stratification and identify high-risk individuals who may benefit from

targeted interventions.

Key words:

TG/HDL-C ratio; cardiovascular high-risk; cross-sectional study;

public health.

Introduction

Cardiovascular diseases (CVD) continue to be a leading cause of

mortality and morbidity worldwide, imposing a significant burden on

healthcare systems and society as a whole. Early identification of

individuals at high risk of developing CVD is crucial for implementing

targeted prevention and management strategies.(2) Traditional risk factors

such as hypertension, dyslipidemia, smoking, and diabetes have been

widely recognized, but there is a growing interest in exploring novel

biomarkers that may enhance risk stratification in the general

population.(3)

In recent years, the triglyceride to high-density lipoprotein

cholesterol (TG/HDL-C) ratio has emerged as a potential marker for

cardiovascular risk assessment.(4–8) This ratio reflects the balance

between atherogenic and anti-atherogenic lipid fractions. and has been

proposed as a simple and cost-effective tool to identify individuals at

increased risk of CVD. Some studies have found that this indicator is

associated with morbidity and mortality from cardiovascular disease.(4,9)

However, there is still a lack of relevant studies on whether this indicator

can identify high-risk groups of cardiovascular disease in the general

population.

Our study aims to address this gap in knowledge by conducting a

cross-sectional observational study to evaluate the utility of the

TG/HDL-C ratio in identifying high-risk individuals for cardiovascular

diseases in the general population. By examining this novel biomarker in

a large and diverse sample, we seek to elucidate its potential as a

screening tool for early risk stratification and intervention. If we can

effectively identify individuals at high risk of cardiovascular disease in

the general population using this indicator, we can intervene at earlier

stages of cardiovascular disease, thereby maximizing the prevention of

major adverse cardiovascular events.

Methods

Study population and data collection

The current study is a branch of the China Health Evaluation And

risk Reduction through nationwide Teamwork (ChinaHEART) in Luohe

City, involving 6,859 community-dwelling adults recruited from two

township hospitals in Luohe City between November 2, 2021 and

February 20, 2022. The ChinaHEART cohort is a population cohort

launched by the China National Center for Cardiovascular Diseases

(NCCD) initially aiming at cardiovascular disease (CVD) risk screening

and management.(10) Study methods has been described previously.(11)

The study strictly adhered to ethical guidelines, with approval granted by

the Ethics Committee of Fuwai Hospital and the Ethics Committee of

Luohe Central Hospital. Participants received detailed information on the

study's purpose, significance, and risks before participant, in accordance

with the Declaration of Helsinki. Written informed consent was obtained

from all participants. The inclusion criteria for this study are as follows:

(1) Age: 35-75 years (born between January 1, 1946, and December 31,

1986); (2) Residing near the survey site for at least 6 months in the 12

months prior to screening; (3) Voluntarily participating in this project and

signing the informed consent form. The exclusion criteria are as follows:

(1) Survey participants with missing data; (2) Survey participants who

answered "unclear" to questionnaire items; (3) Survey participants with

fasting time less than 8 hours before blood collection.

We utilized standardized questionnaires to collect data on

socio-demographic characteristics (e.g., age, gender), lifestyle factors

(e.g., smoking, alcohol consumption), medical history (including

hypertension, diabetes, dyslipidemia, coronary heart disease, myocardial

infarction, stroke), and medication usage (such as antihypertensive,

hypoglycemic, and lipid-lowering drugs) of the study participants. Data

were collected through an information system provided by the

ChinaHEART study and was uploaded directly to the national study

center in the Fuwai Hospital. The anonymous data used in the current

study was applied from the national study center in the Fuwai Hospital

after they performed data washing. Cardiovascular disease (CVD) was

defined as a diagnosis of coronary heart disease, previous myocardial

infarction, or stroke. Body mass index (BMI) was calculated as body

mass in kilograms divided by the square of height in meters. Waist

circumference was measured by placing a measuring tape horizontally

around the abdomen at the midpoint between the anterior superior iliac

crest and the lower edge of the twelfth rib. Systolic blood pressure (SBP)

and diastolic blood pressure (DBP) were measured in a seated resting

position at least twice, and the average values were recorded. Lipid

profiles were assessed using fingertip blood samples.

According to the risk assessment prediction map in the Guidelines

for Cardiovascular Risk Assessment and Management issued by the

World Health Organization in 2008, the risk assessment of cardiovascular

disease was conducted.(12) If the screening subjects had 10 years of risk

of cardiovascular disease 20%, they were judged as high risk.

Statistics

The study population was categorized based on quartiles of the

TG/HDL-C ratio for basic characteristic description. Normality of

continuous variables was assessed using the Kolmogorov-Smirnov test,

with mean ± standard deviation for normally distributed data, and median

(25%-75% interval) for non-normally distributed data. Categorical

variables were described using percentages. Distribution differences in

high cardiovascular risk proportions across TG/HDL-C ratio quartile

populations were illustrated with a percentage stacking bar graph.

Continuous variable comparisons between groups involved t-tests or

analysis of variance for normally distributed data, and rank sum tests for

non-normally distributed data. Chi-square tests were conducted for

categorical variable comparisons. The association between the

TG/HDL-C ratio and cardiovascular high risk was examined using

logistic regression analysis, with two models utilized: one without

confounder adjustments for univariate analysis, and another adjusting for

confounding factors including gender, age, marital status, BMI, waist

circumference, fasting blood glucose, smoking history, alcohol

consumption history, diabetes, and systolic and diastolic blood pressure.

Nonlinear associations between the TG/HDL-C ratio and cardiovascular

risk were explored through restricted cubic spline analysis, utilizing key

points at the 10th, 25th, 50th, 75th, and 90th percentiles of the

TG/HDL-C ratio. The ability of the TG/HDL-C ratio to identify high

cardiovascular risk was evaluated using receiver operating characteristic

(ROC) curve analysis. While conducting the ROC curve analysis, we

calculated the appropriate cutoff value, and this cutoff value was used to

divide the participants into the high and low TG/HDL-C groups for

further logistic regression analysis. Furthermore, we used net

reclassification improvement (NRI) index to quantify the added value of

TG/HDL-C for the discriminative performance.

Results

A total of 6,593 participants with sufficient data were included in the

study, comprising 1,417 individuals categorized as cardiovascular

high-risk and 5,176 individuals in the non-high-risk category. The study

cohort was stratified into four segments based on the TG/HDL-C ratio,

highlighting notable variations in demographic and clinical characteristics

such as age, gender, height, weight, BMI, waist circumference, smoking

habits, alcohol consumption, history of hypertension, diabetes, as well as

levels of glucose, lipids, and blood pressure (refer to Table 1). The

distribution of high-risk individuals across these quartiles was 16%, 15%,

18%, and 38% respectively (view Figure 1).

To explore the nonlinear relationship between the TG/HDL-C ratio

and cardiovascular high-risk, a restricted cubic spline analysis was

conducted. The analysis revealed a positive association between elevated

TG/HDL-C ratio levels and an increased likelihood of cardiovascular

high-risk (see Figure 2).

The TG/HDL-C ratio's discriminatory ability in identifying

individuals at cardiovascular high-risk was evaluated through ROC

analysis (refer to Figure 3). Initially, in model 1, the AUC for the

TG/HDL-C ratio alone was 0.562 (95% CI 0.544-0.580) for all

participants. However, upon incorporating all covariates into the analysis

in model 2, the AUC significantly increased to 0.819 (95% CI

0.805-0.834), suggesting the enhanced effectiveness of combining the

TG/HDL-C ratio with traditional cardiovascular risk factors for high-risk

identification. Notably, the NRI index showed a significant improvement

in discriminative performance, with a value of 0.107 ± 0.030 (p = 0.016).

Moreover, the ROC analysis identified the optimal cutoff value for the

TG/HDL-C ratio as 1.18, enabling the classification of participants into

high and low TG/HDL-C ratio groups.

The logistic regression analysis results examining the relationship

between the TG/HDL-C ratio and cardiovascular high-risk are detailed in

Table 2. In Model 1, a 1 unit increase in the TG/HDL-C ratio was

associated with a 1.40-fold increase in the odds of cardiovascular

high-risk (95% CI 1.30-1.51, p < 0.001). Comparatively, the high

TG/HDL-C ratio group exhibited 1.61 times higher odds of high-risk than

the low TG/HDL-C ratio group (95% CI 1.43-1.81, p < 0.001), while the

highest quartile of the TG/HDL-C ratio had 1.79 times higher odds

compared to the lowest quartile (95% CI 1.52-2.10, p < 0.001). In Model

2, each 1 unit rise in the TG/HDL-C ratio was associated with a 1.21-fold

increase in the odds of cardiovascular high-risk (95% CI 1.11-1.32, p <

0.001). Additionally, the high TG/HDL-C ratio group demonstrated 1.27

times higher odds of high-risk compared to the low TG/HDL-C ratio

group (95% CI 1.10-1.47, p = 0.001), with the highest quartile of the

TG/HDL-C ratio displaying 1.49 times higher odds than the lowest

quartile (95% CI 1.21-1.83, p < 0.001).

Discussion

Cardiovascular disease (CVD) remains a significant global health

burden, necessitating the identification of individuals at high risk for

effective prevention and intervention strategies. The TG/HDL-C ratio has

emerged as a potential marker for assessing cardiovascular risk in the

general population. It has been observed that the ratio of TG/HDL-C ratio

not only provides an approximation of insulin resistance but also assists

in identifying patients with an atherogenic lipoprotein profile.(13) The

typical lipoprotein phenotype in insulin-resistant/hyperinsulinemic

patients is typified by high plasma TG and low HDL cholesterol levels,

smaller and denser LDL particles, as well as an increased accumulation of

remnant lipoproteins after a meal. This lipoprotein phenotype is not only

associated with insulin resistance, but it is also linked with an elevated

risk of cardiovascular disease.(14) Although plasma TG and HDL

cholesterol levels are commonly tested, the same cannot be said for the

LDL particle diameter or postprandial remnant concentrations.

Nonetheless, a previous study observed a robust correlation between the

TG/HDL cholesterol ratio and the LDL peak diameter.(15) Moreover, it is

suggested that a TG/HDL cholesterol concentration ratio of 3.5 can

accurately predict the presence of the small dense LDL phenotype (LDL

phenotype B). Therefore, by assessing plasma TG and HDL cholesterol

levels and computing their ratio, it is possible to obtain understanding

into three out of the four modifications in lipoprotein metabolism that

enhance the risk of cardiovascular disease in insulin-resistant patients.

Numerous epidemiological studies have examined the relationship

between the TG/HDL ratio and CVD risk. Most of these studies have

reported a positive association between a higher TG/HDL ratio and

increased CVD risk. For instance, a prospective cohort study by Gaziano

et al. found the ratio of triglycerides to HDL was a strong predictor of

myocardial infarction;(16) Urbina et al. suggests the TG/HDL ratio may

be a useful marker for identifying young individuals at risk for

developing arterial stiffness, an early sign of CVD;(17) Li et al. found a

positive correlation between the TG/HDL-C ratio and carotid intima

thickness, a marker of early atherosclerosis, in adolescents with type 2

diabetes;(18) Du et al. found that the TG/HDL ratio is associated with

insulin resistance, a major risk factor for CVD, in a Chinese

population;(19) Ren et al. found that the TG/HDL ratio is positively

associated with insulin resistance in newly diagnosed type 2 diabetes

patients, indicating an increased risk of CVD;(20) Squillace et al. found

that a higher TG/HDL ratio is predictive of the development of diabetes, a

major risk factor for CVD;(21) Hong et al. found that a high TG/HDL

ratio is associated with an increased risk of cardiovascular events in

patients with diabetes and coronary artery disease.(22)

    The TG/HDL ratio holds clinical significance as a potential marker

for CVD prevention. It provides valuable insights into the balance

between TG and HDL levels, representing the interplay between

dyslipidemia and atherosclerosis. A higher TG/HDL ratio reflects an

increased atherogenic lipid profile and has been associated with arterial

stiffness, carotid intima thickness, insulin resistance and diabetes, all of

which contribute to CVD development. By incorporating the TG/HDL

ratio into risk assessment models, clinicians can more accurately identify

individuals at high risk for CVD and implement targeted interventions.

Despite the clinical significance of the TG/HDL ratio, several

limitations must be acknowledged. First, the cut-off values for defining

high TG/HDL ratios vary across studies, leading to heterogeneity in

research findings. Standardization of these thresholds is necessary to

ensure consistency and comparability between studies. Additionally,

confounding factors, including age, sex, lifestyle factors, and

comorbidities, may influence the association between the TG/HDL ratio

and CVD risk. Although some studies have attempted to account for these

factors, further research is needed to elucidate their precise impact.

Moreover, variations in laboratory methods for lipid profile assessment

may introduce measurement errors and affect the accuracy of TG and

HDL cholesterol estimations.

In conclusion, the TG/HDL ratio demonstrates potential as a

valuable clinical tool for CVD prevention. Its ability to capture the

interplay between triglycerides and HDL cholesterol provides a

comprehensive assessment of the atherogenic lipid profile. By

incorporating the TG/HDL ratio into risk assessment models, healthcare

professionals can enhance risk stratification and identify high-risk

individuals who may benefit from targeted interventions. However,

standardization of cut-off values, better control of confounding factors,

and harmonization of laboratory methods are needed to further validate

the clinical significance of the TG/HDL ratio. With these advancements,

the TG/HDL ratio could significantly contribute to the prevention and

control of cardiovascular disease.

Availability of data and materials

The data that support the findings of this study are available from the

corresponding author Haoran Wang, upon reasonable request.

Declaration of Interest statement

The authors declared that there is no conflict of interest.

Author Contributions statement

Conceptualization and Methodology: Haoran Wang, Pengcheng Li;

Formal analysis: Pengcheng Li, Jirui Cai, Shuaifang Yuan, Yapeng Li,

Huiting Shi, Cui Liang, Bing He; Data Curation: Qiaotao Xie, Baocang

Lei,Jing Bai, Nan Wang, Dongliang Liu, Qichao Wang, Jianwei Xiong,

Jin Wang, Haoran Wang; Writing: Pengcheng Li, Jirui Cai, Shuaifang Yuan, Yapeng Li, Huiting Shi, Cui Liang; Funding acquisition: Qiaotao Xie, Haoran Wang, Pengcheng Li.

Fundings

The study is supported by grants from the National Natural Science Foundation of China (81803318, 82304229), Henan Provincial Science and Technology Research Project (232300420069, 232102310231, 232300420289), and the Henan Ministry of Education (22B320003). The funders have no role in the design and implementation of the current study.

Ethical statement

This study was conducted with approval from the Ethics Committee of Fuwai Hospital (2014–574) as the Central Ethic, and was approved by the Ethics Committee of Luohe Central Hospital as the local ethic. This study was conducted in accordance with the declaration of Helsinki. Written informed consent was obtained from all participants.

References

1. Roth GA, Mensah GA, Johnson CO, Addolorato G, Ammirati E, Baddour LM, Barengo NC, Beaton AZ, Benjamin EJ, Benziger CP, et al. Global Burden of Cardiovascular Diseases and Risk Factors, 1990-2019: Update From the GBD 2019 Study. *J Am Coll Cardiol* (2020) 76:2982–3021. doi: 10.1016/j.jacc.2020.11.010

2. Zhao D, Liu J, Xie W, Qi Y. Cardiovascular risk assessment: a global perspective. *Nat Rev Cardiol* (2015) 12:301–311. doi: 10.1038/nrcardio.2015.28

3. Thupakula S, Nimmala SSR, Ravula H, Chekuri S, Padiya R. Emerging biomarkers for the detection of cardiovascular diseases. *Egypt Heart J* (2022) 74:77. doi: 10.1186/s43044-022-00317-2

4. Yang M, Rigdon J, Tsai SA. Association of triglyceride to HDL cholesterol ratio with cardiometabolic outcomes. *J Investig Med Off Publ Am Fed Clin Res* (2019) 67:663–668. doi: 10.1136/jim-2018-000869

5. Kosmas CE, Rodriguez Polanco S, Bousvarou MD, Papakonstantinou EJ, Peña Genao E, Guzman E, Kostara CE. The Triglyceride/High-Density Lipoprotein Cholesterol (TG/HDL-C) Ratio as a Risk Marker for Metabolic Syndrome and Cardiovascular Disease. *Diagnostics* (2023) 13:929. doi: 10.3390/diagnostics13050929

6. Che B, Zhong C, Zhang R, Pu L, Zhao T, Zhang Y, Han L. Triglyceride-glucose index and triglyceride to high-density lipoprotein cholesterol ratio as potential cardiovascular disease risk factors: an analysis of UK biobank data. *Cardiovasc Diabetol* (2023) 22:34. doi: 10.1186/s12933-023-01762-2

7. Chen C, Dai J-L. Triglyceride to high-density lipoprotein cholesterol (HDL-C) ratio and arterial stiffness in Japanese population: a secondary analysis based on a cross-sectional study. *Lipids Health Dis* (2018) 17:130. doi: 10.1186/s12944-018-0776-7

8. Chen Y, Chang Z, Liu Y, Zhao Y, Fu J, Zhang Y, Liu Y, Fan Z. Triglyceride to high-density lipoprotein cholesterol ratio and cardiovascular events in the general population: A systematic review and meta-analysis of cohort studies. *Nutr Metab Cardiovasc Dis NMCD* (2022) 32:318–329. doi: 10.1016/j.numecd.2021.11.005

9. Chen Z, Chen G, Qin H, Cai Z, Huang J, Chen H, Wu W, Chen Z, Wu S, Chen Y. Higher triglyceride to high-density lipoprotein cholesterol ratio increases cardiovascular risk: 10-year prospective study in a cohort of Chinese adults. *J Diabetes Investig* (2020) 11:475–481. doi: 10.1111/jdi.13118

10. Wang R, Yang Y, Lu J, Cui J, Xu W, Song L, Wu C, Zhang X, Dai H, Zhong H, et al. Cohort Profile: ChinaHEART (Health Evaluation And risk Reduction through nationwide Teamwork) Cohort. *Int J Epidemiol* (2023) 52:e273–e282. doi: 10.1093/ije/dyad074

11. Lu J, Xuan S, Downing NS, Wu C, Li L, Krumholz HM, Jiang L. Protocol for the China PEACE (Patient-centered Evaluative Assessment of Cardiac Events) Million Persons Project pilot. *BMJ Open* (2016) 6:e010200. doi: 10.1136/bmjopen-2015-010200

12. WHO CVD Risk Chart Working Group. World Health Organization cardiovascular disease risk charts: revised models to estimate risk in 21 global regions. *Lancet Glob Health* (2019) 7:e1332–e1345. doi: 10.1016/S2214-109X(19)30318-3

13. Reaven GM. The insulin resistance syndrome. *Curr Atheroscler Rep* (2003) 5:364–371. doi: 10.1007/s11883-003-0007-0

14. Lorenzatti AJ, Toth PP. New Perspectives on Atherogenic Dyslipidaemia and Cardiovascular Disease. *Eur Cardiol* (2020) 15:1–9. doi: 10.15420/ecr.2019.06

15. Dobiásová M, Frohlich J. The plasma parameter log (TG/HDL-C) as an atherogenic index: correlation with lipoprotein particle size and esterification rate in apoB-lipoprotein-depleted

plasma (FER(HDL)). *Clin Biochem* (2001) 34:583–588. doi:
10.1016/s0009-9120(01)00263-6

16.   Gaziano JM, Hennekens CH, O'Donnell CJ, Breslow JL,
Buring JE. Fasting triglycerides, high-density lipoprotein, and risk of
myocardial infarction. *Circulation* (1997) 96:2520–2525. doi:
10.1161/01.cir.96.8.2520

17.   Urbina EM, Khoury PR, McCoy CE, Dolan LM, Daniels
SR, Kimball TR. Triglyceride to HDL-C ratio and increased arterial
stiffness in children, adolescents, and young adults. *Pediatrics* (2013)
131:e1082-1090. doi: 10.1542/peds.2012-1726

18.   Li X, Deng Y-P, Yang M, Wu Y-W, Sun S-X, Sun J-Z.
Triglyceride to high-density lipoprotein cholesterol ratio and carotid
intima-medial thickness in Chinese adolescents with newly
diagnosed type 2 diabetes mellitus. *Pediatr Diabetes* (2016)
17:87–92. doi: 10.1111/pedi.12250

19.   Du T, Yuan G, Zhang M, Zhou X, Sun X, Yu X. Clinical
usefulness of lipid ratios, visceral adiposity indicators, and the
triglycerides and glucose index as risk markers of insulin resistance.
Cardiovasc Diabetol (2014) 13:146. doi:
10.1186/s12933-014-0146-3

20.   Ren X, Chen ZA, Zheng S, Han T, Li Y, Liu W, Hu Y.
Association between Triglyceride to HDL-C Ratio (TG/HDL-C) and
Insulin Resistance in Chinese Patients with Newly Diagnosed Type 2
Diabetes Mellitus. *PloS One* (2016) 11:e0154345. doi:
10.1371/journal.pone.0154345

21.   Squillace N, Lorenzini P, Lapadula G, Bandera A,
Cozzi-Lepri A, Rusconi S, Puoti M, Castagna A, Antinori A, Gori A,
et al. Triglyceride/HDL ratio and its impact on the risk of diabetes
mellitus development during ART. *J Antimicrob Chemother* (2016)
71:2663–2669. doi: 10.1093/jac/dkw185

22.   Yang S-H, Du Y, Li X-L, Zhang Y, Li S, Xu R-X, Zhu
C-G, Guo Y-L, Wu N-Q, Qing P, et al. Triglyceride to High-Density

394   Lipoprotein Cholesterol Ratio and Cardiovascular Events in

395   Diabetics With Coronary Artery Disease. *Am J Med Sci* (2017)

396   354:117–124. doi: 10.1016/j.amjms.2017.03.032

Tables

Table 1. Clinical characteristics of the study population by quartiles of TG/HDL-C.

	Q1	Q2	Q3	Q4	p-value
Number	N=1,648	N=1,648	N=1,648	N=1,649	
Age (median (IQR))	59.00 (50.00-67.00)	58.00 (50.00-67.00)	58.00 (51.00-66.00)	58.00 (52.00-66.00)	0.78
Male (%)	617 (37%)	613 (37%)	599 (36%)	667 (40%)	0.081
BMI (median (IQR))	24.12 (22.14-26.32)	24.80 (22.70-26.83)	25.15 (23.16-27.44)	26.04 (24.01-28.36)	<0.001
Waist (median (IQR))	83.00 (77.00-89.95)	85.00 (79.00-91.00)	86.00 (80.50-93.00)	89.00 (82.00-95.00)	<0.001
SBP (median (IQR))	135.00 (123.25-148.50)	136.00 (124.50-149.50)	136.00 (125.50-149.50)	141.50 (129.50-152.00)	<0.001
DBP (median (IQR))	82.50 (76.00-89.50)	82.50 (76.00-89.50)	82.50 (76.00-90.00)	85.00 (77.00-92.50)	<0.001
Current smoke (%)	338 (21%)	325 (20%)	300 (18%)	358 (22%)	0.083
Drinking (%)	86 (5%)	68 (4%)	98 (6%)	107 (6%)	0.018
HTN (%)	335 (20%)	338 (21%)	367 (22%)	467 (28%)	<0.001
DM (%)	72 (4%)	96 (6%)	91 (6%)	147 (9%)	<0.001
CHD (%)	36 (2%)	25 (2%)	19 (1%)	13 (1%)	0.006
Stroke (%)	63 (4%)	37 (2%)	41 (2%)	40 (2%)	0.021
TC (median (IQR))	4.08 (3.47-4.65)	4.53 (4.03-5.11)	4.97 (4.36-5.43)	5.62 (4.98-6.39)	<0.001
HDL (median (IQR))	1.71 (1.49-1.97)	1.50 (1.33-1.69)	1.39 (1.23-1.52)	1.21 (1.09-1.37)	<0.001
TG (median (IQR))	1.21 (0.95-1.62)	1.41 (1.11-1.74)	1.58 (1.25-2.00)	1.94 (1.47-2.69)	<0.001
LDL (median (IQR))	1.96 (1.45-2.44)	2.48 (2.09-2.88)	2.71 (2.32-3.17)	3.35 (2.71-4.03)	<0.001
Glu (median (IQR))	5.40 (5.10-5.80)	5.40 (5.10-5.90)	5.40 (5.11-6.00)	5.70 (5.30-6.40)	<0.001
TG/HDL-C (median (IQR))	2.43 (2.20-2.59)	3.02 (2.88-3.15)	3.55 (3.41-3.72)	4.50 (4.17-4.97)	<0.001
CVD_highrisk (%)	257 (16%)	251 (15%)	290 (18%)	620 (38%)	<0.001

Data are presented as median (IQR) for continuous variables, and number (percentage) for categorical variables. HTN, hypertension; DM, diabetes mellitus; CHD, coronary heart disease; BMI, body mass index; TC, total cholesterol; TG, triglyceride; LDL, low density lipoprotein; HDL, high density lipoprotein; SBP, systolic blood pressure; DBP, diastolic blood pressure.

Table 2. Association between TG/HDL-C ratio and cardiovascular high-risk by logistic regression.

	Model 1		Model 2	
	OR (95% CI)	p	OR (95% CI)	p
All participants				
TG/HDL-C (per 1 unit)	1.40 (1.30-1.51)	<0.001	1.21 (1.11-1.32)	<0.001
TG/HDL-C (high vs low)	1.61 (1.43-1.81)	<0.001	1.27 (1.10-1.47)	0.001
TG/HDL-C (quartile)				
Q1	Ref		Ref	
Q2	0.90 (0.75-1.07)	0.214	1.04 (0.85-1.29)	0.68
Q3	0.84 (0.71-1.01)	0.06	0.95 (0.77-1.18)	0.634
Q4	1.79 (1.52-2.10)	<0.001	1.49 (1.21-1.83)	<0.001
Female				
TG/HDL-C (per 1 unit)	1.36 (1.24-1.50)	<0.001	1.14 (1.01-1.28)	0.028
TG/HDL-C (high vs low)	1.52 (1.30-1.77)	<0.001	1.16 (0.96-1.40)	0.114
TG/HDL-C (quartile)				
Q1	Ref		Ref	
Q2	0.94 (0.74-1.19)	0.6	0.97 (0.74-1.28)	0.838
Q3	0.91 (0.72-1.15)	0.447	1.00 (0.76-1.32)	0.998
Q4	1.78 (1.44--2.20)	<0.001	1.33 (1.02-1.74)	0.034
Male				
TG/HDL-C (per 1 unit)	1.50 (1.33-1.69)	<0.001	1.31 (1.14-1.52)	<0.001
TG/HDL-C (high vs low)	1.82 (1.51-2.19)	<0.001	1.45 (1.15-1.84)	<0.001
TG/HDL-C (quartile)				
Q1	Ref		Ref	
Q2	0.85 (0.66-1.11)	0.238	1.14 (0.84-1.57)	0.432
Q3	0.78 (0.59-1.03)	0.076	0.90 (0.64-1.27)	0.557
Q4	1.94 (1.51-2.49)	<0.001	1.67 (1.20-2.32)	0.002

Figure legends

Figure 1. Percentage stacked histogram to visually represent the distribution of cardiovascular high-risk individuals across quartiles of the TG/HDL-C ratio.

Figure 2. Distribution histogram of the TG/HDL-C ratio in the study population and restricted cubic spline to assess the relationship between TG/HDL-C ratio and cardiovascular high-risk.

Figure 3. ROC curve to illustrate the discriminatory power of TG/HDL-C ratio for cardiovascular high-risk.

Figure 1. Percentage stacked histogram to illustrate the distribution of cardiovascular high-risk according to quartiles of TG/HDL-C ratio.

Figure 2. Distribution histogram of the TG/HDL-C ratio in the study population and restricted cubic spline to assess the relationship between TG/HDL-C ratio and cardiovascular high-risk.

Figure 3. ROC curve to illustrate the discriminatory power of TG/HDL-C for cardiovascular high-risk.