

1 **Clinical characteristics and valve lesions in rheumatic heart disease among children admitted to a selected**  
2 **tertiary teaching hospital in eastern Ethiopia.**

3 **Authors**

4 **Temesgen Teferi Libe<sup>1\*</sup>, Yunus Edris Kelil<sup>1</sup>, Samrawit Abebaw Tegene<sup>2</sup>, Faisal Abdi<sup>2</sup>, Kibrom**

5 **Hailemariam Mesfin<sup>3</sup>**

6 **Affiliations:**

7 <sup>1</sup> Department of Pediatrics and Child Health, College of Health and Medical Sciences, Haramaya University,  
8 Harar, Ethiopia.

9 <sup>2</sup>Hararghe Health Research, Haramaya University, Harar, Ethiopia.

0 <sup>3</sup>Maedot Charity Organization, Mekelle, Ethiopia

1 **\*Corresponding author:** Temesgen Teferi Libe (MD), School of Medicine, College of Health and Medical  
2 Sciences, Haramaya University Harar, Ethiopia, Email: [libe.temesgen.temesgen@gmail.com](mailto:libe.temesgen.temesgen@gmail.com), Tel:  
3 +251913730243, P.O Box 235 Harar, Ethiopia.

## 4 Abstract

5 **Background:** Rheumatic heart disease remains a significant public health concern, especially among children in  
6 resource-limited settings like Ethiopia. Despite effective prevention strategies, RHD persists due to factors such  
7 as poverty and limited healthcare access. Understanding the clinical characteristics and valve lesions of  
8 rheumatic heart disease is crucial for improving diagnosis and management.

9 **Objective:** This study aimed to characterize the clinical features and valve lesions in children with rheumatic  
0 heart disease admitted to a tertiary teaching hospital in Eastern Ethiopia.

1 **Methods:** A hospital-based cross-sectional study was conducted from September 1, 2020, to August 31, 2021, at  
2 Hiwot Fana Comprehensive Specialized Hospital. Data were collected from medical records, clinical  
3 assessments, echocardiography reports, and laboratory tests. Descriptive statistics were used for analysis.

4 **Results:** A total of 39 children with rheumatic heart disease were included, predominantly females (71.8%).  
5 Shortness of breath (53.9%) and cough (38.5%) were common presenting symptoms. Acute decompensated  
6 heart failure (ADHF) was prevalent (89.7%). Mitral regurgitation (94.9%) and aortic regurgitation (66.7%) were  
7 frequent valve lesions. Other findings included mitral stenosis (56.4%) and left atrial enlargement (86.1%).  
8 Laboratory results showed mean hemoglobin of 10.29 g/dL and mean ESR of 45 mm/hr.

9 **Conclusions:** This study highlights the burden of RHD in Eastern Ethiopia, with advanced disease at  
0 presentation. Mitral and aortic valve lesions were predominant, emphasizing the need for early detection and  
1 comprehensive management strategies. Collaboration among healthcare providers and policymakers is essential  
2 to address the challenges of RHD in resource-limited settings.

3 **Keywords:** Rheumatic heart disease, clinical characteristics, valve lesions, Eastern Ethiopia

## 34 **Introduction**

## 35 **Background**

36 Rheumatic heart disease (RHD) remains a significant public health threat, particularly for  
37 children in developing countries like Ethiopia. It is a debilitating condition primarily affecting  
38 the heart valves, particularly the mitral and aortic valves, resulting from acute rheumatic fever  
39 (ARF), a complication of untreated streptococcal pharyngitis(1). The global burden of RHD is  
40 immense, with developing nations disproportionately affected due to factors like poverty,  
41 overcrowding, and limited access to healthcare(2, 3, 4, 5, 6, 7, 8, 9).

42 RHD is a significant contributor to the overall burden of acquired heart diseases worldwide,  
43 accounting for nearly half of all cardiovascular problems(10). In 2015 alone, it was responsible  
44 for over 10 million disability-adjusted life-years and over 300,000 premature deaths(2). Despite  
45 the existence of effective prevention strategies, RHD continues to disproportionately affect  
46 developing countries. Here, untreated ARF often progresses to RHD(3, 4, 5, 6, 7, 8, 9, 10).

47 Patients with RHD often present with symptoms of heart failure, including shortness of breath,  
48 fatigue, and chest pain. In severe cases, RHD can lead to complications such as pulmonary  
49 arterial hypertension, arrhythmias, and infective endocarditis(11, 12). Echocardiography plays a  
50 crucial role in diagnosing RHD, determining the location, type, and severity of valve lesions, and  
51 identifying associated complications(11, 12, 13, 14).

52 Despite the substantial national burden of RHD, there is a scarcity of studies exploring the  
53 clinical characteristics and patterns of valve lesions in Ethiopia, particularly in the Eastern  
54 region. To our knowledge, no comprehensive research has been conducted in this specific

55 geographic area regarding valve lesion patterns and clinical characteristics. This study aims to  
56 address this critical knowledge gap by investigating RHD in Ethiopian children.

57 The lack of understanding concerning the specific clinical features and valve lesion patterns in  
58 this population hinders the development of tailored interventions, early detection, and optimal  
59 management of RHD. Healthcare providers, policymakers, and public health practitioners  
60 require a clearer picture to implement effective strategies for reducing the disease burden,  
61 associated morbidity, and mortality.

62 Therefore, this study aims to characterize the clinical features and valve lesion patterns in  
63 children with RHD. This knowledge will ultimately inform the development of evidence-based  
64 interventions and improved healthcare for this vulnerable population.

## 65 **Objectives**

### 66 **General Objective**

67 To determine the clinical characteristics and valve lesions in children with rheumatic heart  
68 disease admitted to a tertiary teaching hospital in eastern Ethiopia, from September 1, 2020, to  
69 August 31, 2021.

### 70 **Specific objectives**

71 To describe the clinical characteristics of children admitted to a tertiary teaching hospital in  
72 eastern Ethiopia with a diagnosis of rheumatic heart disease.

73 To evaluate the patterns of valve lesions in children admitted to a tertiary teaching hospital in  
74 eastern Ethiopia with a diagnosis of rheumatic heart disease.

## 75 **Methods and Materials**

### 76 **Study area, design and period**

77 A hospital-based cross-sectional study design was used to prospectively study children admitted  
78 from September 01, 2020 to August 31, 2021 to pediatric ward and pediatrics ICU of Hiwot Fana  
79 Comprehensive Specialized Hospital (HFCSH). Hiwot Fana Comprehensive Specialized  
80 Hospital (HFCSH), Haramaya University's tertiary teaching hospital, is located in Harar and  
81 serves 5.8 million people, including the populations of Harari Region, Diredawa City  
82 Administration, Somali Region, and Eastern Hararghe Zone of Oromia Regional State. At  
83 HFCSH, Pediatric Ward and ICU accommodate 30 beds (6 in PICU and 24 in the Pediatric  
84 ward) with about 1,110 total admissions per year, among which about 79 had RHD diagnosis.  
85 (Health Management Information System data 2019; unpublished).

### 86 **Patient involvement**

87 Patients were not directly involved in the design of this study.

### 88 **Population and eligibility**

89 All children with RHD admitted to the pediatric ward and pediatric ICU of HFCSH from  
90 September 01, 2020 to August 31, 2021, by excluding repeat admissions and disappeared cases  
91 before obtaining adequate information.

### 92 **Sample size determination and sampling technique**

93 Study population, and Sample size was calculated Using OpenEpi software by taking P=50%),  
94 with a 95% confidence level and the margin of error d=5% (0.05). The sample size was  
95 determined to be 384. However, over the study period of one year, there were about 67 children

96 with RHD admitted to pediatric ward and ICU. Thus, we decided to enroll all of these children to  
97 have more representative data. [Fig 1]

98 Fig 1. Sampling procedure for the study Clinical characteristics and valve lesions in rheumatic  
99 heart disease.

## 100 **Data Collection Methods**

101 A comprehensive review of medical records for children with RHD admitted to the pediatric  
102 ward and ICU at Hiwot Fana Comprehensive Tertiary Teaching Hospital was conducted. This  
103 review included demographic information, clinical history, presenting symptoms, laboratory test  
104 results, chest X-ray findings, and echocardiography reports. A team of experienced pediatricians  
105 and cardiologists conducted standardized clinical assessments on each enrolled child, including a  
106 thorough physical examination, for cardiac symptom evaluation, vital sign measurement, and  
107 collection of additional clinical data. Pediatric cardiologists performed echocardiographic  
108 assessments adhering to standardized protocols to identify valve lesions and their characteristics.  
109 Additionally, laboratory data (complete blood count, erythrocyte sedimentation rate [ESR]),  
110 chest X-ray, and ECG were obtained.

## 111 **Study Variables and measurements**

112 The outcome variables of this study were patterns of valve lesions and clinical characteristics of  
113 RHD. Independent variables included; socio-demographic variables; age, sex, address, ethnicity.

## 114 **Data quality control**

115 The data collection tool was piloted on two RHD patients in the Jugol Hospital pediatric ward to  
116 assess its consistency, completeness, and ease of understanding. Two days of training were given  
117 to both the data collectors and supervisor. The principal investigator closely monitored the data

118 collection process to ensure completeness, accuracy, and data collection consistency. During  
119 sessions, thorough checking was done before analyzing the filled questionnaires. Two separate  
120 individuals conducted double data entry. When any gap was identified, it was communicated  
121 with data collectors daily. Following data entry, the dataset was reviewed for completeness and  
122 any missing data points were addressed.

### 123 **Data Processing and Analysis**

124 After data collection, each questionnaire was checked thoroughly for completeness. The data was  
125 then coded using a standardized coding scheme and double-entered into EpiData version 3.1.  
126 Subsequently, the data was transferred to Stata version 14.2 for statistical analysis.  
127 Descriptive statistics, including frequency, proportions, means, and standard deviation were  
128 computed.

### 129 **Ethical consideration**

130 Ethical clearance was obtained from Haramaya University College of Health and Medical  
131 Sciences Institution Health Research Ethical Review Committee (with ethical approval ref.no:  
132 IHRERC/245/2020). An official letter was sent to Hiwot Fana Comprehensive Specialized  
133 Hospital. This study was conducted in accordance with the Declaration of Helsinki. Data  
134 collection was started after the study protocol participants (mothers of children) under the age of  
135 18 years were reviewed and approved by the IRB of Haramaya University to provide informed  
136 consent on their own behalf.

137 Permissions were obtained from hospital managers/medical directors and after written or (verbal  
138 for those unable to read and write) informed consent was obtained from mothers of the baby. The  
139 purpose of the study was briefly explained to the children's mothers/care givers. Data that were

140 collected from the participants were kept in secured and order to maintain confidentiality.

141 Voluntary participation was ensured and participants were free to out.

## 142 **Result**

### 143 **Baseline Characteristics**

144 A total of 39 children with RHD were enrolled in the study. The majority (71.8%, n=28) were  
145 females, and the remaining participants (28.2%, n=11) were males. The mean age of the children  
146 was 10 years (SD  $\pm$  2.6 years). [Table 1]

147 Table 1 Baseline

Sex	Frequency	Percentage	Age in years	148
Male	11	28.21%	Mean	10 149
Female	28	71.79%	SD	2.6
Total	39	100%	(Min, Max)	(5, 14) <sup>150</sup>

Characteristics of  
RHD patients

151

152

### 153 **Clinical characteristics**

154 Fourteen children had a known diagnosis of RHD, and 25 were newly diagnosed. Shortness of  
155 breath (53.9%) was the most frequent presenting complaint, followed by cough (38.5%). Among  
156 the 14 children with a known diagnosis of RHD, only 2 (14.3%) were adherent to secondary  
157 prophylaxis, while 6 (42.9%) were non-adherent and the remaining 6 (42.9%) had never taken  
158 prophylaxis. Shortness of breath (53.9%) was the most frequent presenting complaint, followed  
159 by cough (38.5%) and other symptoms. [Table 2]



160 Table 2 clinical characteristics of RHD patients

<b>Chief Complaint</b>	Freq	Percent	<b>RHD status</b>	Freq	Percent	<b>Adherence to prophylaxis</b>	Freq	Percent
Cough	15	38.46	New	25	64.1	Adherent	2	14.29
Shortness of breath	21	53.85	Known	14	35.9	Not Adherent	6	42.86
Other	3	7.69				Not taking prophylaxis	6	42.86
Total	39	100	Total	39	100	Total	14	100

161

162 Ten (25.6%) children had at least one comorbidity. The most frequent comorbidities were severe  
 163 acute malnutrition (5; 12.8%) and severe anemia (Hb < 7 gm/dl; 5; 12.8%).

164 A significant proportion of patients (35 out of 39, or 89.7%) presented with acute decompensated  
 165 heart failure (ADHF) upon admission.

166 The most frequent identified trigger was pneumonia, affecting 21 patients (60% of ADHF  
 167 patients). One patient (3% of ADHF patients) had atrial fibrillation as a possible contributing  
 168 factor. Four patients (11% of ADHF patients) experienced a recurrence of rheumatic disease,  
 169 potentially contributing to their heart failure. Five patients (14% of ADHF patients) had  
 170 confirmed or suspected infective endocarditis, which could have exacerbated their heart failure.

## 171 **Vital signs and Anthropometry**

172 The mean pulse rate was 115.25 beats per minute (bpm) with a standard deviation (SD) of 19.44  
 173 bpm (range: 60-178 bpm). The mean respiratory rate was 37.46 breaths per minute (bpm) with  
 174 an SD of 13.49 bpm (range: 18-72 bpm). The mean body temperature was 36.94°C (SD ±  
 175 1.05°C; range: 32-38.7°C). Mean oxygen saturation on room air was 94.3% (SD ± 3.5%; range:

176 86-99%). The mean body mass index (BMI) was 14.56 kg/m<sup>2</sup> (SD ± 3.1 kg/m<sup>2</sup>; range: 11.3-28.2  
 177 kg/m<sup>2</sup>). [Table 3]

178 Table 3

Variable	Mean	SD	Min	Max
Pulse rate	115.25	19.44	60	178
Respiratory Rate	37.46	13.49	18	72
Temperature	36.94	1.05	32	38.7
Saturation	94.3	3.5	86	99
BMI	14.56	3.1	11.3	28.2

179 Vital sign  
 180 and BMI  
 181 distribution  
 182 of RHD  
 183 patients

184  
 185  
 186  
 187  
 188 According to WHO Z-scores for age, 5 (12.8%) children were severely wasted (< -3 Z-score), 12  
 189 (30.8%) were moderately wasted (-3 to -2 Z-score), 21 (53.8%) had normal weight (-2 to +2 Z-  
 190 score), and 1 (2.6%) was overweight (> +2 Z-score). [Fig 2]

191 Fig 2. BMI for age distribution of RHD patients

## 192 Chest examination

193 At least one abnormal chest examination finding was present in 24 (61.5%) children. The most  
 194 common findings were crepitations (41.0%), decreased or absent air entry (39.5%), dullness  
 195 (39.5%), and wheezing (9.3%). [Table 4]

196 Table 4 Chest examination findings of RHD patients

<b>Chest findings (N=39)</b>	<b>At least one finding</b>	<b>Crepitation</b>	<b>Decreased or absent air entry</b>	<b>Dullness</b>	<b>Wheezing</b>
Frequency	24	16	17	17	4
Percent	61.54	41.03	39.53	39.53	9.3

197

## 198 **Cardiac examination**

199 The point of maximal impulse (PMI) was displaced downward and to the left in 28 (71.8%)  
 200 children, while it was in the normal position in 9 (23.1%) and on the left lower sternal border in  
 201 2 (5.1%). Apical heave (61.5%) and parasternal heave (56.4%) were the most frequent abnormal  
 202 cardiac examination findings. P2 heart sound was accentuated in 28 (77.8%) children. A third  
 203 heart sound (S3 gallop) was present in 5 (12.8%) children. Almost all children (97.4%) had a  
 204 significant murmur on cardiac auscultation. [Table 5]

205 Table 4 Cardiac examination findings of RHD patients

		<b>Freq.</b>	<b>Percent</b>
<b>Point of Maximum impulse</b>	Normal position	9	23.08
	On left lower sternal	2	5.13
	Deviated to left and/or downward	28	71.79
	Total	39	100
<b>Apical Heave</b>	Yes	24	61.54
	No	15	38.46
	Total	39	100
<b>Parasternal heave</b>	Yes	22	56.41

	No	17	43.59
	Total	39	100
<b>S3 gallop</b>	Yes	5	12.82
	No	34	87.18
	Total	39	100

206

### 207 **Abdomen and integumentary examination**

208 Hepatomegaly was present in 17 (43.6%) children, while ascites (fluid collection in the  
209 abdomen) was detected in 14 (35.9%). Peripheral edema was observed in 8 (20.5%).

### 210 **Laboratory findings**

211 Laboratory analysis revealed the following: mean hemoglobin (Hb) concentration was 10.29  
212 g/dL (SD  $\pm$  2.67 g/dL; range, 3.9-17.6 g/dL), mean erythrocyte sedimentation rate (ESR) was 45  
213 mm/hr (SD  $\pm$  33.58 mm/hr; range, 1-140 mm/hr), and mean serum creatinine level was 0.53  
214 mg/dL (SD  $\pm$  0.265 mg/dL; range, 0.1-1.1 mg/dL).

### 215 **Chest x ray findings**

216 Chest X-ray revealed abnormal findings in a significant proportion of participants. Left atrial  
217 enlargement was the most frequent abnormality, observed in 86.1% (31/36) of patients. Other  
218 findings included cardiomegaly (80.6%), pulmonary edema (26.5%), signs suggestive of  
219 pulmonary arterial hypertension (PAH; 36.7%), and pleural effusion (24.2%). [Table 6]

220 Table 6 Chest X-Ray Findings of RHD patients

Chest X-Ray Finding	Frequency (%)	Number of Patients (n=36)
---------------------	---------------	---------------------------

Left Atrial Enlargement (LAE)	86.1	31
Cardiomegaly	80.6	29
Pulmonary Edema (PE)	26.5	9
Signs suggestive of Pulmonary Arterial Hypertension (PAH)	36.7	11
Pleural Effusion	24.2	8
Pneumonia	21.2	7

221

222 **Echocardiographic Findings in RHD Patients**

223 Echocardiography revealed significant valvular involvement in a vast majority of patients with  
 224 RHD. All 39 patients had evidence of mitral valve involvement. The most frequent finding was  
 225 mitral regurgitation (MR), present in 37 patients (94.9%). The severity of MR varied: 3 patients  
 226 (7.7%) had mild MR, 7 patients (17.9%) had moderate MR, and 27 patients (69.2%) had severe  
 227 MR. Mitral stenosis (MS) was identified in a subset of patients (22/39, 56.4%). The severity of  
 228 MS also varied: 9 patients (23.1%) had mild MS, 6 patients (15.4%) had moderate MS, and 7  
 229 patients (17.9%) had severe MS. Notably, 20 patients (51.3%) had both MR and MS present  
 230 concurrently. However, it's important to note that 2 patients (5.1%) had isolated MS without MR.

231 [Table 7]

232 Table 7 Mitral Valve Findings (n=39)

Feature	Frequency	Percent (%)
<b>Mitral Regurgitation (MR)</b>		
No	2	5.13
Mild	3	7.69

Moderate	7	17.95
Severe	27	69.23
<b>Mitral Stenosis (MS)</b>		
No	17	43.59
Mild	9	23.08
Moderate	6	15.38
Severe	7	17.95
<b>Combined MR &amp; MS</b>	20	51.3

233

234 A substantial portion of patients (26 out of 39, or 66.7%) had aortic valve involvement. Similar  
 235 to the mitral valve, the severity of aortic regurgitation (AR) varied. Fourteen patients (35.9%)  
 236 had mild AR, 11 patients (28.2%) had moderate AR, and only 1 patient (2.6%) had severe AR.  
 237 Interestingly, only 1 patient (2.6%) had aortic stenosis (AS), and it was severe. It's important to  
 238 note that this severe AS co-existed with MR, highlighting a potential link between these valve  
 239 dysfunctions in some cases. [Table 8]

240 Table 8 Aortic Valve Findings (n=39)

Feature	Frequency	Percent (%)
<b>Aortic Regurgitation (AR)</b>		
No	13	33.33
Mild	14	35.9
Moderate	11	28.21
Severe	1	2.56
<b>Aortic Stenosis (AS)</b>		

No	38	97.44
Severe	1	2.56

241

242 Pulmonic valve involvement was less frequent compared to the mitral and aortic valves. Only 7  
243 patients (17.9%) had pulmonic regurgitation, and no patients had pulmonic stenosis.

244 The pressure gradient measured across the tricuspid valve averaged 27.17 mmHg, with some  
245 variation (standard deviation of 19.734 mmHg).

246 Right atrial size varied among patients. Normal size was observed in nearly 40% (15 patients),  
247 while mild, moderate, and severe dilation were identified in 21%, 37%, and 2.6% of patients,  
248 respectively. Similarly, right ventricular size varied, with normal dimensions in almost half  
249 (47%) of patients. Mild and moderate dilation were seen in 18% and 34% of patients,  
250 respectively. Data for one patient's right ventricle was missing.

251 Left atrial size also showed variation. Most patients had either moderate (47%) or severe (45%)  
252 dilation. Only a small portion (8%) had mildly dilated left atria, and normal size was uncommon  
253 (observed in just 4 patients). Left ventricular size followed a similar pattern, with the majority of  
254 patients having moderate (34%) or severe (39%) dilation. Normal size was seen in less than 20%  
255 of patients, and mild dilation was identified in another 8%.

256 One important finding was the presence of a thrombus in a single patient (2.56%). Notably, left  
257 ventricular function remained normal in all cases, despite the observed chamber size variations.

## 258 **Discussion**

259 This study investigated the clinical characteristics and valve lesion patterns of rheumatic heart  
260 disease (RHD) in a resource-limited setting in Eastern Ethiopia. Our findings offer valuable  
261 insights into the burden and presentation of RHD in this understudied population.

262 The results highlight the persistent burden of RHD, with a majority of affected children being  
263 young and predominantly female. Shortness of breath and cough were the most frequent  
264 presenting symptoms, indicating a significant impact on their quality of life. Furthermore, a  
265 substantial proportion of patients presented with advanced disease, indicated by acute  
266 decompensated heart failure (ADHF) at admission. These observations emphasize the urgent  
267 need for improved strategies for early detection and management of RHD in resource-limited  
268 settings like Eastern Ethiopia. This includes health education initiatives and promoting better  
269 healthcare utilization.

270 Our findings are consistent with previous studies in Ethiopia and other low-resource settings,  
271 which consistently report a high burden of RHD in children, often presenting at advanced  
272 stages(6, 9, 11, 12, 15). The predominance of mitral regurgitation and aortic regurgitation as the  
273 most prevalent valve lesions aligns with existing literature, underlining the extensive valvular  
274 damage associated with pediatric RHD(9, 15, 16). However, this study provides novel insights  
275 into the specific clinical characteristics and prevalent valve lesions in Eastern Ethiopia,  
276 contributing to a more comprehensive understanding of RHD epidemiology in this region.

277 The high prevalence of ADHF and significant valvular involvement observed in our study have  
278 significant implications for clinical practice and public health policies in Ethiopia. Healthcare  
279 providers need to be alert for symptoms suggestive of RHD, particularly in children from



280 vulnerable populations. Early screening and diagnosis should be prioritized to facilitate timely  
281 intervention. Additionally, efforts to improve access to secondary prophylaxis and implement  
282 comprehensive RHD control programs are crucial to reduce disease burden and improve long-  
283 term outcomes for affected children. These findings underscore the need for multidisciplinary  
284 collaboration between healthcare professionals, policymakers, and community stakeholders. This  
285 collaboration is essential to develop and implement effective strategies for RHD prevention,  
286 diagnosis, and management in resource-limited settings.

### 287 **Limitations**

288 Several limitations should be considered when interpreting the findings of this study. Firstly, the  
289 relatively small sample size and single-center design may limit the generalizability of our results  
290 to other settings within Ethiopia and beyond. Furthermore, the lack of long-term follow-up data  
291 precludes a comprehensive assessment of treatment outcomes and disease progression in this  
292 population. Future studies with larger sample sizes, prospective designs, and longer follow-up  
293 periods are needed to address these limitations and provide a more robust understanding of RHD  
294 epidemiology and outcomes in Eastern Ethiopia.

### 295 **Conclusion:**

296 In conclusion, this study provides valuable insights into the clinical characteristics and valve  
297 lesions in RHD among children in Eastern Ethiopia. The findings underscore the persistent  
298 burden of RHD in this region and highlight the urgent need for improved early detection,  
299 diagnosis, and management strategies. Healthcare providers, policymakers, and researchers must  
300 collaborate to develop and implement comprehensive RHD control programs tailored to the  
301 unique challenges faced by resource-limited settings like Eastern Ethiopia. By addressing these

302 challenges and prioritizing RHD prevention and management, we can mitigate the morbidity and  
303 mortality associated with this preventable yet devastating disease among vulnerable pediatric  
304 populations.

## 305 **Data Sharing Statement**

306 Any of the data used for analysis in the study is available from the corresponding author and  
307 ready to be provided up on reasonable request.

## 308 **Contributors**

309 Conceptualization: Temesgen T Libe, Yunus E Kelil, Samrawit Abebaw, Faisel, and Kibrom  
310 Mesfin; formal analysis: Temesgen T Libe, Faisel, and Yunus E Kelil; investigation: Temesgen  
311 T Libe, Faisel, Yunus E Kelil, Samrawit Abebaw, and Kibrom; methodology: Temesgen T Libe,  
312 Faisel, and Kibrom; Validation: Temesgen T Libe, Faisel, and Kibrom, Yunus E Kelil and  
313 Samrawit Abebaw; writing the original draft: Temesgen T Libe, and Samrawit Abebaw.

## 314 **Acknowledgments**

315 We are very grateful to the Haramaya University College of Health and Medical Sciences for its  
316 ethical clearance provision and financial support. We also acknowledge our study participants,  
317 data collectors and supervisors for enabling us to realize this research work.

## 318 **Funding**

319 The thesis was funded by Haramaya University.

320 **Disclosure**

321 The authors declared that they have no conflicts of interest.

## 322 **References**

- 323 1. Kliegman RM, St Geme, J. W. Nelson Textbook of Pediatrics E-Book: Elsevier Health  
324 Sciences; 2019.
- 325 2. Watkins DA, Johnson CO, Colquhoun SM, Karthikeyan G, Beaton A, Bukhman G, et al.  
326 Global, Regional, and National Burden of Rheumatic Heart Disease, 1990-2015. *N Engl J Med.*  
327 2017;377(8):713-22.
- 328 3. Cilliers AM. Rheumatic fever and rheumatic heart disease in Africa. *S Afr Med J.*  
329 2015;105(5):361-2.
- 330 4. Arvind B, Ramakrishnan S. Rheumatic Fever and Rheumatic Heart Disease in Children.  
331 *Indian J Pediatr.* 2020;87(4):305-11.
- 332 5. Carapetis JR, Beaton A, Cunningham MW, Guilherme L, Karthikeyan G, Mayosi BM, et  
333 al. Acute rheumatic fever and rheumatic heart disease. *Nat Rev Dis Primers.* 2016;2:15084.
- 334 6. Zühlke L, Engel ME, Karthikeyan G, Rangarajan S, Mackie P, Cupido B, et al.  
335 Characteristics, complications, and gaps in evidence-based interventions in rheumatic heart  
336 disease: the Global Rheumatic Heart Disease Registry (the REMEDY study). *Eur Heart J.*  
337 2015;36(18):1115-22a.
- 338 7. Condemi F, Rossi G, Lupiz M, Pagano A, Zamatto F, Marini S, et al. Screening of  
339 asymptomatic rheumatic heart disease among refugee/migrant children and youths in Italy.  
340 *Pediatr Rheumatol Online J.* 2019;17(1):12.
- 341 8. Roberts K, Cannon J, Atkinson D, Brown A, Maguire G, Remenyi B, et al.  
342 Echocardiographic Screening for Rheumatic Heart Disease in Indigenous Australian Children: A  
343 Cost-Utility Analysis. *J Am Heart Assoc.* 2017;6(3).

- 344 9. Sani MU, Karaye KM, Borodo MM. Prevalence and pattern of rheumatic heart disease in  
345 the Nigerian savannah: an echocardiographic study. *Cardiovasc J Afr.* 2007;18(5):295-9.
- 346 10. Karthikeyan G, Connolly SJ, Ntsekhe M, Benz A, Rangarajan S, Lewis G, et al. The  
347 INVICTUS rheumatic heart disease research program: Rationale, design and baseline  
348 characteristics of a randomized trial of rivaroxaban compared to vitamin K antagonists in  
349 rheumatic valvular disease and atrial fibrillation. *Am Heart J.* 2020;225:69-77.
- 350 11. Animasahun BA, Madise Wobo AD, Itiola AY, Adekunle MO, Kusimo OY, Thomas FB.  
351 The burden of rheumatic heart disease among children in Lagos: how are we fairing? *Pan Afr*  
352 *Med J.* 2018;29:150.
- 353 12. Marijon E, Ou P, Celermajer DS, Ferreira B, Mocumbi AO, Jani D, et al. Prevalence of  
354 rheumatic heart disease detected by echocardiographic screening. *N Engl J Med.*  
355 2007;357(5):470-6.
- 356 13. Guilherme L, Cury P, Demarchi LM, Coelho V, Abel L, Lopez AP, et al. Rheumatic  
357 heart disease: proinflammatory cytokines play a role in the progression and maintenance of  
358 valvular lesions. *Am J Pathol.* 2004;165(5):1583-91.
- 359 14. Guilherme L, Köhler KF, Postol E, Kalil J. Genes, autoimmunity and pathogenesis of  
360 rheumatic heart disease. *Ann Pediatr Cardiol.* 2011;4(1):13-21.
- 361 15. Tantchou Tchoumi JC, Butera G. Rheumatic valvulopathies occurrence, pattern and  
362 follow-up in rural area: the experience of the Shisong Hospital, Cameroon. *Bull Soc Pathol Exot.*  
363 2009;102(3):155-8.
- 364 16. Koirala PC, Sah RK, Sharma D. Pattern of rheumatic heart disease in patients admitted at  
365 tertiary care centre of Nepal. *Nepalese Heart Journal.* 2018;15(1):29-33.

366