Decoding ruminative states from neurophysiological patterns

Jana Welkerling[1], Patrick Schneeweiss[2], Sebastian Wolf[1,*], Tim Rohe[3,*]

[1] Faculty of Economics and Social Sciences, Institute of Sports Science, Department of

Education & Health Research, University of Tübingen, Tübingen, Germany

[2] Department of Sports Medicine, University Hospital Tübingen, Tübingen, Germany

[3] Institute of Psychology, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen,

Germany

Corresponding Author: jana.welkerling@uni-tuebingen.de

*: These authors contributed equally.

Keywords: Rumination, Depression, EEG, Multivariate pattern decoding

Author contributions

Jana Welkerling: Methodology, Software, Formal Analysis, Investigation, Data Curation,
Writing Original Draft, Visualization, Funding acquisition

Patrick Schneeweiss: Resources, Writing Review & Editing

Sebastian Wolf: Conceptualization, Methodology, Funding Acquisition, Writing Review &
Editing, Supervision

Tim Rohe: Conceptualization, Methodology, Formal analysis, Funding Aquisition, Writing
Review & Editing, Supervision

Funding

The Robert Enke Foundation provided partial funding for this project. The foundation was
neither involved in the study design nor in the collection, analysis and interpretation of data
nor in the writing of the report and in the decision to submit the article for publication.

Declarations of interest: none

Abstract

Individuals with depression often engage in iterative "rumination" about challenging

situations and potential outcomes. Although the state of rumination has been associated with

diverse univariate neurophysiological features, the potential to use multivariate patterns to

decode it remains uncertain. In this study, we trained linear support vector machines to

differentiate state rumination from distraction using patterns in the alpha, beta, and theta

bands, as well as inter-channel connectivity. We used validated tasks to induce rumination or

distraction for eight minutes in 24 depressed individuals in six runs over three sessions.

During inductions, we recorded 64-channel EEG data and measured self-reported levels of

rumination. Participants reported strongly increased rumination, and we decoded state

rumination from EEG patterns with significant accuracy. However, the informative features

were not consistent across participants, demonstrating that while ruminative states can indeed

be decoded from EEG data, these states appear to reflect processes unique to each individual.

1 Introduction

When faced with an emotionally challenging situation, such as social exclusion, humans often

start to think about the situation, potential solutions, and their own negative emotional

reaction to it. This process of "chewing" on negative thoughts long beyond the eliciting

situation is known as "rumination" and has been defined as "a mode of responding to distress

that involves repetitively and passively focusing on symptoms of distress and on the possible

causes and consequences of these symptoms" (Nolen-Hoeksema, Wisco, & Lyubomirsky,

2008). Typically, ruminative thoughts are accompanied by negative emotional reactions.

While some situations induce rumination in most people, there is variability in individuals'

general tendency to ruminate. Thus, rumination can be conceptualized both as a transient state

experienced in response to a distressing situation and as a stable personality trait.

Trait rumination is a key clinical feature of depression, as well as other psychiatric

disorders such as anxiety and eating disorders (Aldao, Nolen-Hoeksema, & Schweizer, 2010).

Depression, one of the most burdensome diseases worldwide (GBD 2019 Mental Disorders

Collaborators, 2022), is characterized by persistent low mood, loss of interest, and low energy.

Trait rumination is linked to the onset of depression, acting as a risk factor (Aldao et al., 2010;

Kuyken, Watkins, Holden, & Cook, 2006; Watkins & Roberts, 2020), as well as to the

duration and the severity of depression, with a stronger tendency to ruminate being associated

with longer episodes (Nolen-Hoeksema, 1991; Timm et al., 2017) and an increased risk of

suicide attempts (Chiang et al., 2022; Eshun, 2000).

Individuals who exhibit heightened ruminative thought patterns during emotionally

challenging situations are more likely to demonstrate higher levels of trait rumination (Fang,

Marchetti, Hoorelbeke, & Koster, 2019). In fact, neurocognitive models, such as the

attentional scope model of rumination (Whitmer & Gotlib, 2013), have identified state

rumination as a key cognitive factor that perpetuates momentary depressive symptoms (Timm

et al., 2017; Watkins & Roberts, 2020) through a vicious cycle: When in a sad mood,

individuals with depression narrow their attentional focus to their mood, its causes, and

consequences, impeding the retrieval of alternative thoughts. By maintaining information

related to the sad mood in their working memory, patients are unable to access anti-depressive

strategies stored in their long-term memory, thus perpetuating their sad mood. To disrupt this

cycle of repetitive negative thoughts and feelings, individuals attempt to interrupt rumination,

either physiologically, for example by abusing alcohol (Devynck, Rousseau, & Romo, 2019),

or psychologically, by distracting themselves from negative thoughts. Employing distraction

as a cognitive coping strategy involves deliberately redirecting one's attentional focus from

the distressing thoughts towards neutral or enjoyable activities (Nolen-Hoeksema, 1991),

offering transient relief from the sad mood.

In individuals with depression, rumination appears to be a highly stable recursive

attractor state of brain dynamics (Rolls, 2016) accompanied by specific cognitive and

negative affective mental representations. Rumination has been associated with numerous

aspects of brain dynamics, such as activity in the default mode network as observed through

fMRI (Zhou et al., 2020) and neural oscillations indicating functional connectivities as

evident in spectral EEG features (see below). However, it is unknown whether an

experimentally induced ruminative brain state, as measured by neurophysiological patterns,

can be distinguished from alternative brain states such as distraction. Should this be

achievable, it would enable the classification of an individual's ruminative state based on his

or her dynamic brain activity. This capability would not only contribute to neurocognitive

(e.g., Whitmer & Gotlib, 2013) and clinical models of the neural dynamics of state

rumination, but also offer a means for measuring ruminative states through

neurophysiological data online e.g., during intervention studies.

In this study, we therefore aimed to decode state rumination from neurophysiological

EEG patterns as a proof-of-principle. We chose EEG for its ability to capture brain dynamics

93 with high temporal resolution, presenting a more practical option than fMRI, which is also

94 often not feasible in interventional studies, such as those involving exercise. EEG studies have

95 linked both trait and state rumination with numerous univariate neurophysiological features.

96 Trait rumination has been shown to be positively associated with functional beta-band

97 connectivity between the posterior cingulate cortex (PCC) and subgenual prefrontal cortex

98 (sgPFC) (Benschop et al., 2021). Resting-state EEG measurements have demonstrated that

99 higher levels of trait rumination are associated with asymmetrically increased alpha power

100 over the left parietal-occipital cortex (Umemoto et al., 2021) and decreased alpha power over

101 the right frontal cortex (Keune, Bostanov, Hautzinger, & Kotchoubey, 2011). To investigate

102 neurophysiological correlates of state rumination, previous studies, such as Laicher et al.

103 (2022) and Huffziger and Kuehner (2009), have experimentally induced rumination (mostly

104 in contrast to distraction) in depressed patients to elicit increased self-reported rumination and

105 negative affect. These studies have demonstrated that state rumination correlates with higher

106 EEG beta power in the left temporal cortex compared to a positive and a neutral condition

107 (Ferdek, van Rijn, & Wyczesany, 2016), and with an increased EEG alpha power over the

108 prefrontal cortex (Putnam & McSweeney, 2008). Moreover, internal attention, which is

109 closely linked to rumination (Magosso, Ricci, & Ursino, 2021), was found to increase EEG

110 theta power over frontal midline regions. External attention, however, which is closely linked

111 to distraction, was found in the same study to increase theta power while decreasing alpha

112 power over parieto-occipital regions (Magosso et al., 2021).

113   Overall, the body of research examining the neurophysiological correlates of state

114 rumination through EEG is very limited, with most studies to date having focused on trait

115 rumination. Those studies that have investigated state rumination have been subject to

116 methodological shortcomings, such as the absence of self-reported state rumination

117 measurements (Ferdek et al., 2016). Moreover, prior studies examining the

118 neurophysiological correlates of state rumination have used group-based analysis. Such an

approach may not be ideal given the results of meta-analyses of brain activation patterns

associated with depression, which have shown a lack of statistical significance when

aggregated across individuals (Horato, Quagliato, & Nardi, 2022). Thus, it seems very likely

that complex mental states such as state rumination are characterized by substantial

interindividual variability in neurophysiological patterns.

Two recent studies used a *multivariate* decoding approach to decode *trait* rumination

from neurophysiological patterns that suggests that trait rumination is reflected in consistent

patterns across participants: Aydın and Akın (2022) were able to distinguish between high and

low trait ruminators, as defined by self-report questionnaires, through spectral EEG patterns.

Similarly, Kim et al. (2023) predicted *trait* rumination, as measured by a rumination

questionnaire, from connectivity patterns in resting-state fMRI data (Kim et al., 2023).

However, while both of these studies successfully trained their *trait rumination* decoding

models on data aggregated *across* participants, they did not consider individual variability in

the mapping between rumination and neurophysiological patterns. Additionally, it remains

unknown whether *state* rumination is associated with neurophysiological patterns unique to

each individual and how temporal dynamics of state rumination behave.

In a proof-of-concept study, we therefore addressed the following research questions:

1) Do inductions of rumination or distraction lead to differential brain states that can be

decoded from neurophysiological patterns)? 2) Are the neurophysiological patterns encoding

rumination consistent across individuals, or are they unique to each individual? 3) Are

decoded and self-reported levels of rumination correlated across individuals? 4) Which

features of the neurophysiological patterns are most informative for decoding state

rumination? Answers to these question will not only provide fundamental insights which

neurophysiological states of the brain correlate with ruminative states, but they will also

provide a measurement tool to quantify rumination from neurophysiological data, for example

to assess intervention effects on rumination (Welkerling et al., 2024).

To answer our research questions, we experimentally induced state rumination versus

state distraction and used multivariate pattern analysis to decode the two states from

neurophysiological patterns (Fig. 1). During the inductions, we collected EEG data and

measured self-reported rumination and affect before, during, and after the inductions. To

classify rumination and distraction, we trained support vector machines (SVM) to decode both

states from spectral EEG features in the alpha, beta, and theta bands, as well as functional

connectivity, which have been linked to rumination in previous research. As alpha and beta

connectivity have been linked to rumination, we decided to calculate whole-spectrum

connectivity to ensure that no potentially relevant information, such as from other frequency

bands, is overlooked. (Benschop et al., 2021; Ferdek et al., 2016; Keune et al., 2011; Magosso

et al., 2021; Putnam & McSweeney, 2008; Umemoto et al., 2021).

Figure 1. Experimental design and decoding approach. a) The experimental design measured

self-reported rumination and affect (i.e., Perseverative Thinking Questionnaire (PTQ-

S),Positive and Negative Affect Schedule (PANAS); for further details please refer to the

Methods section), as well as one-minute resting state EEG data, before visual items induced

rumination or distraction. During the eight-minute inductions, participants reported their

ruminative and affective state on a seven-point visual analogue scale (VAS) every minute. b)

To decode rumination versus distraction from neurophysiological patterns, we computed

feature vectors comprising alpha (60 features), beta (60 features), and theta (60 features)

power, as well as connectivity between electrodes (1770) from eight one-minute bins of 64-

channel EEG data. c) Next, we trained linear SVM decoders to distinguish between induced

rumination and distraction from neurophysiological feature vectors (feature values are shown

on the x and y axis and exemplars are the dots lying in x/y space) using cross-validation. To

evaluate decoder performance in each one-minute bin, we computed the proportion of

correctly predicted state labels.

172   **2 Methods**

2.1 Participants

Twenty-four moderately or severely depressed individuals (17 female, M_{age} = 24.6 years)

were included in the study after providing written informed consent. A pilot study with $n = 6$

participants was conducted to calculate statistical power and sample size for the main study.

For a proof-of-concept study such as ours, the primary goal was to show that decoding

accuracy is significantly larger than a chance level of 50%. Thus, we converted decoding

accuracy from the pilot sample into Cohen's d to compute the required sample size to

statistically safeguard above-chance decoding accuracy. In the pilot study the mean decoding

accuracy across all participants differed with a medium effect size (d = 0.60) from chance-

level. Based on this effect, power analysis using the open-source programme G*Power (3.1)

resulted in a minimum sample size of 24 people (two-tailed t-test for dependent samples, α-

level of .05, power of 80%, not taking into account possible drop-outs). In addition to the six

data sets that had already been collected, data from an additional 18 participants were also

gathered. The pilot study additionally involved a positive mood induction and fNIRS

measurements. Because there were no other changes in the experimental design or setup

between the pilot and main studies, we included data from these six individuals in the main

sample. Sociodemographic and clinical information of the participants can be found in Tab.

S10. Sensitivity analyses including only the main sample are described in the supplementary

results (Fig. S4).

Inclusion criteria were a diagnosis of moderate or severe depression according to the

Structured Clinical Interview for DSM-5 Disorders, Clinician Version (SCID-V; Beesdo-

Baum, Zaudig, & Wittchen, 2019); F32.1-F32.2, F33.1-F33.2 DSM-V), a Beck Depression

Inventory II (BDI-II; Beck, Steer, & Brown, 1996) score greater than 19, age between 18 and

40 years, fluency in German, no current psychopharmacological treatment, and no current

psychological treatment. To ensure that rumination and distraction inductions reliably altered

the mental states in each participant, we only included participants who showed an increase in

rumination after a rumination induction pretest (PTQ-S$_{post-pre}$ > 0) and a decrease in

rumination after the distraction induction pretest (i.e., PTQ-S$_{post-pre}$ < 0). Exclusion criteria

were acute suicidality, schizophrenia or other psychotic disorders (F23, F20.81, F20.9 DSM-

V), acute substance misuse excluding tobacco (F10, F11, F12, F13, F14, F15, F16, F18, F19

DSM-V), acute eating disorders (F50.01, F50.02, F50.2, F50.81, F50.82 DSM-V) acute or

chronic diseases or conditions influencing brain metabolism, such as diabetes mellitus (E10-

E14 ICD-10), kidney insufficiency of stage 3 according to the Kidney Disease Outcome

Quality Initiative Guidelines (National Kidney Foundation, 2002), non-adjusted hypertension

(I10 ICD-10), and moderate to severe craniocerebral trauma (Glasgow Coma Scale; Teasdale

& Jennett, 1974; GSC 3-12) or craniocerebral trauma second or third degree with loss of

consciousness for more than 30 minutes. Additionally, we excluded participants who engaged

in regular physical exercise, as this was an exclusion criterion pertinent to the second part of

the study, details of which will be reported in a separate publication (Welkerling et al., 2024);

for more information, refer to our pre-registration document available online at osf.io/qf2ha.

Participants were recruited through local hospital departments and ambulatory

psychotherapeutic practices. Participants were compensated financially and provided with

additional mental health treatment information. The study was approved by the local Ethics

Committee for Medical Research of the University of Tübingen (133/2020BO2) and was

conducted in accordance to the tenets of the Declaration of Helsinki of 2010.

2.2 Procedure

Individuals interested in taking part in our study were first screened for eligibility in a

telephone call based on the pre-defined inclusion and exclusion criteria. Those found eligible

were then invited to an in-person appointment. After providing informed written consent, they

underwent a clinical interview (SCID-V) conducted by a psychologist trained in clinical

diagnostics. They also completed questionnaires to measure the severity of their depression

(BDI-II) and their general tendency towards rumination (PTQ). Additionally, this initial

appointment included pretests for the rumination and distraction induction protocols to

determine if these procedures led to a sufficient change in individuals' levels of rumination, as

required for inclusion in the study. At a subsequent appointment, participants were medically

examined by sports physicians for eligibility to engage in the physical exercises involved in

the second part of the study, which will be reported in a separate publication (Welkerling et

al., 2024). In the second week of the study, participants attended three sessions during which

they underwent both rumination and distraction inductions consecutively to generate training

data for the decoder. The order of these inductions was randomized within each session but

counterbalanced across participants.

2.3 Experimental tasks

Before each induction, a baseline resting-state measurement was taken, with participants

closing their eyes for 60 seconds while EEG data were collected. For the rumination induction

protocol, participants were exposed to 16 different items, which were adapted from Huffziger

and Kuehner (2009) and designed to direct participants' attention towards their own

symptoms and emotions (e.g., "Think about the way you feel inside."). Each item appeared on

the screen for seven seconds, and participants were instructed to reflect on it. Following this, a

blank screen was displayed for 30 seconds to avoid reading artifacts, and participants were

asked to continue focusing on the item. After every two items, participants assessed their level

of rumination and their affective state using two consecutive visual analogue scales (VAS; see

supplemental methods for VAS). The VAS appeared on the screen and participants rated their

current levels of rumination and affect by pressing the corresponding button on the keyboard.

Before the induction protocols started, participants were familiarized with the definition of

rumination. We explicitly instructed participants that rumination does not refer to general

thinking, but rather means brooding about their own emotions and symptoms.

For the distraction induction protocol, participants were exposed to 64 items from four

different tasks, all designed to shift their attentional focus away from their own symptoms and

distress. In the first task, adapted from Huffziger and Kuehner (2009), 16 different prompts

(randomly selected from the original set of 28 items) were presented, aiming to elicit neutral

mental images (e.g., "Think of raindrops flowing down a window"). The second task was a

knowledge-based exercise, requiring participants to evaluate the correctness of 16 randomly

selected statements from the Distraction Questionnaire by Ehring, Fuchs, and Kläsener (2009)

(e.g., "Spain is not a neighboring country of Germany!"), with the response options "true" and

"false". The third task, also adapted from Ehring et al. (2009), was a word-generation activity

requiring participants to think of a new word beginning with each letter from given five-letter

words (e.g., from "ocean": oven, concentration, elephant, apple, night). The fourth task,

developed by the authors of this paper, aimed to redirect participants' attention towards self-

related topics unrelated to symptoms, similar to the prompts of Huffziger and Kuehner (2009)

(e.g., "Think about your next weekly shopping"). In each trial, an item from one of the four

tasks was presented for two to five seconds, with their order randomized within and across

participants. After each item, a blank screen appeared for 7.5 seconds. Participants were

instructed to maintain their focus on the previously presented item. After every eight items,

participants reported their level of rumination and affective state using the VAS. Before and

after each induction, levels of rumination and affect were assessed with the PTQ-S and the

PANAS, respectively. Further details concerning the PANAS and VAS affect can be found in

the supplementary methods. To accommodate potential shifts in attentional focus away from

the screen, a beep was used to signal the onset of the next item or the VAS (Fig. 1).

271   The experiments were conducted in a laboratory room using Psychtoolbox 3.0.16

(Brainard, 1997) running under MATLAB 2020b (The MathWorks Inc., 2020) on a laptop

connected to an LCD screen (Acer B246HYL; 60.5 cm diagonal display size). Acoustic

stimuli were delivered at approximately 77 dB SPL through speakers (Amazon basics

V216Custom1) positioned to the right and left of the screen. Participant responses were

collected with a keyboard connected to the laptop via USB. Participants were seated

approximately 75 cm away from the screen.

2.4 Self-report measurements

2.4.1 Perserverative Thinking Questionnaire (PTQ)

The German version of the PTQ comprises 15 items to assess the core processes of repetitive

negative thinking: repetitive, intrusive, difficult to disengage from, unproductive and

capturing mental capacity. Nine items load on the first three processes and three on each of

the last two. Furthermore, a total sum score for all 15 items can be computed and assesses the

general tendency to ruminate. In the PTQ-S, the state version, instructions change to how

participants thought about negative events and problems in the last moments. Higher values

correspond to higher levels of rumination (Ehring et al., 2011). Compared to the widely used

Response Styles Questionnaire (RSQ) by Nolen-Hoeksema and Morrow (1991), the PTQ

focuses on key processes of rumination without specifying the content of those thoughts.

Since we were interested in the underlying processes of rumination compared to distraction,

we measured rumination with the PTQ. Internal consistency for the total sum score amounted

to $\alpha = .95$ in a clinical sample and construct validity to the RSQ amounted to $r = .72$ (Ehring

et al., 2011). Of a total of 288 timepoints seven were missing (2.43%). Missing values were

handled by averaging the total sum score of the PTQ-S per induction and timepoint over the

three sessions.

2.4.2 Visual Analogue Scales (VAS)

Adapted to the self-assessment manikins from Bradley and Lang (1994) the authors designed

manikins assessing rumination and affect. The VAS range from 1 to 7. For rumination one

corresponds to *"no rumination"* and seven to *"a lot"* (see supplemental methods). 54 from

1296 datapoints of self-reported VAS-rumination were missing (4.17%). Missings were

handled by averaging the VAS responses per induction and timepoint over the three sessions.

2.5 EEG measurements

EEG data were recorded using a 64-electrode actiCHamp Plus system (Brain Products GmbH,

Gilching, Germany) with a sampling rate of 1000 Hz. Fifty-eight of these electrodes were

positioned according to the international 10-20 system (Jasper, 1958). Five of the 64

electrodes (i.e., O1, O2, TP9, TP10, T8) deviated from the 10-20 system and were placed

around the eyes and mouth to monitor eye movement and facial motion (adjacent to both the

right and left eyes; below and above the left eye; on the risorius muscle). FCz served as the

reference and AFz as the ground electrode. Additionally, to control for muscle artifacts from

neck and shoulder movements, four pairs of bipolar electrodes were applied: two pairs on

either side of the trapezius muscle, and two pairs on either side of the neck targeting the

sternocleidomastoid muscles. To monitor head movements, an accelerometer was attached to

the electrode cap to the left of the Cz electrode. In the case of the six pilot study participants,

59 electrodes were placed according to the 10-20 system. However, for the pilot participants,

a separate reference electrode channel was used, which was part of the electrode setup for the

main sample: for them, the Iz electrode was placed in FCz, which then served as the reference

electrode in the main sample.

2.5.1 EEG preprocessing

The pre-processing of EEG data was carried out using Brainstorm software (Tadel, Baillet,

Mosher, Pantazis, & Leahy, 2011) (version 14 October 2022) in six steps. First, to remove

muscle- and head-motion artifacts, the raw EEG data were corrected for muscle and

movement artifacts using sliding-window regressions (van der Meer et al., 2016) implemented

in EEGLab (Delorme & Makeig, 2004) (O1, O2, TP9, TP10, T8, bipolar electrodes from the

trapezius and sternocleidomastoid muscles, as well as the accelerometer). Second, the EEG

data were bandpass filtered with a frequency range of 0.25 to 45 Hz. Third, eye blinks were

automatically detected by Brainstorm's standard algorithms using data from the electrode

above the left eye. Signal-space projectors (SSPs) were subsequently created from 400 ms

segments, band-pass filtered between 1.5 and 15 Hz and centered on the detected blinks. The

first spatial component of the SSPs was then used to correct for blink artifacts in the

continuous EEG data. Fourth, all data were visually inspected to identify and exclude any

segments showing artifacts from blinks, saccades, motion, electrode drifts, or jumps. Fifth,

whole channels were marked as bad if visual inspections revealed suspicious time courses or

spectral power distributions resulting from artifacts. Bad channels were interpolated as the

distance-weighted average of neighboring channels. Lastly, data were re-referenced to the

PO7 and PO8 electrodes (PO9 and PO10 for the pilot sample) and resampled to a frequency

of 250 Hz.

To ensure that EEG data used for training the decoder were not contaminated by

reading artifacts, analysis was restricted to data from the blank-screen sequences. Because the

shortest blank-screen sequence lasted for 7.5 seconds, all blank-screen sequences were binned

into 7.5-second epochs for consistency. Because the electrode set-up differed slightly between

the pilot ($n = 6$) and main ($n = 18$) samples, channels that were missing (AFz in the pilot

sample and PO9 and PO10 in the main sample) were interpolated similarly to previously

marked bad channels, resulting in 60 channels for all participants. To compute feature vectors

for the decoding algorithm, 7.5-second epochs were transformed using a Fast Fourier

transformation. Feature vectors included the power of the alpha (i.e., averaged over 8 to 13

Hz), beta (i.e., averaged over 13 to 30 Hz), and theta (i.e., averaged over 4 to 7 Hz) bands

across the 60 channels and averaged within each 7.5s epoch. Additionally, connectivity

features comprised the pairwise connectivity between the 60 electrodes computed as pairwise

correlation of time courses within the 7.5-second epochs (i.e., 1770 correlations per epoch:

[60*60 channels – 60 values from the diagonal] / 2). Each feature vector was z-standardized

(*zscore* function Matlab) to normalize scaling differences between feature vectors (Pereira,

Mitchell, & Botvinick, 2009). Overall, the feature vector per 7.5-second epoch comprised

1950 features including 60 z-standardized alpha power values, 60 z-standardized beta power

values, 60 z-standardized theta power values, and 1770 z-standardized connectivity values

(Fig. 1). Shorter epochs of 3.75 seconds length did not change our results, we therefore chose

7.5 s to align the definition of features to the duration of the blank screen sequences from the

distraction induction.

2.5.2 Decoding procedure

To decode rumination versus distraction, we trained linear support-vector machine (SVM)

classification models using LIBSVM (Chang & Lin, 2011). For training and prediction, we

employed a leave-one-session-out cross-validation (CV) procedure: The SVMs learned the

mapping between feature vectors and the two classes (rumination vs. distraction) using data

from two of the three induction sessions. Subsequently, these models were applied to predict

the class labels of the independent feature vectors from the remaining session. This CV

procedure, recommended by best practice guidelines (Scheinost et al., 2019), ensured that

training and testing of the decoders was performed on independent data sets with balanced

classes within each cross-validation fold. Because we implemented a within-participant

experimental design, the data from the induced experimental conditions (i.e., rumination vs.

distraction) were independent in all three sessions at the level of individual participants. The

SVMs' nu parameter (i.e., from [0.0001,0.1:0.025:0.7]) was optimized through nested cross-

validations. To explore the time course of rumination across the eight one-minute bins, the CV

procedure was repeated for each bin. Importantly, the data of each bin were divided into eight

7.5-second epochs (see above) that were treated as data segments for decoding, resulting in an

effective bin size of one minute. In other words, in each cross-validation fold for a bin, the

decoder was trained on 32 data points (= 2 sessions x 2 inductions x 8 epochs). We chose

eight one-minute bins, each consisting of eight 7.5-second epochs to align the decoding

approach with the VAS reports that were requested every minute during inductions. This

approach facilitated an exploratory comparison between the SVM decoding performance and

the VAS reports, analyzing decoding probability estimates against the rumination scales. The

decoder predicted class labels (i.e., rumination vs. distraction) and corresponding probability

estimates for each class. The decoded probability estimates represented the probability p that

a data segment belongs to a rumination state (if $p > 0.5$) or to a distraction state (if $p < 0.5$) as

defined in the induction protocols for decoder training (i.e., the probability for the classes of

rumination and distraction add up to one). Additionally, to establish a baseline decoding

accuracy before the inductions, we applied the same CV procedure to the one-minute resting-

state data collected before each induction, epoched similarly to the induction data. The chance

level for the decoder performance was computed by repeating the CV procedure on

randomized class labels within each bin (i.e., 1000 random allocations of the labels

rumination versus distraction). Out of the 10368 data segments obtained from all participants

and sessions, 447 (4.31%) were missing or excluded from the decoder training due to

technical reasons. The decoder for minute 0 before the inductions served only to establish

baseline performance; hence, all inferential analyses concerning the performance of the

decoder excluded this one-minute baseline (Fig. 1).

2.6 Statistical analyses

2.6.1 PTQ-S and VAS

To assess whether the inductions increased or decreased state rumination as intended, we

averaged the total score of the PTQ-S across sessions for each induction type, timepoint (pre

vs. post induction), and participant. Subsequently, we computed a linear mixed model with the

averaged total score of the PTQ-S as the outcome variable. Fixed effects included induction

(rumination vs. distraction), timepoint (pre vs. post), and their interaction. Random intercepts

per participant were implemented. The random slopes per participant did not converge

because the observed data were insufficient compared to what should have been estimated.

Post-hoc two-sided pairwise *t*-tests based on estimated marginal means were used to assess

differences in self-reported rumination levels for every type of induction. Additionally,

estimated marginal means of the total scores were compared at every timepoint (pre vs. post)

between both inductions. The significance levels for the post-hoc tests were adjusted using the

Bonferroni correction to account for four comparisons (Fig. 2a).

407   VAS responses were averaged across sessions for each induction type, timepoint

(minute 0 to minute 8 during induction), and participant. A linear mixed model was then

calculated with these averaged responses as the dependent variable. The model included fixed

effects for the type of induction (rumination vs. distraction), timepoint (minute 0 to 8 during

induction), and their interaction as well as random intercepts for participants. The random

slopes per participant did not converge because the observed data were insufficient compared

to what should have been estimated. To compare the two conditions at each timepoint, post-

hoc two-sided pairwise *t*-tests based on estimated marginal means were conducted and

subsequently Bonferroni corrected for nine tests (Fig. 2b). Detailed results can be found in

supplementary Tab. S1 and in figure 2. Additionally, to verify whether the inductions reliably

induced rumination and distraction, the averaged estimated marginal means of VAS responses

over sessions for both conditions were calculated for each timepoint. Each timepoint

beginning from minute 1 was compared to baseline rumination at minute 0 using a paired

two-sided *t*-test (Tab. S2). These tests were also Bonferroni corrected, this time for 16

comparisons.

2.6.2 Decoding analysis

To evaluate the accuracy of the person-specific decoders, the predicted class labels

(rumination vs. distraction) from the independent left-out session were compared with the

actual label of the data point. Decoding accuracy for each participant was quantified as a

percentage of correct predictions for each bin, across all CVs and both induction types. To

establish a baseline decoding accuracy for each participant in every bin, a chance level was

computed by repeating the CV procedure on randomized class labels (i.e., 1000 randomized

allocations of the labels to the data). Subsequently, the mean of a participant's randomization

distribution was subtracted from the original decoding accuracy for that participant to obtain

chance-level-corrected decoding accuracies. For a global test of decoding accuracy, chance-

level-corrected accuracies were averaged across the eight bins from the inductions for each

participant, excluding the baseline measurement from the one-minute resting bin. A one-sided

one-sample *t*-test was conducted to test if these chance-corrected decoding accuracies across

bins and participants were significantly greater than zero. This approach allows for the

assessment of whether the decoders could distinguish between rumination and distraction

states beyond what would be expected by chance alone. 95%-Confidence Intervals (CI) and

Bayes Factor Analysis *bf10* were calculated for the primary analysis, quantifying evidence for

a difference to chance level (H□) relative to the null hypothesis of no difference (H□) using

the BayesFactor toolbox (Krekelberg, 2024).

To analyze the progression of decoding accuracies throughout the eight minutes of

induction, we fitted a linear mixed model with decoding accuracy as dependent variable. Bins

were defined as fixed effects and random intercepts for participants were included. The

random slopes per participant did not converge because the observed data were insufficient

compared to what should have been estimated. To correct for multiple comparisons across

bins, we derived p-values for the averaged accuracies across participants for each bin using

randomization tests that applied the max-stat approach from Blair and Karniski (1993). In this

procedure, the maximum decoding accuracies across the nine bins for each of the 1000

permutations were taken as the max-stat randomization distribution. For each bin, multiple-

comparison-corrected p-values were then calculated as the proportion of the sum of values

from the max-stat distribution that were higher than or equal to the original decoding accuracy

of that bin averaged across participants (Fig. 3). Additionally, pairwise two-sided t-tests with

the estimated marginal means of the accuracies compared accuracies of bin 0 to all other

timepoints. These tests were Bonferroni corrected for eight tests.

For a global test of the performance of the decoder applied to each participant, we

averaged the decoding accuracies across all bins for every participant and assessed whether

the resulting mean decoding accuracies were significantly different from zero. This was

determined using a randomization test in which the p-value was calculated based on the

proportion of decoding accuracies from 1000 random permutations that were equal to or

greater than each participant's actual decoding accuracy (Tab. S4). All of these analyses were

conducted using custom-written code in MATLAB (2020b).

Additionally, to investigate whether rumination-related EEG patterns were consistent

across participants, we trained a 'group decoder' using combined data segments from all

participants. This decoder was subjected to the same cross-validation process used for

evaluating individual-level participant data. In conducting a global assessment of the accuracy

of this group decoder, we averaged the decoding accuracies across the eight applicable data

segments (i.e., omitting the one-minute resting period used as a temporal baseline) per

participant. A one-sided one-sample t-test was then performed to determine if these accuracies

were significantly greater than 50%, suggesting a performance above chance level.

All linear mixed models conducted for PTQ-S, VAS and decoders' accuracies were

fitted with the lmer function from the package "lme4" as well as the anova function in R (R

Core Team, 2017) (version 4.4.0). Model assumptions, including linearity, normal

distribution, homoscedasticity, and absence of outliers, were checked visually. Additionally

for outlier analysis Cook's Distance was calculated and if values were greater than one,

sensitivity analyses were conducted as described in the following. If assumptions were

violated sensitivity analyses were calculated without influential cases and if results did not

differ results of the complete sample were reported. Effect sizes for the linear models were

estimated using partial eta squared (Cohen, 1973) (F_to_eta2 function in R) and Cohen's d

was calculated using the R function t_to_d for post-hoc tests. Both functions belong to Rs

package "effectsize".

2.6.3 Post-hoc and exploratory analysis

In our post-hoc analysis exploring the convergence or divergence of self-reported rumination

and EEG-decoded rumination, we correlated the self-reported rumination levels from VAS

scales and PTQ-S questionnaires with the decoders' probability estimates of rumination

versus distraction. Individual VAS values for rumination, as well as the decoders' probability

estimates, were averaged across sessions for each bin per induction type. We then performed

linear correlations between the 18 VAS values (resulting from two inductions across nine

timepoints) and the corresponding probability estimates. Fisher *z*-transformed individual

correlations were tested against zero at the group level using one-sided one-sample *t*-tests.

The analysis was complemented with the Bayes Factor *bf10*. Additionally, individual PTQ-S

rumination change scores and probability estimates were averaged for each induction type and

assessed through linear correlation (Fig. 4).

Adopting the "virtual lesion approach" developed by Kohoutová et al. (2020) and

applied by Kim et al. (2023), we assessed the contribution of different feature sets (i.e., alpha,

beta, theta band power, and connectivity) to decoder performance for those participants with

significant decoders ($n = 10$): First, we trained decoders excluding one feature set at a time to

observe the impact on decoding accuracy compared to using the full feature sets.

Subsequently, we trained a decoder on only a single feature set to assess the feature set's

decoding performance in isolation. A change in decoding performance related to the omission

or isolated use of a feature set indicates the amount of rumination-related information

represented by that feature set (Tab. 1). For the "virtual lesion" approach, decoders were

trained as described above and baseline resting-state data were not included in the

performance statistic. Comparisons between decoder performance with one feature set

omitted and the full feature set decoders were made using paired two-sided t-tests, adjusted

for multiple comparisons with Bonferroni correction for four tests. Similarly, the performance

of decoders trained on single feature sets was compared to a chance level of 50% using paired

one-sided t-tests, Bonferroni corrected for four tests. Bayes Factor Analysis (*bf10*) was

additionally conducted for all t-tests.

509   Additionally, we analyzed the feature weights of the SVMs in the subsample of 10

participants whose decoder performance exceeded chance level. To determine whether feature

weights contributed significantly to the decoders' performance, the feature weights were

averaged across cross-validations and bins for these participants and then absolute values

were taken. To determine the significance of these averaged feature weights at the group level

while correcting for multiple comparisons across the various features, we calculated p-values

as the proportion of maximum mean feature weights averaged across the aforementioned

participants in the feature set from randomized data ($n = 1000$ randomizations) that were

equal to or larger than the original feature weights (cf. max-stat approach above). Similarly,

feature weights from the ten participants were analyzed at the individual level (Fig. 5). It

should be noted that the analysis based on the ten participants with significant decoder

accuracies is exploratory in nature and therefore not circular. Consequently, it does not

constitute double dipping (Kriegeskorte, Simmons, Bellgowan, & Baker, 2009).

2.7 Data availability statement

The datasets and scripts generated and used for analysis in the current study are available in

the OSF repository: https://doi.org/10.17605/OSF.IO/QF2HA

525   **3 Results**

3.1 Self-reported rumination (PTQ-S and VAS)

Self-reported rumination increased after the rumination induction and decreased after the

distraction induction (Fig. 2a; linear mixed model with significant induction x timepoint

interaction for PTQ-S; $F(1, 69) = 121.77$, $p < .0001$, $\eta_p = 0.64$). At post-measurements,

rumination was significantly higher in the rumination condition ($t(69) = 10.76$, $p < .0001$, $d =$

1.30), whereas at pre-measurements it was significantly lower ($t(69) = -4.85$, $p < .0001$, $d =$

-0.58).

Similarly, continuous VAS-reported rumination quickly increased during the

rumination induction to a stable plateau and slightly decreased during the distraction

induction (Fig. 2B; significant interaction in the linear mixed model between induction and

timepoint; $F(8, 391) = 17.02$, $p < .0001$, $\eta_p = 0.26$). Thus, VAS-reported rumination differed at

each timepoint between both induction conditions (see Fig. 2b). Unexpectedly, but consistent

with the PTQ-S pre-measurements, rumination levels already differed slightly, yet

significantly, before the induction. This difference resulted from carry-over effects from the

previous inductions despite our counter-balanced experimental design: When we analyzed

only the first run of each session, we found no difference in rumination between the two

induction conditions (see supplemental results, Fig. S1). Consistent with the pre- and post-

questionnaire measurements, VAS-reported data indicated that rumination was successfully

induced in our sample already from the first minute of rumination induction (i.e., significant

difference in self-reported VAS rumination in the rumination induction from each timepoint

compared to t0; supplemental results Tab. S2).

Figure 2. Self-reported rumination from the questionnaire and VAS scales for the rumination

and distraction conditions. a) The total score of the PTQ-S for pre and post timepoints for

each induction (across-participant mean +/- SEM and individual data in the background). b)

Time course of VAS self-reported rumination (across-participant mean +/- SEM and

individual-level data in the background). Asterisks indicate significant ($p < .05$) differences

from post-hoc pairwise *t*-tests between rumination and distraction. VAS were recorded after

the resting-state baseline measurement before the inductions (minute 0) and at minutes 1 to 8

during inductions (see supplemental results Tab. S1).

3.2 Decoding rumination from individual neurophysiological patterns

Having established robust inductions of state rumination, we proceeded to train decoders (i.e.,

linear SVMs) to distinguish between states of induced rumination and distraction using

multivariate EEG patterns, analyzed in one-minute bins across the eight-minute induction

periods. The decoders were trained to classify rumination and distraction from training data

and then predicted the two mental states from independent test data using cross validation,

which allowed to assess the accuracy of decoding.

Decoding accuracy, when averaged **across** all one-minute timepoints, was

significantly greater than the chance level of 50% (one-sided one sample t-test, $t(23) = 1.74$, p

$=.048$, $d = 0.35$ [0.30; 0.40], $bf10 = 1.50$), with a mean accuracy of 54.39% ($SD = 8.76\%$).

Analyzing the accuracies across the bins, we found no significant main effect of timepoint

(linear mixed model for accuracies, $F(8,184) = 1.84$, $p = .073$, $\eta_p = 0.01$). Post-hoc analyses

for each timepoint showed that the decoding accuracies before the induction, as expected, did

not differ from chance level ($p > .05$ and Bayes Factor < 1). Descriptively minutes 1, 2, 6 and

7 showed highest decoding accuracies (Fig. 3 and Tab. S4). In line with these findings,

accuracies from minutes 1, 6 and 7 differed significantly from minute 0 (supplemental

results). However, there was considerable variation in decoding accuracies across participants:

Individual accuracies averaged over timepoints ranged from 27.6% to 82.6%. Decoding

accuracies differed significantly from chance at the individual level for only 10 participants

(Fig. 3; supplemental results Tab. S4). In summary, our decoding results indicate that

multivariate EEG patterns contain weak but statistically significant information distinguishing

between induced ruminative and distractive states.

Classifier based on EEG data

Figure 3. Decoding accuracies of the decoder classifying rumination versus distraction from EEG patterns across the eight-minute induction phase. Decoding accuracies (across-participant mean +/- SEM; individual data dotted in gray) for every one-minute bin and averaged over bins. Chance-level decoding accuracies (black dashed line) were calculated from randomized data (i.e., 1000 random permutations of rumination vs. distraction labels). Asterisks indicate decoding accuracies significantly larger than chance level ($p < .05$, average accuracy tested against chance-level).

3.3 Group decoder

The decoders described thus far were trained using individual-level neurophysiological EEG patterns, which may include both generic patterns that are consistent across participants and patterns specific to individuals. To determine whether these decoders primarily exploit person-specific EEG patterns to predict rumination, we trained a 'group decoder'. This decoder was designed to identify commonalities in the mapping between induced states and EEG patterns across all participants. However, the group decoder achieved a mean decoding accuracy of only 49.59%, which does not differ significantly from the chance level of 50%

($t(23)$ = -0.23, p = .591, $bf10$ = 0.18). Therefore, it appears that the neurophysiological

patterns informative for rumination, as used by our individual decoders, are specific to each

individual.

3.4 Exploratory analysis of the relationship between reported and decoded rumination

Subsequently, we explored the relationship between the neural state of rumination, as

quantified using the decoders' probability estimates of rumination versus distraction, and self-

reported rumination throughout the induction phase. At the group level, we observed that the

decoders' probability estimates for rumination diverged from the 50% mark immediately after

the onset of the induction, coinciding with an increase in VAS-reported rumination during the

rumination induction compared to the distraction induction (Fig. 4a). To analyze this

neurobehavioral link more formally, we examined the relationship between the decoders'

probability estimates and the VAS scale ratings over time for each participant. The *z*-

transformed linear correlations between the VAS values and the probability estimates differed

significantly from zero at the group level ($t(23)$ = 2.57 p = .009, $bf10$ = 6.17; Fig. 4b).

Participant-specific correlations between self-reported and decoded rumination are provided

in supplementary Tab. S5.

Figure 4. *Neurobehavioral link between decoded rumination and reported rumination*. a) Self-

reported rumination measured with the VAS (solid lines) and decoders' probability estimates

for rumination (dashed lines) across timepoints of both induction phases. VAS values, which

range from 1 to 7, were standardized between 0 and 1 (divided through six). b) Self-reported

VAS rumination and decoders' probability estimates over inductions and timepoints from all

individuals. The averaged correlation between self-reported VAS rumination and decoded

rumination ($r = .24$) is shown as an average regression slope.

In contrast to the VAS reports, which account for individual time courses of induced

rumination, the differences observed pre- and post-induction in the PTQ-S scores – when

averaged per induction across sessions and runs – and the mean probability estimates of

rumination for each induction did not show a significant correlation ($r = .10$, $p = .500$).

Overall, our analyses indicated that there was a weak but significant relationship between

decoded and self-reported rumination when taking the participant-specific time courses of

induced ruminations into account.

3.5 Exploratory analyses of the informativeness of neurophysiological patterns

Our decoding results demonstrated that multivariate EEG feature patterns contain information

on ruminative state, and that this information correlates with self-reports. However, it was not

clear whether this information was specifically encoded in patterns of the alpha, beta, or theta

band, or rather in the connectivity between electrodes, as has been suggested by previous

studies on univariate neurophysiological correlates of rumination (Benschop et al., 2021;

Ferdek et al., 2016; Keune et al., 2011; Magosso et al., 2021; Putnam & McSweeney, 2008;

Umemoto et al., 2021). Therefore, we analyzed the feature weights of decoders from

participants whose decoder performance significantly exceeded chance level ($n = 10$

participants with individually significant decoding accuracies). To do so, we chose a 'virtual

lesion' approach, which evaluates the contribution of each feature set (i.e., alpha, beta, and

theta power, as well as connectivity) to overall decoding performance (Kohoutová et al.,

2020). This approach trained 'reduced' decoders on EEG patterns, which involved omitting

one feature set at a time or focusing exclusively on one feature set. We found that the

exclusion and inclusion of connectivity between all channels influenced the decoding of

rumination: The performance of the 'reduced' decoders, which lacked connectivity patterns,

was lower than that of a decoder using all patterns, although not significantly (Tab. 1). The

decoder trained solely on connectivity showed the best decoding performance, although

decoders trained on alpha, beta and theta power also achieved accuracies significantly above

chance level.

Table 1. Results of a virtual lesion approach (Kohoutová et al., 2020) to attribute decoding

performance to feature sets from the alpha, beta, and theta bands, as well as connectivity.

Feature Set	One feature removed for prediction			One feature used for prediction		
	Features used	Accuracy (%)	Test statistics	Features used	Accuracy (%)	Test statistics
Alpha	1890	65.31 (9.34)	$t(9) = -0.81$, $p = .440$, $bf10 = 0.41$	60	60.22 (8.83)	$t(9)= 2.56$, $p = .015$, $bf10 = 5.03$
Beta	1890	66.24 (9.09)	$t(9) = 1.22$, $p = .254$, $bf10 = 0.56$	60	61.78 (10.32)	$t(9) = 3.59$, $p = .003$, $bf10 = 19.06$
Theta	1890	66.32 (8.87)	$t(9) = 1.35$, $p = .211$, $bf10 = 0.63$	60	53.58 (9.12)	$t(9) = 0.95$, $p = .185$, $bf10 = 0.73$
Connectivity	180	61.55 (9.02)	$t(9) = -1.24$, $p = .247$, $bf10 = 0.57$	1770	66.58 (9.34)	$t(9) = 5.42$, $p < .001$, $bf10 > 100$
All together	1950	65.81 (9.14)				

Decoders were trained either by leaving out one feature set or by using solely one feature set.

Performance was averaged across eight minutes and all participants whose decoder

performance significantly exceeded chance level (n = 10). Performance of reduced decoders

was compared to the one encompassing all and those exclusively focusing on one feature with

chance level of 50%. Tests were Bonferroni corrected for four tests. Bf10 represents the Bayes

Factor.

Additionally, we evaluated the importance of distinct features of the

neurophysiological patterns to predict rumination. To do so, we analyzed, at the group level,

whether the SVM feature weights among participants with significant decoding accuracies (*n*

= 10) differed significantly from chance level using a max-stat (Blair & Karniski, 1993)

approach for multiple-comparison correction across features. However, only the beta band

activity in channel FP2, F5, F6 and AF8 and the connectivity between channel CP2 and AF3

showed a significant difference from chance, reinforcing the notion that informative features

were not consistent across participants. Consistent with this finding, individual-level analysis

showed highly variable weight patterns: for example, we found significant results for one

participant in the alpha and beta bands in channel P6, while another participant showed

significant weights in the beta band, especially in the frontal channels (for detailed results, see

supplemental results Tab. S6). Additionally, visual inspection of the feature weight maps (i.e.

importance maps) for the alpha, beta and theta bands, as well as the connectivity feature

weights, suggested substantial differences across participants (Figure 5).

Figure 5. Post-hoc analysis of participant-specific decoder feature weights by channel for

participants whose decoder performance significantly exceeded chance level; SVM weights

for each feature were extracted. These weights were averaged across bins, cross-validation

folds and experimental runs. Absolute weights are presented for each participant for a) alpha,

b) beta, c) theta and, d) connectivity. In a), b) and c), brighter colors indicate higher values.

Please note that feature weight maps were scaled individually. Each color in the connectivity

plot represents one electrode, while the outer circle summarizes electrode subsets. The 20

most informative features from inter-channel connectivity are plotted.

4 Discussion

In this proof-of-principle study, we trained linear decoders to distinguish between states of

experimentally induced rumination and distraction based on EEG patterns comprising alpha,

beta, and theta power, as well as connectivity between electrodes. The inductions led to

significant increases in self-reported rumination as measured in pre versus post assessments

(PTQ-S). Continuous self-reported VAS-rumination showed that high rumination levels were

quickly achieved within one minute and remained stable across the eight-minute induction

period. In contrast, during the distraction induction, VAS-rumination remained at baseline

levels throughout the induction period. Across all participants and timepoints, decoding

performance significantly differed from chance level (i.e., 50%), albeit at overall rather low

decoding accuracies of around 54%. As expected, the EEG patterns did not allow for

differentiation between rumination and distraction during resting state before the inductions.

Highest decoding accuracies occurred at one, six and seven minutes (all above 55.0%) of the

induction period. Post-hoc analyses across all participants and timepoints showed that

continuous self-reported VAS-rumination was significantly correlated with decoded

rumination. A group decoder trained on patterns across all participants did not exceed chance-

level performance. Based on this finding and in line with the notion that rumination patterns

are individually different, we found large variability (27.6% to 82.6%) in decoding accuracies

between participants, with only 10 out of 24 showing decoding performance above chance

level on an individual basis. Additionally, feature weights from the individual significant

decoders varied substantially among participants, indicating that these decoders mapped the

induced mental states to individually specific neurophysiological patterns. When comparing

EEG patterns from different frequency bands and connectivity across participants in more

detail, we found that information on the mental states could be mostly attributed to

connectivity.

701   Although our overall decoder performance for state rumination differed significantly

702 from chance level, the accuracies were slightly lower than those observed in studies decoding

703 trait rumination (Aydın & Akın, 2022; Kim et al., 2023). This difference could stem from

704 several factors other than the fundamental distinctions between mental states and traits. For

705 example, our study trained decoders to distinguish between rumination and the

706 complementary neutral state of distraction (Nolen-Hoeksema, 1991), whereas previous

707 research has compared rumination-related EEG features with those of a positive-affect

708 condition (Ferdek et al., 2016), possibly resulting in more pronounced neurophysiological

709 effects. However, despite incorporating a positive-affect condition in pilot tests, we did not

710 observe higher decoding accuracies (supplemental results Tab. S7). The low decoding

711 accuracies for state rumination in our study might also be attributable to the multiple

712 neurocognitive and affective processes involved in rumination, such as executive functions,

713 attention, memory, and abstract semantic reasoning (Watkins & Roberts, 2020; Whitmer &

714 Gotlib, 2013). During rumination, these neurocognitive processes differ from moment to

715 moment within and between individuals, even though the recursive dynamics may form a

716 stable attractor state on a larger timescale. Although these processes may have clear and

717 consistent neural correlates individually, their variable interplay during state rumination likely

718 results in weak and noisy neural signatures of rumination at a global level. Controlling for

719 these processes experimentally is difficult, even with targeted inductions. Additionally,

720 individuals may respond to rumination through cognitive reappraisal, a distinct regulatory

721 strategy from distraction. However, similar to distraction, reappraisal is expected to reduce

722 negative affect by maintaining attention on ruminative thoughts while actively reinterpreting

723 their content (Gross & John, 2003). However, our findings indicate that negative affect

724 remained consistently higher during rumination inductions compared to distraction, whereas

725 positive affect was consistently lower during rumination relative to distraction.

Our findings of interindividual variability in neurophysiological rumination patterns

indicate that rumination cannot be conceptualized as a stable, canonical brain state. Instead,

the ruminating brain appears to exhibit highly dynamic and individualized activity: Decoding

performance significantly exceeded chance level in only 10 out of 24 participants. This

relatively small proportion of successful individual decoding outcomes suggests that

rumination did not induce strong and reliable neurophysiological signatures within individuals

that could be detected through EEG patterns in the alpha, beta, and theta bands and

connectivity measures. Additionally, we observed high variability in rumination-related EEG

patterns between participants. In the subsample of 10 participants with significant accuracies,

we found at group level only very little state rumination patterns in the beta band and the

inter-channel connectivity as determined by the decoders' feature weights; instead, there were

highly variable weight patterns across individuals. While previous studies have decoded trait

rumination from multivariate patterns across individuals (Aydın & Akın, 2022; Kim et al.,

2023) or identified univariate neural features of state rumination that were consistent across

individuals (e.g., Ferdek et al., 2016), they did not take account of the neural correlates of

rumination within individuals. Our results suggest, however, that the neural basis of state

rumination is highly specific to the individual. For example, we observed significant weights

in the alpha and theta bands over the temporal cortex in one participant, but increased weights

in the beta bands over frontal electrodes in another. Future studies should therefore analyze

how individual brain activation correlates with state rumination, speaking to personalized

approaches in cognitive and clinical neuroscience (Sui, Jiang, Bustillo, & Calhoun, 2020), as

well as computational and precision psychiatry (e.g., Bzdok & Meyer-Lindenberg, 2018;

Passos, Ballester, Rabelo-da-Ponte, & Kapczinski, 2021).

Previous studies have rarely examined the dynamics of state rumination. For example,

it is unclear over what timescales brain dynamics shift from a non-ruminative to a stable

attractor state (Rolls, 2016), a transition that is characteristic of rumination as a recursive

process. Given this gap in the evidence, our study was designed to discern how quickly an

experimental induction might elicit state rumination, which could occur rapidly (e.g., within a

minute) or develop more gradually (e.g., over several minutes). To capture these dynamics,

we trained the decoders using one-minute segments of EEG patterns. Within this temporal

resolution, our results indicate that the process of rumination begins swiftly, noticeable

already within the first minute following the start of induction, but then increases only

gradually, eventually reaching a plateau over the induction period of eight minutes. Decoding

performance over all participants yielded highest results for minutes 6 and 7, during which

time continuously self-reported rumination was also at its peak. Interestingly, our results

indicate that self-reported rumination – both in pre-post measurements and continuous

monitoring – showed differences right from the start of the induction periods. By comparing

only the first induction in each session across participants we were able to demonstrate that

these differences arise due to spill-over effects from the previous induction (Fig. S1). This

finding indicates that rumination is a stable state over timescales of several minutes.

Based on previous EEG studies investigating neural correlates of trait and state

rumination (Benschop et al., 2021; Ferdek et al., 2016; Keune et al., 2011; Magosso et al.,

2021; Putnam & McSweeney, 2008; Umemoto et al., 2021), we decided a priori to focus our

decoding analysis on spectral EEG patterns in the alpha, beta, and theta bands, as well as on

connectivity at the electrode level. Using a 'virtual lesion' approach, we found that patterns of

connectivity in particular, and to some extent alpha, beta and theta power, provided

information on ruminative states. However, our approach did not yield significant decoding

results for 14 of the 24 participants, even though this subsample showed strong increases in

self-reported rumination (Supplemental Fig. S2). This observation suggests that the few EEG

studies that have examined neurophysiological patterns of state rumination (Ferdek et al.,

2016; Putnam & McSweeney, 2008) to date may not have fully captured the

neurophysiological correlates of state rumination in EEG data. Furthermore, the ability to

measure neural signatures of rumination depends on the neuroimaging method. In this study,

we used EEG to identify neural signatures of rumination; however, other techniques like

fMRI might be better at capturing rumination-related neural activity, such as that within the

DMN. For example, fMRI studies (Zhou et al., 2020) have consistently linked the default

mode network with rumination, a type of connectivity rarely measured with EEG (although

there are exceptions; see e.g., Das, de los Angeles, & Menon, 2022). Despite these limitations,

EEG offers advantages in ease of use and affordability compared to fMRI, potentially

facilitating the translation of our findings to field studies and clinical settings. The low

decoding accuracies seen in a majority of our participants may also result from our choice of a

linear SVM decoder. However, our sensitivity analyses indicate that switching to non-linear

SVMs did not improve the decoding results (Tab. S8 and Tab. S9).

Our study has at least five limitations: First, we included only moderately to severely

depressed participants who exhibited robust changes in self-reported rumination during pre-

tests. Thus, our results may not generalize to all individuals with depression, nor to those with

other psychiatric disorders in which rumination is an important psychopathological symptom

(e.g., Aldao et al., 2010). Second, the accuracy of our decoding of rumination may have been

negatively affected by peripheral-physiological signals that were correlated with rumination

and thus contaminated the EEG measurements. However, during preprocessing, we cleaned

the EEG signals of artifacts related to facial muscles, head movements, and eye movements.

Thus, while not absolutely certain, it is highly likely that our decoding results reflect cortical

neurophysiological patterns. Third, we continuously assessed subjective rumination using

VAS scales. Reporting of rumination every minute may have interfered with cognitive

processes, specifically rumination. However, in our proof-of-principle study, we required

subjective data to externally validate the decoding results: Only by simultaneously measuring

rumination with VAS scales, we could correlate the time course of decoded rumination with

subjective rumination. This correlation established an important link between the

psychological and neural levels, showing that decoded rumination is psychologically

meaningful. Moreover, visually presented items such VAS scales are processed within

milliseconds (von der Heydt, 2023) and can quickly be answered. They were applied in the

same fashion in the distraction control condition, so that VAS-response-related cognitive or

motor processing were controlled. Because we have validated decoded rumination, future

studies may use this measure along with or even instead of regular self-reports during the

assessment of state rumination, leading to less potential interferences. Fourth, overall

decoding accuracies were significant but small. Future studies may increase decoding

accuracies. Nevertheless, also small decoding accuracies can help to measure rumination in

clinical studies that investigate how interventions affect rumination (Welkerling et al., 2024).

Fifth, the large number of features, especially for the connectivity, may have led to variable

feature weight maps because individual features are not informed by sufficient data sizes.

Therefore, post-hoc analyses regarding the feature weight maps have to be interpreted with

caution.

To conclude, our study has shown that neurophysiological spectral and connectivity

EEG patterns represent information about individual state rumination. Although the

information was weak and encoded in highly variable patterns across individuals, this proof-

of-principle study demonstrates that individual dynamic state rumination can be decoded from

neurophysiological patterns on a timescale of minutes. These findings may lay the

groundwork for future studies investigating the dynamic evolution of neurocognitive and

affective processes during state rumination.

Declaration of Generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used ChatGPT (version 4) and DeepL write in

order, to improve the readability and clarity of the manuscript. After using this tool/service,

the authors reviewed and edited the content as needed and take full responsibility for the

content of the publication.

831   **References**

832   Aldao, A., Nolen-Hoeksema, S., & Schweizer, S. (2010). Emotion-regulation strategies across

833   psychopathology: A meta-analytic review. *Clinical Psychology Review, 30*(2), 217–

834   237. doi:10.1016/j.cpr.2009.11.004

835   Aydın, S., & Akın, B. (2022). Machine learning classification of maladaptive rumination and

836   cognitive distraction in terms of frequency specific complexity. *Biomedical Signal*

837   *Processing and Control, 77*, 103740. doi:10.1016/j.bspc.2022.103740

838   Beck, A. T., Steer, R. A., & Brown, G. K. (1996). Manual for the beck depression inventory-

839   II. *San Antonio, TX: Psychological Corporation, 1*(82), 10.1037.

840   Beesdo-Baum, K., Zaudig, M., & Wittchen, H.-U. (2019). *Strukturiertes Klinisches Interview*

841   *für DSM-5®-Störungen – Klinische Version (SCID-5-CV)*. Göttingen: Hogrefe.

842   Benschop, L., Poppa, T., Medani, T., Shahabi, H., Baeken, C., Leahy, R. M., . . .

843   Vanderhasselt, M.-A. (2021). Electrophysiological scarring in remitted depressed

844   patients: Elevated EEG functional connectivity between the posterior cingulate cortex

845   and the subgenual prefrontal cortex as a neural marker for rumination. *Journal of*

846   *Affective Disorders, 281*, 493-501. doi:10.1016/j.jad.2020.12.081

847   Blair, R. C., & Karniski, W. (1993). An alternative method for significance testing of

848   waveform   difference   potentials.   *Psychophysiology,   30*(5),   518-524.

849   doi:10.1111/j.1469-8986.1993.tb02075.x

850   Bradley, M. M., & Lang, P. J. (1994). Measuring emotion: The self-assessment manikin and

851   the semantic differential. *Journal of Behavior Therapy and Experimental Psychiatry,*

852   *25*(1), 49-59. doi:10.1016/0005-7916(94)90063-9

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spat Vis, 10*(4), 433-436.

Bzdok, D., & Meyer-Lindenberg, A. (2018). Machine Learning for Precision Psychiatry:

Opportunities and Challenges. *Biological Psychiatry: Cognitive Neuroscience and*

Neuroimaging, 3(3), 223-230. doi:https://doi.org/10.1016/j.bpsc.2017.11.007

Chang, C.-C., & Lin, C.-J. (2011). LIBSVM: A library for support vector machines. *ACM*

Transactions on Intelligent Systems and Technology, 2(3), Article 27.

doi:10.1145/1961189.1961199

Chiang, Y. H., Ma, Y. C., Lin, Y. C., Jiang, J. L., Wu, M. H., & Chiang, K. C. (2022). The

Relationship between Depressive Symptoms, Rumination, and Suicide Ideation in

Patients with Depression. *International Journal of Environmental Research and*

Public Health, 19(21). doi:10.3390/ijerph192114492

Cohen, J. (1973). Eta-Squared and Partial Eta-Squared in Fixed Factor Anova Designs.

Educational and Psychological Measurement, 33(1), 107-112.

doi:10.1177/001316447303300111

Das, A., de los Angeles, C., & Menon, V. (2022). Electrophysiological foundations of the

human default-mode network revealed by intracranial-EEG recordings during resting-

state and cognition. *NeuroImage, 250*, 118927.

doi:https://doi.org/10.1016/j.neuroimage.2022.118927

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-

trial EEG dynamics including independent component analysis. *Journal of*

Neuroscience Methods, 134(1), 9-21. doi:10.1016/j.jneumeth.2003.10.009

Devynck, F., Rousseau, A., & Romo, L. (2019). Does Repetitive Negative Thinking Influence

Alcohol Use? A Systematic Review of the Literature. *Frontiers in Psychology, 10*.

doi:10.3389/fpsyg.2019.01482

Ehring, T., Fuchs, N., & Kläsener, I. (2009). The effects of experimentally induced rumination

versus distraction on analogue posttraumatic stress symptoms. *Behavior Therapy,*

40(4), 403–413. doi:10.1016/j.beth.2008.10.001

Ehring, T., Zetsche, U., Weidacker, K., Wahl, K., Schönfeld, S., & Ehlers, A. (2011). The

881   Perseverative Thinking Questionnaire (PTQ): validation of a content-independent

882   measure of repetitive negative thinking. *Journal of Behavior Therapy and*

883   *Experimental Psychiatry, 42*(2), 225-232. doi:10.1016/j.jbtep.2010.12.003

884   Eshun, S. (2000). Role of Gender and Rumination in Suicide Ideation: A Comparison of

885   College Samples From Ghana and the United States. *Cross-Cultural Research, 34*(3),

886   250-263. doi:10.1177/106939710003400303

887   Fang, L., Marchetti, I., Hoorelbeke, K., & Koster, E. H. W. (2019). Do daily dynamics in

888   rumination and affect predict depressive symptoms and trait rumination? An

889   experience sampling study. *Journal of Behavior Therapy and Experimental*

890   *Psychiatry, 63*, 66-72. doi:https://doi.org/10.1016/j.jbtep.2018.11.002

891   Ferdek, M. A., van Rijn, C. M., & Wyczesany, M. (2016). Depressive rumination and the

892   emotional control circuit: An EEG localization and effective connectivity study.

893   *Cognitive, affective & behavioral neuroscience, 16*(6), 1099–1113.

894   doi:10.3758/s13415-016-0456-x

895   GBD 2019 Mental Disorders Collaborators. (2022). Global, regional, and national burden of

896   12 mental disorders in 204 countries and territories, 1990-2019: a systematic analysis

897   for the Global Burden of Disease Study 2019. *The Lancet Psychiatry, 9*(2), 137-150.

898   doi:10.1016/S2215-0366(21)00395-3

899   Gross, J. J., & John, O. P. (2003). Individual differences in two emotion regulation processes:

900   implications for affect, relationships, and well-being. *J Pers Soc Psychol, 85*(2), 348-

901   362. doi:10.1037/0022-3514.85.2.348

902   Horato, N., Quagliato, L. A., & Nardi, A. E. (2022). The relationship between emotional

903   regulation and hemispheric lateralization in depression: a systematic review and a

904   meta-analysis. *Translational Psychiatry, 12*(1), 162. doi:10.1038/s41398-022-01927-9

905 Huffziger, S., & Kuehner, C. (2009). Rumination, distraction, and mindful self-focus in

906     depressed patients. *Behavior Research and Therapy, 47*(3), 224–230.

907     doi:10.1016/j.brat.2008.12.005

908 Jasper, H. H. (1958). The ten-twenty electrode system of the International Federation.

909     *Electroencephalography and Clinical Neurophysiology, 10*, 370–375.

910     doi:10.1016/0013-4694(58)90053-1

911 Keune, P. M., Bostanov, V., Hautzinger, M., & Kotchoubey, B. (2011). Mindfulness-based

912     cognitive therapy (MBCT), cognitive style, and the temporal dynamics of frontal EEG

913     alpha asymmetry in recurrently depressed patients. *Biological psychology, 88*(2-3),

914     243–252. doi:10.1016/j.biopsycho.2011.08.008

915 Kim, J., Andrews-Hanna, J. R., Eisenbarth, H., Lux, B. K., Kim, H. J., Lee, E., . . . Woo, C.-

916     W. (2023). A dorsomedial prefrontal cortex-based dynamic functional connectivity

917     model of rumination. *Nature Communications, 14*(1), 3540. doi:10.1038/s41467-023-

918     39142-9

919 Kohoutová, L., Heo, J., Cha, S., Lee, S., Moon, T., Wager, T. D., & Woo, C.-W. (2020).

920     Toward a unified framework for interpreting machine-learning models in

921     neuroimaging. *Nature Protocols, 15*(4), 1399-1435. doi:10.1038/s41596-019-0289-5

922 Krekelberg, B. (2024). klabhub/bayesFactor: Bayes only (v3.0). *Zenodo.*

923     doi:https://doi.org/10.5281/zenodo.13744717

924 Kriegeskorte, N., Simmons, W. K., Bellgowan, P. S. F., & Baker, C. I. (2009). Circular

925     analysis in systems neuroscience: the dangers of double dipping. *Nature Neuroscience,*

926     *12*(5), 535-540. doi:10.1038/nn.2303

927 Kuyken, W., Watkins, E., Holden, E., & Cook, W. (2006). Rumination in adolescents at risk

928     for depression. *Journal of Affective Disorders, 96*(1), 39-47.

929     doi:10.1016/j.jad.2006.05.017

Laicher, H., Int-Veen, I., Torka, F., Kroczek, A., Bihlmaier, I., Storchak, H., . . . Rosenbaum,

D. (2022). Trait rumination and social anxiety separately influence stress-induced

rumination and hemodynamic responses. *Scientific Reports, 12*(1), 5512.

doi:10.1038/s41598-022-08579-1

Magosso, E., Ricci, G., & Ursino, M. (2021). Alpha and theta mechanisms operating in

internal-external attention competition. *Journal of Integrative Neuroscience, 20*(1), 1-

19. doi:10.31083/j.jin.2021.01.422

National Kidney Foundation. (2002). K/DOQI Clinical Practice Guidelines for Chronic

Kidney Disease: Evaluation, Classification and Stratification. *American Journal of*

Kidney Diseases, 39, S1-S266.

Nolen-Hoeksema, S. (1991). Responses to depression and their effects on the duration of

depressive episodes. *Journal of Abnormal Psychology, 100*(4), 569-582.

doi:10.1037//0021-843x.100.4.569

Nolen-Hoeksema, S., & Morrow, J. (1991). A prospective study of depression and

posttraumatic stress symptoms after a natural disaster: the 1989 Loma Prieta

Earthquake. *Journal of Personality and Social Psychology, 61*(1), 115-121.

doi:10.1037//0022-3514.61.1.115

Nolen-Hoeksema, S., Wisco, B. E., & Lyubomirsky, S. (2008). Rethinking Rumination.

Perspectives on Psychological Science, 3(5), 400-424. doi:10.1111/j.1745-

6924.2008.00088.x

Passos, I. C., Ballester, P., Rabelo-da-Ponte, F. D., & Kapczinski, F. (2021). Precision

Psychiatry: The Future Is Now. *The Canadian Journal of Psychiatry, 67*(1), 21-25.

doi:10.1177/0706743721998044

Pereira, F., Mitchell, T., & Botvinick, M. (2009). Machine learning classifiers and fMRI: A

tutorial overview. *NeuroImage, 45*(1, Supplement 1), S199-S209.

doi:https://doi.org/10.1016/j.neuroimage.2008.11.007

Putnam, K. M., & McSweeney, L. B. (2008). Depressive symptoms and baseline prefrontal

EEG alpha activity: a study utilizing Ecological Momentary Assessment. *Biological*

Psychology, 77(2), 237-240. doi:10.1016/j.biopsycho.2007.10.010

R Core Team. (2017). R: A language and environment for statistical computing. Vienna: R

Foundation for Statistical Computing.

Rolls, E. T. (2016). A non-reward attractor theory of depression. *Neuroscience and*

Biobehavioral Reviews, 68, 47-58. doi:https://doi.org/10.1016/j.neubiorev.2016.05.007

Scheinost, D., Noble, S., Horien, C., Greene, A. S., Lake, E. M. R., Salehi, M., . . . Constable,

R. T. (2019). Ten simple rules for predictive modeling of individual differences in

neuroimaging.   *NeuroImage,*   *193*,   35-45.

doi:https://doi.org/10.1016/j.neuroimage.2019.02.057

Sui, J., Jiang, R., Bustillo, J., & Calhoun, V. (2020). Neuroimaging-based Individualized

Prediction of Cognition and Behavior for Mental Disorders and Health: Methods and

Promises.   *Biological*   *Psychiatry,*   *88*(11),   818-828.

doi:https://doi.org/10.1016/j.biopsych.2020.02.016

Tadel, F., Baillet, S., Mosher, J. C., Pantazis, D., & Leahy, R. M. (2011). Brainstorm: A User-

Friendly Application for MEG/EEG Analysis. *Computational Intelligence and*

Neuroscience, 2011, 879716. doi:10.1155/2011/879716

Teasdale, G., & Jennett, B. (1974). ASSESSMENT OF COMA AND IMPAIRED

CONSCIOUSNESS: A Practical Scale. *The Lancet, 304*(7872), 81-84.

doi:https://doi.org/10.1016/S0140-6736(74)91639-0

The MathWorks Inc. (2020). MATLAB (Version version 9.9.0 (R2020b)). Natick,

Masachusetts.

Timm, C., Ubl, B., Zamoscik, V., Ebner-Priemer, U., Reinhard, I., Huffziger, S., . . . Kuehner,

C. (2017). Cognitive and affective trait and state factors influencing the long-term

symptom course in remitted depressed patients. *PloS One, 12*(6), e0178759.

doi:10.1371/journal.pone.0178759

Umemoto, A., Panier, L. Y. X., Cole, S. L., Kayser, J., Pizzagalli, D. A., & Auerbach, R. P.

(2021). Resting posterior alpha power and adolescent major depressive disorder.

Journal of Psychiatric Research, 141, 233-240. doi:10.1016/j.jpsychires.2021.07.003

van der Meer, J. N., Pampel, A., Van Someren, E. J. W., Ramautar, J. R., van der Werf, Y. D.,

Gomez-Herrero, G., . . . Walter, M. (2016). Carbon-wire loop based artifact correction

outperforms post-processing EEG/fMRI corrections—A validation of a real-time

simultaneous EEG/fMRI correction method. *NeuroImage, 125*, 880-894.

doi:10.1016/j.neuroimage.2015.10.064

von der Heydt, R. (2023). Visual cortical processing—From image to object representation.

Frontiers in Computer Science, 5, 1136987.

Watkins, E. R., & Roberts, H. (2020). Reflecting on rumination: Consequences, causes,

mechanisms and treatment of rumination. *Behaviour Research and Therapy, 127*,

103573. doi:10.1016/j.brat.2020.103573

Welkerling, J., Niess, A., Schneeweiss, P., Sudeck, G., Rohe, T., & Wolf, S. (2024). Preprint:

Single bout of exercise reduces self-reported and decoded rumination in favor of

distraction in patients with major depression. doi:https://doi.org/10.31219/osf.io/rhg3s

Whitmer, A. J., & Gotlib, I. H. (2013). An attentional scope model of rumination.

Psychological Bulletin, 139(5), 1036-1061. doi:10.1037/a0030923

Zhou, H.-X., Chen, X., Shen, Y.-Q., Li, L., Chen, N.-X., Zhu, Z.-C., . . . Yan, C.-G. (2020).

Rumination and the default mode network: Meta-analysis of brain imaging studies and

implications for depression. *NeuroImage, 206*, 116287.

doi:https://doi.org/10.1016/j.neuroimage.2019.116287

1005