Full title: Comparing the Value of Left Atrial Strain and HFA-PEFF Score in Diagnosing Heart Failure with Preserved Ejection Fraction in Vietnamese Population: A Cross-Sectional Study

Short Title: Left Atrial Strain in Heart Failure With Preserved Ejection Fraction

Hai Dang Nguyen Ngoc[1][¶], Thang Viet Luong[2][¶], Bang Hai Ho[2], Tien Anh

Hoang[2][*]

[1]The Faculty of Medicine, Duy Tan University, Da Nang, Viet Nam

[2]Department of Internal Medicine, Hue University of Medicine and Pharmacy,

Hue, Viet Nam

*Corresponding author:

Email: hatien@huemed-univ.edu.vn or hatien@hueuni.edu.vn

¶Hai Nguyen Ngoc Dang and Thang Viet Luong contributed equally to this work.

Author contributions:

Hai Nguyen Ngoc Dang: Conceptualization, Methodology, Acquisition,

Investigation, Data curation, Writing – original draft, Writing – review &

editing.

Thang Viet Luong: Conceptualization, Methodology, Acquisition, Investigation,

Data curation, Writing – original draft, Writing – review & editing.

Bang Hai Ho: Conceptualization, Acquisition, Writing – original draft.

Tien Anh Hoang: Conceptualization, Acquisition, Writing – original draft.

All authors have read and agreed to the published version of the manuscript.

Abstract

Background: Heart failure with preserved ejection fraction (HFpEF) has

a high hospitalization rate. While recent guidelines recommend specific

parameters like E/e', e' velocity, and left atrial volume index (LAVI) for

diagnosing HFpEF, their clinical accuracy remains limited. Left atrial (LA)

strain has emerged as a potential diagnostic parameter, yet its role in the

Vietnamese population is unclear. This study aims to evaluate LA strain's

diagnostic role in HFpEF among Vietnamese patients, exploring its relationship

with established parameters of left ventricle (LV) diastolic function to determine

its potential utility as a diagnostic tool.

Methods: A cross-sectional study was conducted from 15/04/2022 to

01/12/2023, involving 118 subjects, including 49 patients with HFpEF and 69

individuals without cardiac dysfunction. The study subjects were evaluated for

LA strain and HFA-PEFF.

Results: LA strain including LA reservoir (LASr), conduit (LAScd), and contractile (LASct) function, in the HFpEF group were 19.97% ± 9.18%, 9.08% ± 6.18%, and 10.89% ± 5.16%, respectively. The control group had corresponding LASr, LAScd, and LASct values of 34.70% ± 5.38%, 17.33% ± 5.72%, and 17.38% ± 4.41% (p < 0.001). The area under the curve (AUC) for LASr, LAScd, LAScr, HFA-PEFF, LAVI, and GLS to diagnose HFpEF was 0.877, 0.836, 0.749, 0.975, 0.587, and 0.717, respectively. Comparing the AUCs for diagnosing HFpEF between LASr and HFA-PEFF, no difference was found with p = 0.062.

Conclusion: LA strain has a diagnostic value equivalent to the HFA-PEFF score in diagnosing HFpEF. These indices could be incorporated into the existing guidelines to enhance the diagnosis of HFpEF.

Key words: Speckle tracking echocardiography, left atrial, strain, heart failure, left ventricular ejection fraction

Introduction

Heart failure with preserved ejection fraction (HFpEF) is defined as heart failure with an ejection fraction of 50% or higher at diagnosis, affecting approximately 32 million people worldwide. Patients with HFpEF are hospitalized about 1.4 times per year and have an annual mortality rate of around 15% [1]. Left ventricular (LV) diastolic dysfunction plays a fundamental role, overarching in the pathophysiology of HFpEF [2].

The recent recommendations of the American Society of Echocardiography/European Society of Cardiovascular Imaging in 2016 use parameters such as E/e', septal e' velocity, lateral e' velocity, maximum tricuspid valve regurgitation flow (TRV) and left atrial volume index (LAVI) to evaluate LV diastolic dysfunction as well as to diagnose HFpEF [3]. However, in clinical practice, the recommendations parameters have limited precision in diagnosing HFpEF [4].

The assessment of left atrial (LA) function has recently gained prominence as a critical parameter in cardiac evaluation [5]. The LA function encompasses three primary aspects: blood storage (reservoir function), blood conduction (conduit function), and ejection function (contractile function) [6]. Commonly used indices to assess LA function include LAVI and LA size. Increased LAVI is associated with prolonged chronic LV/LA pressure overload. However, LA size takes time to change, often significantly dilating in later stages, making LAVI less sensitive in the early stages [7].

LA strain is a novel echocardiographic technique that comprehensively evaluates reservoir, conduit, and contractile functions. This method proves particularly valuable when changes are subtle and challenging to detect using conventional parameters such as LA dimensions and LAVI [8]. While LA dimensions have been previously utilized, the role of LA function as a biomarker is increasingly under evaluation, both independently and in conjunction with LA size. LA strain serves as a tool to assess LA function and can be measured

throughout the cardiac cycle, enabling a comprehensive evaluation of LA reservoir, conduit, and contractile functions [7]. Additionally, LA strain offers the advantage of being a technique mostly independent of angle and less susceptible to influences from mitral annulus calcification and bundle branch block effects [9].

Notably, impaired LA strain has been observed in HFpEF patients, indicating its potential diagnostic value [10–12]. Studies conducted in the United Kingdom and China have demonstrated the utility of LA strain in assessing and diagnosing HFpEF [9,13]. However, its role in the Vietnamese population remains unexplored. Therefore, this study aims to evaluate the diagnostic role of LA strain in HFpEF among Vietnamese individuals, contributing to a deeper understanding of its applicability in clinical practice.

Materials and methods

Ethics approval statement

The study received approval from The Institutional Ethics Committee of Hue University of Medicine and Pharmacy (Approval number: H2022/034) and adhered to the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline, conducted under the principles outlined in the Declaration of Helsinki 2013. Prior to participation, written informed consent was obtained from each patient or their legal representative.

Study population

During the period from 15/04/2022 to 01/12/2023, a cross-sectional study was undertaken. The study population consisted of adults aged 18 and above, who provided informed consent prior to their inclusion. A total of 118 subjects were recruited, comprising 49 individuals diagnosed with HFpEF (Heart Failure with preserved Ejection Fraction) in the disease group, and 69 individuals without cardiac dysfunction in the control group. These participants sought examination and treatment at Hue University Hospital within the aforementioned timeframe. Patients with HFpEF are evaluated according to the standards of AHA/ACC/HFSA in 2022 [4]: (1) Symptoms of heart failure (pulmonary congestion or systemic congestion); (2) Normal LV ejection fraction LVEF ≥ 50%; (3) Objective evidence of structural and/or functional cardiac abnormalities consistent with LV diastolic dysfunction; (4) NT-proBNP ≥ 125 pg/mL. The four recommended variables for identifying diastolic dysfunction and their abnormal cutoff values are septal $e' < 7$ cm/s or lateral $e' < 10$ cm/s, average E/e' ratio > 14, LAVI > 34 mL/m^2, and TRV > 2.8 m/s. LV diastolic dysfunction was considered present if more than half of the available parameters met these cutoff values. In cases where half of the parameters met the cutoff values (indeterminate), diastolic dysfunction was assumed to be present if patients had E/A > 2.0 or if ≥ 2 of the three parameters (E/e' ratio, TRV, and LAVI) met the cutoff values [14]. Exclusion criteria included patients who declined participation, severe valvular heart disease, heart failure with EF < 50%, and arrhythmias. Patients with unclear echocardiography images or images lacking clear visualization of the myocardial endocardial layer were also excluded. The

control group comprised 69 healthy adults undergoing health screening with no history of heart failure.

Clinical data collection, laboratory tests, and transthoracic echocardiogram

The clinical data collected included personal and family medical histories, as well as clinical variables obtained through direct interviews and medical records. N-Terminal Pro-B-Type Natriuretic Peptide (NT-proBNP) levels were measured using a Cobas 8000 analyzer.

The echocardiographic procedure followed the guidelines of the American Society of Echocardiography for performing a comprehensive transthoracic echocardiographic examination in adults. The machine recorded the electrocardiogram alongside the echocardiographic images during the echocardiography procedure. All echocardiograms included in the final data analysis were performed on patients with normal sinus rhythm [15].

Left atrial and left ventricular strain analysis

Echocardiography images in DICOM format, meeting acceptable image quality standards, were uploaded to Philips QLAB Cardiovascular ultrasound quantification software Cardiac Analysis version 15. LA strain assessment was conducted in both the two-chamber and four-chamber views, with reference points set at the onset of the P wave in the cardiac cycle. Measurements of LA strain were

acquired during the reservoir, conduit, and contractile phases of LA function,

designated as LA strain reservoir function (LASr), LA strain conduit function

(LAScd), and LA strain contractile function (LASct), respectively. For LV strain

analysis, endocardial borders were traced on the end-systolic frame in three apical

views (4-chamber, 2-chamber, and 3-chamber), with end-systole defined by the

QRS complex or as the smallest LV volume during the cardiac cycle. The software

tracked speckles along the endocardial border and myocardium throughout the

cardiac cycle, automatically computing peak longitudinal strain and generating

regional data from six segments, as well as an average value for each view. For

patients in sinus rhythm, analyses were performed on a single cardiac cycle; for

those in atrial fibrillation, strain values were calculated as the average of three

cardiac cycles [16]. All strain measurements were performed by one strain

specialist in the core laboratory who was blinded to the patients' other data.

Results of LV and LA strain are conventionally represented as negative

values. However, for convenience in analysis and display, we utilized the absolute

values of these results. The detailed methodology is illustrated in **Fig 1**.

Fig 1. Parameters of LA strain on speckle tracking echocardiography. LASr

(Reservoir) in red, LAScd (Conduit) in green, and LASct (Contractile) in yellow.

LAScd: left atrial conduit function; LASct: left atrial contractile function; LASr:

left atrial reservoir function.

Calculation of the Heart Failure Association-PEFF Score

The Heart Failure Association-PEFF (HFA-PEFF) score comprises

functional, morphological, and biomarker domains (**Supplementary**). A patient

can score zero, minor (1 point), or major (2 points) for each domain, and then those

subscores are summed to produce a total score that ranges from 0 to 6 points. The

total score is classified as low likelihood (0 – 1 point), intermediate likelihood (2 –

4 points), and high likelihood (5 – 6 points) [17].

Statistics

All statistical analyses were conducted using SPSS Version 26 (IBM, New

York, United States), MedCalc Software Version 22.019 (MedCalc Software,

Ostend, Belgium), and GraphPad Prism Version 10 (GraphPad Software, Boston,

United States). Continuous variables were presented as mean ± standard deviation

for normally distributed variables, as determined by the Kolmogorov-Smirnov test

and the Shapiro-Wilk test. Non-normally distributed variables were expressed as

median values with interquartile ranges (25^{th}-75^{th} percentile). Categorical

variables were reported as frequencies and percentages. Intergroup differences in

categorical variables were assessed using Fisher's exact test, while differences in

continuous variables were analyzed using the unpaired T-test or the Mann-Whitney

U test, as appropriate.

Missing data were excluded from the analyses. Pearson's correlation

coefficient (r) and its corresponding p-value were calculated to elucidate the

correlation between continuous variables. The area under the curve (AUC) was

determined using the Wilson/Brown method to predict HFpEF. AUC comparison

was conducted to assess the diagnostic value of strain compared to existing

guideline criteria, employing the DeLong method [18]. All statistical tests were

two-sided, with a P-value of < 0.05 considered significant.

Results

The study was conducted on 118 subjects, of whom 49 patients with HFpEF

and 69 people qualified as the control group, meeting the disease selection criteria

according to the research criteria. Our research obtained the following results:

Table 1 displays the mean age of the HFpEF and control groups as $71.45 \pm$

13.92 and 60.39 ± 6.58, respectively. The average body mass index (BMI) for the

case and control group was $21.92 \pm 2.03 \, \text{kg/m}^2$ and $21.71 \pm 3.54 \, \text{kg/m}^2$. The results

indicated no statistically significant differences in BMI between the two study

groups ($p > 0.05$). The NT-proBNP concentration in the HFpEF group exhibited a

non-normal distribution, with a median of 663.20 pg/mL (Quartile: 286.55 pg/mL

$- 1417.00$ pg/mL).

Table 1. General Characteristics of Study Subjects.

Characteristics	Control group (n = 69)	HFpEF group (n = 49)	p-value
Baseline demographic and clinical features			
Age (years)	60.39 ± 6.58	71.45 ± 13.92	< 0.001

Female	37 (53.62)	27 (55.10)	0.512
BSA (m²)	1.55 ± 0.13	1.52 ± 0.16	0.357
BMI (kg/m²)	21.92 ± 2.03	21.71 ± 3.54	0.710
Heart rate (bpm)	74.94 ± 10.92	80.47 ± 15.11	0.032
SBP (mmHg)	121.09 ± 8.17	130.43 ± 13.94	< 0.001
DBP (mmHg)	74.49 ± 7.48	76.10 ± 9.08	0.295
Hypertension	0	26 (53.06)	< 0.001
Diabetes	0	8 (16.33)	< 0.001
Coronary disease	0	29 (59.18)	< 0.001
NT-ProBNP (pg/mL)	64.00 [43.50 - 77.95]	663.20 [286.55 - 1417.00]	< 0.001
Left ventricular structure and function			
LVMI (g/m²)	96.75 ± 24.48	130.45 ± 35.66	< 0.001
LVIDd (cm)	4.61 ± 0.48	5.02 ± 0.68	0.001
LVIDs (cm)	2.77 ± 0.38	3.29 ± 0.63	< 0.001
IVSd (cm)	0.94 ± 0.15	1.07 ± 0.19	< 0.001
IVSs (cm)	1.28 ± 0.15	1.41 ± 0.20	< 0.001
LVPWd (cm)	0.92 ± 0.14	0.99 ± 0.20	0.023
LVPWs (cm)	1.37 ± 0.15	1.46 ± 0.21	0.013
RWT	0.400 ± 0.067	0.403 ± 0.098	0.891
LV EF (%)	70.22 ± 6.38	63.38 ± 8.22	< 0.001
FS (%)	39.90 ± 5.22	35.27 ± 6.68	< 0.001

LV GLS (%)	20.67 ± 2.44	15.95 ± 4.76	< 0.001
Doppler echocardiography			
E velocity (cm/s)	65.61 ± 14.21	84.52 ± 25.12	0.001
A velocity (cm/s)	80.86 ± 14.34	93.79 ± 30.71	0.041
e' septal velocity (cm/s)	7.35 [6.29 - 7.93]	6 [4.57 - 7.38]	0.003
e' lateral velocity (cm/s)	8.80 [8.10 - 10.70]	7.11 [5.32 - 9.38]	0.008
E/A ratio	0.78 [0.69 - 0.93]	0.87 [0.65 - 1.24]	0.249
Average E/e ratio	7.96 ± 1.98	13.15 ± 8.12	0.002
TRV (cm/s)	129.45 ± 44.43	244.73 ± 115.50	< 0.001
Left atrial structure and function			
LAd (cm)	3.07 ± 0.48	3.55 ± 0.62	< 0.001
LAVI (mL/m²)	18.30 [16.70 - 22.15]	24.00 [16.65 - 33.50]	0.002
LASr (%)	34.70 ± 5.38	19.97 ± 9.18	< 0.001
LASct (%)	17.33 ± 5.72	9.08 ± 6,18	< 0.001
LAScd (%)	17.38 ± 4.41	10.89 ± 5.16	< 0.001
LASr/LAVI	1.89 ± 0.53	0.95 ± 0.89	< 0.001
LASr/E/e'	4.71 ± 1.42	2.13 ± 1.29	< 0.001
HFA-PEFF score			
HFA-PEFF score	2 [2 − 3]	4 [3 − 5]	<0.001

195   Values are presented as mean ± standard deviation or number (%) or median [IQR,

25th-75th percentile]. A: A-wave velocity; BMI: body mass index; BSA: body surface area; DBP: diastolic blood pressure; E: E-wave velocity; e': e'-wave velocity; HFA-PEFF: Heart Failure Association-PEFF; IVSd: interventricular septum thickness in diastole; IVSs: interventricular septum thickness in systole; Lad: left atrial diameter; LAScd: left atrial conduit function; LASct: left atrial contractile function; LASr: left atrial reservoir function; LAVI: left atrial volume index; LV EF: left ventricular ejection fraction; LV GLS: left ventricular global longitudinal strain; LVMI: left ventricular mass index; LVIDd: left ventricular internal diameter in diastole; LVIDs: left ventricular internal diameter in systole; LVPWd: left ventricular posterior wall thickness in diastole; LVPWs: left ventricular posterior wall thickness in systole; NT-proBNP: N-terminal pro-B-type natriuretic peptide; RWT: relative wall thickness; SBP: systolic blood pressure; TRV: tricuspid regurgitation velocity.

Additionally, LV structural and functional indices, including LVMI, RWT, and LV EF of the HFpEF group were 130.45 ± 35.66 g/m^2, 0.403 ± 0.098, and $63.38 \pm 8.22\%$, respectively; while those of the control group were 96.75 ± 24.48 g/m^2, 0.400 ± 0.067, and $70.22 \pm 6.38\%$, respectively. On echocardiography, LA strain indices in the HFpEF group were lower than in the control group, with all differences being statistically significant (< 0.001). Further details are illustrated in **Table 1** and **Fig 2**.

Fig 2. The HFA-PEFF score, LASr, LAScd, and LASct of both HFpEF and control groups. LA functional indices including LASr, LAScd, and LASct in the

control group are higher than those in the HFpEF group. Additionally, HFpEF

patients exhibit higher HFA-PEFF scores with a median of 4. HFA-PEFF: Heart

Failure Association-PEFF; LAScd: left atrial conduit function; LASct: left atrial

contractile function; LASr: left atrial reservoir function; ***p < 0.001.

Additionally, **Fig 3** depicts the correlation between echocardiography

indices, NT-proBNP, and HFA-PEFF. LASr, LAScd, and LASct showed an inverse

correlation with HFA-PEFF score and NT-proBNP, while also correlating with

measured indices of cardiac function. Detailed parameters are demonstrated in **Fig**

3.

Fig 3. Heatmap depicts the correlation between echocardiography indices, NT-

ProBNP, and HFA-PEFF. A: A-wave velocity; E: E-wave velocity; e': e'-wave

velocity; HFA-PEFF: Heart Failure Association-PEFF; LAScd: left atrial conduit

function; LASct: left atrial contractile function; LASr: left atrial reservoir function;

LAVI: left atrial volume index; LV GLS: left ventricular global longitudinal strain;

NT-proBNP: N-terminal pro-B-type natriuretic peptide; TRV: tricuspid

regurgitation velocity.

Table 2 and **Fig 4** illustrate that the indices LASr (AUC = 0.877), LAScd

(AUC = 0.836), LASct (AUC = 0.749), HFA-PEFF score (AUC = 0.975) exhibit

high accuracy in diagnosing HFpEF. The cutoff points, sensitivity, and specificity

of LASr are 29.93%, 82.61%, and 91.38%, respectively. Further detailed

information is presented in **Table 2** and **Fig 4**.

Table 2. The performance of the LA strain parameters and existing criteria to discriminate HfpEF.

Parameters	AUC	95%Cl		p-value	Cut-off point	Sensitivity (%)	Specificity (%)
LAVI (mL/m²)	0.587	0.440	0.734	0.224	34	17.39	100.00
Average E/e'	0.578	0.432	0.724	0.274	14	17.39	98.26
Septal e' velocity (cm/s)	0.615	0.478	0.751	0.109	7	60.87	62.07
TRVmax (m/s)	0.565	0.419	0.711	0.362	2.8	13.04	100.00
LV GLS (%)	0.717	0.576	0.836	< 0.001	16	43.48	100.00
LASr (%)	0.877	0.776	0.979	< 0.001	29.93	82.61	91.38
LAScd (%)	0.836	0.723	0.949	< 0.001	11.63	69.57	91.38
LASct (%)	0.749	0.618	0.880	0.001	12.35	56.52	87.93
HFA-PEFF	0.975	0.942	1.000	< 0.001	5	56.52	100.00
LASr/LAVI	0.850	0.736	0.965	< 0.001	1.23	69.57	96.55
LASr/E/e'	0.894	0.809	0.978	< 0.001	3.12	82.61	89.66

E: E-wave velocity; e': e'-wave velocity; HFA-PEFF: Heart Failure Association-PEFF; LAScd: left atrial conduit function; LASct: left atrial contractile function;

LASr: left atrial reservoir function; LAVI: left atrial volume index; LV GLS: left ventricular global longitudinal strain; TRV: tricuspid regurgitation velocity.

Fig 4. Receiver operating characteristic curves for predicting HFpEF. E: E-wave velocity; e': e'-wave velocity; HFA-PEFF: Heart Failure Association-PEFF; LAScd: left atrial conduit function; LASct: left atrial contractile function; LASr: left atrial reservoir function; LAVI: left atrial volume index; TRV: tricuspid regurgitation velocity.

Fig 5 illustrates the area under the ROC curve for diagnosing HFpEF of HFA-PEFF and LASr, showing no difference ($p = 0.062$). Furthermore, the AUC of conventional echocardiographic indices (LAVI, E/e', e', TRV) with HFA-PEFF and LASr differed significantly with $p < 0.001$.

Fig 5. Comparison of the AUC values of LASr, HFA-PEFF score, and existing parameters in predicting HFpEF. The analysis reveals no significant difference in diagnostic value between LASr and HFA-PEFF score ($p > 0.05$). However, significant disparities were observed between LASr and LAVI, E/e', and between HFA-PEFF score and LAVI, E/e', TRV, and e' ($p < 0.05$). E: E-wave velocity; e': e'-wave velocity; HFA-PEFF: Heart Failure Association-PEFF; LASr: left atrial reservoir function; LAVI: left atrial volume index; TRV: tricuspid regurgitation velocity.

Discussion

In this study, we examined 118 subjects to evaluate the role of LA strain in

diagnosing HFpEF.

LA function indices such as LASr, LAScd, and LASct decreased significantly

in the HFpEF group compared to the control group, with a p-value < 0.001. This

finding is consistent with Aung et al.'s (2017) study on 83 patients, which also

reported decreased LASr and LASct in the HFpEF group compared to the control

group [19]. Similar results were observed in several other studies where LASr,

LAScd, and LASct were all reduced in HFpEF patients compared to the control

group [9,13,20,21]. Therefore, our study confirms a significant decline in LA

function in the HFpEF group compared to healthy subjects. Reddy et al. also

highlighted the important role of the LA in the progression of HFpEF. They

suggested that LA strain reflects the overall LA function, which progressively

deteriorates in the presence of chronic LV diastolic dysfunction, such as in patients

with HFpEF [9].

The American Echocardiography Association and the European Association

of Cardiovascular Imaging (ASE/EACVI) recommend using echocardiography to

diagnose HFpEF in patients with sinus rhythm using E/A ratio, LAVI, TRV, and

E/e' [3]. The ESC guidelines for the diagnosis of HFpEF are based on evidence of

functional and structural alterations of the heart using the parameters E/e', LVM,

LAVI, septal e' velocity, and lateral e' velocity [14]. However, these classic

indicators still have many limitations, and many clinical practice cases encounter

difficulties when the diagnosis falls into the "undetermined" state, when Doppler

measurements cannot be made, such as tachycardia or severe mitral valve disease

[22]. Compared with Doppler echocardiography, the advantage of speckle tracking

echocardiography when compared to conventional echocardiography is that it is

independent of angle and less affected by mitral valve disease. On the other hand,

the LA strain evaluates the function of the LA throughout the entire cardiac cycle

rather than the functional state of a certain time point in the cardiac cycle [11]. In

our study, LA strain indices including LASr (AUC = 0.877), LAScd (AUC =

0.836), and LASct (AUC = 0.749) have high values in diagnosing HFpEF

equivalent to the score HFA-PEFF (AUC = 0.975). At the same time, the ability of

LASr to diagnose HFpEF was superior to the LAVI (AUC = 0.670), E/e' (AUC =

0.578), septal e' velocity (AUC = 0.615), and TRV (AUC = 0.565) with p < 0.05.

Many studies have also shown the superiority of LA strain indices in the ability to

diagnose HFpEF compared to commonly used classical indices [9,23,24].

HFA-PEFF is a widely used scoring system for diagnosing HFpEF. However,

evaluating this HFA-PEFF is a widely used scoring system for diagnosing HFpEF.

However, evaluating this scoring system requires numerous parameters, including

echocardiography, NT-proBNP, and atrial fibrillation diagnosis [17,25]. Through

our study, LA strain indices have demonstrated an AUC equivalent to that of the

HFA-PEFF score. When comparing the AUC of LASr and HFA-PEFF, our study

found no significant difference in the HFpEF diagnostic value of LASr and HFA-

PEFF score with p = 0.062. The 5-point HFA-PEFF score has 100% specificity,

however, the sensitivity is low at only 56.52%, which can cause difficulties in

applying this score in clinical practice.

In conclusion, LA parameters on speckle tracking echocardiography may be useful in diagnosing HFpEF and integration into the 2016 EACVI/ASE criteria will improve diagnostic effectiveness with accuracy not inferior to conventional echocardiographic parameters for diagnosing HFpEF.

Limitations of the study

Our study only compares non-invasive indices in diagnosing HFpEF and does not utilize invasive interventions for evaluation or comparison with other invasive indices. We only used one software for strain analysis and did not compare different software programs. Furthermore, although our sample size is larger than some studies, it is still relatively small, necessitating larger studies to establish cutoff points applicable to clinical central Vietnam and was conducted at a single location, which may limit the generalizability of the findings to other populations or settings. Different populations or geographical locations may exhibit variations in disease prevalence and characteristics, potentially affecting the study outcomes. During sample collection, technical limitations of Doppler echocardiography may have prevented us from obtaining all possible Doppler echocardiography indices for comparison with LA strain indices. This study only focused on sinus rhythm subjects; however, atrial fibrillation is a significant risk factor for HFpEF, necessitating further research to assess the best way to combine LA strain parameters with conventional parameters to determine HFpEF.

Conclusions

LA strain has a diagnostic value equivalent to the HFA-PEFF score in diagnosing HFpEF. These indices could be incorporated into the existing guidelines to enhance the diagnosis of HFpEF.

Acknowledgements

Nil

References

1. Redfield MM, Borlaug BA. Heart Failure With Preserved Ejection Fraction: A Review. JAMA. 2023;329: 827. doi:10.1001/jama.2023.2020

2. Obokata M, Reddy YNV, Borlaug BA. Diastolic Dysfunction and Heart Failure With Preserved Ejection Fraction. JACC: Cardiovascular Imaging. 2020;13: 245–257. doi:10.1016/j.jcmg.2018.12.034

3. Nagueh SF, Smiseth OA, Appleton CP, Byrd BF, Dokainish H, Edvardsen T, et al. Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography: An Update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. Journal of the American Society of Echocardiography. 2016;29: 277–314. doi:10.1016/j.echo.2016.01.011

4.   Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM, et al.

2022 ACC/AHA/HFSA Guideline for the Management of Heart Failure. Journal

of Cardiac Failure. 2022;28: e1–e167. doi:10.1016/j.cardfail.2022.02.010

5.   Boe E, Smiseth OA. Left atrial strain imaging: ready for clinical implementation

in heart failure with preserved ejection fraction. European Heart Journal -

Cardiovascular Imaging. 2022;23: 1169–1170. doi:10.1093/ehjci/jeac059

6.   Rosca M, Lancellotti P, Popescu BA, Pierard LA. Left atrial function:

pathophysiology, echocardiographic assessment, and clinical applications.

Heart. 2011;97: 1982–1989. doi:10.1136/heartjnl-2011-300069

7.   Gan GCH, Ferkh A, Boyd A, Thomas L. Left atrial function: evaluation by strain

analysis. Cardiovasc Diagn Ther. 2018;8: 29–46. doi:10.21037/cdt.2017.06.08

8.   Jarasunas J, Aidietis A, Aidietiene S. Left atrial strain - an early marker of left

ventricular diastolic dysfunction in patients with hypertension and paroxysmal

atrial fibrillation. Cardiovasc Ultrasound. 2018;16: 29. doi:10.1186/s12947-018-

0147-6

9.   Reddy YNV, Obokata M, Egbe A, Yang JH, Pislaru S, Lin G, et al. Left atrial

strain and compliance in the diagnostic evaluation of heart failure with preserved

ejection fraction. European J of Heart Fail. 2019;21: 891–900.

doi:10.1002/ejhf.1464

10. Brand A, Romero Dorta E, Wolf A, Blaschke-Waluga D, Seeland U, Crayen C,

et al. Phasic left atrial strain to predict worsening of diastolic function: Results

from the prospective Berlin Female Risk Evaluation follow-up trial. Front

Cardiovasc Med. 2023;10: 1070450. doi:10.3389/fcvm.2023.1070450

11. Ma C, Liao Y, Fan J, Zhao X, Su B, Zhou B. The novel left atrial strain

parameters in diagnosing of heart failure with preserved ejection fraction.

Echocardiography. 2022;39: 416–425. doi:10.1111/echo.15304

12. Kim D, Seo JH, Choi KH, Lee SH, Choi J-O, Jeon E-S, et al. Prognostic

Implications of Left Atrial Stiffness Index in Heart Failure Patients With

Preserved Ejection Fraction. JACC: Cardiovascular Imaging. 2023;16: 435–445.

doi:10.1016/j.jcmg.2022.11.002

13. Rimbas RC, Visoiu IS, Magda SL, Mihaila-Baldea S, Luchian ML, Chitroceanu

AM, et al. New insights into the potential utility of the left atrial function analysis

in heart failure with preserved ejection fraction diagnosis. Johnson DM, editor.

PLoS ONE. 2022;17: e0267962. doi:10.1371/journal.pone.0267962

14. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JGF, Coats AJS, et al.

2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart

failure: The Task Force for the diagnosis and treatment of acute and chronic heart

failure of the European Society of Cardiology (ESC). Developed with the special

contribution of the Heart Failure Association (HFA) of the ESC. European J of

Heart Fail. 2016;18: 891–975. doi:10.1002/ejhf.592

15. Mitchell C, Rahko PS, Blauwet LA, Canaday B, Finstuen JA, Foster MC, et al.

Guidelines for Performing a Comprehensive Transthoracic Echocardiographic

Examination in Adults: Recommendations from the American Society of

Echocardiography. Journal of the American Society of Echocardiography.

2019;32: 1–64. doi:10.1016/j.echo.2018.06.004

16. Park JJ, Hwang I, Kang S, Park J, Park J, Cho G. Myocardial strain for heart

failure with preserved ejection fraction but without diastolic dysfunction. ESC

Heart Failure. 2022;9: 3308–3316. doi:10.1002/ehf2.14078

17. Pieske B, Tschöpe C, De Boer RA, Fraser AG, Anker SD, Donal E, et al. How

to diagnose heart failure with preserved ejection fraction: the HFA–PEFF

diagnostic algorithm: a consensus recommendation from the Heart Failure

Association (HFA) of the European Society of Cardiology (ESC). European

Heart Journal. 2019;40: 3297–3317. doi:10.1093/eurheartj/ehz641

18. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two

or more correlated receiver operating characteristic curves: a nonparametric

approach. Biometrics. 1988;44: 837–845.

19. Aung SM, Güler A, Güler Y, Huraibat A, Karabay CY, Akdemir I. Left atrial

strain in heart failure with preserved ejection fraction. Herz. 2017;42: 194–199.

doi:10.1007/s00059-016-4456-y

20. Obokata M, Negishi K, Kurosawa K, Arima H, Tateno R, Ui G, et al. Incremental

Diagnostic Value of LA Strain With Leg Lifts in Heart Failure With Preserved

Ejection Fraction. JACC: Cardiovascular Imaging. 2013;6: 749–758.

doi:10.1016/j.jcmg.2013.04.006

21. Ye Z, Miranda WR, Yeung DF, Kane GC, Oh JK. Left Atrial Strain in Evaluation

of Heart Failure with Preserved Ejection Fraction. Journal of the American

Society of Echocardiography. 2020;33: 1490–1499.

doi:10.1016/j.echo.2020.07.020

22. Dzhioeva O, Belyavskiy E. Diagnosis and Management of Patients with Heart

Failure with Preserved Ejection Fraction (HFpEF): Current Perspectives and

Recommendations. TCRM. 2020;Volume 16: 769–785.

doi:10.2147/TCRM.S207117

23. Braunauer K, Düngen H-D, Belyavskiy E, Aravind-Kumar R, Frydas A, Kropf

M, et al. Potential usefulness and clinical relevance of a novel left atrial filling

index to estimate left ventricular filling pressures in patients with preserved left

ventricular ejection fraction. European Heart Journal - Cardiovascular Imaging.

2020;21: 260–269. doi:10.1093/ehjci/jez272

24. Lin J, Ma H, Gao L, Wang Y, Wang J, Zhu Z, et al. Left atrial reservoir strain

combined with E/E' as a better single measure to predict elevated LV filling

pressures in patients with coronary artery disease. Cardiovasc Ultrasound.

2020;18: 11. doi:10.1186/s12947-020-00192-4

25. Choi KH, Yang JH, Seo JH, Hong D, Youn T, Joh HS, et al. Discriminative Role

of Invasive Left Heart Catheterization in Patients Suspected of Heart Failure

With   Preserved   Ejection   Fraction.   JAHA.   2023;12:   e027581.

doi:10.1161/JAHA.122.027581

Supplementary materials

Supplementary 1. Calculation of HFA-PEFF score

	HFA-PEFF score				
Parameter	**Minor**		**Major**		
	Value	**Point**	**Value**	**Point**	
Functional	Average E/e' 9 - 14 or GLS < 16%	1	Septal e' < 7 cm/s or lateral e' < 10 cm/s or Average E/e' ≥ 15 or TR velocity > 280 cm/s (PASP > 35 mmHg)	2	
Morphological	LAVI 29 – 34 mL/m^2 or LVMI ≥ 115/95 g/m^2 (female/male) or RWT > 0.42 or LV wall thickness ≥	1	LAVI > 34mL/m^2 or LVMI ≥ 149/122 g/m^2 (female/male) and RWT > 0.42	2	

	12 mm			
Biomarker (SR)	NT-proBNP 125 - 220 pg/mL or BNP 35 - 80 pg/mL	1	NT-proBNP > 220 pg/mL or BNP > 80 pg/mL	2
Biomarker (AF)	NT-proBNP 365 - 660 pg/mL or BNP 105 - 240 pg/mL	1	NT-proBNP > 660 pg/mL or BNP > 240 pg/mL	2

Adapted from Heart Failure Association of the European Society of Cardiology

Abbreviations: AF: atrial fibrillation; BNP: B-type natriuretic peptide; E: early transmitral flow velocity; e': early diastolic mitral annular velocity; GLS: left ventricular global longitudinal strain; HFA-PEFF: Heart Failure Association-PEFF; LAVI: left atrial volume index; LV: left ventricle; LVMI: left ventricular mass index; NT-proBNP: N-terminal pro-B-type natriuretic peptide; PASP: pulmonary artery systolic pressure; RWT: relative wall thickness; SR: sinus rhythm; TRV: tricuspid regurgitation velocity.

Fig1

Fig2

Fig3

Fig4

Fig5