1 **Assessing the impact of Enhanced-Case-Finding on tuberculosis case**
2 **notifications and transmission in The Gambia using epidemiological and**
3 **phylodynamic approaches**
4

Florian Gehre*[1,2,3,4], Francis Oko*[3], Boatema Ofori-Anyinam*[2,3,5], Conor J. Meehan*
[2,6], Etthel M. Windels[7,8,9], Ken Joof[3], Tutty Faal[3], Francis Mendy[3], Tijan Jobarteh[3],
Ensa Gitteh[3], Abi Janet Riley[3], Binta Sarr-Kuyateh[3], Catherine Okoi[3,10], Edward
Demba[3], William Del-Alorse[3], Kodjovi Mlaga[3], Basil Sambou[3], Fatoumatta Kanuteh[3],
Koduh Lette[3], Abdul Khalie Muhammad[3], Shadrac Agbla[3], Wim Mulders[2], Simon
Donkor[3], David Jeffries[3], Anna Roca[3], Umberto D'Alessandro[3], Martin Antonio*[3],
Ifedayo Adetifa*[3,10,11], Bouke C. de Jong*[2]
12

Affiliations:
[1] Bernhard-Nocht-Institute for Tropical Medicine (BNITM), Hamburg, Germany
[2] Institute for Tropical Medicine (ITM), Antwerp, Belgium
[3] Medical Research Council (MRC) Unit The Gambia Unit at the London School
of Hygiene and Tropical Medicine, Fajara, The Gambia.
[4] East African Community (EAC), Arusha, Tanzania
[5] Ruy V. Lourenço Center for Emerging and Re-Emerging Pathogens, Rutgers
University New Jersey Medical School, New Jersey, USA.
[6] Nottingham Trent University, Nottingham, UK.
[7] ETH Zurich, Basel, Switzerland
[8] Swiss Tropical and Public Health Institute, Allschwil, Switzerland
[9] Swiss Institute of Bioinformatics, Lausanne, Switzerland
[10] National Reference Laboratory, Nigeria Centre for Disease Control and
Prevention, Abuja, Nigeria.
[11] Department of Paediatrics and Child Health, College of Medicine, University of
Lagos, Lagos, Nigeria.

* Equal contributions

Running Head: TB Enhanced Case Finding in The Gambia
Key words: tuberculosis, phylodynamics, TB-ECF trial, InterrupTB study
Corresponding author: Conor Meehan, email: conor.meehan@ntu.ac.uk
33
34
35

Abstract:

Background: Most tuberculosis (TB) cases in The Gambia are notified in the Greater Banjul Area (GBA). We conducted an Enhanced-Case-Finding (ECF) intervention in the GBA and determined its effect on TB case notifications and ongoing TB transmission.

Methods: This was a cluster randomized trial in which randomly assigned intervention areas of grouped settlements received three rounds of an ECF strategy consisting of sensitization followed by auramine microscopy, whereas people with TB in control areas continued to be identified through passive case finding. People with TB were recruited at the TB diagnostic and treatment centers serving both the intervention- and control areas. The primary outcome was TB case notification rate. To exclude that an increase in notified cases, followed by a decrease in notified cases, would hide the future impact of the intervention, we tested for changes in transmission dynamics using both genetic clustering and phylodynamic methods.

Results: 3,047 people living with TB were recruited in the study, evenly split between intervention and control regions. No significant difference in TB case notification rates, transmission clustering or effective reproductive number was detected between intervention and control areas using either a case notification rate or phylodynamic approach.

Conclusion: Although we did not find evidence for decreased TB case notification nor TB transmission through the ECF strategy used, this approach is an examplar of how both classical epidemiology and genomic phylodynamics approaches can be integrated to better assess public health intervention outcomes.

Introduction:

Tuberculosis (TB) remains endemic in most countries worldwide. Despite ongoing efforts to meet Sustainable Development Goals, there has been an extremely slow decline of TB incidence (1) sustained by ongoing transmission from the 'missing millions' i.e., the estimated three of the ten million incident people with TB worldwide who are not diagnosed nor treated.

To improve TB detection, case finding efforts have been scaled up through several initiatives, such as 'TB reach' (2). Nevertheless, the evidence-base for the various approaches to detect additional people with TB is scarce. Although people with TB become rapidly non-infectious with effective treatment (3), there is little concrete evidence that increased case detection interrupts TB transmission. Historical evidence showed that mass tuberculosis screening by chest x-ray in Europe and North America was effective in reducing tuberculosis incidence (4). More recent evidence includes mass Xpert screening to have reduced IGRA positivity among children (5), supporting that the intervention interrupted tuberculosis transmission. Landmark studies in Zimbabwe (6) and Vietnam (7) showed an impact of case finding strategies on TB prevalence, and the TREATS study in Zambia and South Africa found marked reduction in incidence after community-wide screening (8). However, other studies such as ZAMSTAR (9) found little impact from household and community interventions and systematic comparisons of active case finding approaches have found moderate quality evidence that active case-finding, when implemented with sufficient coverage and intensity in high-prevalence settings, can positively affect the community epidemiology of tuberculosis(10).

Measuring the impact of case finding interventions on TB transmission and incidence is complicated by several factors including the long latent period of *Mycobacterium tuberculosis*(11) and the dynamics of case finding, which is expected to initially *increase* TB case notifications (TB case detection rate) by decreasing the diagnostic gap. If sustained, such intervention efforts are expected to eventually *decrease* TB case notifications and thus TB incidence. It is expected that the effect of case finding would be more reliably measured as a decline in transmission, inferred from the phylogenetic relatedness of the people with TB's mycobacterial isolates.

In The Gambia, previous spatial analyses identified the Greater Banjul Area, a suburban region bordering the capital city of Banjul, as the area with most TB notifications (12). Therefore, this area was chosen to carry out a cluster randomized trial to determine the effect of enhanced case finding (ECF) through community

96 sensitization. The impact of the intervention on TB transmission was modelled
97 through the analysis of whole genome sequence data from bacterial isolates
98 collected during the trial.
99

Methods

Study population and overall study design: The cluster randomized trial on
Enhanced Case Finding (ECF-TB study, ClinicalTrials.gov Identifier: NCT01660646)
was conducted from 1st March 2011 to 31st December 2014 in the Greater Banjul
Area (GBA) with an estimated population size of 680,000 people (based on the
available 2004 census at the time of the study design). The GBA was geographically
divided into 20 clusters which included 121 settlements with population ≥ 100, which
were randomized (10 each) to either the intervention (Enhanced Case Finding) or the
control arm. The primary endpoint was TB case notification rate. To understand the
effect of the intervention on TB transmission, all mycobacterial isolates collected
between 2011 and 2014 underwent culturing and whole genome sequencing to
estimate changes in TB transmission between intervention and control clusters
through genetic clustering and phylodynamics (InterrupTB sub-study).

Enhanced Case Finding study:

Design and cluster assignment:

An unrestricted randomization of the study settlements would have resulted in a
higher risk of 'contamination' from intervention to control clusters, without controlling
for spatial proximity or population size. Therefore, a stratified randomization was
used to minimize the chance of contamination between settlements, to balance
population size between the study arms and balance the number of settlements and
settlement sizes in each arm and produce a design that allowed efficient
implementation of ECF. Randomization was performed by many runs of an algorithm
that used a clustering algorithm (k-means) to spatially cluster settlements with a
population of at least 100. For each run the following operations were performed: 1)
The clusters of settlements were randomly assigned to two arms (which would later
be randomised to the ECF or the control arm). 2) The distance between the two
closest villages in different clusters was estimated. 3) The balance of large, medium
and small settlements in each arm was quantified. 4) The balance of the total
numbers of settlements in each arm was quantified, including the overall population
in each arm. Out of 100,000 runs, the assignment maximizing the closest distance
(1.1km) between settlements from different clusters, with the best settlement number

(60 vs. 61) and overall population balance (imbalance of 708 out of 660,000) was
chosen. The two groups of 10 clusters (note that there was an intervention cluster
with only one settlement as it was remote from the other settlements) were then
randomly allocated to either ECF or control.

Community Intervention: ECF was implemented over several days. On day 1,
trained field workers informed the village elder (alkalo) and community leaders of the
upcoming intervention and sought their agreement. This was followed by a
community meeting to inform and sensitize about TB symptoms (persistent cough,
night sweats, weight loss) through a public speech and by showing an educational
movie on a portable TV. The participants also had the opportunity to ask questions
about the disease during a Q&A session. Before their departure, the research team
left sputum collection cups for the inhabitants. These were collected on day 3,
provided the participants had signed a written informed consent. Epidemiological
data (ethnicity, reason for presenting at Health Centre, symptoms, risk factors) were
collected at the same time as collection. The intervention was repeated 3 times
(every 6 months) over a period of 18 months. Acid Fast Bacilli (AFB) microscopy
positive samples identified in this manner were attributed to the direct effect of the
ECF. The difference in people living with TB presenting in any of the seven chest
clinics in the GBA from intervention clusters versus from non-intervention clusters
was considered as the indirect effect of the ECF, as they represented sensitized,
symptomatic individuals or their indirectly sensitized contacts unable or unwilling to
directly return their sputum sample just after the sensitization. All health staff in the
seven chest clinics were trained in the use of the case report form to (i) confirm
whether a person living with TB had heard of or participated in the ECF strategy, (ii)
assess risk factors, and (iii) document the person's village of residence. To confirm
such information, the research team visited people living with TB presenting during
the study period at home and collected the geographical coordinates through GPS,
allowing the correct allocation of each person to either the intervention or the control
arm.

Primary TB Diagnostics: All sputum cups collected in the field during ECF
campaigns were brought to the MRC TB laboratory and immediately analyzed using
direct fluorescent microscopy prior to decontamination. Sputum samples from people
living with TB recruited at the Chest Clinics were processed by Fluorescent
Microscopy and MGIT culture (BACTEC MGIT 960; Becton Dickinson Microbiology
Systems, Sparks, Maryland, USA). Positive MGIT cultures were confirmed as

Mycobacterium tuberculosis (*Mtb*) by MPT64 antigen test and subcultured on
Löwenstein Jensen (LJ) medium. All isolates were stored at -80°C for whole genome
sequencing. People with AFB positive sputa were informed and referred for TB
treatment at the nearest facility.
173

TB case notification rate difference (CNRD)
One of the major read-outs of the ECF study was to understand whether an ECF
strategy would reduce case notifications in intervention clusters when compared to
control areas. Therefore, we calculated the 60-day moving average TB case
notification rate difference (CNRD) between intervention and control clusters (for all
clusters with more than 30 screened people).
180

InterrupTB study:
Genome sequencing, Bioinformatics and estimation of TB transmission
Cultured TB isolates were retrieved from -80°C storage and cultured in the BACTEC
MGIT 960 System (MGIT 960; Becton Dickinson Microbiology Systems, Sparks,
Maryland, USA) at 37°C following manufactures' instructions. Positive cultures were
plated on blood agar to check for purity. Pure cultures were subjected to ZN
microscopy to confirm the presence of AFB; AFB positive cultures were sub-cultured
on LJ slants, one with pyruvate and another with glycerol. If the stored isolate did not
regrow in the MGIT, leftovers of processed sputa of AFB-positive and MGIT culture
negative clinical samples detected within the study period were retrospectively
retrieved from freezer storage at -20°C and cultured on LJ slants with pyruvate, to
lessen culture bias against *Mtb* L5 and L6 (*M. africanum*) (13). Cultures were
incubated at 37°C for up to 8 weeks. From loopfuls of pure *Mtb* colonies grown on
LJ, samples for whole genome analysis had genomic DNA extracted using the
Maxwell 16 DNA Purification Kit (Promega) (14).
196

Sequencing was undertaken on Illumina HiSeq and Miseq platforms. TBprofiler was
used (Pipeline version: 3.0.3, access date: December 2020)(15) on the resulting raw
reads to assign lineages and drug resistance profiles. Mixed infections were
excluded based on whether the TBprofiler strain calling algorithm detected 2 or more
(sub)lineages. Data filtering, quality control and SNP calling was performed as
previously reported (16). Briefly, non-*Mtb* reads were filtered out using Centrifuge
v1.0.3 (17) and then SNPs were called using MTBseq (18) with default settings and
the inferred ancestor genome as the reference (19). Samples with less than 95% of
the reference genome covered or with a mean coverage <30x were then excluded.

SNP distance matrices were created using the MTBseq join and amend functions

using default settings (only sites with >95% of samples with an unambiguous

nucleotide call and SNPs within 12bp of each other filtered out). Custom python

scripts were then used to create loose transmission clusters (as previously defined

(20)) at 5 and 1 SNP cut-offs. These two cut-offs represent best the likely divergence

during the 3-year sampling period (20).

Transmission clusters were assigned to either the intervention or control arm using

two methods: majority counts and earliest sampling (Supplementary Figure 1). For

majority counts, the arm (intervention or control area) of each sample in a

transmission cluster was extracted from the database and the total of each arm

summed up. Transmission clusters where >50% of samples were from an

intervention area were labelled as intervention transmission clusters. The same

approach was applied to the control clusters. Clusters where the number of

intervention and control samples were equal (i.e., 50% each) were excluded. For

earliest sampling, the date of sampling of each isolate in the transmission cluster was

extracted from the database and a cluster was assigned to either intervention or

control arm according to the origin of the earliest sample. Clusters where two

samples were taken on the same day but from different trial arms were excluded.

To look for significant differences between intensity of transmission in intervention

and control arms based on the majority clustered and earliest sample methods, a chi-

squared test on the raw counts of clusters assigned to each arm was used as

implemented in R v3.6.3 (21) using a p-value cut-off of 0.001 with trial arm vs

clustered/unclustered as the variables. To create the graph of clusters, the Wilson

confidence interval for each method was calculated using the binom (22), plyr (23)

and ggplot2 (24) packages in R.

Phylodynamic analyses
235

To further investigate the effect of the intervention on *Mtb* transmission, we employed

a phylodynamic approach. For computational feasibility, we randomly subsampled

400 sequences belonging to *Mtb* lineage 4 (as it was the most prominent lineage in

the dataset) and either the intervention or control arm. On this subset, we fit a

structured birth-death model (25) with two compartments: one containing people

living with TB in the control arm of the study and the other containing people living

with TB in the intervention arm. In this birth-death model, the birth rate corresponds

to the *Mtb* transmission rate, i.e., the number *Mtb* transmission events occurring from
one person living with TB per unit of time. The death rate corresponds to the rate of
becoming uninfectious for *Mtb* due to recovery or death. People living with TB can
infect people in the same trial arm as well as in the other arm, with probabilities
determined by the probabilities of contact with people from the same versus the other
trial arm. We assumed no further structuring of the population by village.

The effective reproductive number (R_e), corresponding to the ratio of the birth and
death rate, was assumed equal for both trial arms before the start of the study ($R_{e,1}$).
After that time, a different R_e was inferred, which was also allowed to be different for
the control and intervention group:

$$R_{e,cc} = (1 - p) * R_{e,2}$$
$$R_{e,ii} = (1 - p) * f * R_{e,2}$$
$$R_{e,ci} = p * R_{e,2}$$
$$R_{e,ic} = p * f * R_{e,2}$$

with $R_{e,cc}$, $R_{e,ii}$, $R_{e,ci}$, and $R_{e,ic}$ the effective reproductive numbers within the control
group, within the intervention group, from the control group to the intervention group,
and from the intervention group to the control group, respectively. $R_{e,2}$ represents the
base R_e after the start of the study, *p* is the probability that a contact belongs to a
different trial arm, and *f* represents the multiplicative effect of the intervention on the
reproductive number. We assumed that the transmission rate was equal for both trial
arms due to the random selection and allocation of clusters to study arms. We also
assumed that any intervention effect on R_e was the result of an effect on the
infectious period. The reproductive numbers for samples in the control and
intervention arm ($R_{e,c}$ and $R_{e,i}$, respectively) are then as follows:

$$R_{e,c} = R_{e,cc} + R_{e,ci} = R_{e,2}$$
$$R_{e,i} = R_{e,ii} + R_{e,ic} = f * R_{e,2}$$

People living with TB are sampled with sampling proportion *s*, which was set to zero
before the onset of sampling and assumed equal for both intervention arms
afterwards. Upon sampling an infected person, the person was assumed to become
uninfectious with probability *r* (26). A strict molecular clock was assumed, and a
general time-reversible nucleotide substitution model with four gamma rate
categories to account for site-to-site rate heterogeneity (GTR+Γ_4). Phylodynamic

inference was performed using the bdmm package (25) in BEAST2 (27,28). A
variable SNP alignment was generated and augmented with a count of invariant A,
C, G, and T nucleotides to avoid ascertainment bias (29). Three independent Markov
Chain Monte Carlo chains were run for each analysis, with states sampled every
1,000 steps. Tracer (30) was used to assess convergence and confirm that the
effective sample size was at least 200 for the parameters of interest. 10% of each
chain was discarded as burn-in, and the remaining samples across the three chains
were pooled using LogCombiner (27), resulting in at least 200,000,000 iterations in
combined chains.
284

Prior distributions
All parameters and their corresponding prior distributions are listed in Supplementary
Table 1. For the sampling proportion, a uniform prior was chosen, with lower bound
set to zero and upper bound set equal to the ratio of the number of sequences,
corrected for down sampling, and the total number of reported cases during the
sampling period.
291

Sensitivity analyses
The robustness of the phylodynamic inference to potential sampling biases was
assessed by estimating a different sampling proportion for the control and
intervention group. Furthermore, we assessed the sensitivity of the results to prior
assumptions by setting the clock rate prior to a Lognormal(-17,1) distribution and the
prior on f to a Lognormal(0,2) distribution.
298

Ethical Approval: For the ECF study ethical approval was granted by the Gambia
Government / MRC Joint Ethics Committee. In brief, any newly diagnosed person
initiating TB therapy at any of the Gambia Government TB diagnostic and treatment
centres in the Greater Banjul Area, regardless of age, residency, HIV status, or type
of TB, with permanent residence in the Greater Banjul Area was eligible for inclusion.
Individuals unable to understand the implications of study participation, whether
through cognitive impairment or insurmountable language barrier, were excluded
from the study.
For the transmission blocking study, ethical approval was provided by the Gambia
Government/ MRC Joint Ethics Committee and the Institutional Review Board of the
Institute of Tropical Medicine, Antwerp, Belgium. Isolates from all ECF participants
(except minors, i.e., <18 years old) were included.
311

312 **Data accessibility**

313 Raw reads are stored on the ENA under accession number PRJEB53138. Custom

314 python scripts used for SNP clustering can be found at

315 https://github.com/conmeehan/pathophy. XML files for BEAST analyses can be

316 found at https://github.com/EtthelWindels/tb-ecf-gambia.

Results and Discussion:

Design set up, cluster assignment and recruitment: We chose a clustered

randomized trial design over a before-and-after design as this most stringent study

setup allows randomization and adjustment for longitudinal external effects that

influence the whole study population. We positioned our study in the GBA, with a

population of ~700.000 inhabitants where ~77% of all TB cases reported in the

Gambia occur (personal communication, Gambian National Tuberculosis and

Leprosy Control Program, NTLP).

A total of 121 villages in the Greater Banjul Area, with a total population of 648,835,

were assigned to either 10 intervention (331,800 people) or 10 control clusters

(317,035 people) (Fig. 1). During three ECF rounds, 10,219 individuals were

screened for TB and provided a sputum sample. Thirty-five individuals (0.34%) were

AFB+ and therefore confirmed to be people living with TB and attributable to the

direct effect of the ECF strategy (Fig. 2). A total of 3,047 people living with TB were

recruited through the seven Chest Clinics in the GBA, 1,314 from intervention, 1,375

from control clusters (Fig. 2), and 358 not residing in a cluster. For an

epidemiological overview of the ECF study population, see Supplementary Table 2.

About 43% of isolates from intervention (563/1,314) and from control (588/1,375)

clusters were retrieved for sub-culture and extraction of genomic DNA. The people

whose isolate was retrieved did not differ significantly from those who were not

included in this analysis, except for higher rates of positive cultures (Supplementary

Table 2).

Dynamics of TB case notifications: If an intervention successfully reduces TB

incidence, you would expect the case notification rate in the intervention areas to

begin to drop compared to the control areas. This would result in the comparative

case notification rate of intervention to control dropping below zero. Although the

CNRD tended to decrease around the interventions, the upper limit of the 95% CI

never crossed the zero-difference line. This suggests that, based on case reporting,

we did not observe a significant reduction of case notifications in the intervention
clusters at any time of the study period (Fig. 3).

As the health facilities where people with TB were recruited served both the
intervention and control arm, selection bias due to different access was probably
limited. Therefore, assuming the study design and study location were appropriate, it
is unclear why the intervention did not have any effect on the study outcomes. The
direct effect of the intervention was unexpectedly small, with auramine microscopy
on sputum collected the next morning only detecting 35 cases (0.34% of submitted
sputum samples). Auramine microscopy has a reported sensitivity around 60% for
the detection of pulmonary tuberculosis in symptomatic persons, although recent
studies suggest that the majority of culture positive people with TB do not report a
cough (31).

Another possible explanation is the lower-than-expected TB prevalence. Indeed,
based on WHO prevalence estimates at the time of study design, the TB prevalence
in The Gambia was assumed to be 327/100,000. However, a nationwide TB
prevalence survey (GAMSTEP) reported a much lower prevalence, i.e., 128/100,000.
Indeed, in the latest 2021 WHO guidelines systematic screening for TB disease
among the general population is recommended in areas with an estimated TB
prevalence of 500/100,000 or higher, albeit with low certainty of evidence (32).
Based on these prevalence estimates, the intervention retrieved above expected
numbers in the prevalence survey (32/10,219 vs the expected ~13/~10,000
samples), yet below the expected yield after sensitizing on TB symptoms.

Estimating TB transmission using genomic clustering proportion and changes
in R_e: An alternative way to assess the impact of an intervention on TB incidence is
to look at its effect on transmission rates. An effective intervention should reduce the
number of new TB cases in the region, thus resulting in longer term reduction in case
notifications. Using a pathogen genome-based approache, this should result in fewer
transmission clusters arising in intervention areas compared to control area. In a
phylodynamics-based approach this should result in a reduction of the R_e (effective
secondary case contact rate) in intervention areas compared to the control.

We sequenced the whole genome of 536 and 555 isolates (post QC and removal of
mixed infections) from intervention and control areas, respectively. Most sequenced
strains came from L4 (67.39%) or L6 (26.43%). Only 10 MDR-TB samples were
predicted in this dataset based on genotyping, suggesting a low prevalence of drug
resistance in this setting.

Using a 5-SNP cut-off, 150 transmission clusters were detected in the data
containing a total of 568 (52%) of people with TB. A total of 119 clusters were
detected using the more stringent 1-SNP cut-off containing 349 (32%) of people with
TB. We determined the clustering proportion of isolates belonging to intervention in
comparison to the control group (Fig 4) but did not find significant differences
between the two groups, even after considering various SNP thresholds as cut-offs
for clustering (Chi-Squared test p-value 0.82).
Further transmission dynamics analyses were performed using BEAST to estimate
the effective reproductive number (R_e) in the two arms. If the intervention worked, the
$R_{e\ in}$ the intervention group should be lower than the R_e in the control group,
indicating a reduction in transmission. However, all of our phylodynamic models
showed no significant effect of the intervention on R_e (posterior mean of the relative
R_e = 0.58-0.72), even with different prior assumptions on sampling evenness and
molecular clock rates (Figure 5; Table 2). This correlates with what is observed in the
case notification data from the entire ECF sample, indicating that the intervention did
not significantly reduce transmission nor incidence in The Gambia.
This phylodynamic approach serves as an example of how molecular data,
especially whole genome sequencing, can be incorporated into such interventions to
better evaluate reductions in incidence during the study period and predict if future
transmission and incidence will also be reduced (based on projected parameters).
Such genomic epidemiology approaches have been used previously to indicate when
small outbreaks in low-incidence settings are over (33). However, here we show the
additional power of phylodynamics to enhance outcome measures in higher
incidence settings as well.
408

Lessons learnt and limitations:
As the overall transmission-blocking effect is expected to be lower in this setting than
in medium prevalence settings, a longer follow-up period after the last intervention
round may have shown a lower-case notification rate and transmission in the
intervention arm. Compounding this with disease latency of months to years(11) may
mean that the impact of interventions may not be seen for several years after studies
have finished. In addition, the intervention may not have had the expected reach,
although the research team tried to be as inclusive as possible. During the
community interventions, attendees were educated on various aspects of TB (using a
film and presentation), with a special focus on three symptoms (weight loss, night
sweats, persistent cough), which were in line with WHO recommendations (32), even

if our current understanding suggests poor sensitivity of such symptom screen (31).
All community meetings were interactive, with the research team encouraging the
audience to ask any question they may have. Moreover, depending on the location of
the village/cluster, the campaign was held in at least one of the four main languages,
namely Wolof, Mandinka, Jola and Fula, but also any other local language spoken by
the field team. However, we cannot fully dismiss the possibility that the delivered
messages were not disseminated wide enough within the recipient population, thus
not reaching the incident people with TB we aimed to invite for testing.

It is also possible that the overall awareness of TB in the Gambian population was
already high, and that people with TB in the control area would have attended the
health facilities for TB diagnosis even without the sensitization campaigns, with
minimal incremental knowledge gained with ECF. This seems to be the case as a
study carried out before and after the intervention confirmed that background
knowledge on TB was already high (34). After all, the Gambia has a long-lasting,
ongoing TB case-contact platform (TBCC) in which thousands of people with TB and
their household contacts had been enrolled and sensitized over the years (35). Such
baseline knowledge studies should be done in areas where sensitization
interventions are planned to be carried out, to ensure that they will have an impact,
or are indeed necessary in the first place.

Another consideration is the structuring of populations which may also affect
transmission dynamics beyond what can be modelled as done here. We assumed no
further structuring at the village level beyond assignment to trial arm. This may not be
the case as interactions may be subdivided based on habits within villages. A better
understanding of the anthropological aspects of *Mtb* transmission would help
improve such phylodynamic models going forward.

There are some diagnostic limitations which can lead to issues with further studies in
such settings. For instance, people with TB may have been missed due to the limited
sensitivity of the diagnostic tests used, explaining why the research team identified
only 35 people living with TB among 10,291 individuals who provided a sputum
sample. For instance, a molecular test such as GenXpert, which became available in
The Gambia during the trial implementation, would have probably (more than)
doubled the sensitivity of fluorescent microscopy, especially in (pre) symptomatic
people with incipient TB. However, it is unlikely GenXpert would have led to a
significant increase in case notifications from intervention clusters. Moreover,

applying GenXpert rather than Fluorescent Microscopy would have required a high
throughput set-up, with equipment and cartridges at a much higher cost (7).

Another diagnostic shortcoming was the use of the MTP64-based rapid test for
confirmation of positive MGIT cultures. After completing the trial, it was found that the
globally used MPT64-based rapid tests do not have a sufficiently high sensitivity to
confirm cultures of *Mtb* L5 and L6, (previously termed *M. africanum*), two lineage
endemic to West Africa and The Gambia. Therefore, it is possible people living with
TB were missed during primary diagnosis when a positive MGIT culture with (false)
negative MPT64-based rapid test would have been interpreted as a non-tuberculous
mycobacterium. This would not have affected the primary outcome of the ECF trial
which was based on TB diagnosed by the TB diagnostic and treatment centers rather
than on the outcome of primary isolation. In the future, whole genome sequencing
directly conducted on primary sputum could be a solution to avoid any culture bias
towards certain mycobacterial lineages.

Moreover, the high workload for field teams and laboratory staff warrants evaluation
of the cost-effectiveness of such intervention in conditions as seen in The Gambia. A
strength of our study is that we demonstrate for the first time that a combination of
whole genome sequencing of bacterial populations followed by Bayesian
phylodynamic analysis can be successfully used to estimate key transmission
parameters in TB ECF studies. We suggest that such phylodynamic approaches be
used in conjunction with classic epidemiology approaches to assess the impact of
interventions on transmission dynamics. This could be beneficial, especially if
combined with or assessing other types of active case finding such as community
worker referrals(36). Such molecular epidemiological approaches can add an
additional outcome measure for clinical trials and studies such as this in other
settings.

Conclusions: As ECF in a Gambian low TB prevalence setting did not reduce case
notification nor block transmission, we recommend for any future study to carefully
consider the underlying TB prevalence, and to use a more sensitive diagnostic
algorithm. Molecular studies confirmed that the ECF strategy did not impact TB
transmission, an effect that could have been masked when solely monitoring case
notifications.

Acknowledgements: In memoriam, Jacob Kweku Otu was the Higher Scientific Officer, MRC TB laboratory and was very dedicated to providing high quality diagnostic services for people living with tuberculosis and study participants.

Funding: This work was supported by an ERC grant [INTERRUPTB; no. 311725] to BdJ and by the Global Fund for Tuberculosis, AIDS and Malaria. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authorship contributions:
FG, BCdJ, MA developed and conducted the InterrupTB study; IA, MA, AR, UDA designed and conducted the TB-ECF study; FO, KJ coordinated all field work and conducted the ECF intervention, CJM, WM, BOA, and EMW conducted the bioinformatics component of the InterrupTB; FG, BOA, TF, FM, TJ, EG, AJR, BSK, KO, ED, WDA, KM, BS, FK were responsible for all mycobacteriology activities in both TB-ECF and InterrupTB; KL, AKM, SA, SD, DJ were part of data management and statistics team in both studies. All authors critically reviewed the manuscript.

Competing interests: The authors declare no competing interests.

References

1. Silva S, Arinaminpathy N, Atun R, Goosby E, Reid M. Economic impact of tuberculosis mortality in 120 countries and the cost of not achieving the Sustainable Development Goals tuberculosis targets: a full-income analysis. Lancet Glob Health. 2021 Oct 1;9(10):e1372–9.

2. Creswell J, Khan A, Bakker MI, Brouwer M, Kamineni VV, Mergenthaler C, et al. The TB REACH Initiative: Supporting TB Elimination Efforts in the Asia-Pacific. Trop Med Infect Dis. 2020 Dec;5(4):164.

3. Rouillon A, Perdrizet S, Parrot R. Transmission of tubercle bacilli: The effects of chemotherapy. Tubercle. 1976 Dec;57(4):275–99.

4. MacPherson P, Stagg HR, Schwalb A, Henderson H, Taylor AE, Burke RM, et al. Impact of active case finding for tuberculosis with mass chest X-ray screening in Glasgow, Scotland, 1950–1963: An epidemiological analysis of historical data. PLOS Med. 2024 Nov 5;21(11):e1004448.

5. Detjen AK, DiNardo AR, Leyden J, Steingart KR, Menzies D, Schiller I, et al.
Xpert MTB/RIF assay for the diagnosis of pulmonary tuberculosis in
children: a systematic review and meta-analysis. Lancet Respir Med. 2015
June 1;3(6):451–61.

6. Corbett EL, Bandason T, Duong T, Dauya E, Makamure B, Churchyard GJ, et al.
Comparison of two active case-finding strategies for community-based
diagnosis of symptomatic smear-positive tuberculosis and control of
infectious tuberculosis in Harare, Zimbabwe (DETECTB): a cluster-
randomised trial. The Lancet. 2010 Oct 9;376(9748):1244–53.

7. Marks GB, Nguyen, Nhung V., Nguyen, Phuong T.B., Nguyen, Thu-Anh, Nguyen
Hoa B., Tran Khoa H., et al. Community-wide Screening for Tuberculosis in a
High-Prevalence Setting. N Engl J Med. 2019 Oct 3;381(14):1347–57.

8. Klinkenberg E, Floyd S, Shanaube K, Mureithi L, Gachie T, de Haas P, et al.
Tuberculosis prevalence after 4 years of population-wide systematic TB
symptom screening and universal testing and treatment for HIV in the HPTN
071 (PopART) community-randomised trial in Zambia and South Africa: A
cross-sectional survey (TREATS). PLoS Med. 2023 Sept;20(9):e1004278.

9. Ayles H, Muyoyeta M, Toit ED, Schaap A, Floyd S, Simwinga M, et al. Effect of
household and community interventions on the burden of tuberculosis in
southern Africa: the ZAMSTAR community-randomised trial. The Lancet.
2013 Oct 5;382(9899):1183–94.

10. Burke RM, Nliwasa M, Feasey HRA, Chaisson LH, Golub JE, Naufal F, et al.
Community-based active case-finding interventions for tuberculosis: a
systematic review. Lancet Public Health. 2021 May;6(5):e283–99.

11. Behr MA, Kaufmann E, Duffin J, Edelstein PH, Ramakrishnan L. Latent
Tuberculosis: Two Centuries of Confusion. Am J Respir Crit Care Med. 2021
July 15;204(2):142–8.

12. Touray K, Adetifa IM, Jallow A, Rigby J, Jeffries D, Cheung YB, et al. Spatial
analysis of tuberculosis in an urban west African setting: is there evidence of
clustering? Trop Med Int Health TM IH. 2010 June;15(6):664–72.

13. Sanoussi CN, Affolabi D, Rigouts L, Anagonou S, de Jong B. Genotypic
characterization directly applied to sputum improves the detection of
Mycobacterium africanum West African 1, under-represented in positive
cultures. Picardeau M, editor. PLoS Negl Trop Dis. 2017
Sept;11(9):e0005900–e0005900.

14. Lempens P, Meehan CJ, Vandelannoote K, Fissette K, de Rijk P, Van Deun A, et
al. Isoniazid resistance levels of Mycobacterium tuberculosis can largely be
predicted by high-confidence resistance-conferring mutations. Sci Rep. 2018
Dec;8(1):3246–3246.

15. Phelan JE, O'Sullivan DM, Machado D, Ramos J, Oppong YEA, Campino S, et al.
Integrating informatics tools and portable sequencing technology for rapid

detection of resistance to anti-tuberculous drugs. Genome Med. 2019 June
24;11(1):41.

16. Ngabonziza JCS, Rigouts L, Torrea G, Decroo T, Kamanzi E, Lempens P, et al.
Multidrug-resistant tuberculosis control in Rwanda overcomes a successful
clone that causes most disease over a quarter century. J Clin Tuberc
Mycobact Dis. 2022 Jan 24;27:100299.

17. Kim D, Song L, Breitwieser FP, Salzberg SL. Centrifuge: rapid and sensitive
classification of metagenomic sequences. Genome Res. 2016
Dec;26(12):1721–9.

18. Kohl TA, Utpatel C, Schleusener V, De Filippo MR, Beckert P, Cirillo DM, et al.
MTBseq: a comprehensive pipeline for whole genome sequence analysis of
Mycobacterium tuberculosis complex isolates. PeerJ. 2018;6:e5895.

19. Comas I. Genome of the inferred most recent common ancestor of the
Mycobacterium tuberculosis complex [Internet]. Zenodo; 2019 [cited 2024
May 17]. Available from: https://zenodo.org/records/3497110

20. Meehan CJ, Moris P, Kohl TA, Pečerska J, Akter S, Merker M, et al. The
relationship between transmission time and clustering methods in
Mycobacterium tuberculosis epidemiology. EBioMedicine. 2018
Aug;37(0):410–6.

21. R Core Team. R: A language and environment for statistical computing
[Internet]. 2021. Available from: https://www.R-project.org/

22. Dorai-Raj, Sundar. binom: Binomial Confidence Intervals for Several
Parameterizations [Internet]. 2022. Available from: https://CRAN.R-
project.org/package=binom

23. Wickham, Hadley. The Split-Apply-Combine Strategy for Data Analysis
[Internet]. 2011. Available from: https://www.jstatsoft.org/v40/i01/

24. Wickham, Hadley. ggplot2: Elegant Graphics for Data Analysis [Internet].
2016. Available from: https://ggplot2.tidyverse.org

25. Kühnert D, Stadler T, Vaughan TG, Drummond AJ. Phylodynamics with
Migration: A Computational Framework to Quantify Population Structure
from Genomic Data. Mol Biol Evol. 2016 Aug 1;33(8):2102–16.

26. Gavryushkina A, Welch D, Stadler T, Drummond AJ. Bayesian Inference of
Sampled Ancestor Trees for Epidemiology and Fossil Calibration. PLOS
Comput Biol. 2014 Dec 4;10(12):e1003919.

27. Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M,
Gavryushkina A, et al. BEAST 2.5: An advanced software platform for
Bayesian evolutionary analysis. PLOS Comput Biol. 2019 Apr
8;15(4):e1006650.

28. Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu CH, Xie D, et al. BEAST 2: A
Software Platform for Bayesian Evolutionary Analysis. PLOS Comput Biol.
2014 Apr 10;10(4):e1003537.

29. Leaché AD, Banbury BL, Felsenstein J, de Oca A nieto-Montes, Stamatakis A.
Short Tree, Long Tree, Right Tree, Wrong Tree: New Acquisition Bias
Corrections for Inferring SNP Phylogenies. Syst Biol. 2015 Nov 1;64(6):1032–
47.

30. Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. Posterior
Summarization in Bayesian Phylogenetics Using Tracer 1.7. Syst Biol. 2018
Sept 1;67(5):901–4.

31. Stuck L, Klinkenberg E, Abdelgadir Ali N, Basheir Abukaraig EA, Adusi-Poku
Y, Alebachew Wagaw Z, et al. Prevalence of subclinical pulmonary
tuberculosis in adults in community settings: an individual participant data
meta-analysis. Lancet Infect Dis [Internet]. 2024 Mar 12 [cited 2024 May 17];
Available from:
https://www.sciencedirect.com/science/article/pii/S1473309924000112

32. World Health Organization. WHO operational handbook on tuberculosis:
module 2: screening: systematic screening for tuberculosis disease
[Internet]. World Health Organization; 2021 [cited 2024 May 17]. Available
from: https://www.who.int/publications-detail-redirect/9789240022614

33. Hatherell HA, Didelot X, Pollock SL, Tang P, Crisan A, Johnston JC, et al.
Declaring a tuberculosis outbreak over with genomic epidemiology. Microb
Genomics. 2016 May;2(5):e000060.

34. Price, S. The Effect of a Community Sensitization Intervention on Knowlegde
of TB in The Gambia. 2013.

35. Hill PC, Ota MOC. Tuberculosis case-contact research in endemic tropical
settings: design, conduct, and relevance to other infectious diseases. Lancet
Infect Dis. 2010 Oct;10(10):723–32.

36. Adejumo AO, Azuogu B, Okorie O, Lawal OM, Onazi OJ, Gidado M, et al.
Community referral for presumptive TB in Nigeria: a comparison of four
models of active case finding. BMC Public Health. 2016 Feb 23;16(1):177.

Tables

Table 1. Overview of WGS data. A total of 1294 samples were sequenced with 84%

coming from intervention or control villages. The vast majority of samples were from

L4 or L6.

Lineage	Control	Intervention	Corridor	Total	Percentage
Animal	0	1	0	N/A	N/A
L1	7	11	4	22	1.7

L2	21	14	7	42	3.25
L3	8	3	1	12	0.93
L4	358	370	144	872	67.39
L5	0	2	1	3	0.23
L6	161	135	46	342	26.43
Total:	555	536	203	1294	
Percentage	42.89	41.42	15.69		

Table 2. Effect of intervention on effective reproduction number (R_e) of Mtb.
Phylodynamic analyses were used to examine if there was a reduction in the R_e in
the intervention arm of the trial vs the control. Although the mean calculated effect of
the intervention across the Bayesian analysis run is below 1, the 95% HPD interval
includes 1, indicating no significant effect. *see Supplementary Table 1 and Methods
for default settings and priors

Approach	Effect mean	Effect 95% HPD	Control R_e mean	Control R_e 95% HPD	Intervention R_e mean	Intervention R_e 95% HPD
Default*	0.72	[0.39,1.06]	1.30	[0.99,1.62]	0.90	[0.64,1.17]
Unequal sampling proportions	0.58	[0.16,1.03]	1.52	[0.94,2.24]	0.80	[0.46,1.16]
Lognormal(-17,1) prior on clock rate	0.72	[0.39,1.06]	1.29	[0.99,1.63]	0.90	[0.64,1.17]
Lognormal(0,2) prior on effect parameter f	0.70	[0.38,1.06]	1.31	[0.99,1.64]	0.89	[0.62,1.17]

Figure Legends:

Figure 1: An overview of locations of ECF intervention and control clusters in

the Greater Banjul Area. Map created using OpenStreetMap and distributed under

CC-BY license.

Figure 2: Flowchart of study population recruitment into intervention and

control areas within the TB-ECF trial as well as in InterrupTB. Samples for the

TB-ECF trial were collected from patients in intervention areas via sensitisation

clinics and encouragement to present at health clinics and from control areas through

passive presentation at health clinic only. The breakdown of these groups is outlined

in Supplementary Table 2. A subset of these isolates were whole genome

sequenced for the transmission interruption assessment in InterrupTB.

Figure 3: Longitudinal development of the study case notification rate

difference in relation to three rounds of ECF intervention. Displayed is the 60-

day moving average TB case notification rate difference (CNRD, red line) and 95%CI

(light red lines) between intervention and control clusters (for all clusters with more

than 30 screened people).

Figure 4: Estimating TB transmission in intervention and control clusters

based on whole genome sequencing of bacterial populations and genetic

clustering based on varying SNP thresholds. Clusters are assigned to an arm

(intervention and control) using two methods: >50%: if over 50% of the samples in

cluster come from a specific arm, the clustered is designated as being from that arm

(and clusters that are 50:50 to each arm are discounted); time: a cluster is assigned

to an arm based on the assignment of the sample with the earliest sampling date.

Bar charts indicate the proportion of clusters assigned to each arm base don the

related method (excluding those clusters not assigned to either arm due to being

50:50). Error bars on each group indicate the Wilson confidence interval around the

estimate.

Figure 5: Estimating the impact of the intervention on the effective

reproductive number (R_e) of *Mycobacterium tuberculosis*. A) Prior and posterior

distribution for the multiplicative effect of enhanced case finding on the effective

reproductive number (*f* parameter in the phylodynamic model). The genomic data

was found to have enough signal for analysis, as shown by the posterior distribution

(prior + data) being different from the prior distribution. The intervention did not result

in a significant decrease in the R_e compared to the control arm, as shown by the

posterior distribution overlapping with 1 (= no effect). B) Posterior distributions for the

R_e of cases in the two study arms, again showing no significant difference.

Supplementary information legends

Supplementary Figure 1. Approaches for assigning clusters to control or

intervention arms. A mock phylogenetic tree showing a transmission cluster based

on a SNP cut-o; is shown above. It contains five samples, three from control areas

and two from intervention areas. The year of isolation is indicated after the

underscore on each sample name. The majority counts cluster assignment approach

would assign this cluster to the control arm as >50% of the samples in the cluster are

from the control arm. The earliest sample cluster assignment method would assign

this cluster to the control arm as the sample taken at the earliest timepoint (C1_2013)

is from the control arm.

Supplementary Table 1. Prior distributions for the parameters of the

phylodynamic model.

Supplementary Table 2. Epidemiological overview of the dataset. Aggregated
associated meta and clinical data for the participants are shown for the main study
(ECF), the sub-study of only samples with an associated genome sample
(InterrupTB). The proportion of the total study/sub-study included in that group is
shown with the difference in proportion between the main and sub-study. Where
these proportions differed by more than 5% are highlighted in bold.
712
713 **Figures**
714 **Figure 1:**
715

716
717
718 **Figure 2:**

720 **Figure 3:**

724   **Figure 4:**

725
726
727   **Figure 5:**

728