Human genetics implicate thromboembolism in the pathogenesis of long COVID in individuals of European ancestry

Art Schuermans[1,2,3,*], Andreas Verstraete[1,4], Vilma Lammi[5], Tomoko Nakanishi[6-10], Maddalena Ardissino[11,12], Jef Van den Eynde[1], Benjamin B. Sun[13,14], Marios K. Georgakis[2,15,16], Beatriz Guillen-Guio[17,18], Louise V. Wain[17,18], Christopher E. Brightling[18,19], PHOSP-COVID Collaborative Group, Johan Van Weyenbergh[20], Adam J. Lewandowski[21], Betty Raman[22], Hugo Zeberg[23,24], Hanna M. Ollila[2,3,5,25], Stephen Burgess[26], Pradeep Natarajan[2,3,27], Michael C. Honigberg[2,3,27], Kathleen Freson[1], Thomas Vanassche[1,4], Peter Verhamme[1,4].

[1] KU Leuven Department of Cardiovascular Sciences, Center for Molecular and Vascular Biology, Leuven, Belgium

[2] Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[3] Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA

[4] UZ Leuven Department of Cardiovascular Diseases, Leuven, Belgium

[5] Institute for Molecular Medicine Finland, Helsinki Institute of Life Science, University of Helsinki, Helsinki, Finland

[6] Department of Human Genetics, McGill University, Montréal, QC, Canada

[7] Lady Davis Institute, Jewish General Hospital, McGill University, Montréal, QC, Canada

[8] Kyoto-McGill International Collaborative Program in Genomic Medicine, Graduate School of Medicine, Kyoto University, Kyoto, Japan

[9] Department of Genome Informatics, Graduate School of Medicine, the University of Tokyo, Tokyo, Japan

[10] Japan Society for the Promotion of Science, Tokyo, Japan

[11] BHF Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

[12] National Heart and Lung Institute, Imperial College London, London, UK

[13] Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

[14] Human Genetics, Informatics and Predictive Sciences, Research, Bristol Myers Squibb, USA

[15] Institute for Stroke and Dementia Research, Ludwig Maximilian University of Munich, Munich, Germany

[16] Munich Cluster for Systems Neurology (SyNergy), Munich, Germany

[17] Department of Population Health Sciences, University of Leicester, Leicester, UK

[18] NIHR Leicester Biomedical Research Centre, Leicester, United Kingdom

[19] Institute for Lung Health, Department of Respiratory Sciences, University of Leicester, Leicester, UK

[20] Laboratory of Clinical and Epidemiological Virology, Rega Institute for Medical Research, KU Leuven, Leuven, Belgium

[21] Nuffield Department of Population Health, University of Oxford, Oxford, UK

[22] Division of Cardiovascular Medicine, Radcliffe Department of Medicine, University of Oxford, Oxford, UK

[23] Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, Sweden

[24] Department of Evolutionary Genetics, Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany

[25] Anesthesia, Critical Care, and Pain Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, USA

[26] MRC Biostatistics Unit, University of Cambridge, Cambridge, UK

[27] Department of Medicine, Harvard Medical School, Boston, MA, USA

[*]**Correspondence to:**

Art Schuermans

KU Leuven Department of Cardiovascular Sciences

Center of Molecular and Vascular Biology

Herestraat 49 (box 911)

3000 Leuven, Belgium

Tel: +32 16 343491

Fax: +32 16 343493

Email: aschuerm@broadinstitute.org

Short title: Thromboembolism and long COVID

Abstract word count: 150

Word count (excluding methods): 3,965

Figure/table count: 5

Abstract

SARS-CoV-2 infection can result in long COVID, characterized by post-acute

symptoms from multiple organs. Current hypotheses on mechanisms underlying long COVID

include persistent inflammation and thromboembolism; however, compelling evidence from

humans is limited and causal associations remain unclear. Here, we tested the association of

thromboembolism-related genetic variants with long COVID in the Long COVID Host

Genetics Initiative (n_{cases}=3,018; $n_{controls}$=994,582). Primary analyses revealed that each unit

increase in the log-odds of genetically predicted venous thromboembolism risk was

associated with 1.21-fold odds of long COVID (95%CI: 1.08-1.35; P=1.2×10^{-3}). This

association was independent of acute COVID-19 severity, robust across genetic instruments

and methods, and replicated in external datasets for both venous thromboembolism and long

COVID. Downstream analyses using gene-specific instruments, along with protein and gene

expression data, suggested the protease-activated receptor 1 (PAR-1) as a potential

molecular contributor to long COVID. These findings provide human genetic evidence

implicating thromboembolism in long COVID pathogenesis.

Main

Infection with SARS-CoV-2 can lead to different acute clinical manifestations, ranging
from mild respiratory disease to multi-organ dysfunction. It is becoming increasingly clear
that infection with SARS-CoV-2 also has chronic effects, with epidemiological data
suggesting that 5-10% of acutely infected individuals experience persistent complaints three
months after infection[1]. These post-acute symptoms—referred to as "long COVID"—are not
limited to the respiratory tract but involve various organ systems[2]. Common manifestations
include physical (e.g., fatigue, post-exertional malaise, and shortness of breath), cognitive
(e.g., memory loss and cognitive impairment), and psychological (e.g., anxiety and
depression) symptoms[3]. Up to 85% of long COVID patients report ongoing complaints even
after one year[4,5]. Despite its high prevalence and impact on quality of life, clinical strategies
for the prevention and treatment of long COVID remain limited.

The lack of disease-specific therapies for long COVID can be partially ascribed to an
incomplete understanding of its mechanistic drivers. Currently hypothesized mechanisms
underlying long COVID include immune dysregulation, virus-induced inflammation, and
endothelial dysfunction[3]. For instance, in a prospective cohort study of 1,837 adults
hospitalized with COVID-19, selected blood markers of thromboinflammation (i.e., fibrinogen
and D-dimer relative to C-reactive protein) predicted cognitive defects at 6-12 months after
infection[6]. Another recent study of 113 individuals infected with SARS-CoV-2 revealed that
the development of long COVID was associated with changes to blood proteins indicative of
persistent complement activation and altered coagulation[7]. These primarily observational
findings have prompted some physicians to offer treatments like apheresis and
anticoagulation therapy to long COVID patients[8]. However, the evidence supporting these
treatments is limited, and it remains unclear whether thromboinflammation plays a causal
role in the development of long COVID.

Large-scale genome-wide association studies (GWAS) have identified numerous
genetic variants associated with human diseases[9], facilitating the identification of causal
mechanisms using Mendelian randomization (MR)[10,11]. MR is an epidemiological method that
uses the random assortment of alleles at conception, which leads to an effective
randomization of individuals to different levels of genetic predisposition for specific traits or
conditions[12,13]. This randomization limits the risk of bias due to confounding and reverse
causation and can therefore provide support for a causal relationship between an exposure
and outcome of interest[10,11]. Previous MR analyses have identified causal risk factors (e.g.,
body mass index) and molecular players (e.g., the interleukin-6 receptor) for acute COVID-
19[14–16], with clinical trial data confirming some of these as effective therapeutic targets in
human patients[17,18]. Given the limitations and high costs associated with traditional methods

for identifying causal mediators of diseases, genetic approaches such as MR may also help

prioritize new therapeutic targets for long COVID[19].

Here, we used MR to test the hypothesis that thromboembolism contributes causally

to long COVID. In primary analyses, we constructed a genetic instrument for venous

thromboembolism and estimated its causal effect on the development of long COVID. We

assessed the specificity of this association and tested its robustness using different

methodological approaches and external datasets. Finally, we performed gene-focused

analyses to identify potential molecular mediators of long COVID and gain further insights

into the possible biological mechanisms driving this condition.

Results

Overall analysis approach

The study design is shown in **Fig. 1**. In brief, we used a two-sample MR approach to

test the association of thromboembolism-related genetic variants with long COVID. Two-

sample MR leverages genetic data from two separate GWAS: the first one is used to

construct a genetic instrument for the exposure of interest (e.g., venous thromboembolism),

while the second is used to test the association of this instrument with the outcome of

interest (e.g., long COVID)[10,20]. When an MR analysis adheres to specific assumptions (**Fig.**

1; see **Methods**), it provides a robust framework for testing the causal association of a given

exposure with the corresponding outcome[10,20].

In line with this framework, we first identified 35 genetic variants associated with

venous thromboembolism in FinnGen. These variants were then combined into a single

statistical entity, i.e., a genetic instrument. Primary analyses tested the association of this

genetic instrument (referred to as "genetically predicted venous thromboembolism risk") with

long COVID in the Long COVID Host Genetics Initiative. Downstream analyses (1) assessed

the robustness of this association using different methods and external datasets; (2) tested

the effects of genetically predicted venous thromboembolism risk on specific conditions with

symptom profiles shared with long COVID (referred to as "long COVID-resembling

phenotypes"); and (3) explored molecular mediators of long COVID using gene-focused MR

approaches (**Fig. 1**).

Cohort characteristics

The main genetic instrument for venous thromboembolism was constructed in

FinnGen[21], using data from 412,181 participants (venous thromboembolism: n_{cases}=21,021

and $n_{controls}$=391,160) from ten contributing Finnish biobanks. Venous thromboembolism

cases were identified using diagnostic codes from hospital discharge and death cause

registers (see **Methods**). Of the included participants, 44.1% were male, and the median age

was approximately 62.9 years[22,23].

90   The association of genetically predicted venous thromboembolism risk with long

COVID was tested in the Long COVID Host Genetics Initiative[24], characteristics of which are

provided in **Supplementary Table 1**. While the Long COVID Host Genetics Initiative

included 25 cohorts (long COVID: n_{cases}=6,450 and $n_{controls}$=1,093,995), primary analyses

excluded participants who met the criteria for long COVID but did not have a prior test-

verified SARS-CoV-2 infection and therefore only used data from 12 cohorts (n_{cases}=3,018;

$n_{controls}$=994,582). Of these 12 cohorts, the majority (83.3%) used questionnaires rather than

electronic health data to ascertain long COVID (**Supplementary Table 1**). Almost all

participants (99.1%) included in these cohorts had European ancestry, 48.0% were male,

and the weighted mean age across cohorts was 62.3 years[24]. Around 30.5% of long COVID

cases had a history of hospitalization due to COVID-19.

Thromboembolism-promoting genetic variants increase long COVID risk

103   Our main genetic instrument included 35 independent genetic variants (linkage

disequilibrium R^2<0.001) strongly associated with venous thromboembolism (P<5×10⁻⁸) in

FinnGen (**Supplementary Table 2**). The mean F-statistic was 114.5, suggesting adequate

strength[10]. In primary analyses, we tested the effect of this genetic instrument on long COVID

risk in the Long COVID Host Genetics Initiative using the inverse-variance weighted MR

method. Each unit increase in the log-odds of genetically predicted venous thromboembolism

risk was associated with 1.21-fold odds of developing long COVID (95% confidence interval

[CI]: 1.08-1.35; P=1.2×10⁻³; **Fig. 2**), indicating a significant positive association of

thromboembolism-promoting genetic variants with long COVID.

112   To contextualize these findings, we assessed whether genetic instruments for other

conditions were associated with long COVID using a similar approach. First, to corroborate

the observed association of venous thromboembolism with long COVID, we tested the

effects of genetic instruments for the most common manifestations of venous

thromboembolism—deep vein thrombosis and pulmonary embolism—on long COVID risk.

Next, to evaluate the specificity of these associations, we extended our analysis to a range of

conditions representing different organ systems and pathophysiological pathways (e.g., type

2 diabetes, dementia, rheumatoid arthritis). These analyses showed that only genetically

predicted risk for deep vein thrombosis (odds ratio [OR], 1.10 [95%CI, 0.98-1.22] per log-

odds increase in genetically predicted deep vein thrombosis risk; P=9.3×10⁻²) and pulmonary

embolism (OR, 1.17 [95%CI, 1.02-1.34] per log-odds increase in genetically predicted

pulmonary embolism risk; P=2.2×10⁻²) had effect estimates within the 95%CI of the venous

thromboembolism-long COVID association. None of the conditions less closely related to

venous thromboembolism had effect estimates within this 95%CI or a positive association

reaching nominal significance (**Fig. 2** and **Supplementary Table 3**). These findings support

a specific association of venous thromboembolism with long COVID rather than a

generalized association across pathophysiologically distinct conditions.

Sensitivity and replication analyses

To evaluate the potential role of horizontal pleiotropy (i.e., effects of a genetic

instrument on the outcome outside of its effects on the exposure), we tested the association

of genetically predicted venous thromboembolism risk with long COVID using MR-Egger[25].

MR-Egger showed no evidence of horizontal pleiotropy ($P_{intercept}$>0.99), with an effect

estimate closely aligned with our primary analysis (OR, 1.21 [95%CI, 1.00-1.46] per log-odds

increase in genetically predicted venous thromboembolism risk; P=5.1×10^{-2}). This suggests

that genetic variants associated with venous thromboembolism affect long COVID risk via

thromboembolic mechanisms rather than alternative pathways. Consistent findings were

observed with the weighted median (OR, 1.26 [95%CI, 1.06-1.49] per log-odds; P=7.4×10^{-3})

and mode-based (OR, 1.33 [95%CI, 1.12-1.56] per log-odds; P=8.5×10^{-4}) estimators,

providing additional support for a pleiotropy-robust association.

We also performed sensitivity analyses using different linkage disequilibrium R^2

(R^2<0.001/R^2<0.001/R^2<0.01/R^2<0.1) and P-value thresholds (P<5×10^{-4}/P<5×10^{-6}/P<5×10^{-8}/P<5×10^{-10}) to construct our genetic instrument for venous thromboembolism. These

analyses showed that the association of venous thromboembolism with long COVID was

directionally consistent across all tested parameters (**Fig. 3**; **Supplementary Table 4**).

Additionally, we tested for reverse causation (i.e., long COVID increasing venous

thromboembolism risk) by applying Steiger filtering. While Steiger filtering can be used to

identify variants explaining more variance in the outcome than in the exposure of interest[26],

we did not identify any such reverse causal variants. Correspondingly, MR analyses in the

opposite direction (see **Methods**)—evaluating the effect of genetically predicted long COVID

on venous thromboembolism risk—yielded null results (OR, 1.01 [95%CI, 0.94–1.09] per log-

odds increase in genetically predicted long COVID risk; P=0.79), providing no evidence for

reverse causation.

Replication analyses evaluated whether the observed associations were consistent

across different exposure and outcome datasets (see **Methods**). First, we constructed an

externally derived genetic instrument for venous thromboembolism using data from the

Million Veteran Program[27] and tested its association with long COVID in the Long COVID

Host Genetics Initiative (**Supplementary Table 5**). Second, we tested the genetic

association of venous thromboembolism with long COVID in a subcohort of the Long COVID

Host Genetics Initiative that applied stricter criteria for controls, restricting to participants with

proven SARS-CoV-2 infection[24] (**Supplementary Table 6**). Third, we tested this association

in the Post-Hospitalization COVID-19 (PHOSP-COVID) study, a prospective cohort study

designed to investigate the medium- and long-term sequelae of hospitalized COVID-19[28,29]

(**Supplementary Table 7**). All three replication analyses demonstrated a significant positive

association of genetically predicted venous thromboembolism risk with long COVID (**Fig. 3**).

Additional sensitivity analyses, performed using the same replication datasets with varying

linkage disequilibrium R^2 and P-value thresholds for genetic instrument construction, yielded

consistent results throughout (**Fig. 3**; **Supplementary Tables 5-7**).

The thromboembolism-long COVID link does not reflect acute COVID-19 severity

Given that thrombotic events are frequently observed in individuals with COVID-19,

correlating partly with disease severity[30,31], and that severe COVID-19 is a thought to be a

causal risk factor for long COVID[24], we investigated whether acute COVID-19 severity might

mediate the genetic association of venous thromboembolism with long COVID. Specifically,

we tested the effects of genetically predicted venous thromboembolism risk across three

categories of acute COVID-19 severity in the COVID-19 Host Genetics Initiative[16,32]: SARS-

CoV-2 infection, hospitalized COVID-19, and critical COVID-19 (defined as hospitalization

with respiratory support or COVID-19-related death). Our results showed that genetically

predicted venous thromboembolism risk had effect estimates within the 95%CI for its

association with long COVID for both hospitalized COVID-19 (OR, 1.09 [95%CI, 1.02-1.16]

per log-odds increase in genetically predicted venous thromboembolism risk; P=1.6×10^{-2})

and critical COVID-19 (OR, 1.08 [95%CI, 0.99-1.17] per log-odds; P=6.6×10^{-2}), but not for

SARS-CoV-2 infection alone (**Fig. 4a**). Only the association with hospitalized COVID-19

reached nominal significance.

Building on these findings, we performed multivariable MR[33] to estimate the direct

effects of genetic predisposition to venous thromboembolism on long COVID while adjusting

for its impact on acute COVID-19 severity. Consistent with previous findings[24], all acute

COVID-19 phenotypes had significant genetic associations with long COVID in these

multivariable MR models (**Supplementary Table 8**). However, genetically predicted venous

thromboembolism was also significantly associated with long COVID across all models

(**Supplementary Table 8**), suggesting that its effects on long COVID are not mediated by

acute COVID-19 severity alone. These findings imply that a predisposition to

thromboembolism contributes to the risk of developing long COVID through mechanisms that

are not solely dependent on the severity of acute infection.

Predisposition to thromboembolism does not promote symptoms shared with long COVID

To gain additional insights into the mechanisms driving the genetic association of

venous thromboembolism with long COVID, we tested the associations of genetically

predicted venous thromboembolism risk with specific conditions with symptom profiles

shared with long COVID. Genetically predicted risk of venous thromboembolism was not

associated with a higher risk of long COVID-resembling conditions or symptoms such as

memory loss or mood disorders (**Fig. 4b**; **Supplementary Tables 9-10**). Although there was

no significant association with postviral fatigue (OR, 1.13 [95%CI, 0.88-1.44] per log-odds

increase in genetically predicted venous thromboembolism risk; P=0.35), confidence

intervals were very wide, suggesting this association was potentially underpowered due to a

small number of cases in FinnGen's GWAS of postviral fatigue (n_{cases}=195; $n_{controls}$=382,198).

Nevertheless, these findings suggests the genetic association of venous thromboembolism

with long COVID represents a specific association, wherein thromboembolism plays a role in

the pathophysiology of long COVID rather than the occurrence of its individual, non-specific

symptoms.

Exploration of potential molecular mediators

Finally, we aimed to identify potential molecular mediators of the thromboembolism-

long COVID association by evaluating the effects of individual genes and their associated

proteins on long COVID risk. First, we constructed "gene-specific genetic instruments" by

identifying variants strongly associated with venous thromboembolism within or near different

coagulation-related genes (see **Methods**). Gene-specific instruments consisting of one or

multiple independent variants were available for a total of 40 genes (**Supplementary Tables**

11-12). Most genes' instruments included only one genetic variant (n=32/40 [80.0%]). All

mean F-statistics of the gene-specific instruments were greater than 10 (mean, 65.2; range,

12.9-377.4), indicating adequate instrument strength[10]. Among the different gene-specific

instruments, *F2R*'s instrument (rs73131633) had the strongest effect on long COVID risk

(**Fig. 5a**). Each unit increase in the log-odds of genetically predicted venous

thromboembolism risk mediated through *F2R* was nominally associated with 4.50-fold odds

of developing long COVID (95%CI, 1.12-18.2; P=3.5×10^{-2}).

The *F2R* gene encodes the protease-activated receptor 1 (PAR-1; i.e., the thrombin

receptor). PAR-1 is a G-protein-coupled receptor that is activated predominantly by thrombin

and drives thrombosis and inflammation through involvement in platelet aggregation,

endothelial activation, and leukocyte recruitment[34]. To verify its association with long COVID,

we tested the effects of genetically predicted circulating levels of PAR-1 on long COVID risk.

Genetically predicted circulating levels of PAR-1 were instrumented using a single intronic

F2R variant (rs168753; in moderate linkage disequilibrium [R^2=0.36] with rs73131633)

associated with modified PAR-1 expression on platelets leading to altered procoagulant

activity[35,36]; this variant was also the lead *F2R* signal for circulating PAR-1 levels in the UK

Biobank Pharma Proteomics Project[37]. MR analyses using this instrument revealed that each

unit increase in genetically predicted PAR-1 levels was associated with 1.80-fold odds of

developing long COVID (95%CI, 1.07-3.04; P=2.7×10^{-2}). This association was consistent

across genetic instrument construction parameters (**Fig. 5b**; **Supplementary Table 13**).

Colocalization analyses supported the hypothesis of a causal variant for trait 1 (i.e.,

circulating PAR-1 levels) but not trait 2 (i.e., long COVID) as the most likely (75.8%),

suggesting that statistical power was insufficient to discriminate between whether the genetic

association was due to shared causal variants or variants in linkage disequilibrium (i.e.,

horizontal pleiotropy)[38,39]. The probability for shared causal variants, conditional on the

presence of a causal variant for both traits, was 60.9%.

We also evaluated the association between genetically predicted *F2R* expression and

long COVID risk. Using data from the Genotype-Tissue Expression (GTEx) project[40], we

identified the tissues with the highest levels of *F2R* expression and constructed genetic

instruments for those most relevant to long COVID pathophysiology. As GTEx lacks data on

blood platelets—an important site of *F2R* expression[34]—we supplemented our analysis with

genetic instruments for platelets from the GeneSTAR Research Study[41]. Each tissue-specific

instrument was constructed using the variant most strongly associated with *F2R* expression

within a 200-kilobase *cis*-region of the gene. As *F2R*'s *cis*-region partially overlaps with those

of other genes (e.g., *F2RL1* and *F2RL2*)[42], we also performed analyses using only intragenic

variants. Our analyses revealed that genetically predicted *F2R* expression had positive effect

estimates on long COVID risk across all tissues, with nominal significance for *F2R*

expression in platelets, lung, and spleen tissue (**Fig. 5c**; **Supplementary Table 14**). The

strongest effect was observed in platelets, indicating that genetic variants increasing *F2R*

expression in platelets are linked to higher long COVID risk, further supporting a mechanistic

link between thromboembolism and long COVID.

Discussion

Using genetic data from multiple cohorts, we tested the association of genetically

predicted risk of venous thromboembolism with long COVID. Our primary analysis

demonstrated that a higher genetically predicted risk of venous thromboembolism was

significantly associated with an increased risk of developing long COVID. This association

was robust across various sensitivity analyses, replicated using data from external cohorts,

and was independent of acute COVID-19 severity. Additional analyses revealed no genetic

associations of venous thromboembolism with any long COVID-resembling conditions (e.g.,

memory loss or mood disorders), suggesting a specific role for thromboembolism-related

pathways in long COVID. Furthermore, gene-focused analyses identified PAR-1—a thrombin

receptor strongly involved in platelet and coagulation activation—as a potential contributor to
its pathophysiology. Collectively, our findings provide new mechanistic insights and identify a
potential therapeutic target for long COVID.
These results have several implications. First, our findings suggest that
thromboembolic pathways are causally involved in the development of long COVID. This
extends earlier observational research linking thromboinflammatory biomarkers to long
COVID[6,7,43]. For instance, prior analyses have shown that blood biomarkers of coagulation
and inflammation (i.e., fibrinogen and D-dimer relative to C-reactive protein) can help predict
cognitive defects in patients hospitalized for COVID-19 up to 12 months after infection[6].
Similarly, a recent prospective study of 113 COVID-19 patients found that individuals who
developed long COVID exhibited changes to blood proteins indicative of persistent
thromboinflammation at 6 months post-infection, compared to those who did not develop
long COVID[7]. However, these studies relied predominantly on traditional observational
approaches to identify molecular risk factors for long COVID. While such approaches are
often well-suited to identify biomarkers for the detection and prediction of diseases, they
cannot always discern true causal associations from those influenced by confounding or
reverse causation[10]. MR is an approach that can help deal with these challenges by using
genetic variants as instruments for specific exposures. Because an individual's genetic
makeup is fixed at conception, MR analyses are less directly susceptible to external
influences and reverse causation than those performed using more traditional methods[10].
Consequently, this study provides human genetic evidence supporting a causal association
of thromboembolism with long COVID.
Second, while thrombotic events are common in individuals with acute COVID-19 and
correlate partially with disease severity[30,31], our findings indicate that thromboembolic
pathways may contribute to long COVID independently of acute disease severity. During the
early stages of the pandemic, one in four COVID-19 patients admitted to the intensive care
unit experienced venous thromboembolism despite standard dose thromboprophylaxis[44].
Mechanistic research suggests that this heightened risk of venous thromboembolism partially
reflects a systemic COVID-19-associated coagulopathy involving immune response
dysregulation, endothelial cell dysfunction, and hypercoagulability[45]. For instance, early work
demonstrated that SARS-CoV-2 infection is associated with platelet hyperreactivity (e.g., in
response to PAR-1 stimulation)[46] potentially contributing to the formation of microthrombi
across organs (e.g., the lungs and brain)[45,47–49]. Of note, epidemiological research indicates
that individuals with a genetic predisposition to venous thromboembolism are more likely to
experience acute thrombotic complications due to COVID-19[50], which aligns with our
observation that genetic predisposition to venous thromboembolism is associated with an
increased risk of hospitalized COVID-19. However, we also found that the genetic

association of venous thromboembolism with long COVID was not mediated by acute

disease severity. This finding is consistent with *in vitro* evidence showing that the SARS-

CoV-2 spike protein binds human fibrin, promoting persistent thromboinflammation and

neuropathology independent of active infection[51]. Together with recent reports of circulating

SARS-CoV-2 spike in individuals with long COVID[52], our findings support the hypothesis that

SARS-CoV-2 creates a substrate for sustained thromboinflammation post-infection,

potentially leading to chronic symptoms from multiple organ systems.

Third, gene-focused analyses identified PAR-1 as a potential contributor to the

pathogenesis of long COVID. PAR-1 is encoded by the *F2R* gene and represents the

prototype of a small family of G-protein-coupled protease-activated receptors (PARs)

involved in thrombin signaling[34,53]. Thrombin cleaves the amino-terminal extracellular domain

of PAR-1, resulting in the formation of a new amino terminus that binds intramolecularly to

the body of the receptor and leads to platelet activation[53]. While PAR-1 predominantly

functions as a receptor and not conventionally as a secreted protein in the circulation, we

found that higher genetically predicted plasma levels of this protein (e.g., through receptor

shedding or in extracellular vesicles) were associated with a higher risk of long COVID. As

PAR-1 is considered the main mediator of thrombin's effects on platelet aggregation, it has

gained interest as a potential drug target for the prevention of thrombotic events[54,55]. Large

trials found that inhibition of PAR-1 with vorapaxar (i.e., a competitive PAR-1 antagonist)

reduced the risk of cardiovascular death or ischemic events in selected populations[54];

however, it also increased the risk of major bleeding including intracranial hemorrhage[54,55]. In

addition to mediating thrombin-induced platelet aggregation, PAR-1 also promotes

coagulation-driven inflammation. For instance, in an experimental mouse model of sepsis,

genetic and pharmacological inhibition of PAR-1 led to reduced inflammation and coagulation

in comparison with control treatment, as did pharmacological inhibition of thrombin[56].

Consistent with these findings, data from recent mechanistic work suggest that virus-

exposed endothelium promotes thromboinflammation through a positive feedback loop

sustained and enhanced by thrombin and PAR-1, implicating PAR-1 in virus-induced

inflammation and thrombosis[57]. Together with the results from the present study, these

findings support the idea that PAR-1 is a mediator of virus-induced thromboinflammation,

potentially contributing to chronic symptoms associated with SARS-CoV-2 infection. Future

animal experiments and clinical trials are needed to evaluate the effectiveness of drugs

targeting PAR-1 or other thromboinflammatory mediators to prevent and/or treat long COVID.

Although this study benefitted from multiple large-scale and well-profiled datasets, as

well as a robust framework for causal inference using human genetics, certain limitations

must be considered when interpreting the study findings. First, while MR analyses can be

used to infer causality in exposure-outcome pairs, causal inference relies on the justification

of the underlying MR assumptions[10]. However, the present study used a robust framework

with multiple sensitivity and replication analyses and assessed the key MR assumptions

where possible (see **Methods**), minimizing the chance of systematically violating the MR

assumptions. Second, we only tested genetic instruments constructed using data from

European-ancestry cohorts, limiting generalizability to other ancestries. Third, certain

phenotypes such as (post-viral) post-exertional malaise did not have GWAS data available

and could therefore not be evaluated in relation to genetically predicted venous

thromboembolism risk. Fourth, the use of partially overlapping cohorts can lead to bias

towards the observational exposure-outcome association; however, given the limited number

of long COVID cases in FinnGen (n_{cases}=263), the binary nature of the outcome phenotype

(i.e., long COVID), and the strong genetic instrument for venous thromboembolism (mean F-

statistic, 114.5), any bias due to sample overlap was expected to be minimal[58]. Furthermore,

replication analyses with non-overlapping exposure and outcome datasets (i.e., the Million

Veteran Program and PHOSP-COVID cohorts) yielded highly consistent results, suggesting

sample overlap did not strongly affect our findings. Fifth, although the gene-specific

association of venous thromboembolism with long COVID in *F2R* had a large effect size, this

association did not reach Bonferroni-significance due to limited power. Similarly,

colocalization analyses to support the association of circulating PAR-1 and long COVID had

limited power. Nevertheless, the probability for shared causal variants, conditional on the

presence of a causal variant for long COVID, was 61.0%, which means that the probability

for shared causal variants was greater than that of distinct causal variants and suggests no

confounding due to linkage disequilibrium. Finally, while the gene-focused analyses from the

present study used a robust *cis*-MR framework facilitating adherence to the core

assumptions of MR[59–61], the identification of PAR-1 as a potential therapeutic target should

be validated in prospective intervention trials.

In conclusion, this study found a strong and robust genetic association of venous

thromboembolism with long COVID, pointing to thromboembolic pathways as potential

drivers of post-acute sequelae following SARS-CoV-2 infection. The thrombin receptor PAR-

1 was identified as a putative molecular contributor to long COVID pathogenesis. These

findings offer human genetic evidence for a causal link between thromboembolism and long

COVID and support clinical research into anticoagulant and antiplatelet therapies as

strategies for its prevention and treatment.

Methods

MR and instrumental variable assumptions

MR analyses can be used to infer causality in certain exposure-outcome

relationships, given that these analyses adhere to three key instrumental variable

assumptions[10] (**Fig. 1**). First, the tested genetic instrument must comprise genetic variants

that are associated with the exposure of interest (i.e., relevance assumption). Second, there

can be no confounding pathway between the genetic variants comprising the tested genetic

instrument and the outcome (i.e., independence assumption). Third, there can be no

horizontal pleiotropy in the association of the tested genetic instrument with the outcome,

meaning that the tested genetic instrument can only affect the outcome through its effects on

the exposure of interest (i.e., exclusion restriction assumption).

The relevance assumption was tested by estimating the *F*-statistics (indicating

instrument strength) for the variants included in the genetic instrument for venous

thromboembolism[10]. The independence and exclusion restriction assumptions were

evaluated by testing the associations of the genetic instrument for venous thromboembolism

with a range of long COVID-resembling conditions[10]. We also performed replication analyses

using a wide range of genetic instruments and pleiotropy-robust methods (e.g., MR-Egger[25])

to further evaluate the robustness of the observed associations against the aforementioned

MR assumptions.

GWAS of venous thromboembolism

Genetic association data for venous thromboembolism were obtained from

FinnGen[21]. The FinnGen study is a large-scale genomics initiative that has analyzed over

500,000 Finnish biobank samples and correlated genetic variation with health data to

understand disease mechanisms and predispositions. The present study used data from

FinnGen data release 10, which includes 412,181 participants with genetic and phenotype

data. The Coordinating Ethics Committee of the Hospital District of Helsinki and Uusimaa

(HUS) approved the FinnGen study protocol, and all participants provided informed

consent[21].

The genetic instrument for venous thromboembolism (I9_VTE) was constructed using

a GWAS in 21,021 venous thromboembolism cases and 391,160 controls[23]. Venous

thromboembolic events were defined using diagnostic codes from hospital discharge and

death cause registers. In brief, qualifying events included diagnoses of pulmonary embolism

and deep vein thrombosis. The exact codes used to construct the venous thromboembolism

phenotype are listed in **Supplementary Table 15**. Participants were genotyped using

different Illumina (Illumina Inc., San Diego, CA, USA) and Affymetrix arrays (Thermo Fisher

Scientific, Santa Clara, CA, USA)[62]. Samples with ambiguous gender, high genotype

missingness (>5%), excess heterozygosity (more than 4 standard deviations from the mean)

and non-Finnish ancestry were excluded[62]. Samples were imputed to the 1000 Genomes

Project reference panel (phase 3)[63] prior to association analyses adjusted for age, sex, ten

principal components of ancestry, FinnGen chip version 1 or 2, and legacy genotyping batch

as covariates[62]. Variants with high missingness (>2%) and Hardy-Weinberg equilibrium

$P<1\times10^{-6}$ were excluded[62]. Variants with minor allele frequency <1% were also excluded.

GWAS of long COVID

Genetic association data for long COVID were obtained from the Long COVID Host

Genetics Initiative[24]. Data from a total of 3,018 long COVID cases and 994,582 controls from

12 individual cohorts were included for primary analyses (**Supplementary Table 1**).

Consistent with the genetic ancestry of the exposure cohort, more than 99% of participants

had European ancestry. The maximal overlap in effective sample size (calculated as [4 ×

$n_{case} \times n_{control}$] / $[n_{case} + n_{control}]^{24}$) between the Long COVID Host Genetics Initiative and

FinnGen[21] cohorts was 10.3%. Given the limited number of long COVID cases in FinnGen

(n_{cases}=263), the binary nature of the outcome phenotype (i.e., long COVID), and the strong

genetic instrument for venous thromboembolism (mean *F*-statistic, 114.5), any bias due to

sample overlap is expected to be minimal[58]. Furthermore, we performed replication analyses

with different exposure and outcome datasets (i.e., the Million Veteran Program and PHOSP-

COVID) without any sample overlap (see **Replication analyses**). Given that these datasets

were smaller—and less well-powered—than the FinnGen and Long COVID Host Genetics

Initiative datasets, we chose to use the latter set of datasets for primary analyses and the

other for replication analyses. Long COVID Host Genetics Initiative participants provided

informed consent for participation in their respective studies, with recruitment and ethics

following study-specific protocols approved by their respective institutional review boards[24].

Study participants were classified as long COVID cases if they had an earlier test-

verified SARS-CoV-2 infection and met long COVID criteria at least three months since

SARS-CoV-2 infection or COVID-19 onset[24]. Aligning with the World Health Organization

guidelines[2], individuals meeting long COVID criteria were defined as those who (1) reported

presence of COVID-19 symptoms that could not be explained by alternative diagnoses; (2)

reported ongoing significant impact on day-to-day activities; or (3) had any diagnosis codes

of long COVID in their electronic health records. Primary analyses used population controls

(i.e., all non-cases) as controls; these individuals were defined as those who were not

identified as having long COVID using the above-mentioned criteria. As such, data from a

total of 3,018 long COVID cases and 994,582 controls from 12 cohorts were included for

primary analyses. All cohort-specific criteria to ascertain long COVID are shown in

Supplementary Table 1. Each cohort study applied their own methods for genotyping,

genotype and sample quality control, imputation, and association analyses (**Supplementary**

Table 1), according to a central analysis plan[24]. Each cohort's GWAS was minimally adjusted

for age, age², sex, age × sex, and the first 10 genetic principal components. Prior to analysis,

all variants with imputation INFO score <0.6 were excluded. Meta-analysis was performed

using a fixed-effects inverse-variance weighted method.

Primary analyses used population controls to increase statistical power and limit

selective inclusion of certain groups within the population, which can introduce collider bias in

MR studies. Population controls were defined as those who were not identified as having

long COVID using the above-mentioned criteria[24]. Replication analyses used a stricter

control definition (i.e., all individuals with a history of SARS-CoV-2 infection but without long

COVID criteria [n_{cases}=2,975; $n_{controls}$=37,935]). Additional details on the long COVID case and

control phenotypes, including diagnostic codes and study-specific criteria, can be found in

the **Supplementary Table 1** and in the main Long COVID Host Genetics Initiative paper[24].

MR analyses

In our primary analysis, we tested the association of genetically predicted venous

thromboembolism risk with long COVID. To do so, we constructed a genetic instrument (i.e.,

a statistical entity comprising one or more genetic variants) for venous thromboembolism in

FinnGen[21]. All genetic instrument construction steps were performed using genetic variants

that were present in both the exposure and outcome GWAS datasets. To ensure that our

genetic instrument was strongly associated with venous thromboembolism, we only included

variants that were associated with venous thromboembolism at genome-wide significance

(P<5×10^{-8}). In addition, to limit bias caused by genetic variants in linkage disequilibrium, we

only retained independent variants in our genetic instrument (R^2<0.001; clumped within

regions of 10 megabases using PLINK [version 1.90][64]). The linkage disequilibrium reference

panel for clumping was derived from the European panel of phase 3 of the 1000 Genomes

Project[63].

Primary analyses used the inverse-variance weighted method with multiplicative

random effects[12] to estimate the effects of genetically predicted venous thromboembolism

risk on long COVID. In downstream analyses, unless specified otherwise, all MR analyses

testing the effects of a genetic instrument with more than three variants used this method.

MR analyses testing the effects of a genetic instrument with two or three variants used the

inverse-variance weighted method with fixed effects[12]; those testing the effects of a genetic

instrument with only one variant used the Wald ratio estimator[65].

All associations were presented as odds ratios (ORs) and 95% confidence intervals

(CIs) per log-odds increase in genetically predicted risk for the exposure condition, along with

their corresponding *P*-values. Two-sided *P*<0.05 indicated statistical significance in primary

analyses (i.e., the inverse-variance weighted estimate of genetically predicted venous

thromboembolism risk on long COVID). All MR analyses were performed using the

TwoSampleMR[66] (version 0.5.7) and *MendelianRandomization*[67] (version 0.7.0) packages in

R (version 4.1.0) were used for analyses.

Genetic instruments for non-venous thromboembolism conditions

495   To contextualize the genetic association of venous thromboembolism with long

COVID (i.e., the association observed in our primary analysis), we also tested the genetic

associations of different other conditions with long COVID. These conditions were chosen

from FinnGen[21] to represent a variety of organ systems and pathophysiological pathways

with well-powered GWAS and included: atrial fibrillation (FinnGen phenotypee code: I9_AF),

asthma (J10_ASTHMA_EXMORE), chronic kidney disease (N14_CHRONKIDNEYDIS),

chronic lower respiratory disease (J10_LOWCHRON), coronary artery disease

(I9_REVASC), dementia (F5_DEMENTIA), diabetes mellitus (T2D), hypertension

(I9_HYPTENS), migraine (MIGRAINE_TRIPTAN), multiple sclerosis (G6_MS), osteoarthritis

(M13_ARTHROSIS), and rheumatoid arthritis (M13_RHEUMA) (**Supplementary Table 15**).

505   Consistent with the primary analysis, all genetic instruments for these conditions were

constructed using GWAS in FinnGen (release 10)[21]. We also tested the associations of

genetically predicted risk for deep vein thrombosis (I9_PHLETHROMBDVTLOW) and

pulmonary embolism (I9_PULMEMB) with long COVID to corroborate the genetic association

of venous thromboembolism with long COVID. All tested phenotypes were defined using

nationwide registries that were linked to the biological samples and harmonized across

diagnosis-, procedure-, and drug-specific codes. The exact codes used to construct the

different phenotypes are listed in **Supplementary Table 15**. The GWAS for these

phenotypes (including genotyping, variant- and sample-wise quality control steps, and

association analyses) were identical to those used for the GWAS of venous

thromboembolism in FinnGen (see **GWAS of venous thromboembolism**). All GWAS were

well-powered with an adequate number of cases per condition ($n>2,000$) and multiple

independent ($R^2<0.001$) variants reaching genome-wide significance ($P<5\times10^{-8}$).

518   All genetic instruments for these conditions were constructed using a *P*-value

threshold of 5×10^{-8} and a linkage disequilibrium R^2 threshold of 0.001. The associations of

these genetic instruments with long COVID were tested using the inverse-variance weighted

method. As the main aim of these analyses was to compare the association observed for

venous thromboembolism (i.e., our primary analysis) with the associations observed for

other, unrelated conditions, statistical significance for these analyses was set at the same

threshold (i.e., two-sided $P<0.05$). In addition, to compare effect estimates, we also

evaluated which genetic instruments had effect estimates within the 95%CI of the venous

thromboembolism-long COVID association.

Sensitivity analyses

We performed sensitivity analyses using different MR approaches and genetic

instrument selection parameters to test the robustness of the association of genetically

predicted venous thromboembolism risk with long COVID. First, to evaluate the possibility of

horizontal pleiotropy (i.e., effects of a genetic instrument on the target outcome beyond its

effects on the exposure of interest) affecting the observed associations, we performed

sensitivity analyses using MR methods that allow for horizontal pleiotropy (including MR-

Egger, the weighted median estimator, and the weighted mode estimator). MR-Egger is an

MR method used to test and correct for horizontal pleiotropy within a genetic instrument at

the cost of lower precision and statistical power[25]. The weighted median estimator is an

alternative MR method with preserved statistical power that allows for horizontal pleiotropy

as long as the majority of the genetic instrument's weight consists of valid instrumental

variables (i.e., genetic variants)[68]. The weighted mode-based estimator allows for horizontal

pleiotropy as long as a plurality of the genetic instrument's weight consists of valid

instrumental variables[69]. Second, to explore the potential effects of residual correlation

between the genetic variants in our genetic instrument, we conducted sensitivity analyses

with genetic instruments constructed using various linkage disequilibrium R^2 thresholds

($R^2 < 0.0001$/$R^2 < 0.001$/$R^2 < 0.01$/$R^2 < 0.1$). Third, we also conducted additional sensitivity

analyses with genetic instruments constructed using various P-value thresholds ($P < 5 \times 10^{-4}$

/$P < 5 \times 10^{-6}$/$P < 5 \times 10^{-8}$/$P < 5 \times 10^{-10}$). These sensitivity analyses were performed in line with

previous reports[59,60,70]. Although there is no consensus on the exact values to use for these

sensitivity analyses, we chose the aforementioned R^2 and P-value thresholds to cover a

reasonable range of parameters at fixed intervals. As MR analyses using multiple genetic

variants from across the genome conventionally use a genome-wide significance threshold

($P < 5 \times 10^{-8}$) and a relatively stringent R^2 threshold ($R^2 < 0.001$), the different R^2 and P-value

parameters for sensitivity analyses were centered around these values. Fourth, we

performed Steiger filtering, which removes genetic variants that explain more variation in the

outcome than in the exposure of interest[26], to evaluate the possibility of reverse causation

affecting the observed results. Fifth, we performed MR analyses in the opposite direction

(i.e., with long COVID as the exposure and venous thromboembolism as the outcome) to

further explore the possibility of reverse causation. Genetic instruments were constructed

using independent ($R^2 < 0.001$) genome-wide significant ($P < 5 \times 10^{-8}$) variants associated with

long COVID in the Long COVID Host Genetics Initiative (**Supplementary Table 16**), using

the same GWAS data as those used for the outcome in primary analyses. These reverse MR

analyses used the inverse-variance weighted method. Because the genetic instrument for

long COVID constructed using a *P*-value threshold of $5.0×10^{-8}$ (mean *F*-statistic, 35.2) only

included two variants, we also performed an additional Mendelian randomization analysis

using a *P*-value threshold of $5.0×10^{-6}$ for instrument construction (mean *F*-statistic, 23.8),

yielding similar results (OR, 0.99 [95%CI, 0.95-1.03] per log-odds increase in long COVID

risk; *P*=0.52).

568   Sensitivity analyses were considered robust if (1) MR-Egger suggested no horizontal

pleiotropy (*P*≥0.05 for the intercept test or *P*<0.05 for the intercept test with *P*<0.05 for the

causal test) and all pleiotropy-robust methods (i.e., the MR-Egger, weighted median, and

weighted mode-based estimators) had a consistent direction of effect; (2) MR estimates were

directionally consistent across all linkage disequilibrium *R*² thresholds for instrument

construction; (3) MR estimates were directionally consistent across all *P*-value thresholds for

instrument construction; (4) MR analyses with Steiger filtering yielded directionally consistent

results (or Steiger filtering did not identify any "reverse causal" variants); and (5) there was

no significant positive association of genetically predicted long COVID risk on venous

thromboembolism.

Replication analyses

580   Replication analyses evaluated whether the observed associations were consistent

across different exposure and outcome datasets. First, we constructed a genetic instrument

for venous thromboembolism using a data source external to the FinnGen[21] dataset (i.e., the

data source used to construct the genetic instrument used for our primary analyses). To this

end, we used data from 8,929 venous thromboembolism cases and 181,337 controls with

European ancestry from the Million Veteran Program[27]. In brief, the Million Veteran Program

recruited individuals aged 19 to >100 years from more than 50 Veterans Affairs Medical

Centers across the USA since 2011[71]. Participants with at least two qualifying diagnostic

codes for deep vein thrombosis or pulmonary embolism in their electronic health record were

classified as cases[27]. Participants were genotyped using a customized Affymetrix Axiom

array (Thermo Fisher Scientific, Santa Clara, CA, USA). Samples with excess

heterozygosity, excess missingness (>2.5%), or discordance between genetically inferred

sex and phenotypic gender were excluded[27]. Samples were imputed to the 1000 Genomes

Project reference panel (phase 3)[63] prior to association analyses adjusted for age, sex, and

five principal components of ancestry[27]. Variants with ancestry-specific Hardy-Weinberg

equilibrium $P<1×10^{-20}$, posterior call probability <0.9, imputation INFO score <0.3, or call rate

<97.5% were excluded, as were variants deviating >10% from their expected allele

frequency based on reference data from the 1000 Genomes Project[63]. Variants with minor

allele frequency <1% were also excluded. The genetic instrument for venous

thromboembolism used for replication was constructed using the same approach as for the

instrument used in our primary analysis (e.g., using a P-value threshold of 5×10^{-8} and a

linkage disequilibrium R^2 threshold of 0.001). The Million Veteran Program received ethical

and study protocol approval by the Veterans Affairs Central Institutional Review Board, and

informed consent was obtained from all participants[27].

Second, we evaluated whether the observed associations persisted when we used

stricter criteria for controls in the Long COVID Host Genetics Initiative GWAS[24]. Unlike our

primary analyses, which used population controls (i.e., all study participants without long

COVID criteria; $n_{controls}$=994,582), these replication analyses only included proven controls

(i.e., all study participants with proven SARS-CoV-2 infection but without long COVID criteria;

$n_{controls}$=37,935).

Third, we tested the association of genetically predicted venous thromboembolism

risk (i.e., the FinnGen instrument) with long COVID in the Post-Hospitalization COVID-19

(PHOSP-COVID) study, a prospective cohort study designed to investigate the medium- and

long-term sequelae of hospitalized COVID-19[28,29]. PHOSP-COVID recruited adults who were

discharged from one of 53 hospitals across the United Kingdom between March 5 and

November 30, 2020, after confirmed or clinician-diagnosed COVID-19[28,29]. A total of 1,097

participants with genetic data and information on long COVID status (n_{cases}=697; $n_{controls}$=400)

were included in this analysis. Long COVID status was primarily based on the question "Do

you feel fully recovered from COVID-19?" during the a visit occurring at least 3 months after

COVID-19 diagnosis. Participants who answered "no" were classified as long COVID cases.

For participants who responded with "not sure" or who had a missing response, long COVID

status was determined by their answers to the EuroQol five-dimension five-level (EQ-5D-5L)

questionnaire. At their first visit (i.e., at least 3 months after COVID-19 diagnosis),

participants completed the EQ-5D-5L questionnaire for their health state at that moment and

retrospectively for their health state before COVID-19 diagnosis. Patients were categorized

as having long COVID if the difference between the current and pre-COVID-19 EQ-5D-5L

scores was ≤-0.1. Samples were genotyped using the Illumina Global Screening Array v3.0

(Illumina Inc., San Diego, CA, USA). Samples with call rates <95%, discordance between

genetically imputed and recorded sex, excess relatedness (third degree of kinship), or

excess heterozygosity (more than 4 standard deviations from the mean) were excluded, as

were duplicates. Genetic variants with Hardy-Weinberg equilibrium $P<1\times10^{-6}$, call rates

<99%, or minor allele frequencies <1% were also excluded. Genetic association analyses for

long COVID were adjusted for age, age², sex, age × sex, and ten principal components of

genetic ancestry. The PHOSP-COVID study was approved by the Leeds West Research

Ethics Committee, and all participants provided informed consent[28,29].

Two-sided $P<0.05$ indicated statistical significance for all replication analyses. In

addition to the three main replication analyses, which used the same methods as those used

for our primary analyses (i.e., genetic instruments with $P<5\times10^{-8}$ and $R^2<0.001$ tested using

the inverse-variance weighted method), we also performed additional analyses using

different P-value ($P<5\times10^{-4}$/$P<5\times10^{-6}$/$P<5\times10^{-8}$/$P<5\times10^{-10}$) and linkage disequilibrium R^2

($R^2<0.0001$/$R^2<0.001$/$R^2<0.01$/$R^2<0.1$) thresholds to construct genetic instruments. Analyses

were considered robust if they were directionally consistent across all genetic instrument

construction parameters.

MR analyses of acute COVID-19 phenotypes

We tested the associations of genetically predicted venous thromboembolism risk

with acute COVID-19 in the COVID-19 Host Genetics Initiative (release 7)[16,32]. The COVID-

19 Host Genetics Initiative includes GWAS data for three acute COVID-19 categories based

on disease severity: SARS-CoV-2 infection (defined as any SARS-CoV-2 infection with or

without symptoms of any severity); hospitalized COVID-19 (defined as hospitalization due to

symptoms associated with SARS-CoV-2 infection); and critical COVID-19 (defined as

hospitalization with respiratory support or death due to symptoms associated with SARS-

CoV-2 infection)[16,32]. All three acute COVID-19 severity categories were tested as separate

outcomes. We only used genetic data from individuals with European ancestry to match the

ancestry of the outcome cohort (i.e., the COVID-19 Host Genetics Initiative[16,32]) with that of

the exposure cohort (i.e., FinnGen[21]). All protocols for studies included in the COVID-19 Host

Genetics Initiative followed local ethics recommendations, and informed consent was

obtained when required[16,32].

The different GWAS meta-analyses—excluding the 23andMe samples—included

122,616 cases and 2,475,240 controls for SARS-CoV-2 infection; 32,519 cases and

2,062,805 controls for hospitalized COVID-19; and 13,769 cases and 1,072,442 controls for

critical COVID-19[16,32]. Each contributing cohort genotyped the samples, performed variant-

wise and sample-wise quality control, and conducted association analyses independently

following a central analysis plan. The different cohort-specific GWAS were filtered for

variants with imputation INFO scores >0.6 and meta-analyzed using the inverse-variance

weighted method. Variants with minor allele frequency <1% were excluded. Consistent with

our primary analysis, we tested the associations of our main genetic instrument for venous

thromboembolism with the three aforementioned acute COVID-19 phenotypes using the

inverse-variance weighted MR method.

We performed multivariable MR[33] analyses to estimate the direct effects of genetic

predisposition to venous thromboembolism on long COVID while adjusting for its impact on

acute COVID-19. Multivariable MR is an extension of conventional (i.e., univariable) MR that

allows for multiple exposures to be modelled at once[33]. We performed three sets of

multivariable MR analyses; in each analysis, venous thromboembolism was modelled as an

exposure together with one acute COVID-19 phenotype (i.e., SARS-CoV-2 infection,

hospitalized COVID-19, and critical COVID-19) while long COVID was considered the

outcome. Genetic instruments were constructed using independent variants ($R^2<0.001$)

reaching genome-wide significance ($P<5\times10^{-8}$) in either the GWAS of venous

thromboembolism or the GWAS of the acute COVID-19 phenotype under study. The

independent genetic effects of each exposure on long COVID were estimated using the

inverse-variance weighted method for multivariable MR studies, implemented using the

mr_mvivw() function from the *MendelianRandomization*[67] package (version 0.7.0) in R.

As the main aim of the conventional (i.e., univariable) MR analyses was to compare

the genetic effects of venous thromboembolism on long COVID (i.e., our primary analysis)

with those on acute COVID-19 phenotypes, statistical significance for these analyses was set

at the same threshold (i.e., two-sided $P<0.05$). We also evaluated which associations had

effect estimates within the 95%CI of the venous thromboembolism-long COVID association.

Results from multivariable MR analyses were considered robust if venous thromboembolism

had a significant independent association (i.e., two-sided $P<0.05$) with long COVID across

models.

MR analyses of long COVID-resembling conditions

We tested the genetic associations of venous thromboembolism with a range of "long

COVID-resembling" phenotypes. These long COVID-resembling phenotypes included

various conditions sharing similar symptoms—yet likely distinct mechanisms—to those

observed in long COVID[3]. The phenotypes were chosen based on prior evidence suggesting

overlapping symptoms with long COVID[3] and were tested if they had GWAS data available in

FinnGen (release 10)[21].

All phenotypes were defined using nationwide registries that were linked to the

biological samples and harmonized across diagnosis-, procedure-, and drug-specific codes[21].

The tested phenotypes included: chronic lower respiratory disease (FinnGen phenotype

code: J10_LOWCHRON), functional dyspepsia (K11_FUNCDYSP), irritable bowel syndrome

(K11_IBS), endometriosis (N14_ENDOMETRIOSIS), fibromyalgia (M13_FIBROMYALGIA),

anxiety disorders (F5_ALLANXIOUS), memory loss (MEMLOSS), sleep disorders (SLEEP),

mood disturbances (F5_MOOD), depression (F5_DEPRESSIO), and postviral fatigue

(G6_POSTVIRFAT). Additional information on the tested long COVID-resembling

phenotypes can be found in **Supplementary Table 10**. The GWAS for these phenotypes

(including genotyping, variant-wise and sample-wise quality control, and association

analyses) were identical to those used for the GWAS of venous thromboembolism in

FinnGen (see **GWAS of venous thromboembolism**).

To avoid sample overlap, we used the genetic instrument from the Million Veteran

Program[27] (see **Replication analyses**) to test the effects of genetically predicted venous

thromboembolism risk on long COVID-resembling phenotypes. These analyses used the

inverse-variance weighted MR method. Statistical significance was set at the same threshold

as in our primary analysis (i.e., two-sided $P<0.05$). We also evaluated which associations

had effect estimates within the 95%CI of the venous thromboembolism-long COVID

association.

Gene-specific genetic instruments for venous thromboembolism

We constructed "gene-specific genetic instruments" by identifying variants associated

with venous thromboembolism within or near different coagulation-related genes.

Coagulation-related genes were defined as those included in the "*Complement and*

coagulation cascades" gene set from the KEGG database[72] (**Supplementary Table 17**).

Each gene-specific genetic instrument was constructed using genetic variants associated

with venous thromboembolism within a *cis*-region 200 kilobases of the indicated coagulation-

related gene in FinnGen[21]. Because a *cis*-region only comprises a small region within the

genome, and because sensitivity analyses showed that genetic instruments for venous

thromboembolism were still associated with long COVID when using more lenient *P*-value

thresholds (i.e., $P<5\times10^{-4}$ or $P<5\times10^{-6}$) (**Supplementary Table 4**), we also included genetic

variants associated with venous thromboembolism at a sub-genome-wide significance

threshold to increase the number of genes with valid gene-specific genetic instruments. More

specifically, if a gene's *cis*-region included variants associated with venous thromboembolism

at $P<5\times10^{-8}$, we used this threshold to construct the genetic instrument; if not, we used

$P<5\times10^{-6}$ or $P<5\times10^{-4}$, depending on the presence of *cis*-variants surpassing those *P*-value

thresholds. Independent genetic variants were selected using a linkage disequilibrium R^2

threshold of 0.001.

MR analyses testing the effects of gene-specific genetic instruments with more than

three variants used the inverse-variance weighted method with multiplicative random

effects[12]; those instruments with two or three variants used the inverse-variance weighted

method with fixed effects[12]; those testing the effects of a genetic instrument with only one

variant used the Wald ratio estimator[65]. Considering that genetically predicted venous

thromboembolism was strongly associated with long COVID and that this association was

robust across multiple genetic instrument *P*-value thresholds, and that gene-specific genetic

instruments inherently include fewer genetic variants than genetic instruments constructed

using the entire genome, we did not formally correct these analyses for multiple

comparisons. Instead, we prioritized associations based on effect size and used two-sided

$P<0.05$ to indicate statistical significance.

MR analyses using genetically predicted protein levels

Because *F2R*'s gene-specific genetic instrument had largest effect on long COVID

across all gene-specific genetic instruments, downstream analyses focused on this gene and

its gene products (i.e., *F2R* RNA and the PAR-1 protein). First, we tested the effects of

genetically predicted circulating levels of PAR-1 on long COVID risk using a genetic

instrument from the UK Biobank Pharma Proteomics Project (UKB-PPP)[37].

The UKB-PPP is a collaboration between 13 biopharmaceutical companies that

funded the measurement of 2,923 circulating proteins in participants from the UK Biobank, a

population-based cohort of individuals aged 40–69 years recruited between 2006 and 2010

across the United Kingdom[37]. A total of 54,306 UK Biobank participants who donated blood

samples at baseline or follow-up study visits were selected for proteomic profiling and were

included in the UKB-PPP cohort. Blood samples were analyzed using proximity extension

assay technology (Olink Explore 3072 platform [Olink Proteomics, Inc; Waltham, MA, USA]).

Sun *et al.*[37] performed a GWAS of circulating proteins measured using Olink technology in a

discovery cohort of 35,571 UKB-PPP participants of European ancestry. Genotyping,

imputation, and quality control of the UK Biobank samples were performed as described

previously[37,73]. In addition to checking for sex mismatch, sex chromosome aneuploidy, and

heterozygosity, imputed genetic variants were filtered for imputation INFO score >0.7.

Individual protein levels were inverse-rank normalized prior to association analyses, which

were adjusted for age, age², sex, age × sex, age² × sex, batch, assessment center, genetic

array, time between blood sampling and measurement, and 20 principal components of

genetic ancestry. The UK Biobank received approval from the North West Multi-center

Research Ethics Committee, and all participants provided informed consent.

We constructed a genetic instrument for circulating PAR-1 levels using independent

(linkage disequilibrium $R^2<0.001$) genetic variants within 200 kilobases of *F2R* strongly

associated with circulating PAR-1 levels ($P<5\times10^{-8}$). Such variants are also called *cis*-protein

quantitative trait loci (*cis*-pQTLs) and have favorable effects on the assumptions of MR

analyses[59–61]. The aforementioned thresholds resulted in only one variant (i.e., rs168753) to

be included in the genetic instrument for circulating PAR-1; therefore, the association of this

instrument with long COVID was tested using the Wald ratio estimator[65]. Sensitivity analyses

used different *P*-value ($P<5\times10^{-4}$/$P<5\times10^{-6}$/$P<5\times10^{-8}$/$P<5\times10^{-10}$) and linkage disequilibrium

R^2 ($R^2<0.0001$/$R^2<0.001$/$R^2<0.01$/$R^2<0.1$) thresholds to construct genetic instruments. In

addition, because *F2R*'s *cis*-region partially overlaps with those of neighboring genes (i.e.,

F2RL1 and *F2RL2*)[42], we also performed analyses with genetic instruments selectively

including intragenic variants to limit pleiotropic effects of extragenic variants potentially

influencing other genes' expression patterns. Two-sided P<0.05 indicated statistical

significance for this corroboratory analysis, and directional consistency was used as the

criterium for robustness in sensitivity analyses.

Colocalization analyses

To complement the MR, we performed colocalization analyses to test for shared

causal variants between circulating PAR-1 levels and long COVID. Analyses were carried out

with variants within 200 kilobases of *F2R*, using the *coloc.abf()* function from the R package

coloc (version 5.2.2)[74]. The output of these analyses was expressed as test statistics that

estimate the posterior probabilities of five hypotheses: (1) no causal variant for either trait; (2)

a causal variant for the first but not the second trait; (3) a causal variant for the second but

not the first trait; (4) distinct causal variants underlying both traits; and (5) one shared causal

variant underlying both traits[38,39]. A high posterior probability for the fifth hypothesis (>80%)

indicates strong evidence of colocalization, suggesting a high likelihood of a shared causal

variant driving both traits. A high posterior probability for the fourth hypothesis (>80%)

indicates strong evidence of distinct causal variants, suggesting a high likelihood of

confounding by linkage disequilibrium (or horizontal pleiotropy) in the corresponding MR

analysis. If there is no high probability for the fourth or fifth hypothesis (<80% for each) but

the MR association is statistically significant and not a false positive finding, this suggests

that the colocalization analysis is likely not adequately powered to distinguish between

whether the MR association is due to a shared causal variant or a variant in linkage

disequilibrium (i.e., horizontal pleiotropy)[38,39]. In the latter situation, we also estimated the

probability for shared causal variants, conditional on there being at least one variant

associated with each trait under study, calculated as: $(PP_{H4}) / (PP_{H3} + PP_{H4})$[38,39], where PP_{H3}

indicates the posterior probability of the fourth hypothesis (i.e., distinct causal variants for

both traits) and PP_{H4} that of the fifth hypothesis (i.e., one shared causal variant for both

traits).

MR analyses using genetically predicted transcript levels

In addition to analyses focused on the PAR-1 protein, we also performed MR

analyses testing the effects of genetically predicted *F2R* RNA levels on long COVID risk. To

this end, we selected five tissues or cell types with (1) high *F2R* expression (queried using

the Genome-Tissue Expression [GTEx] database[40]); (2) biological relevance to long COVID;

and (3) at least one valid variant near *F2R* associated with *F2R* expression (i.e., *cis-*

expression quantitative trait locus [*cis*-eQTL]). The highest median *F2R* RNA levels in GTEx

were observed for the following tissues: aorta, coronary artery, tibial artery, skin (not sun-

exposed), skin (sun-exposed), minor salivary gland, lung, and spleen[40]. Because we

assumed that the skin (both sun-exposed and not sun-exposed) and minor salivary gland

had less direct biological relevance to long COVID in comparison with the other tissues, we

did not carry these tissues forward for further analyses. Given high *F2R* expression in

platelets[34]—a cell type not included in the GTEx database[40]—we added platelets to our list of

tissues and cell types of interest. Genetic instruments for platelets were obtained from the

GeneSTAR Research Study[41], while those for the other tissues (i.e., aorta, coronary artery,

tibial artery, lung, and spleen) were queried in GTEx (v8)[40].

We used the GTEx (v8) dataset[40] to construct genetic instruments for *F2R* expression

in the aforementioned tissues. In brief, the GTEx project has created a resource of gene

expression levels from normal (i.e., non-diseased) tissues from deceased human donors.

GTEx included 838 donors with available RNA sequences and genotypes, collectively

providing 17,382 samples from 52 tissues and two cell lines[75]. A total of 715 (85.3%)

participants had European ancestry and were used to construct genetic instruments in the

present analysis. RNA sequencing libraries were generated using the Illumina TruSeq

protocol (Illumina Inc., San Diego, CA, USA); gene-level expression levels were quantified as

reads per kilobase of transcript per million mapped reads[40]. Participants were genotyped

using the Illumina Human Omni 2.5M and 5M Beadchips and imputed using the multi-ethnic

reference panel from the 1000 Genomes Project (phase 3)[63]. Samples with fewer than 10

million mapped reads or outlier expression measurements were removed[40]. Variants with call

rates <95%, minor allele frequencies <1%, Hardy-Weinberg equilibrium $P<1\times10^{-6}$, or

imputation INFO score <0.4 were excluded. Genetic association analyses of transcript levels

were adjusted for PEER factors, sex, genotyping platform, and three principal components of

genetic ancestry. The protocol for the GTEx project was reviewed by Chesapeake Research

Review Inc., Roswell Park Cancer Institute's Office of Research Subject Protection, and the

institutional review board of the University of Pennsylvania. For all donors, consent was

obtained via next-of-kin consent[40].

We used data from the GeneSTAR Research Study[41] to construct genetic instruments

for *F2R* expression in platelets. In brief, the GeneSTAR Research Study was designed to

examine gene-environment determinants of platelet reactivity in response to low-dose aspirin

therapy[76]. GeneSTAR included siblings identified from probands with coronary artery disease

at age <60 years, as well as the spouses and adult offspring of the siblings. A total of 307

apparently healthy individuals were included for platelet eQTL analyses, of whom 180

(58.6%) had European ancestry and were used to construct genetic instruments in the

present analysis[41]. Once platelets were isolated from whole blood samples, RNA sequencing

libraries were generated using the Illumina TruSeq protocol (Illumina Inc., San Diego, CA,

USA). Transcript abundances were quantified as fragments per kilobase of transcript per

million sequenced reads. Participants underwent whole genome sequencing as part of the

NHLBI's Trans-Omics for Precision Medicine (TOPMed) program[77]. Low-expression outliers

and samples with poor quality or unexpected familial relationships were excluded[41]. Variants

were filtered for autosomal single-nucleotide polymorphisms with at least two samples per

genotype and a call rate of >80%. Genetic association analyses of transcript levels were

adjusted for sex, age, RNA sequencing batch, 15 principal components of the filtered and

log-transformed gene expression matrix, and three principal components of genetic ancestry.

The GeneSTAR Research Study was approved by the institutional review board of the Johns

Hopkins Medical Institutions, and all participants provided informed consent[76].

We constructed a genetic instrument for tissue-specific *F2R* expression using

independent (linkage disequilibrium R^2<0.001) genetic variants within 200 kilobases of *F2R*

associated with *F2R* RNA levels from the GeneSTAR Research Study[41] (for platelets) or

GTEx project[40] (for the other tissues). Similar to our analyses using gene-specific genetic

instruments for venous thromboembolism (see **Gene-specific genetic instruments for**

venous thromboembolism), each tissue-specific genetic instrument's *P*-value threshold

depended on the association strength of the different variants near *F2R*. If *F2R*'s *cis*-region

included variants associated with tissue-specific *F2R* expression at $P<5\times10^{-8}$, we used this

threshold to construct the genetic instrument; if not, we used $P<5\times10^{-6}$ or $P<5\times10^{-4}$,

depending on the presence of *cis*-variants surpassing those *P*-value thresholds. This

approach resulted in one genetic variant (i.e., the strongest *cis*-eQTL) for each tissue-specific

genetic instrument; therefore, the association of these instruments with long COVID was

tested using the Wald ratio estimator[65]. The only tissue without adequate *cis*-eQTLs for *F2R*

was coronary arterial tissue, which was excluded from the analyses. Similar to the analyses

using *cis*-pQTLs for PAR-1, we also performed analyses that only included intragenic

variants. The genetic instruments for *F2R* expression in tibial arterial tissue included

rs253072 (using variants within 200 kilobases) and rs250730 (using only intragenic variants),

respectively; those for splenic tissue included rs253072 and rs250750; those for pulmonary

tissue included rs2227750 and rs12735; and those for platelets included rs253072 and

rs32934. We could only construct an adequate instrument for aortic tissue using variants

within 200 kilobases (i.e., rs9293695) and not using intragenic variants alone. Two-sided

$P<0.05$ indicated statistical significance for these confirmatory analyses.

Data availability

This study used summary statistics from genome-wide association studies performed

in FinnGen (release 10; available from https://r10.finngen.fi/), the Long COVID Host Genetics

Initiative (available from https://my.locuszoom.org/gwas/793752/;

https://doi.org/10.1101/2023.06.29.23292056), the Million Veteran Program (available from

dbGAP, accession code no. phs001672.v2.p1), the PHOSP-COVID study

(https://www.phosp.org/), the COVID-19 Host Genetics Initiative (release 7; available from

https://www.covid19hg.org/results/r7/), the UK Biobank (available from http://ukb-

ppp.gwas.eu/), the Genotype-Tissue Expression project (v10; available from

https://gtexportal.org/home/downloads/adult-gtex/qtl), and the GeneSTAR Research Study

(available from http://www.biostat.jhsph.edu/~kkammers/GeneSTAR/).

Code availability

901   Code used for the main analyses of this study can be accessed at [link to be

902 published upon publication].

Acknowledgements

The authors acknowledge the participants and investigators of the FinnGen study, the

Long COVID Host Genetics Initiative, the Million Veteran Program, the Post-Hospitalization

COVID-19 study, the UK Biobank, the Genotype-Tissue Expression project, and the

GeneSTAR Research Study. A.S. is supported by Life Sciences Research Partners (LSRP).

T.N. is supported by a research fellowship of the Japan Society for the Promotion of Science

for Young Scientists (22J30004) and by a Grant-in-Aid for Scientific Research (B)

(23H02917). M.K.G. is funded by the German Research Foundation in the form of an Emmy

Noether grant (GZ: GE 3461/2-1, ID 512461526). B.G.G. is supported by the Wellcome Trust

(221680/Z/20/Z). L.V.W. holds a GlaxoSmithKline Asthma + Lung UK Chair in Respiratory

Research (C17-1). B.R. is supported by a Wellcome Career Development Award fellowship

(302210/Z/23/Z). H.Z. is supported by the Swedish Research Council (2021-03050), the Knut

and Alice Wallenberg Foundation (2023.0141), Groschinskys Minnesfond, the Max Planck

Society as well as Cornell's, Philip-Sörensen's, and Hedlund's foundations. H.M.O. is

supported by the Academy of Finland (#1350181) and the NIH (R01AI170850). K.F. is

supported by the National Institute of Health (R01Hl161365), KU Leuven (C14/23/121),

research foundation Flanders (FWO; G072921N). T.V. is supported by a grant from FWO

(1843423N). PHOSP-COVID is jointly funded by a grant from the MRC-UK Research and

Innovation and the Department of Health and Social Care through the National Institute for

Health Research (NIHR) rapid response panel to tackle COVID-19 (grant references:

MR/V027859/1 and COV0319). The views expressed in the publication are those of the

author(s) and not necessarily those of the National Health Service (NHS), the NIHR or the
Department of Health and Social Care. This study would not be possible without all the
participants who have given their time and support. We thank all the participants and their
families. We thank the many research administrators, health-care and social-care
professionals who contributed to setting up and delivering the study at all of the 65 NHS
trusts/Health boards and 25 research institutions across the UK, as well as all the supporting
staff at the NIHR Clinical Research Network, Health Research Authority, Research Ethics
Committee, Department of Health and Social Care, Public Health Scotland, and Public
Health England, and support from the ISARIC Coronavirus Clinical Characterisation
Consortium. We thank Kate Holmes at the NIHR Office for Clinical Research Infrastructure
(NOCRI) for her support in coordinating the charities group. The PHOSP-COVID industry
framework was formed to provide advice and support in commercial discussions, and we
thank the Association of the British Pharmaceutical Industry as well NOCRI for coordinating
this. We are very grateful to all the charities that have provided insight to the study: Action
Pulmonary Fibrosis, Alzheimer's Research UK, Asthma + Lung UK, British Heart Foundation,
Diabetes UK, Cystic Fibrosis Trust, Kidney Research UK, MQ Mental Health, Muscular
Dystrophy UK, Stroke Association Blood Cancer UK, McPin Foundations, and Versus
Arthritis. We thank the NIHR Leicester Biomedical Research Centre patient and public
involvement group and Long Covid Support.
944

Competing interests

B.S. is a full-time employee of Bristol Myers Squibb. M.K.G. has consulted for
Tourmaline Bio and serves in the Editorial Board of Neurology, both not relevant to this work.
L.V.W. reports research funding from GlaxoSmithKline, Genentech and Orion Pharma, and
consultancy for Galapagos and GlaxoSmithKline, outside of the submitted work. C.E.B. has
received grants and consultancy fees from 4D Pharma, AstraZeneca, Chiesi, Genentech,
GSK, Mologic, Novartis, Regeneron Pharmaceuticals, Roche and Sanofi. M.C.H. reports
research funding from Genentech, site principal investigator work for Novartis, consulting
fees from Comanche Biopharma, and advisory board service for Miga Health, all unrelated to
the present work. K.F. received research grants from Novo Nordisk and Swedish Orphan
Biovitrum AB (Sobi). T.V. has participated in advisory boards and/or acted as a speaker on
behalf of Bayer, BMS/Pfizer, Boehringer Ingelheim, Daiichi Sankyo, Leo Pharma, Sanofi
Aventis. P.V. has received research funding from Bayer, BMS, Pfizer, and Leo Pharma, and
honoraria from Bayer, Pfizer, BMS, Daiichi-Sankyo, Sanofi-Aventis, Leo Pharma, Anthos
Therapeutics, and Astra-Zeneca.
960

Inclusion and ethics

962   Inclusion and ethics standards have been reviewed where applicable.

964   **Corresponding authors**

965   Correspondence and requests for materials should be addressed to A.S.

References

1. Vos T., Hanson SW., Abbafati C., Aerts JG., Al-Aly Z., Ashbaugh C., et al. Estimated Global Proportions of Individuals With Persistent Fatigue, Cognitive, and Respiratory Symptom Clusters Following Symptomatic COVID-19 in 2020 and 2021. *JAMA* 2022;328:1604–15. https://doi.org/10.1001/JAMA.2022.18931.

2. Soriano JB., Murthy S., Marshall JC., Relan P., Diaz J V. A clinical case definition of post-COVID-19 condition by a Delphi consensus. *Lancet Infect Dis* 2022;22:e102–7. https://doi.org/10.1016/S1473-3099(21)00703-9.

3. Davis HE., McCorkell L., Vogel JM., Topol EJ. Long COVID: major findings, mechanisms and recommendations. *Nat Rev Microbiol* 2023;21:133–46. https://doi.org/10.1038/s41579-022-00846-2.

4. Seeble J., Waterboer T., Hippchen T., Simon J., Kirchner M., Lim A., et al. Persistent Symptoms in Adult Patients 1 Year After Coronavirus Disease 2019 (COVID-19): A Prospective Cohort Study. *Clin Infect Dis* 2022;74:1191–8. https://doi.org/10.1093/CID/CIAB611.

5. Tran VT., Porcher R., Pane I., Ravaud P. Course of post COVID-19 disease symptoms over time in the ComPaRe long COVID prospective e-cohort. *Nat Commun* 2022;13:1–6. https://doi.org/10.1038/s41467-022-29513-z.

6. Taquet M., Skorniewska Z., Hampshire A., Chalmers JD., Ho LP., Horsley A., et al. Acute blood biomarker profiles predict cognitive deficits 6 and 12 months after COVID-19 hospitalization. *Nat Med* 2023;29:2498–508. https://doi.org/10.1038/s41591-023-02525-y.

7. Cervia-Hasler C., Brüningk SC., Hoch T., Fan B., Muzio G., Thompson RC., et al. Persistent complement dysregulation with signs of thromboinflammation in active Long Covid. *Science (1979)* 2024;383:1–18. https://doi.org/10.1126/SCIENCE.ADG7942.

8. Davies M. Long covid patients travel abroad for expensive and experimental "blood washing." *BMJ* 2022;378. https://doi.org/10.1136/BMJ.O1671.

9. Uffelmann E., Huang QQ., Munung NS., de Vries J., Okada Y., Martin AR., et al. Genome-wide association studies. *Nat Rev Methods Primers* 2021;1:1–21. https://doi.org/10.1038/s43586-021-00056-9.

10. Davies NM., Holmes M V., Davey Smith G. Reading Mendelian randomisation studies: a guide, glossary, and checklist for clinicians. *BMJ* 2018;362:601. https://doi.org/10.1136/BMJ.K601.

11. Sanderson E., Glymour MM., Holmes M V., Kang H., Morrison J., Munafò MR., et al. Mendelian randomization. *Nat Rev Methods Primers* 2022;2:1–21. https://doi.org/10.1038/s43586-021-00092-5.

12.   Burgess S., Butterworth A., Thompson SG. Mendelian Randomization Analysis With
Multiple Genetic Variants Using Summarized Data. *Genet Epidemiol* 2013;37:658–65.
https://doi.org/10.1002/GEPI.21758.

13.   Ardissino M., Morley AP., Slob EAW., Schuermans A., Rayes B., Raisi-Estabragh Z.,
et al. Birth weight influences cardiac structure, function, and disease risk: evidence of
a causal association. *Eur Heart J* 2023.
https://doi.org/10.1093/EURHEARTJ/EHAD631.

14.   Gaziano L., Giambartolomei C., Pereira AC., Gaulton A., Posner DC., Swanson SA.,
et al. Actionable druggable genome-wide Mendelian randomization identifies
repurposing opportunities for COVID-19. *Nat Med* 2021;27:668–76.
https://doi.org/10.1038/s41591-021-01310-z.

15.   Bovijn J., Lindgren CM., Holmes M V. Genetic variants mimicking therapeutic inhibition
of IL-6 receptor signaling and risk of COVID-19. *Lancet Rheumatol* 2020;2:e658–9.
https://doi.org/10.1016/S2665-9913(20)30345-3.

16.   Niemi MEK., Karjalainen J., Liao RG., Neale BM., Daly M., Ganna A., et al. Mapping
the human genetic architecture of COVID-19. *Nature* 2021;600:472–7.
https://doi.org/10.1038/s41586-021-03767-x.

17.   Interleukin-6 Receptor Antagonists in Critically Ill Patients with Covid-19. *N Engl J Med*
2021;384:1491–502. https://doi.org/10.1056/NEJMOA2100433.

18.   Group TWREA for C-19 T (REACT) W., Domingo P., Mur I., Mateo GM., Gutierrez M
del M., Pomar V., et al. Association Between Administration of IL-6 Antagonists and
Mortality Among Patients Hospitalized for COVID-19: A Meta-analysis. *JAMA*
2021;326:499–518. https://doi.org/10.1001/JAMA.2021.11330.

19.   Trajanoska K., Bhérer C., Taliun D., Zhou S., Richards JB., Mooser V. From target
discovery to clinical drug development with human genetics. *Nature* 2023;620:737–45.
https://doi.org/10.1038/s41586-023-06388-8.

20.   Pierce BL., Burgess S. Efficient Design for Mendelian Randomization Studies:
Subsample and 2-Sample Instrumental Variable Estimators. *Am J Epidemiol*
2013;178:1177–84. https://doi.org/10.1093/AJE/KWT084.

21.   Kurki MI., Karjalainen J., Palta P., Sipilä TP., Kristiansson K., Donner KM., et al.
FinnGen provides genetic insights from a well-phenotyped isolated population. *Nature*
2023;613:508–18. https://doi.org/10.1038/s41586-022-05473-8.

22.   Leinonen JT., Pirinen M., Tukiainen T. Disentangling the link between maternal
influences on birth weight and disease risk in 36,211 genotyped mother–child pairs.
Commun Biol 2024;7:1–9. https://doi.org/10.1038/s42003-024-05872-9.

23.   Access results | FinnGen. Available at: https://www.finngen.fi/en/access_results.
Accessed October 24, 2024.

24.   Lammi V., Nakanishi T., Jones SE., Andrews SJ., Karjalainen J., Cortés B., et al.
Genome-wide Association Study of Long COVID. MedRxiv.
https://doi.org/10.1101/2023.06.29.23292056.

25.   Burgess S., Thompson SG. Interpreting findings from Mendelian randomization using
the MR-Egger method. *Eur J Epidemiol* 2017;32:377. https://doi.org/10.1007/S10654-
017-0255-X.

26.   Hemani G., Tilling K., Davey Smith G. Orienting the causal relationship between
imprecisely measured traits using GWAS summary data. *PLoS Genet*
2017;13:e1007081. https://doi.org/10.1371/JOURNAL.PGEN.1007081.

27.   Klarin D., Busenkell E., Judy R., Lynch J., Levin M., Haessler J., et al. Genome-wide
association analysis of venous thromboembolism identifies new risk loci and genetic
overlap with arterial vascular disease. *Nat Genet* 2019;51:1574–9.
https://doi.org/10.1038/s41588-019-0519-3.

28.   Evans RA., McAuley H., Harrison EM., Shikotra A., Singapuri A., Sereno M., et al.
Physical, cognitive, and mental health impacts of COVID-19 after hospitalisation
(PHOSP-COVID): a UK multicentre, prospective cohort study. *Lancet Respir Med*
2021;9:1275–87. https://doi.org/10.1016/S2213-2600(21)00383-0.

29.   Elneima O., McAuley HJC., Leavy OC., Chalmers JD., Horsley A., Ho L-P., et al.
Cohort Profile: Post-Hospitalisation COVID-19 (PHOSP-COVID) study. *Int J Epidemiol*
2024;53:165. https://doi.org/10.1093/IJE/DYAD165.

30.   Bilaloglu S., Aphinyanaphongs Y., Jones S., Iturrate E., Hochman J., Berger JS.
Thrombosis in Hospitalized Patients With COVID-19 in a New York City Health
System. *JAMA* 2020;324:799–801. https://doi.org/10.1001/JAMA.2020.13372.

31.   Katsoularis I., Fonseca-Rodríguez O., Farrington P., Jerndal H., Lundevaller EH.,
Sund M., et al. Risks of deep vein thrombosis, pulmonary embolism, and bleeding
after covid-19: nationwide self-controlled cases series and matched cohort study. *BMJ*
2022;377:e069590. https://doi.org/10.1136/BMJ-2021-069590.

32.   Kanai M., Andrews SJ., Cordioli M., Stevens C., Neale BM., Daly M., et al. A second
update on mapping the human genetic architecture of COVID-19. *Nature*
2023;621:E7–26. https://doi.org/10.1038/s41586-023-06355-3.

33.   Burgess S., Thompson SG. Multivariable Mendelian Randomization: The Use of
Pleiotropic Genetic Variants to Estimate Causal Effects. *Am J Epidemiol*
2015;181:251–60. https://doi.org/10.1093/AJE/KWU283.

34.   Angiolillo DJ., Capodanno D., Goto S. Platelet thrombin receptor antagonism and
atherothrombosis. *Eur Heart J* 2010;31:17–28.
https://doi.org/10.1093/EURHEARTJ/EHP504.

35. Smith SMG., Judge HM., Peters G., Armstrong M., Dupont A., Gaussem P., et al.
PAR-1 genotype influences platelet aggregation and procoagulant responses in
patients with coronary artery disease prior to and during clopidogrel therapy. *Platelets*
2005;16:340–5. https://doi.org/10.1080/00207230500120294.

36. Dupont A., Fontana P., Bachelot-Loza C., Reny JL., Bièche I., Desvard F., et al. An
intronic polymorphism in the PAR-1 gene is associated with platelet receptor density
and the response to SFLLRN. *Blood* 2003;101:1833–40.
https://doi.org/10.1182/BLOOD-2002-07-2149.

37. Sun BB., Chiou J., Traylor M., Benner C., Hsu YH., Richardson TG., et al. Plasma
proteomic associations with genetics and health in the UK Biobank. *Nature*
2023;622:329–38. https://doi.org/10.1038/s41586-023-06592-6.

38. Woolf B., Rajasundaram S., Cronjé HT., Yarmolinsky J., Burgess S., Gill D. A drug
target for erectile dysfunction to help improve fertility, sexual activity, and wellbeing:
mendelian randomisation study. *BMJ* 2023;383. https://doi.org/10.1136/BMJ-2023-
076197.

39. Zuber V., Grinberg NF., Gill D., Manipur I., Slob EAW., Patel A., et al. Combining
evidence from Mendelian randomization and colocalization: Review and comparison of
approaches. *Am J Hum Genet* 2022;109:767–82.
https://doi.org/10.1016/J.AJHG.2022.04.001/ATTACHMENT/FC339381-C46B-413F-
A624-9A1B10820BED/MMC1.PDF.

40. Aguet F., Brown AA., Castel SE., Davis JR., He Y., Jo B., et al. Genetic effects on
gene expression across human tissues. *Nature* 2017;550:204–13.
https://doi.org/10.1038/nature24277.

41. Kammers K., Taub MA., Rodriguez B., Yanek LR., Ruczinski I., Martin J., et al.
Transcriptional profile of platelets and iPSC-derived megakaryocytes from whole-
genome and RNA sequencing. *Blood* 2021;137:959–68.
https://doi.org/10.1182/BLOOD.2020006115.

42. Kahn ML., Hammes SR., Botka C., Coughlin SR. Gene and Locus Structure and
Chromosomal Localization of the Protease-activated Receptor Gene Family. *J Biol*
Chem 1998;273:23290–6. https://doi.org/10.1074/JBC.273.36.23290.

43. Pretorius E., Vlok M., Venter C., Bezuidenhout JA., Laubscher GJ., Steenkamp J., et
al. Persistent clotting protein pathology in Long COVID/Post-Acute Sequelae of
COVID-19 (PASC) is accompanied by increased levels of antiplasmin. *Cardiovasc*
Diabetol 2021;20:1–18. https://doi.org/10.1186/S12933-021-01359-7/FIGURES/7.

44. Klok FA., Kruip MJHA., van der Meer NJM., Arbous MS., Gommers DAMPJ., Kant
KM., et al. Incidence of thrombotic complications in critically ill ICU patients with

COVID-19. *Thromb Res* 2020;191:145–7.

https://doi.org/10.1016/J.THROMRES.2020.04.013.

45. Conway EM., Mackman N., Warren RQ., Wolberg AS., Mosnier LO., Campbell RA., et al. Understanding COVID-19-associated coagulopathy. *Nat Rev Immunol* 2022;22:639–49. https://doi.org/10.1038/s41577-022-00762-9.

46. Kanth Manne B., Denorme F., Middleton EA., Portier I., Rowley JW., Stubben C., et al. Platelet gene expression and function in patients with COVID-19. *Blood* 2020;136:1317–29. https://doi.org/10.1182/BLOOD.2020007214.

47. Bryce C., Grimes Z., Pujadas E., Ahuja S., Beasley MB., Albrecht R., et al. Pathophysiology of SARS-CoV-2: the Mount Sinai COVID-19 autopsy experience. *Mod Pathol* 2021;34:1456–67. https://doi.org/10.1038/s41379-021-00793-y.

48. Ackermann M., Verleden SE., Kuehnel M., Haverich A., Welte T., Laenger F., et al. Pulmonary Vascular Endothelialitis, Thrombosis, and Angiogenesis in Covid-19. *N Engl J Med* 2020;383:120–8. https://doi.org/10.1056/NEJMOA2015432.

49. Oxley TJ., Mocco J., Majidi S., Kellner CP., Shoirah H., Singh IP., et al. Large-Vessel Stroke as a Presenting Feature of Covid-19 in the Young. *N Engl J Med* 2020;382. https://doi.org/10.1056/NEJMC2009787.

50. Stevens H., Canovas R., Tran H., Peter K., McFadyen JD. Inherited Thrombophilias Are Associated With a Higher Risk of COVID-19–Associated Venous Thromboembolism: A Prospective Population-Based Cohort Study. *Circulation* 2022;145:940–2. https://doi.org/10.1161/CIRCULATIONAHA.121.057394.

51. Ryu JK., Yan Z., Montano M., Sozmen EG., Dixit K., Suryawanshi RK., et al. Fibrin drives thromboinflammation and neuropathology in COVID-19. *Nature* 2024;633:905–13. https://doi.org/10.1038/s41586-024-07873-4.

52. Swank Z., Senussi Y., Manickas-Hill Z., Yu XG., Li JZ., Alter G., et al. Persistent Circulating Severe Acute Respiratory Syndrome Coronavirus 2 Spike Is Associated With Post-acute Coronavirus Disease 2019 Sequelae. *Clinical Infectious Diseases* 2023;76:e487–90. https://doi.org/10.1093/CID/CIAC722.

53. Coughlin SR. Thrombin signalling and protease-activated receptors. *Nature* 2000;407:258–64. https://doi.org/10.1038/35025229.

54. Morrow DA., Braunwald E., Bonaca MP., Ameriso SF., Dalby AJ., Fish MP., et al. Vorapaxar in the Secondary Prevention of Atherothrombotic Events. *N Engl J Med* 2012;366:1404–13. https://doi.org/10.1056/NEJMOA1200933.

55. Tricoci P., Huang Z., Held C., Moliterno DJ., Armstrong PW., Van de Werf F., et al. Thrombin-Receptor Antagonist Vorapaxar in Acute Coronary Syndromes. *N Engl J Med* 2012;366:20–33.

https://doi.org/10.1056/NEJMOA1109719/SUPPL_FILE/NEJMOA1109719_DISCLOS

URES.PDF.

56.   Niessen F., Schaffner F., Furlan-Freguia C., Pawlinski R., Bhattacharjee G., Chun J.,

et al. Dendritic cell PAR1–S1P3 signalling couples coagulation and inflammation.

Nature 2008;452:654–8. https://doi.org/10.1038/nature06663.

57.   Subramaniam S., Ogoti Y., Hernandez I., Zogg M., Botros F., Burns R., et al. A

thrombin-PAR1/2 feedback loop amplifies thromboinflammatory endothelial responses

to the viral RNA analogue poly(I:C). *Blood Adv* 2021;5:2760–74.

https://doi.org/10.1182/BLOODADVANCES.2021004360.

58.   Burgess S., Davies NM., Thompson SG. Bias due to participant overlap in two-sample

Mendelian randomization. *Genet Epidemiol* 2016;40:597–608.

https://doi.org/10.1002/GEPI.21998.

59.   Schuermans A., Truong B., Ardissino M., Bhukar R., Slob EAW., Nakao T., et al.

Genetic Associations of Circulating Cardiovascular Proteins With Gestational

Hypertension and Preeclampsia. *JAMA Cardiol* 2024.

https://doi.org/10.1001/JAMACARDIO.2023.4994.

60.   Schuermans A., Pournamdari AB., Lee J., Bhukar R., Ganesh S., Darosa N., et al.

Integrative proteomic analyses across common cardiac diseases yield mechanistic

insights and enhanced prediction. *Nature Cardiovascular Research* 2024:1–15.

https://doi.org/10.1038/s44161-024-00567-0.

61.   Swerdlow DI., Kuchenbaecker KB., Shah S., Sofat R., Holmes M V., White J., et al.

Selecting instruments for Mendelian randomization in the wake of genome-wide

association studies. *Int J Epidemiol* 2016;45:1600–16.

https://doi.org/10.1093/IJE/DYW088.

62.   FinnGen Public Documentation. Available at:

https://finngen.gitbook.io/documentation/r10. Accessed November 5, 2024.

63.   The 1000 Genomes Project Consortium. A global reference for human genetic

variation. *Nature* 2015;526:68–74. https://doi.org/10.1038/nature15393.

64.   Purcell S., Neale B., Todd-Brown K., Thomas L., Ferreira MAR., Bender D., et al.

PLINK: A tool set for whole-genome association and population-based linkage

analyses. *Am J Hum Genet* 2007;81:559–75. https://doi.org/10.1086/519795.

65.   Wald A. The Fitting of Straight Lines if Both Variables are Subject to Error. *Ann Math*

Stat 1940;11:284–300. https://doi.org/10.1214/AOMS/1177731868.

66.   Hemani G., Zheng J., Elsworth B., Wade KH., Haberland V., Baird D., et al. The MR-

base platform supports systematic causal inference across the human phenome. *Elife*

2018;7. https://doi.org/10.7554/ELIFE.34408.

67. Yavorska OO., Burgess S. MendelianRandomization: an R package for performing
Mendelian randomization analyses using summarized data. *Int J Epidemiol*
2017;46:1734–9. https://doi.org/10.1093/IJE/DYX034.
68. Bowden J., Davey Smith G., Haycock PC., Burgess S. Consistent Estimation in
Mendelian Randomization with Some Invalid Instruments Using a Weighted Median
Estimator. *Genet Epidemiol* 2016;40:304–14. https://doi.org/10.1002/GEPI.21965.
69. Hartwig FP., Smith GD., Bowden J. Robust inference in summary data Mendelian
randomization via the zero modal pleiotropy assumption. *Int J Epidemiol*
2017;46:1985–98. https://doi.org/10.1093/IJE/DYX102.
70. Henry A., Gordillo-Marañón M., Finan C., Schmidt AF., Ferreira JP., Karra R., et al.
Therapeutic Targets for Heart Failure Identified Using Proteomics and Mendelian
Randomization. *Circulation* 2022;145:1205–17.
https://doi.org/10.1161/CIRCULATIONAHA.121.056663.
71. Gaziano JM., Concato J., Brophy M., Fiore L., Pyarajan S., Breeling J., et al. Million
Veteran Program: A mega-biobank to study genetic influences on health and disease.
J Clin Epidemiol 2016;70:214–23. https://doi.org/10.1016/J.JCLINEPI.2015.09.016.
72. Kanehisa M., Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic*
Acids Res 2000;28:27–30. https://doi.org/10.1093/NAR/28.1.27.
73. Bycroft C., Freeman C., Petkova D., Band G., Elliott LT., Sharp K., et al. The UK
Biobank resource with deep phenotyping and genomic data. *Nature* 2018;562:203–9.
https://doi.org/10.1038/s41586-018-0579-z.
74. Giambartolomei C., Vukcevic D., Schadt EE., Franke L., Hingorani AD., Wallace C., et
al. Bayesian Test for Colocalisation between Pairs of Genetic Association Studies
Using Summary Statistics. *PLoS Genet* 2014;10:e1004383.
https://doi.org/10.1371/JOURNAL.PGEN.1004383.
75. Aguet F., Barbeira AN., Bonazzola R., Brown A., Castel SE., Jo B., et al. The GTEx
Consortium atlas of genetic regulatory effects across human tissues. *Science (1979)*
2020;369:1318–30.
https://doi.org/10.1126/SCIENCE.AAZ1776/SUPPL_FILE/AAZ1776_TABLESS10-
S16.XLSX.
76. Becker DM., Segal J., Vaidya D., Yanek LR., Herrera-Galeano JE., Bray PF., et al.
Sex Differences in Platelet Reactivity and Response to Low-Dose Aspirin Therapy.
JAMA 2006;295:1420–7. https://doi.org/10.1001/JAMA.295.12.1420.
77. Taliun D., Harris DN., Kessler MD., Carlson J., Szpiech ZA., Torres R., et al.
Sequencing of 53,831 diverse genomes from the NHLBI TOPMed Program. *Nature*
2021;590:290–9. https://doi.org/10.1038/s41586-021-03205-y.
1219

1220

 Figures

 Figure 1. Study design.

1224   The flow chart shows the study design and the data sources. The diagram illustrates the design and three core assumptions of Mendelian

1225   randomization analyses: (1) the tested genetic instrument must be associated with the exposure of interest; (2) there must be no confounders

1226   affecting the association between the tested genetic instrument and outcome of interest; (3) the genetic instrument can only affect the outcome

1227   through its effect on the exposure of interest (i.e., no horizontal pleiotropy).

Figure 2. Analyses testing the associations of genetically predicted venous thromboembolism and other conditions with long COVID.

1229
Analyses testing the associations of genetically predicted venous thromboembolism and related (i.e., pulmonary embolism or deep vein

thrombosis) or unrelated conditions (i.e., "negative controls") with long COVID. The *x*-axis highlights the different conditions for which genetic

instruments were constructed and associations with long COVID were tested (i.e., different exposures). All genetic instruments for the indicated

conditions were constructed using genetic variants that were strongly associated with the corresponding condition ($P<5\times10^{-8}$) and not in linkage

disequilibrium with each other (linkage disequilibrium $R^2<0.001$) in up to 412,181 participants from FinnGen. Data for long COVID were obtained

from 997,700 participants from the Long COVID Host Genetics Initiative[24]. All associations were calculated using the inverse-variance weighted

method, visualized using odds ratios (ORs) and 95% confidence intervals (CIs). Genetically predicted venous thromboembolism was

significantly associated with a higher risk of long COVID (depicted in dark blue; the blue band corresponds to the 95% confidence interval of this

association).

1239

Figure 3. Associations of genetically predicted venous thromboembolism with long COVID in sensitivity and replication analyses.

Sensitivity analyses tested the association of genetically predicted venous thromboembolism with long COVID using the inverse-variance

weighted method with genetic instruments constructed using a range of P-value (y-axis) and linkage disequilibrium R^2 (colors) thresholds to

construct genetic instruments. The P-value threshold determines the minimal association strength for genetic variants included in each genetic

instrument, whereas the linkage disequilibrium R^2 threshold determines the maximal allowed correlation between genetic variants in each

genetic instrument. **a**, In primary analyses, associations were calculated using genetic instruments for venous thromboembolism constructed in

412,181 participants from FinnGen[21] and tested against long COVID ascertained in the Long COVID Host Genetics Initiative ("broad control

definition"; N=997,600)[24]. Replication analyses used: **b**, genetic instruments from an external genotyped cohort with ascertainment of venous

thromboembolism events (Million Veteran Program [MVP]; N=190,266)[27]; **c**, an alternative, stricter, control definition for long COVID in the Long

1249 COVID Host Genetics Initiative (N=40,910)[24]; and **d**, data from an external genotyped cohort with long COVID ascertainment (PHOSP-COVID;

1250 N=1,097)[28]. CI indicates confidence interval; OR, odds ratio.

Figure 4. Analyses exploring the associations of genetically predicted venous thromboembolism with acute COVID-19 phenotypes

and long COVID-resembling conditions.

Analyses testing the associations of genetically predicted venous thromboembolism with **a**, long COVID and acute COVID-19 phenotypes and

b, long COVID-resembling symptoms and conditions. For **a**, long COVID was ascertained in 997,700 participants from the Long COVID Host

Genetics Initiative[24] and the acute COVID-19 phenotypes were ascertained in up to 2,597,856 participants from the COVID-19 Host Genetics

Initiative[32]. The genetic instruments for venous thromboembolism were constructed using data from 412,181 FinnGen participants[21]. For **b**, all

long COVID-resembling conditions were ascertained in up to 412,181 participants from FinnGen[21]. The genetic instruments for venous

thromboembolism were constructed using data from 190,266 Million Veteran Program[27] participants to avoid sample overlap. For **a** and **b**, the *x*-

axes highlight the different outcomes that were tested. All genetic instruments for the indicated conditions were constructed using genetic

variants that were strongly associated with venous thromboembolism (*P*<5×10[-8]) and not in linkage disequilibrium with each other (linkage

disequilibrium *R²*<0.001). All associations were calculated using the inverse-variance weighted Mendelian randomization method, visualized

using odds ratios (ORs) and 95% confidence intervals (CIs). As shown in **Fig. 2**, genetically predicted venous thromboembolism was

significantly associated with a higher risk of long COVID (depicted in dark blue; the blue band corresponds to the 95%CI of this association).

Only "hospitalized COVID-19" and "critical COVID-19" (depicted in light blue) had associations that fell within the 95%CI of long COVID's

association.

 Figure 5. Identification of thromboembolism-related molecular mediators of long COVID.

a

b **c**

a, To identify coagulation-related genes driving the association of genetically predicted venous thromboembolism with long COVID, we tested

the effects of gene-specific instruments for venous thromboembolism with long COVID. These gene-specific instruments were constructed

using independent genetic variants associated with venous thromboembolism in FinnGen[21] within 200 kilobases of the indicated gene. The *x*-

axis depicts the different coagulation- and complement-related genes for which gene-specific instruments could be constructed. The *y*-axis and

color gradient depict the effect size (β) of each association. The circle size depicts the negative $\log_{10}$-transformed *P*-value for each association.

The greatest effect size was observed for *F2R*, which encodes the protease-activated receptor 1 (PAR-1) and was carried forward for further

analyses. **b**, We tested the associations of genetically predicted circulating levels of PAR-1 with long COVID. Primary analyses tested a genetic

instrument constructed using independent ($R^2<0.001$) *cis*-variants (within 200 kilobases of *F2R*) associated wih circulating PAR-1 levels

($P<5\times10^{-8}$) in the UK Biobank Pharma Proteomics Project (UKB-PPP)[37]. Sensitivity analyses tested instruments constructed using a range of *P*-

value (*y*-axis) and linkage disequilibrium R^2 (different colors) thresholds as well as those constructed using only intragenic variants (different

shapes). The *P*-value threshold determines the minimal association strength for genetic variants included in each genetic instrument, whereas

the linkage disequilibrium R^2 thresholds determines the maximal allowed correlation between genetic variants in each genetic instrument. **c**,

Additional analyses tested the associations of genetically predicted expression levels of *F2R* in physiologically relevant tissues and blood cell

types. Genetic instruments were constructed using independent genetic variants associated with expression levels of *F2R* in different tissues

and platelets in GTEx[40] and the GeneSTAR Research Study[41], respectively. Analyses were performed using genetic instruments with intragenic

variants or variants within 200 kilobases of *F2R* (different shapes). The *y*-axis and color gradient depict the effect size (β) of each association.

The circle size depicts the negative $\log_{10}$-transformed *P*-value of each association. For **a**, **b**, and **c**, all genetic instruments were tested in the

Long COVID Host Genetics Initiative[24] using the inverse-variance weighted method.

Venous thromboembolism

FINNGEN
FinnGen
(n_{cases}=21,021; $n_{controls}$=391,160)

Long COVID

COVID-19 hg
Long COVID Host Genetics Initiative
(n_{cases}=3,018; $n_{controls}$=994,582)

Construction of a genetic instrument
including 35 genetic variants associated
with venous thromboembolism

Primary analyses

Association of genetically predicted
venous thromboembolism with
long COVID

Schematic illustration of Mendelian randomization

Genetic instrument

Variants associated
with venous
thromboembolism

Exposure

Venous
thromboembolism

Outcome

Long COVID

Confounders

Other risk factors

Venous thromboembolism

Million Veteran Program
(n_{cases}=8,929; $n_{controls}$=181,337)

Long COVID

PHOSP-COVID
Improving long-term health outcomes
PHOSP-COVID
(n_{cases}=697; $n_{controls}$=400)

Sensitivity and replication
analyses

Sensitivity analyses using different
approaches and replication analyses
using different datasets

Analyses evaluating long COVID-
resembling phenotypes

Associations of venous
thromboembolism with clinical
outcomes resembling long COVID

Identification of molecular
mediators of long COVID

Analyses using gene-specific
instruments for thromboembolism,
blood proteins, and gene transcripts

Circulating protein levels

uk**biobank**
UK Biobank Pharma Proteomics
Project (*N*=34,557)

Gene transcript levels
(different tissues)

GTEx
Genotype-Tissue Expression Project
(*N*=715)

Gene transcript levels
(platelets)

GeneSTAR Research Study

GeneSTAR Research Study
(*N*=180)

Associations of different genetically predicted conditions with long COVID:

a Associations of genetically predicted venous thromboembolism with acute COVID−19:

b Associations of genetically predicted venous thromboembolism with long COVID−resembling conditions:

a

Associations of gene−specific genetic instruments for venous thromboembolism with long COVID:

b

Associations of circulating PAR−1 with long COVID:

c

Associations of *F2R* expression with long COVID: