Research article

Prefrontal tDCS for improving mental health and cognitive deficits in patients

with Multiple Sclerosis: a randomized, double-blind, parallel-group study

Nasim Zakibakhsh[1], Sajjad Basharpoor *[1], Hamidreza Ghalyanchi Langroodi[2], Mohammad
Narimani[1] Michael A Nitsche [3,4,5], Mohammad Ali Salehinejad *[3,6]

1 Department of Psychology, Faculty of Educational Sciences and Psychology, University of
Mohaghegh Ardabili, Ardabil, Iran
2 Clinical Development and Research Unit, Ghaem Hospital, Guilan University of Medical
Sciences, Rasht, Iran
3 Department of Psychology and Neurosciences, Leibniz Research Centre for Working
Environment and Human Factors, Dortmund, Germany
4 Bielefeld University, University Hospital OWL, Protestant Hospital of Bethel Foundation,
University Clinic of Psychiatry and Psychotherapy, Bielefeld, Germany
5 German Center for Mental Health (DZPG), Bochum, Germany
6 School of Cognitive Sciences, Institute for Research in Fundamental Sciences, Tehran, Iran
18

19 Number of manuscript words: 4373

20 Number of figures: 4

23 Correspondence: Mohammad Ali Salehinejad & Sajjad Basharpoor
24 salehinejadmohammadali@gmail.com
25 basharpoor_sajjad@uma.ac.ir

Abstract

Background: Multiple Sclerosis (MS) is an autoimmune disease associated with physical

disability, psychological impairment, and cognitive dysfunctions. Consequently, the disease

burden is substantial, and treatment choices are limited. In this randomized, double-blind study,

we used repeated prefrontal electrical stimulation and assessed mental health-related variables

(including quality of life, sleep, psychological distress) and cognitive dysfunctions (psychomotor

speed, working memory, attention/vigilance) in 40 patients with MS.

Methods: The patients were randomly assigned (block randomization method) to two groups of

sham ($n=20$), or 1.5-mA ($n=20$) transcranial direct current stimulation (tDCS) targeting the left

dorsolateral prefrontal cortex (F3) and right frontopolar cortex (Fp2) with anodal and cathodal

stimulation respectively (electrode size: 25 cm^2). The treatment included 10 sessions of 20

minutes stimulation delivered every other day. Outcome measures were quality of life, sleep

quality, psychological distress, and performance on a neuropsychological test battery dedicated to

cognitive dysfunctions in MS (psychomotor speed, working memory, and attention). All outcome

measures were examined pre-intervention and post-intervention. Both patients and technicians

delivering the stimulation were unaware of the study hypotheses and the type of stimulation

being used.

Results: The active protocol significantly improved quality of life and reduced sleep difficulties

and psychological distress compared to the sham group. The active protocol, furthermore,

improved psychomotor speed, attention and vigilance, and some aspects of working memory

performance compared to the sham protocol. Improvement in mental health outcome measures

was significantly associated with better cognitive performance.

Conclusions: Modulation of prefrontal regions with tDCS ameliorates secondary clinical

symptoms and results in beneficial cognitive effects in patients with MS. These results support

applying prefrontal tDCS in larger trials for improving mental health and cognitive dysfunctions

in MS.

Trial registration: ClinicalTrials.gov Identifier: NCT06401928

Keywords: Multiple Sclerosis; transcranial direct current stimulation

1. Introduction

Multiple Sclerosis (MS) is the most common autoimmune disorder of the central nervous system, afflicting more than 2.5 million people worldwide, especially young people (1, 2). It is a progressive chronic disease, caused by an autoimmune attack, which results in the gradual loss of the myelin sheath around neuronal axons of the central nervous system. Depending on the affected neurons, different symptoms are expressed in the course of the disease; nonetheless, physical disability, cognitive impairment, and decreased quality of life are common consequences of the disease (3). MS results in motor, sensory, cognitive, and neuropsychiatric symptoms, all of which can occur independently of one another (4). The disease is associated with a high mental health burden due to primary symptoms of sensory and motor deficits. Some of the most common symptoms include fatigue, vision problems, numbness and tingling, muscle spasms, stiffness and weakness, mobility problems, and pain (2, 5).

In addition to primary symptoms such as fatigue, motor deficits, and visual disturbances, people with MS also experience disability in mental health variables, including quality of life, sleep, and emotional disturbances (6-8). Cognitive deficits are also commonly observed symptoms in MS and include deficits in attention and vigilance, information processing, executive functioning, processing speed, and long-term memory (4) with a profound effect on instrumental activities of daily living (9). Both primary and secondary symptoms can vary in severity and may come and go, depending on disease activity, progression, and the type of MS, which include relapsing-remitting, or chronic-progressive dynamics (8, 10). Individuals with MS need to work closely with healthcare professionals to manage their symptoms and optimize their quality of life. Considering the burden of treatment (11), the complex pathophysiology and psychophysiology of MS, and limited treatment options for secondary symptoms (physical

therapy, psychotherapy, cognitive rehabilitation) (3), there is a need for novel and effective

treatment for both primary and secondary symptoms.

Transcranial direct current stimulation (tDCS) is a safe and easy-to-use noninvasive brain

stimulation intervention for studying and modifying human brain functions (12, 13). It involves

applying a low-intensity direct current to the scalp, which induces alterations of the resting

membrane potential of neurons. At the macroscale level, anodal stimulation depolarizes neurons

at a subthreshold level, making them more likely to fire action potentials, while cathodal

stimulation hyperpolarizes neurons, reducing their excitability (14, 15). By changing the

excitability of the brain and inducing neuroplasticity effects, it is possible to restore functional

brain abnormalities and affect target behavior/cognition. Previous studies have shown functional

brain abnormalities in MS that are partially related to secondary symptoms in MS (16, 17). These

abnormalities can take many different forms, such as modified brain activity patterns, neural

network disturbances, and modifications of cognitive functions. Key elements of functional brain

impairments in MS include changes in pain processing pathways as well as alterations of brain

regions involved in the regulation of mood and arousal, including the limbic system,

hypothalamus, and motor regions (18-20). Furthermore, there is growing evidence of grey matter

changes in MS patients that are linked to disability and other clinical symptoms in these patients

(21, 22).

Previous tDCS studies have mostly focused on examining the efficacy of prefrontal and

motor tDCS on clinical symptoms (e.g., fatigue, pain), motor symptoms, and cognitive deficits of

patients with MS (23-26). There is inconsistency in study results regarding the cognitive effects

of non-invasive brain stimulation, such as tDCS, on MS patients. In addition to cognitive and

motor symptoms, MS patients also suffer from other mental health issues that are less commonly

addressed in tDCS studies. While there have been some studies on the effects of transcranial

direct current stimulation (tDCS) on mental health-related variables, such as quality of life (27),

sleep (28), and emotional difficulties (29), there is still a lack of research on the impact of tDCS

on both mental health-related variables and cognitive performance in patients with MS.

Accordingly, the purpose of this study is to address this research gap by examining 1) the effects

of repeated prefrontal tDCS on mental-health-related variables (i.e., quality of life, sleep

difficulties, psychological distress) in patients with MS and 2) the effects of repeated prefrontal

tDCS on the cognitive performance of patients with MS in a randomized, double-blind, sham-

controlled trial.

2. Methods

2.1. Participants

Eighty MS patients from the local community (Rasht, Iran) were screened for inclusion in

the study. The inclusion criteria were: (1) diagnosis of MS according to the diagnostic criteria for

multiple sclerosis: 2010 Revisions of the McDonald criteria (Polman et al., 2011), certified by a

professional neurologist, (2) being 25–55 years old, (3) being non-smoker, (4) no previous

history of neurological diseases, brain surgery, epilepsy, seizures, brain damage, head injury, or

metal brain implants, (5) absence of other psychiatric disorders except mood and anxiety

disorders, and (6) no relapse of MS symptoms within the last two months before beginning the

experiment. Of those who met the inclusion criteria (n=60), forty patients were randomly

assigned to the experimental (active tDCS) and control (sham tDCS) groups based on a sample

size analysis (f = 0.30, α = 0.05, power = 0.95, mixed-model ANOVA for 2 groups with 2

measurements) which resulted in a sample size of 40 patients. For the group assignment of the

participants, block randomization method was applied. Three patients decided to withdraw from

the study following the first session and thus the final analysis was conducted on 37 patients

(mean age = 37.30, SD = 6.21, 27 females, 10 males) (see Table 1 and Figure 1 for demographics

and study inclusion). This is a retrospectively registered clinical trial (ClinicalTrials.gov

Identifier: NCT06401928) approved by the Ethics Committee of the Mohaghegh Ardabili

University (Ethics code: IR.MAU.REC.1401.94). Participants gave their written informed

consent before participation.

2.2. Measures

2.2.1.   Mental health assessments

The primary outcome measures of the study included quality of life, sleep difficulties, and

psychological distress that were measured with the Multiple Sclerosis Impact Scale (MSIS-29)

(30), the Mini Sleep Questionnaire (MSQ) (31), and the Depression Anxiety Stress Scale-21

(DASS-21) (32) respectively. The MSIS-29 is a measure of the physical and psychological

impact of MS from the patients' perspective consisting of 29 questions with the first 20 items

focusing on the physical impact and the remaining 9 on the psychological impact. The Mini Sleep

Questionnaire (MSQ) is typically used to screen sleep disturbances in clinical populations, and

the DASS-21 is a 21-item self-report measure designed to assess the severity of general

psychological distress and symptoms related to depression, anxiety, and stress in adults and older

adolescents (+17 years). Details about these measures and their psychometric properties can be

found in the supplementary information. A native-language version of each test was used in this

study (33-35).

2.2.2.   Neuropsychological performance

In addition to mental health-related outcome variables, we assessed the

neuropsychological performance and cognitive functioning of the MS patients with several

subtests of a neuropsychological battery designed for MS patients using the Cambridge Cognition

Neuropsychological Battery (CANTAB). The battery included 3 computerized tests to measure

psychomotor speed, working memory and sustained attention / vigilance via the Reaction Time

(RTI), Spatial Working Memory (SWM) and Rapid Visual Information Processing (RVP) tests

respectively. The tests included in this battery are highly sensitive to cognitive impairments in

MS across the disease severity spectrum (36). Details about these measures are available in the

supplementary information.

2.3. tDCS

We used a two-channel Neurostim stimulator device (MadinaTeb, Tehran, Iran) powered

by a 9-volt alkaline battery. Electrical current was applied through a pair of saline-soaked sponge

electrodes (5×5 cm) for a period of 20 minutes (with 30 seconds ramping up and 30 seconds

ramping down) and a stimulation intensity of 1.5 mA. We had two stimulation conditions: active

and sham tDCS. In the active condition, the anodal electrode was placed over the left dorsolateral

prefrontal cortex (left DLPFC) over the F3 electrode position according to the 10-20 EEG

International System and the cathodal electrode was placed over the right frontopolar cortex

(FPC) (electrode position Fp2) which includes orbitofrontal cortex (BA 10, 11) (37, 38). To

reduce shunting of current between the electrodes through the scalp, the edges had a distance of

at least 6 cm. In the sham condition, a sham stimulation was employed with the same electrode

configuration. Here, the electrical current was ramped up for 30 seconds followed by 30 seconds

of stimulation and 30 seconds of ramping down to generate the same sensation as the active

condition, and then was turned off without the participants' knowledge. This method of sham

stimulation has been proven reliable (39). During the study, participants were blinded to the type

of stimulation they received. A survey was conducted after each session to document any

reported side effects, but blinding efficacy (asking participants to guess about the type of

stimulation) was not explored due to the multi-session design of the study. After finalizing the

study protocol, also the patients in the sham group were assigned to active tDCS intervention, but

the latter procedure was beyond the focus of the study protocol.

2.4. Procedure

Prior to the experiment, participants completed a brief questionnaire to evaluate their

suitability for brain stimulation. The active tDCS and sham groups received 10 sessions of

stimulation (three sessions per week, resulting in a total of three weeks plus an additional day) with

24-hour between-session intervals (except for the 4th and 8^{th} sessions that had 48-h interval due to

the weekend). Clinical measures (psychological distress questionnaires and cognitive assessment)

were evaluated before the first tDCS session (pre-intervention), and right after the end of the last

tDCS session (post-intervention). All tDCS sessions were scheduled between 2:00 and 5:00 p.m.

and were monitored for sleep pressure to mitigate a potential impact of circadian variation on

cortical excitability and neuroplasticity induction for all participants across all sessions (40, 41).

Before starting the experiment, the participants were given instructions about the cognitive tasks.

The measurement sessions before the intervention and after the intervention took about 2 hours.

Each session was conducted on two separate days to prevent fatigue. The first day was dedicated

to clinical assessment (quality of life, sleep, psychological distress), and the second day for

cognitive assessment (psychomotor speed, working memory, attention/vigilance). To ensure that

the experiment was double-blinded, independent researchers who were not involved in the

administration of stimulation sessions were responsible for supervising the examination of outcome

measures, data analysis, and group assignment (see Fig. 2).

2.5. Statistical analysis

Data were analyzed with the statistical package SPSS, version 27.0 (IBM, SPSS, Inc.,

Chicago, IL). The normality and homogeneity of data variance were confirmed by Shapiro-Wilk

and Levin tests, respectively. Mixed model ANOVAs were conducted for the dependent variables

(MSIS, DASS, MSQ scores; psychomotor speed, working memory and attention tasks) with

"group" (active vs sham) as the between-subject and time (pre-intervention, post-intervention) as

the within-subject factors. Mauchly's test was used to evaluate sphericity of the data before

performing the respective ANOVAs ($p<0.05$). In case of violation, degrees of freedom were

corrected using Greenhouse-Geisser estimates of sphericity. Post hoc comparisons were

conducted with Fisher's LSD post-hoc tests for individual mean difference comparisons across

groups (active 1.5 mA, sham) and time points (pre-intervention, post-intervention). Pearson's

correlation was also calculated to explore potential associations between mental health-related

variables and cognitive performance. The critical level of significance was 0.05 for all statistical

analyses.

3. Results

3.1. Data overview

Demographic information is summarized in Table 1. Patients tolerated the stimulation

well and no adverse effects were reported during and after stimulation. No significant difference

was found between the group ratings of tDCS side effects except for the burning sensation

($p=0.044$) which was higher in the active group (Table 2). The burning sensation, however, did

not correlate with any outcome measure, thus ruling out respective effects. No significant

between-group differences were observed for baseline measurements of outcome variables (Table

3).

3.2. Efficacy of tDCS on quality of life, sleep quality, and psychological distress,

The results of the 2 (group) $\times$ 2 (time: pre, post) mixed model ANOVA showed a

significant main effect of time (F_1=7.94, p=0.008, ηp2=0.21) and a significant group$\times$time

interaction (F_1=9.98, p=0.004, ηp2=0.25) but no main effect of group (F_1=0.94, p=0.338) on

quality of life scores measured by the MSIS-29. Fisher's LSD post-hoc tests showed a significant

increase in quality of life scores in patients with MS who received active tDCS after the

intervention compared to pre-intervention (t=4.74, p<0.001). When compared to the sham group,

active tDCS significantly improved *quality of life* scores after the intervention (t=4,93, p<0.001).

Baseline between-group comparisons (active groups vs sham) showed no significant differences

in the pre-intervention scores (Table 4, Fig. 3A, B).

With respect to the sleep quality scores measured by the MSQ, the results of a 2 $\times$ 2

mixed model ANOVA showed a significant main effect of time ($F_{1,30}$=30.598, p<0.001,

ηp2=0.505) and a significant group$\times$time interaction ($F_{1,30}$=37.314, p<0.001, ηp2=0.554) but no

main effect of group ($F_{1,30}$=1.065, p=0.31). Fisher's LSD post-hoc tests showed a significant

improvement of sleep quality scores only in patients with MS that received active tDCS after the

intervention compared to pre-intervention ($t_{total\ sleep}$=4.50, p<0.001, $t_{insomnia}$=4.48, p<0.001,

$t_{hypersomnia}$=3.29, p<0.001) but not in the sham group. When compared to the sham group, the

active tDCS group significantly improved with respect to the sleep total score (t=3.38, p=0.001),

insomnia (t=2.90, p=0.005) and hypersomnia (t=2.82, p=0.006). Baseline between-group

comparisons were not significant (p>0.05) (Fig. 3 C-H)

For psychological distress, a 3 (domain: depression, anxiety, stress) $\times$ 2 (time: pre, post) $\times$

2 (group) mixed model ANOVA was conducted. The results showed significant main effects of

time ($F_{1,30}$=6.13, p=0.019, ηp2=0.17), domain ($F_{2,60}$=15.608, p<0.001, ηp2=0.342) and a

significant group$\times$time interaction ($F_{1,30}$=23.505, p<0.001, ηp2=0.439). The main effect of group,

and the domain$\times$time, domain$\times$group and domain$\times$group$\times$time interactions were not significant

(Table 4). Fisher's LSD post-hoc tests showed a significant decrease in the DASS total score only

in patients with MS who received active tDCS after the intervention compared to pre-intervention

(t=2.46, p=0.020) but not in the sham group. Compared to the sham group, the active tDCS group

showed significantly reduced psychological distress ($t_{depression}$=2.73, p=0.008, $t_{anxiety}$=1.99, df=30,

p=0.050, t_{stress}=2.43, p=0.017) after the intervention. Baseline between-group comparisons

(active groups vs sham) showed no significant differences in the pre-intervention scores (p>0.05)

(Fig. 3 M-P).

3.3. Efficacy of tDCS on cognitive functions in MS patients

3.3.1.   Psychomotor speed

Psychomotor speed was evaluated with the RTI task. A significant main effect of time

($F_{1,35}$=127.32, p<0.001, ηp2=0.784) and a significant group$\times$time interaction ($F_{1,35}$=73.491,

p<0.001, ηp2=0.677), but no main effect of group was found for the 5-choice movement time. A

similar significant main effect of time ($F_{1,35}$=11.684, p=0.002, ηp2=0.25) and a significant

group$\times$time interaction ($F_{1,35}$=34.992, p<0.001, ηp2=0.50), but a non-significant main effect of

group was found for the 5-choice reaction time (Table 4). Fisher's LSD post-hoc tests revealed a

significantly faster movement time in patients who received active tDCS after the intervention

compared to pre-intervention (t=5.56, p<0.001). Additionally, in the post-intervention condition,

movement time was significantly shorter in the active group compared to the sham tDCS group (t

$=3.86$, $p<0.001$). For reaction time, neither group responded faster after intervention as compared

to the pre-intervention condition. However, performance after intervention was significantly

faster in the active vs sham group ($t=2.08$, $p=0.043$). Baseline between-group comparisons were

not significant in both outcome variables ($p>0.05$) (Fig. 4 A-H).

Other outcome variables of interest in the RTI task were *simple*-choice movement time

and reaction time. Here, the results of the mixed model ANOVA revealed a significant main

effect of time ($F_{1,35}=17.476$, $p<0.001$, $\eta p2=0.333$) and a significant group×time interaction

($F_{1,35}=20.905$, $p<0.001$, $\eta p2=0.374$) but no main effect of group on movement time, and

significant main effects of time ($F_{1,35}=24.729$, $p<0.001$, $\eta p2=0.414$) and group ($F_{1,35}=6.439$,

$p=0.016$, $\eta p2=0.155$) and a significant interaction between both variables ($F_{1,35}=40.317$, $p<0.001$,

$\eta p2=0.535$) with respect to reaction time. Fisher's LSD post-hoc tests showed significantly faster

movement and reaction times in the psychomotor speed task in patients who received active

tDCS after the intervention compared to pre-intervention ($t=4.09$, $p<0.001$; $t=5.60$, $p<0.001$).

Both movement and reaction times in the psychomotor speed task after intervention were

significantly faster in the active tDCS compared to the sham group ($t=2.92$, $p=0.004$; $t=5.31$,

$p<0.001$). Baseline between-group comparisons were not significant for both outcome variables

($p>0.05$) (Fig. 4 A-H).

3.3.2. Working memory

Here strategy scores and total errors were analyzed. For the strategy scores, the results of

the mixed model ANOVA showed no significant main effects of time, group, or their interaction

(Table 4). For the total errors, however, a significant main effect of time ($F_{1,35}=9.081$, $p=0.005$,

ηp2=0.206) but no main effect of group or the group×time interaction was observed.

Accordingly, no post hoc tests were conducted for errors (Fig. 4- I-L).

3.3.3. Attention and vigilance

The results of the mixed model ANOVA showed a significant main effect of time

($F_{1,35}$=39.44, $p<0.001$, ηp2=0.53) and a significant group×time interaction ($F_{1,35}$=47.032,

$p<0.001$, ηp2=0.573) but no main effect of group ($F_{1,35}$=1.698, $p=0.201$) on Hits (accurate

responses). Patients who received active tDCS had a significantly higher number of hits after

intervention vs pre intervention ($t=2.96$, $p=0.004$), and the number of hits was also significantly

higher in the real intervention than the number of hits in those who received sham tDCS after

intervention ($t=2.79$, $p=0.006$). Similarly, significant main effects of time ($F_{1,35}$=291.414,

$p<0.001$, ηp2=0.893), group ($F_{1,35}$=32.178, $p<0.001$, ηp2=0.479) and a significant group×time

interaction ($F_{1,35}$=292.129, $p<0.001$, ηp2=0.893) emerged for the mean reaction time. Fisher's

LSD post-hoc test showed a faster reaction time after the intervention as compared to the pre-

intervention only in the active tDCS group ($t=7.28$, $p<0.001$). Here, however, mean reaction time

at baseline was significantly different across groups, but not after intervention (Fig. 4 M-P).

3.4. Improved mental health is associated with cognitive task performance after intervention

We calculated Pearson's correlations to assess if changes in mental health outcomes are

linked to cognitive task performance after intervention and found a significant correlation among

specific variables. Notably, enhanced quality of life was significantly associated with faster

performance in the psychomotor task ($r_{5choice-move-time}$=-0.383, $p<0.05$, $r_{5choice-reaction-time}$=-0.481,

$p<0.01$, $r_{simple-move-time}$=-0.587, $p<0.01$). Improved attention/vigilance (i.e., higher performance

accuracy) also showed significant negative correlations with insomnia (r=-0.423, $p<0.05$),

hypersomnia (r=-0.478, p<0.01), and overall sleep difficulties (r=-0.434) (i.e., higher accuracy

was associated with lower sleep disturbances). Additional significant pertinent correlations (e.g.,

faster psychomotor speed and lower sleep issues) are detailed in Table S1 (supplementary

materials).

4. Discussion

As a neuroinflammatory disease, MS impairs neuronal efficacy and neuroplasticity. Non-

pharmacological approaches, such as non-invasive brain stimulation, have the potential to

enhance the outcomes of pharmacological and physical interventions in MS by enhancing

neuroplasticity and functional connectivity (42-44) and potentially have anti-inflammatory effects

as observed in several noninvasive brain stimulation studies (45, 46), although the latter is not

completely clarified. In this randomized, double-blind study, we investigated the effectiveness of

repeated prefrontal tDCS in enhancing mental health and cognitive functions in MS patients to

assess their impact and potential associations between improvements. Patients who received real

tDCS reported a significantly higher quality of life and reduced sleep difficulties after the

intervention. They showed also an overall lower level of psychological distress after the

intervention compared to the sham group. In addition to these mental health-related variables,

patients who received prefrontal tDCS showed superior performance in a neuropsychological test

battery that measured psychomotor speed, working memory, and visual attention and vigilance

which are frequently impaired in MS patients.

The use of tDCS to improve motor and/or cognitive deficits in MS has been less common

compared to other neurological/psychiatric conditions. Moreover, results have been mixed so far.

While some studies showed promising benefits in ameliorating fatigue, pain, and cognitive

symptoms, but inconsistent effects of tDCS on motor symptoms (23, 24), others show a more

promising effect on motor functions (25) which could partially be due to heterogeneous protocols

applied in these studies, but also different patient characteristics. This is the first study that

specifically investigated the efficacy of tDCS on quality of life, mental health-related variables

and cognitive functioning of patients with MS. Our study provides supportive evidence for the

efficacy of tDCS in improving quality of life and amelioration of psychological and cognitive

deficits in patients with MS. Here, the applied protocol and proposed mechanisms of effects are

discussed.

344   Prefrontal tDCS has been extensively used to improve cognitive and emotional

impairments in clinical populations, as well as to enhance cognitive abilities in healthy

individuals. Prefrontal tDCS can have pro-cognitive effects in neuropsychiatric disorders that are

characterized by executive and cognitive deficits, such as mood and anxiety disorders (47, 48),

obsessive-compulsive disorder (49, 50), and substance use disorder (51, 52). Pro-cognitive effects

were also shown in neurological disorders such as Parkinson's disease and stroke (53, 54) and

major neurodevelopmental disorders such as ADHD and autism (55-57). It has been also

effectively used for improving fatigue and quality of life (58, 59). The rationale behind the use of

prefrontal tDCS for enhancing cognition mostly comes from the involvement of different regions

of the prefrontal cortex in various aspects of cognitive functions (60-62) which seems to be at

least partially applicable for MS.

355   Cognitive impairment in MS is the consequence of widespread lesions in the brain and

frequently includes deficits in complex attention, efficiency of information processing, executive

functioning, processing speed, and long-term memory (4, 16). Recent large-scale neuroimaging

studies conducted in MS patients have shown a causal relationship between the pathophysiology

of MS and different brain regions, including the frontal and prefrontal cortices (specifically the

orbitofrontal cortex) and other regions that are connected to these cognition-related areas (such as

the parahippocampal gyrus) (21). Our study applied prefrontal tDCS over prefrontal regions

(DLPFC and right FPC) which are important for cognitive deficits in MS, leading to improved

psychomotor speed and attention/vigilance in patients who received active stimulation. In

addition to the cognitive improvement, quality of life of the patients was improved after the

intervention. The frequency of sleep difficulties and states of depression and anxiety were also

significantly reduced after intervention. Sleep improvement after the active tDCS can be

explained by the more recently stressed role of the cortico-thalamo-cortical feedback loop, which

is a top-down regulatory system related to cortical areas (e.g., PFC), in regulating arousal and

sleep (63). Mood alleviating effects can be partially explained by enhanced cognitive control as a

result of tDCS over the DLPFC, which regulates mood and valence of emotional experiences (64,

65). Importantly, there is a link between the prefrontal cortex, specially the DLPFC, the medial

PFC, and the amygdala network in MS patients who experience emotional difficulties (66), and

reduced depression, anxiety, and distress following intervention might be explained via this

prefrontal-amygdala related emotion regulation. Briefly, Upregulating DLPFC activity can

reduce amygdala activity (67), which is usually hyperactive during negative emotional processing

in MS (66). This effect, coupled with the functional connectivity-enhancing effect of tDCS (68),

may be a possible mechanism of action.

The current study's findings should also be compared with previous tDCS studies in MS

patients. Recent systematic reviews and metanalyses of tDCS studies conducted in MS show

inconsistent results about the cognitive effects of tDCS with some studies reporting clear benefits

in ameliorating cognitive symptoms (23), some partial improvement in specific aspects of

cognition (e.g., vigilance) (25) and some with no strong evidence for effectiveness of tDCS on

cognition (24). Previous tDCS cognition-related studies have commonly targeted the left DLPFC

with the anodal polarity while using the return electrode over the right DLPFC, right shoulder, or

right supraorbital area, and stimulation intensity and duration ranged from 1.5 to 2 mA and from

20 to 30 min respectively (25). The present study aimed to shed light on the effects of tDCS on

specific secondary deficits, but also on the interplay between cognitive functions and mental

health domains, which are both significantly impacted by MS, but have been less studied in

previous works. Importantly, here we observed significant correlations between mental health-

related outcome variables and cognitive performance, suggesting that these two impaired

domains in MS (mental health and cognitive deficits) are interrelated. One consideration

regarding the protocol in the present study (anodal F3- cathodal Fp2) is the use of the reference

electrode over the right FPC instead of the right DLPFC. The bilateral DLPFC protocol can also

be applied if cognitive functions are the main focus, as indicated in previous tDCS studies in MS

(25) and other neuropsychiatric disorders (69, 70). In this study, we also aimed to explore the

impact of the intervention on participants' emotional experiences. FPC, which includes the

orbitofrontal cortex (38), was selected over the right DLPFC due to its significant role in

emotional regulation and direct connections with subcortical regions (60). The relative efficacy of

DLPFC-FPC versus bilateral DLPFC tDCS needs to be compared to determine which is more

effective in enhancing mental health and cognition-related variables.

Limitation

We acknowledge that our study has some limitations. Firstly, although the sample size of

this study is larger than the majority of the previous tDCS studies in MS (24, 25), it is still

relatively small, and therefore, these findings need to be confirmed in larger trials in the future.

Secondly, we did not measure outcome variables over a reasonable follow-up period to determine

how long the observed effects lasted and the intervention was not applied daily but every second

or third day, which might have limited efficacy. Finally, we did not include any physiological

measures, which would have been informative to understand how brain functions and physiology

are affected by the intervention and how they align with the behavioural results. These limitations

should be considered when interpreting the results of our study and for future larger trials.

Conclusions

In conclusion, in this study, we upregulated the activity of the left DLPFC and

downregulated the activity of the right FPC in 37 patients with MS and evaluated the quality of

life, sleep difficulties, psychological distress, and cognitive functions after 10 sessions of brain

stimulation. The group that received real intervention showed significant improvement in quality

of life, and reduced psychological distress and sleep difficulties, as compared to the placebo

group. This improvement was linked to an enhancement of psychomotor speed and

attention/vigilance of the patients in the real intervention group. These findings suggest that

prefrontal tDCS targeting the left DLPFC and right FPC has the potential to be a valuable

intervention for treating secondary symptoms (mental-health-related variables and cognitive

deficits) in patients with MS.

List of abbreviations

MS: Multiple Sclerosis
tDCS: transcranial direct current stimulation
MSIS-29: Multiple Sclerosis Impact Scale
MSQ: Mini Sleep Questionnaire
DASS-21: Depression Anxiety Stress Scale-21
RTI: Reaction Time task
SWM: Spatial Working Memory task
RVP: Rapid Visual Information Processing task
DLPFC: dorsolateral prefrontal cortex
FPC: frontopolar cortex

Declarations

Ethics approval and consent to participate

All patients gave their written consent to participate in the study. The protocol was conducted in

accordance with the latest version of the Declaration of Helsinki and was approved by the

Institutional Review Board and Ethical Committee at Mohaghegh Ardabili University.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding

author upon reasonable request.

Competing interests

Michael Nitsche is a member of the Scientific Advisory Boards of Neuroelectrics and Precisis.

All other authors declare no competing interests

Funding

Not applicable

Authors' contributions

NZ: conceptualization, investigation, data curation, visualization, validation; SB & HGL & MN:

supervision, resources, project administration, data curation; MAN: writing—review & editing;

MAS: supervision, methodology, writing original draft, writing—review & editing, visualization,

formal analysis. All authors read and approved the final manuscript.

Acknowledgments

We would like to thank the MS Association located in the city of Rasht, Iran for their assistance

in recruiting patients.

References

1. Attfield KE, Jensen LT, Kaufmann M, Friese MA, Fugger L. The immunology of multiple sclerosis. Nat Rev Immunol. 2022;22(12):734-50.
2. Compston A, Coles A. Multiple sclerosis. Lancet (London, England). 2008;372(9648):1502-17.
3. McGinley MP, Goldschmidt CH, Rae-Grant AD. Diagnosis and Treatment of Multiple Sclerosis: A Review. JAMA. 2021;325(8):765-79.
4. Chiaravalloti ND, DeLuca J. Cognitive impairment in multiple sclerosis. The Lancet Neurology. 2008;7(12):1139-51.
5. Ajdacic-Gross V, Steinemann N, Horváth G, Rodgers S, Kaufmann M, Xu Y, et al. Onset Symptom Clusters in Multiple Sclerosis: Characteristics, Comorbidities, and Risk Factors. Front Neurol. 2021;12.
6. Marrie RA, Reingold S, Cohen J, Stuve O, Trojano M, Sorensen PS, et al. The incidence and prevalence of psychiatric disorders in multiple sclerosis: A systematic review. Multiple Sclerosis Journal. 2015;21(3):305-17.
7. Laslett LL, Honan C, Turner JA, Dagnew B, Campbell JA, Gill TK, et al. Poor sleep and multiple sclerosis: associations with symptoms of multiple sclerosis and quality of life. Journal of Neurology, Neurosurgery & Psychiatry. 2022;93(11):1162-5.
8. Ben-Zacharia AB. Therapeutics for Multiple Sclerosis Symptoms. Mount Sinai Journal of Medicine: A Journal of Translational and Personalized Medicine. 2011;78(2):176-91.
9. Benedict RHB, Amato MP, DeLuca J, Geurts JJG. Cognitive impairment in multiple sclerosis: clinical management, MRI, and therapeutic avenues. Lancet Neurol. 2020;19(10):860-71.
10. Disanto G, Zecca C, MacLachlan S, Sacco R, Handunnetthi L, Meier UC, et al. Prodromal symptoms of multiple sclerosis in primary care. Ann Neurol. 2018;83(6):1162-73.
11. Ferrari AJ, Santomauro DF, Aali A, Abate YH, Abbafati C, Abbastabar H, et al. Global incidence, prevalence, years lived with disability (YLDs), disability-adjusted life-years (DALYs), and healthy life expectancy (HALE) for 371 diseases and injuries in 204 countries and territories and 811 subnational locations, 1990–2021: a systematic analysis for the Global Burden of Disease Study 2021. The Lancet. 2024.
12. Polanía R, Nitsche MA, Ruff CC. Studying and modifying brain function with non-invasive brain stimulation. Nat Neurosci. 2018;21(2):174-87.
13. Salehinejad MA, Siniatchkin M. Safety of noninvasive brain stimulation in children. Current Opinion in Psychiatry. 2024:10.1097/YCO.0000000000000923.
14. Nitsche MA, Paulus W. Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. J Physiol. 2000;527 Pt 3(Pt 3):633-9.
15. Yavari F, Chhabra H, Polania R, Nitsche MA. Mechanisms of action of transcranial direct current stimulation. In: Quevedo J, Brunoni AR, Hamani C, editors. Interventional Psychiatry: Academic Press; 2024. p. 149-86.
16. Cader S, Cifelli A, Abu-Omar Y, Palace J, Matthews PM. Reduced brain functional reserve and altered functional connectivity in patients with multiple sclerosis. Brain. 2005;129(2):527-37.
17. Pantano P, Mainero C, Caramia F. Functional Brain Reorganization in Multiple Sclerosis: Evidence from fMRI Studies. J Neuroimaging. 2006;16(2):104-14.
18. Rocca MA, Filippi M. Functional MRI in multiple sclerosis. J Neuroimaging. 2007;17 Suppl 1:36s-41s.
19. Rocca MA, Schoonheim MM, Valsasina P, Geurts JJG, Filippi M. Task- and resting-state fMRI studies in multiple sclerosis: From regions to systems and time-varying analysis. Current status and future perspective. NeuroImage: Clinical. 2022;35:103076.

20.   Tahedl M, Levine SM, Greenlee MW, Weissert R, Schwarzbach JV. Functional Connectivity in
Multiple Sclerosis: Recent Findings and Future Directions. Front Neurol. 2018;9.
21.   Sun D, Wang R, Du Q, Zhang Y, Chen H, Shi Z, et al. Causal relationship between multiple
sclerosis and cortical structure: a Mendelian randomization study. J Transl Med. 2024;22(1):83.
22.   Colato E, Stutters J, Tur C, Narayanan S, Arnold DL, Gandini Wheeler-Kingshott CAM, et al.
Predicting disability progression and cognitive worsening in multiple sclerosis using patterns of grey
matter volumes. J Neurol Neurosurg Psychiatry. 2021;92(9):995-1006.
23.   Hiew S, Nguemeni C, Zeller D. Efficacy of transcranial direct current stimulation in people
with multiple sclerosis: a review. Eur J Neurol. 2022;29(2):648-64.
24.   Uygur-Kucukseymen E, Pacheco-Barrios K, Yuksel B, Gonzalez-Mego P, Soysal A, Fregni F.
Non-invasive brain stimulation on clinical symptoms in multiple sclerosis patients: A systematic
review and meta-analysis. Multiple Sclerosis and Related Disorders. 2023;78:104927.
25.   Li S, Zhang Q, Zheng S, Li G, Li S, He L, et al. Efficacy of non-invasive brain stimulation on
cognitive and motor functions in multiple sclerosis: A systematic review and meta-analysis. Front
Neurol. 2023;14.
26.   Saiote C, Goldschmidt T, Timäus C, Steenwijk MD, Opitz A, Antal A, et al. Impact of
transcranial direct current stimulation on fatigue in multiple sclerosis. Restor Neurol Neurosci.
2014;32:423-36.
27.   Mortezanejad M, Ehsani F, Masoudian N, Zoghi M, Jaberzadeh S. Comparing the effects of
multi-session anodal trans-cranial direct current stimulation of primary motor and dorsolateral
prefrontal cortices on fatigue and quality of life in patients with multiple sclerosis: a double-blind,
randomized, sham-controlled trial. Clin Rehabil. 2020;34(8):1103-11.
28.   Chalah MA, Grigorescu C, Kümpfel T, Lefaucheur J-P, Padberg F, Palm U, et al. The effects
of transcranial direct current stimulation on sleep in patients with multiple sclerosis–a pilot study.
Neurophysiol Clin. 2022;52(1):28-32.
29.   Hsu W-Y, Cheng C-H, Zanto TP, Gazzaley A, Bove RM. Effects of Transcranial Direct Current
Stimulation on Cognition, Mood, Pain, and Fatigue in Multiple Sclerosis: A Systematic Review and
Meta-Analysis. Frontiers in Neurology. 2021;12:626113.
30.   Hobart J, Lamping D, Fitzpatrick R, Riazi A, Thompson A. The Multiple Sclerosis Impact Scale
(MSIS-29): a new patient-based outcome measure. Brain. 2001;124(Pt 5):962-73.
31.   Zomer J. Mini Sleep Questionnaire (mSQ) for screening large populations for EDS
complaints. Sleep. 1985.
32.   Lovibond SH, Lovibond PF. Depression Anxiety Stress Scales. 2011.
33.   Manavipour D. Psychometric Properties of the Sleep Questionnaire. Shefaye Khatam.
2015;3(3):15-20.
34.   Ayatollahi P, Nafissi S, Eshraghian M, Tarazi A. Cross-Cultural Adaptation Of The Multiple
Sclerosis Impact Scale (MSIS-29) For Iranian MS Patients, Evaluation Of Reliability And Validity.
Tehran Univ Med J. 2006;64(1):62-8.
35.   Karimi S, Andayeshgar B, Khatony A. Prevalence of anxiety, depression, and stress in
patients with multiple sclerosis in Kermanshah-Iran: a cross-sectional study. BMC Psychiatry.
2020;20:166.
36.   Roque DT, Teixeira RAA, Zachi EC, Ventura DF. The use of the Cambridge
Neuropsychological Test Automated Battery (CANTAB) in neuropsychological assessment:
application in Brazilian research with control children and adults with neurological disorders.
Psychol Neurosci. 2011;4:255-65.
37.   Moayedi M, Salomons TV, Dunlop KA, Downar J, Davis KD. Connectivity-based parcellation
of the human frontal polar cortex. Brain Struct Funct. 2015;220(5):2603-16.
38.   Kringelbach ML. The human orbitofrontal cortex: linking reward to hedonic experience.
Nature Reviews Neuroscience. 2005;6(9):691-702.

39. Gandiga PC, Hummel FC, Cohen LG. Transcranial DC stimulation (tDCS): A tool for double-
blind sham-controlled clinical studies in brain stimulation. Clinical Neurophysiology.
2006;117(4):845-50.
40. Salehinejad MA, Wischnewski M, Ghanavati E, Mosayebi-Samani M, Kuo M-F, Nitsche MA.
Cognitive functions and underlying parameters of human brain physiology are associated with
chronotype. Nature Communications. 2021;12(1):4672.
41. Salehinejad MA, Ghanavati E, Reinders J, Hengstler JG, Kuo M-F, Nitsche MA. Sleep-
dependent upscaled excitability, saturated neuroplasticity, and modulated cognition in the human
brain. eLife. 2022;11:e69308.
42. Kuo M-F, Nitsche MA. Effects of Transcranial Electrical Stimulation on Cognition. Clin EEG
Neurosci. 2012;43(3):192-9.
43. León Ruiz M, Sospedra M, Arce Arce S, Tejeiro-Martínez J, Benito-León J. Current evidence
on the potential therapeutic applications of transcranial magnetic stimulation in multiple sclerosis:
a systematic review of the literature. Neurología (English Edition). 2022;37(3):199-215.
44. Cirillo G, Di Pino G, Capone F, Ranieri F, Florio L, Todisco V, et al. Neurobiological after-
effects of non-invasive brain stimulation. Brain Stimulation. 2017;10(1):1-18.
45. Zuo C, Cao H, Feng F, Li G, Huang Y, Zhu L, et al. Repetitive transcranial magnetic
stimulation exerts anti-inflammatory effects via modulating glial activation in mice with chronic
unpredictable mild stress-induced depression. Int Immunopharmacol. 2022;109:108788.
46. Sasso V, Bisicchia E, Latini L, Ghiglieri V, Cacace F, Carola V, et al. Repetitive transcranial
magnetic stimulation reduces remote apoptotic cell death and inflammation after focal brain injury.
J Neuroinflammation. 2016;13(1):150.
47. Nikolin S, Moffa A, Razza L, Martin D, Brunoni A, Palm U, et al. Time-course of the tDCS
antidepressant effect: An individual participant data meta-analysis. Progress in Neuro-
Psychopharmacology and Biological Psychiatry. 2023;125:110752.
48. Jafari E, Alizadehgoradel J, Pourmohseni Koluri F, Nikoozadehkordmirza E, Refahi M,
Taherifard M, et al. Intensified electrical stimulation targeting lateral and medial prefrontal cortices
for the treatment of social anxiety disorder: A randomized, double-blind, parallel-group, dose-
comparison study. Brain Stimulation. 2021;14(4):974-86.
49. Alizadehgoradel J, Molaei B, Barzegar Jalali K, Pouresmali A, Sharifi K, Hallajian A-H, et al.
Targeting the prefrontal-supplementary motor network in obsessive-compulsive disorder with
intensified electrical stimulation in two dosages: a randomized, controlled trial. Translational
Psychiatry. 2024;14(1):78.
50. Sá KN, Baptista RF, Shirahige L, Razza LB, Nogueira M, Coura MHF, et al. Evidence-based
umbrella review of non-invasive brain stimulation in anxiety disorders. The European Journal of
Psychiatry. 2023;37(3):167-81.
51. Nakamura-Palacios EM, Lopes IB, Souza RA, Klauss J, Batista EK, Conti CL, et al. Ventral
medial prefrontal cortex (vmPFC) as a target of the dorsolateral prefrontal modulation by
transcranial direct current stimulation (tDCS) in drug addiction. J Neural Transm (Vienna).
2016;123(10):1179-94.
52. Alizadehgoradel J, Nejati V, Sadeghi Movahed F, Imani S, Taherifard M, Mosayebi-Samani M,
et al. Repeated stimulation of the dorsolateral-prefrontal cortex improves executive dysfunctions
and craving in drug addiction: A randomized, double-blind, parallel-group study. Brain Stimulation.
2020;13(3):582-93.
53. Flöel A. tDCS-enhanced motor and cognitive function in neurological diseases.
Neuroimage. 2014;85:934-47.
54. Begemann MJ, Brand BA, Ćurčić-Blake B, Aleman A, Sommer IE. Efficacy of non-invasive
brain stimulation on cognitive functioning in brain disorders: a meta-analysis. Psychological
Medicine. 2020;50(15):2465-86.

55.   Salehinejad MA, Ghanavati E, Glinski B, Hallajian A-H, Azarkolah A. A systematic review of
randomized controlled trials on efficacy and safety of transcranial direct current stimulation in
major neurodevelopmental disorders: ADHD, autism, and dyslexia. Brain and Behavior.
2022;n/a(n/a):e2724.
56.   Zemestani M, Hoseinpanahi O, Salehinejad MA, Nitsche MA. The impact of prefrontal
transcranial direct current stimulation (tDCS) on theory of mind, emotion regulation and emotional-
behavioral functions in children with autism disorder: A randomized, sham-controlled, and parallel-
group study. Autism Res. 2022;n/a(n/a).
57.   Salehinejad MA, Ghayerin E, Nejati V, Yavari F, Nitsche MA. Domain-specific Involvement of
the Right Posterior Parietal Cortex in Attention Network and Attentional Control of ADHD: A
Randomized, Cross-over, Sham-controlled tDCS Study. Neuroscience. 2020;444:149-59.
58.   Azarkolah A, Noorbala AA, Ansari S, Hallajian A-H, Salehinejad MA. Efficacy of Transcranial
Direct Current Stimulation on Pain Level and Disability of Patients with Fibromyalgia: A Systematic
Review of Randomized Controlled Trials with Parallel-Group Design. Brain Sciences. 2024;14(1):26.
59.   Jagadish A, Shankaranarayana AM, Natarajan M, Solomon JM. Transcranial direct current
stimulation for fatigue in neurological conditions: A systematic scoping review. Physiother Res Int.
2024;29(1):e2054.
60.   Salehinejad MA, Ghanavati E, Rashid MHA, Nitsche MA. Hot and cold executive functions in
the brain: A prefrontal-cingular network. Brain and Neuroscience Advances.
2021;5:23982128211007769.
61.   Narmashiri A, Akbari F. The Effects of Transcranial Direct Current Stimulation (tDCS) on the
Cognitive Functions: A Systematic Review and Meta-analysis. Neuropsychol Rev. 2023.
62.   Dedoncker J, Brunoni AR, Baeken C, Vanderhasselt M-A. A systematic review and meta-
analysis of the effects of transcranial direct current stimulation (tDCS) over the dorsolateral
prefrontal cortex in healthy and neuropsychiatric samples: influence of stimulation parameters.
Brain stimulation. 2016;9(4):501-17.
63.   Herrero Babiloni A, Bellemare A, Beetz G, Vinet S-A, Martel MO, Lavigne GJ, et al. The effects
of non-invasive brain stimulation on sleep disturbances among different neurological and
neuropsychiatric conditions: A systematic review. Sleep Medicine Reviews. 2021;55:101381.
64.   Salehinejad MA, Ghanavai E, Rostami R, Nejati V. Cognitive control dysfunction in emotion
dysregulation and psychopathology of major depression (MD): Evidence from transcranial brain
stimulation of the dorsolateral prefrontal cortex (DLPFC). Journal of Affective Disorders.
2017;210:241-8.
65.   Nejati V, Majdi R, Salehinejad MA, Nitsche MA. The role of dorsolateral and ventromedial
prefrontal cortex in the processing of emotional dimensions. Sci Rep. 2021;11(1):1971.
66.   Meyer-Arndt L, Kuchling J, Brasanac J, Hermann A, Asseyer S, Bellmann-Strobl J, et al.
Prefrontal-amygdala emotion regulation and depression in multiple sclerosis. Brain
Communications. 2022;4(3).
67.   Vicario CM, Salehinejad MA, Felmingham K, Martino G, Nitsche MA. A systematic review on
the therapeutic effectiveness of non-invasive brain stimulation for the treatment of anxiety
disorders. Neuroscience & Biobehavioral Reviews. 2019;96:219-31.
68.   Polanía R, Paulus W, Nitsche MA. Modulating cortico-striatal and thalamo-cortical
functional connectivity with transcranial direct current stimulation. Hum Brain Mapp.
2012;33(10):2499-508.
69.   Regenold WT, Deng Z-D, Lisanby SH. Noninvasive neuromodulation of the prefrontal cortex
in mental health disorders. Neuropsychopharmacology. 2022;47(1):361-72.
70.   Begemann MJ, Brand BA, Ćurčić-Blake B, Aleman A, Sommer IE. Efficacy of non-invasive
brain stimulation on cognitive functioning in brain disorders: a meta-analysis. Psychological
Medicine. 2020:1-22.

 Tables

Table 1. Demographic data

	Active tDCS	**Sham tDCS**	*p-value**
Sample size (n)	19	18	
Age- M(SD)	36.89(1.47)	37.7(1.45)	0.691
Sex – Male (female)	4(15)	6(12)	0.401
Marital Status – Single (married)	9(10)	7(11)	0.603
OAID yes (No)	4(15)	1(17)	0.340
Family history- yes (No)	9(10)	6(12)	0.508
Duration of disease			0.078
1-10 years	14	7	
11-20 years	4	10	
More than 20 years	1	1	
Type of MS			0.208
CIS	3	5	
RR	14	13	
SR	2	0	
Type of medications IS (IM)	10(7)	12(5)	0.663

655 *Note*: tDCS = transcranial Direct Current Stimulation; M = Mean; SD = Standard Deviation; OAID = other
656 .autoimmune. disease; CIS= clinically isolated syndrome; RR=relapsing-remitting multiple sclerosis; SP=secondary
657 progressive multiple sclerosis; IS= immunosuppressant drug; IM=immunomodulatory drug; * = between-group
658 differences in demographic variables were explored by Chi-square tests or Fisher's exact test for categorical variables
659 and F-tests for continuous variables.

Table 2: Means and SDs of reported tDCS side effects.

		active tDCS	Sham	t	df	p-value*
		M(SD)	M(SD)			
	Skin redness	0.32(1.003)	0.11(0.32)	0.844	21.888	0.408
	sleep	0.32(1.15)	0.0(0.0)	1.189	18	0.25
	fatigue	0.26(1.14)	0.11(0.47)	0.532	24.176	0.599
Reported side effect	pain	0.21(0.53)	0.17(0.38)	0.288	32.642	0.775
	burning	1.32(1.63)	0.44(.705)	2.124	24.747	**0.044**
	itching	0.84(1.25)	0.56(0.85)	0.813	31.822	0.422
	tingling	1.05(1.35)	1(0.907)	0.14	31.608	0.89

661 Values are presented as means ± standard deviation (SD). Note: each value represents the average of side effects
662 reported during all 10 tDCS sessions. tDCS = transcranial Direct Current Stimulation; M = Mean; SD = Standard
663 Deviation; * = between-group differences of side effect items were explored by independent samples t-tests,
664 Significant results are highlighted ($p \leq 0.05$) in **bold**

Table 3: Means and SDs of outcome variables before and after the tDCS.

Measure	Outcome variable	Time	active tDCS M (SD)	sham tDCS M (SD)	p-value
quality of life (MSIS)	Score	Pre-intervention	51.20 (19.19)	51.41 (13.94)	0.972
		Post-intervention	41.93 (11.74)	51.94 (16)	
DASS	Depression	Pre-intervention	3.73 (2.34)	6.11 (6.14)	0.168
		Post-intervention	2.06 (2.25)	6.76 (6.41)	
	Anxiety	Pre-intervention	3(1.73)	3.64(4.10)	0.575
		Post-intervention	1.60(1.88)	4(4.54)	
	Stress	Pre-intervention	6.73(2.81)	7.58(7.89)	0.557
		Post-intervention	4.53(3.64)	8.29(5.35)	
Sleep	total score	Pre-intervention	13.20(4.37)	11.29(3.78)	0.197
		Post-intervention	7.26(1.62)	11.58(3.90)	
	Insomnia	Pre-intervention	6.46(2.79)	5.17(1.55)	0.112
		Post-intervention	3.46(0.83)	5.35(1.65)	
	Oversleep	Pre-intervention	6.73(2.52)	6.11(2.71)	0.513
		Post-intervention	3.80(1.08)	6.23(2.88)	
RTI	5 choice movement time	Pre-intervention	1491.37(313.24)	1384.99(448.08)	0.406
		Post-intervention	825.54(294.49)	1294.09(404.63)	
	5 choice reaction time	Pre-intervention	967.30(415.51)	987.08(346.31)	0.876
		Post-intervention	779.37(392.55)	1037.36(338.37)	
	simple movement time	Pre-intervention	711.15(208.95)	626.94(226.50)	0.247
		Post-intervention	448.67(157.16)	638.68(191.55)	
	simple reaction time	Pre-intervention	457.59(101.29)	428.05(112.48)	0.406
		Post-intervention	273.69(86.51)	450.4(102.97)	
SWM	Strategy	Pre-intervention	31.947(5.83)	32.444(5.90)	0.798
		Post-intervention	30.526(6.73)	32.5(4.91)	
	total error	Pre-intervention	28.84(23.59)	25.11(24.83)	0.642
		Post-intervention	15.47(12.23)	19.94(10.87)	
RVP	Hits	Pre-intervention	14.947(4.08)	15.277(4.05)	0.807
		Post-intervention	18.736(3.61)	15.111(3.99)	
	mean latency	Pre-intervention	837.90(88.39)	470.72(127.81)	
		Post-intervention	538.33(114.37)	470.903(142.56)	<0.001

Note: tDCS = transcranial Direct Current Stimulation; M = Mean; SD = Standard Deviation; MSIS= Multiple Sclerosis Impact Scale (quality of life); DASS= Depression, anxiety, stress scale; MSQ= Mini sleep questionnaire; RTI= Reaction time test; SWM= Spatial Working Memory; RVP= Visual Information Processing; *p* values refer to baseline measurement comparisons using ANOVA tests.

Measure	Outcome variable	Source	df	F	*p*-value	partial eta²
(MSIS) Quality of life	score	Time	1, 30	7.945	0.008	0.209
		Group	1, 30	0.949	0.338	0.031
		time×group	1, 30	9.987	0.004	0.250
DASS	score	Time	1, 30	6.13	0.019	0.170
		Group	1, 30	3.877	0.058	0.114
		Domain	2, 60	15.608	**<0.001**	0.342
		time×group	1, 30	23.505	**<0.001**	0.439
		time× domain	2, 60	0.243	0.785	0.008
		group× domain	2, 60	1.156	0.322	0.037
		time×group× domain	2, 60	1.137	0.328	0.037
Sleep Total score	score	Time	1, 30	30.598	**<0.001**	0.505
		Group	1, 30	1.065	0.31	0.034
		time×group	1, 30	37.314	**<0.001**	0.554
Insomnia	score	Time	1, 30	19.756	**<0.001**	0.397
		Group	1, 30	0.277	0.603	0.009
		time×group	1, 30	25.004	**<0.001**	0.455
Oversleep	score	Time	1, 30	21.861	**<0.001**	0.422
		Group	1, 30	1.265	0.270	0.04
		time×group	1, 30	25.667	**<0.001**	0.461
5 choice movement time	score	Time	1, 35	127.32	**<0.001**	0.784
		Group	1, 35	2.412	0.129	0.064
		time×group	1, 35	73.491	**<0.001**	0.677
5 choice reaction time	score	Time	1, 35	11.684	0.002	0.25
		Group	1, 35	1.299	0.262	0.036
		time×group	1, 35	34.992	**<0.001**	0.50
simple movement time	score	Time	1, 35	17.476	**<0.001**	0.333
		Group	1, 35	0.844	0.365	0.024
		time×group	1, 35	20.905	**<0.001**	0.374
simple reaction time	score	Time	1, 35	24.729	**<0.001**	0.414
		Group	1, 35	6.439	0.016	0.155
		time×group	1, 35	40.317	**<0.001**	0.535
SWM strategy	score	Time	1, 35	0.703	0.407	0.02
		Group	1, 35	0.493	0.487	0.014
		time×group	1, 35	0.823	0.371	0.023
SWM total error	score	Time	1, 35	9.081	0.005	0.206
		Group	1, 35	0.005	**<0.001**	0.946
		time×group	1, 35	1.778	0.191	0.048
RVP Hits	score	Time	1, 35	39.44	**<0.001**	0.530
		Group	1, 35	1.698	0.201	0.046
		time×group	1, 35	47.032	**<0.001**	0.573
RVP mean latency	score	Time	1, 35	291.414	**<0.001**	0.893
		Group	1, 35	32.178	**<0.001**	0.479
		time×group	1, 35	292.129	**<0.001**	0.893

Note: tDCS = transcranial Direct Current Stimulation; M = Mean; SD = Standard Deviation; MSIS= Multiple Sclerosis Impact
Scale(quality of life); DASS= Depression, anxiety, stress scale; MSQ= Mini sleep questionnaire(total sleep score, insomnia and
oversleep); RTI= Reaction time test(RTI Mean simple reaction time, RTI Mean five-choice reaction time, RTI Mean simple
movement time, and RTI Mean five choice movement time); SWM= Spatial Working Memory(SWM Strategy, SWM Total
errors, and SWM Between); RVP= Visual Information Processing (RVP Hits, RVP Miss, RVP mean latency); Significant results
are highlighted (*p* ≤ 0.05) in **bold**.

Figure legends

Figure 1 displays the CONSORT Flow Diagram illustrating the study's inclusion procedures. Thirty-seven patients successfully completed the post-intervention measurement for all CANTAB subtests.

Figure 2: Study Design: (A) This study was conducted as a randomized, double-blind trial. Participants were divided into two groups: active tDCS (n = 19) and sham tDCS (n = 18). Both groups underwent pre- and post-intervention assessments. The anodal electrode was positioned over the left dorsolateral prefrontal cortex (DLPFC), while the cathode was placed over the right frontopolar cortex (FPC). (B) 3D models were utilized to examine the flow of electrical current in the brain following the specified protocol. The MR image was segmented into six tissue types: gray matter (GM), white matter (WM), CSF, skull, scalp, and air cavities using SPM8 from the Welcome Trust Center for Neuroimaging, London, UK, with an enhanced tissue probability map. The segmented images were used to create a 3D model with Simpleware software version 5 from Synopsys, Mountain View, CA, incorporating the electrodes and saline-soaked sponges. The distribution of current flow within the brain was then computed using the finite element method in COMSOL Multiphysics software version 5.2 from COMSOL Inc., Burlington, MA. The electric fields were visualized for stimulation intensities of 2.0 mA with an F3 anodal–Fp2 cathodal montage. Please note: This model illustrates the current flow for 2 mA tDCS for illustrative purposes. The induced electric field of 1.5 mA differs from the 2 mA field.

Figure 3: The effect of prefrontal active and sham tDCS on mental health-related variables. The left panel displays average outcome measures before and after the intervention within each group (i.e., within-group comparisons). The right panels illustrate average outcome measures for each time point (i.e., pre-intervention or post-intervention) across groups (active vs sham). *Note*: tDCS = transcranial Direct Current Stimulation; DASS = Depression Anxiety Stress Scale; Floating asterisks [*] in the left panel represent a significant difference between pre-intervention measurements vs post-intervention measurements in all groups. Floating asterisks [*] in the right panel indicate a significant difference between active stimulation (1.5 mA) vs sham tDCS at each time point. ns non-significant. All pairwise comparisons were conducted using Fisher's LSD multiple comparisons tests. All error bars represent s.e.m.

Figure 4. The effect of the intervention on cognition. The left panel displays average outcome measures before and after the intervention within each group (i.e., within-group comparisons). The right panels illustrate average outcome measures for each time point (i.e., pre-intervention or post-intervention) across groups (active vs sham). *Note*: tDCS = transcranial Direct Current Stimulation; RTI = Reaction Time; SWM = Spatial Working Memory; RVP = Rapid Visual Processing; Floating asterisks [*] in the left panel represent a significant difference between pre-intervention measurements vs post-intervention measurements in all groups. Floating asterisks [*] in the right panel indicate a significant difference between active stimulation (1.5 mA) vs sham tDCS at each time point. ns non-significant. All pairwise comparisons were conducted using Fisher's LSD multiple comparisons tests. All error bars represent s.e.m.

Figure 1.

Figure 2

Figure 3.

RTI

Figure 4.