Synchronous remote-based interventions for suicidal behaviour prevention: a systematic review and meta-analyses of clinical trials

Laura Comendador[1,2], María P. Jiménez-Villamizar[3], Josep-Maria Losilla[4], Juan P. Sanabria-Mazo[3,5,6], Corel Mateo-Canedo[3], Antoni Sanz[3,7], Ana-Isabel Cebrià[2,8,9], Diego Palao[1,2,8]

[1]Department of Psychiatry and Forensic Medicine, Faculty of Medicine, Universitat Autònoma de Barcelona. 08193 Cerdanyola del Vallès, Spain.

[2]Department of Mental Health, Parc Taulí University Hospital. Institut d'Investigació i Innovació Parc Taulí (I3PT-CERCA). Universitat Autònoma de Barcelona. 08208 Sabadell, Spain.

[3]Department of Basic, Developmental, and Educational Psychology, Faculty of Psychology, Universitat Autònoma de Barcelona. 08193 Cerdanyola del Vallès, Spain.

[4]Department of Psychobiology and Methodology of Health Sciences, Faculty of Psychology, Universitat Autònoma de Barcelona. 08193 Cerdanyola del Vallès, Spain.

[5]Teaching, Research & Innovation Unit, Parc Sanitari Sant Joan de Déu. 08830 Sant Boi de Llobregat, Spain.

[6]Centre for Biomedical Research in Epidemiology and Public Health (CIBERESP). 28029 Madrid, Spain

[7]Stress and Health Research Group (GIES). Universitat Autònoma de Barcelona. 08193 Cerdanyola del Vallès, Spain.

[8]Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Instituto de Salud Carlos III. 28029 Madrid, Spain.

[9]Department of Clinical and Health Psychology, Faculty of Psychology, Universitat Autònoma de Barcelona. 08193 Cerdanyola del Vallès, Spain.

Corresponding author
Antoni Sanz, PhD
Department of Basics, Developmental, and Educational Psychology, Faculty of Psychology, Universitat Autònoma de Barcelona.
Carrer de la Fortuna, s/n. Campus de Bellaterra, Universitat Autònoma de Barcelona. 08193 Cerdanyola del Vallès (Spain).
E-mail: antonio.sanz@uab.cat

Author's Note
Laura Comendador Vázquez, MSc, PhD student. E-mail: laura.comendador@autonoma.cat
ORCID 0000-0002-5221-4794
María P. Jiménez-Villamizar, MSc, PhD student. E-mail: mariapaola.jimenez@autonoma.cat
ORCID 0000-0003-2264-7422
Josep-Maria Losilla, PhD. E-mail: josepmaria.losilla@uab.cat
ORCID 0000-0002-5140-5847
Juan P. Sanabria-Mazo, PhD. E-mail: juanpablo.sanabria@sjd.es
ORCID 0000-0003-1688-435X
Corel Mateo-Canedo, PhD. E-mail: corel.mateo@uab.cat
ORCID 0000-0002-0620-9257
Antoni Sanz Ruíz, PhD. E-mail: antonio.sanz@uab.cat
ORCID 0000-0002-7952-4477
Ana Isabel Cebrià Meca, PhD. E-mail: acebria@tauli.cat
ORCID 0000-0002-2632-8130
Diego J. Palao Vidal, MD, PhD. E-mail: dpalao@tauli.cat
ORCID 0000-0002-3323-6568

Word count (excluding title page, abstract, tables, acknowledgements, contributions, and references): 7271 words.
Number of tables: 2.
Number of figures: 5.

Synchronous remote-based interventions for suicidal behaviour prevention: a

systematic review and meta-analyses of clinical trials

ABSTRACT

Objectives Suicide is a leading cause of preventable death worldwide. Evidence supports the

impact of providing active contact for individuals who have attempted suicide. The current

systematic review and meta-analyses aim to investigate the effects of suicide prevention

strategies implemented through remote and synchronous technology-based interventions.

Design Systematic review, narrative synthesis, and meta-analysis.

Data sources Electronic databases (PubMed, PsycInfo, Scopus, and Web of Science) and grey

literature sources (ClinicalTrials.gov and Google Scholar) were searched until December 2024.

Eligibility criteria Eligible articles assessed suicide prevention interventions for participants over

12 years with prior suicidal behaviour. Eligible study design included randomised control trials

and non-randomised clinical trials published in English or Spanish.

Data extraction and synthesis Screening, selection process, data extraction, and risk of bias

assessment were performed independently by two reviewers. Data on suicide-related factors

and adherence to treatment were extracted. Meta-analyses were conducted to determine

effect sizes (Hedges' g) for suicidal ideation, risk ratios (RR) for suicide attempts, and Peto odds

ratios (OR) for suicide. Heterogeneity was assessed using the Cochrane's Q test, tau^2 statistic,

and I^2 value. Publication bias was investigated employing funnel plots and Egger's test.

Results A total of 28 studies, comprising 10,015 participants in the intervention group and

10,726 in the comparison group, were included in the systematic review and meta-analyses.

Synchronous remote-based interventions were effective in preventing repeated suicide

attempts at 1 month (RR 0.73, 95% CI 0.62 to 0.85, I^2=0.0%, Q=0.70, tau^2=0.00), 6 months (RR

0.56, 95% CI 0.34 to 0.95, I^2=85.4%, Q=54.92, tau^2=0.36), and 12 months (RR 0.68, 95% CI 0.49

to 0.96, I^2=87.6%, Q=72.63, tau^2=0.27). Additionally, these interventions were associated with a

reduction in suicide-related deaths at 18 months (Peto OR 0.18, 95% CI 0.08 to 0.44, I^2=0.0%,

Q=0.03, tau^2=0.00). Effects on suicidal ideation were not statistically significant at any time-point

(Hedges' *g* -0.07 to -0.28, I^2=0.0 to 69.3%, Q=1.16 to 7.38, tau^2=0.00 to 0.14).

Conclusions Synchronous remote-based interventions demonstrate a potential benefit in

preventing suicide attempts and deaths by suicide, and may serve as an adjunct to usual

treatment; however, the effect on suicidal ideation appears limited. The observed heterogeneity

warrants caution when interpreting these findings. Future research should prioritise

methodological enhancements to improve the quality and consistency of evidence, as well as

investigate the mediating processes underlying their effectiveness in reducing suicidal

behaviour.

PROSPERO registration number CRD42021275044.

Keywords Suicide, Telemedicine, Preventive Medicine.

STRENGTHS AND LIMITATIONS OF THIS STUDY

●   The protocol was registered in PROSPERO and published in a peer-reviewed journal,

41   ensuring methodological transparency.

42   ●   Data extraction, synthesis, and quality assessment are reported in accordance with the

43   Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA-2020)

44   guidelines.

45   ●   The systematic review and meta-analyses used a comprehensive search strategy to

46   avoid missing published articles.

47   ●   Heterogeneity in the methods makes it difficult to draw definitive conclusions about the

48   effectiveness of interventions.

INTRODUCTION

The World Health Organisation (WHO) estimates that over 703,000 people die annually from

suicide worldwide [1, 2]. Suicide occurs across the lifespan; although, in 2019, it was the fourth

leading cause of death among individuals aged 15 – 29 years worldwide [2, 3]. For each death

by suicide, there are twenty suicide attempts [4], which is one of the major disease burden

causes in the world [5, 6, 7]. Moreover, a history of suicide attempts is considered a robust

predictor of completed suicide, supporting prevention efforts during the acute period following

an episode of suicidal behaviour [8–12].

Increasing evidence suggests that post-discharge follow-up contacts can be an effective

suicide prevention strategy [13–19]. The eHealth-based treatments could be classified into two

categories: (1) real-time synchronous therapy – such as telephone contact or telepsychiatry via

videoconferencing – (2) or digital self-help programmes without human guidance, often

asynchronous [20, 21]. Synchronous communication technologies, including telemedicine and

real-time interactions, have the potential to provide immediate and flexible access to support

during critical periods. As opposed to asynchronous methods, these approaches facilitate direct

and immediate engagement between patients and healthcare professionals, potentially

enhancing the effectiveness of suicide prevention efforts. Accordingly, most post-discharge

suicide prevention programmes reinforce early contact with mental health professionals. An

essential component of the effectiveness of suicide prevention interventions is treatment

adherence [22]. The evaluation of therapeutic adherence in clinical care is critically important

since poor adherence results in adverse outcomes, such as relapses and hospitalisations, among

people with mental health disorders worldwide [22, 23, 24]. Brief contact interventions have

three rationales: firstly, they allow for repeated suicide risk assessments; secondly, they reduce

feelings of social isolation and provide emotional support; and thirdly, they improve treatment

adherence and reduce dropout from mental health care [25–28].

In recent decades, numerous studies have reported promising results from the

implementation of follow-up contact interventions for at-risk individuals as a suicide prevention

strategy [29, 30]. Although previous systematic reviews have addressed this topic [31, 32, 33],

evidence on the specific impact of synchronous technologies remains limited. In a scoping

review, Shin *et al.* [34] identified several trends in information and communication technology

(ICT)-based interventions for suicide prevention applied in clinical settings, such as

computerised interventions or applications, highlighting the need for future research to

strengthen the evidence to improve implementation. This review included 66 ICT-based

interventions, targeted at post-discharge follow-up, screening, and safety planning, for any

spectrum of suicide-related thoughts and behaviours.

Along these lines, Luxton *et al.* [35] suggest that repeated follow-up contacts may help

reduce suicidal behaviour, reporting in their systematic review four studies that showed mixed

results with trends towards a preventive effect and two studies that showed no preventive

effect. A more recent systematic review and meta-analysis [36] examined 16 studies evaluating

the effectiveness of telehealth interventions in suicide prevention, reporting that telepsychiatry

is effective in reducing suicide and reattempt rates, with high acceptance and retention rates.

Nevertheless, this review did not specifically evaluate suicidal ideation, and moreover, the

authors emphasise the need for further research to fully elucidate the potential of telepsychiatry

in suicide prevention.

These interventions' distinctive and rapidly evolving nature necessitates a thorough

exploration of their implications. Our research intends to expand on previous evidence by

updating it with a larger number of studies, a comprehensive analysis of the components of

suicidal behaviour, and a more detailed examination of the temporal dynamics of the effects.

The current systematic review and meta-analyses aim to explore the evidence on remote suicide

prevention strategies implemented through synchronous technology-based interventions (i.e.,

via remote, interactive, real-time communication), considering the effects, barriers, and

facilitators to the implementation.

METHODS

Protocol and registration

The protocol of the systematic review and meta-analyses was registered in the Prospective

International Registry of Systematic Reviews (PROSPERO), with the identification number

CRD42021275044. The review was reported according to the Preferred Reporting Items for

Systematic Reviews and Meta-Analyses (PRISMA) guidelines [37] (see Supplemental File 1: Table

S1). A brief mention of the amendments made to the original protocol article [38] can be found

in the Supplemental File 2.

Eligibility criteria

The main inclusion and exclusion criteria are described below. Full details on the eligibility

criteria can be found in the protocol article [38].

Participants

Adolescents over 12 years of age and adults with reported suicidal ideation or prior suicide

attempts. Individuals with non-suicidal self-injury were excluded.

Interventions

Secondary suicide prevention interventions delivered via synchronous distance communication

technologies– i.e., telephone contact, 24-hour telephone hotlines, videoconferencing therapy,

immediate text messaging, etc–. Interventions based on asynchronous telecommunication

devices were excluded.

Comparisons

The eligible control groups include treatment as usual (e.g., visits to mental health services or

pharmacotherapy), enhanced usual care, no treatment, historical control, or geographical

control. Non-reporting of a control group was a reason for exclusion. Studies were excluded if

both, the intervention and control groups, received synchronous and remote interventions.

Outcomes

Suicidal behaviours– i.e., suicidal ideation, suicide attempts, and suicide deaths– were included.

Suicidal ideation refers to thoughts about the possibility of ending one's life [39], assessed using

validated instruments such as the Columbia Suicide Severity Rating Scale (CSSRS). A suicide

attempt is defined as a self-inflicted, potentially injurious act with a non-fatal outcome and

evidence of intent to die [3], measured by the number of reported attempts within a defined

follow-up period. Finally, suicide death involves a self-inflicted behaviour performed with the

deliberate intention of resulting in death [40], and may be quantified by the number of persons

who die by suicide. Outcomes could be assessed at any point following the intervention, with

no restriction on the length of follow-up, provided the assessment uses a quantitative approach.

Adherence to treatment was also extracted as a secondary outcome.

Study design

Randomised controlled trials (RCTs) and non-randomised clinical trials (nRCTs) were eligible for

inclusion. The term nRCTs refers to studies with a defined intervention and comparison group,

in the absence of random allocation. Observational studies uncontrolled and non-experimental

were excluded (i.e., single-arm studies, interrupted time series, case reports, descriptive non-

comparative studies) [41, 42].

Data collection and analysis

Information sources

Bibliographic searches were conducted in PubMed, PsycInfo, Scopus, and Web of Science for studies published before December 2024. ClinicalTrials.gov and Google Scholar were searched for grey literature and unpublished records. Reference lists of previous systematic reviews on the subject and references cited in eligible articles were consulted for data collection.

Search strategy

Details of the search process applied to databases, registers, and websites can be found in Supplemental File 3. The search was unrestricted by year of publication, limited to English or Spanish, and developed adhering to the Peer Review of Electronic Search Strategies (PRESS) [43].

Study selection

The results of the literature search were imported into Mendeley (version 1.19.8), and Rayyan Systems Inc [44]. Duplicate articles were removed automatically by Rayyan Systems Inc. and manually by the first reviewer (LC). After, two authors (LC and MPJ) blind-screened all articles based on titles, abstracts, and keywords. Then, two reviewers (LC and MPJ) independently evaluated the full-text articles according to the eligibility criteria. Discrepancies were resolved through discussion with a third author (AS) until reaching consensus.

Data collection

Two authors (LC and MPJ) performed the data extraction independently, applying an *ad hoc* extraction form in Microsoft Excel (version 16.65).

Data items

Data were extracted on participants and intervention characteristics, along with outcomes

related to suicidal behaviour for both intervention and comparison groups– for all measures and

time points–. In addition, therapeutic adherence– specifically, the moderating effect of

adherence on the effectiveness of the intervention, the description of measures to maximise

adherence to treatment, and predictor variables of adherence– were examined.

Risk of bias

Two reviewers (LC and MPJ) performed the risk of bias (RoB) assessment independently, using

the Revised Cochrane risk-of-bias tool for randomised trials (RoB 2) [45] and Risk-of-bias In Non-

randomised Studies of Interventions (ROBINS-I) [46] for the outcome of suicidal behaviours.

Disagreements were resolved by consensus with a third blinded reviewer (AS).

Effect measures

Systematically examined all types of effect measures used in the selected trials (e.g., risk ratio,

odds ratio, mean difference, frequencies) for the effects of each intervention group on suicidal

ideation, repeated suicide attempts, and death by suicide.

Synthesis methods

Firstly, a systematic review was employed to collect and synthesise the empirical evidence that

met the eligibility criteria. Secondly, meta-analyses were conducted using a random-effects

model. Due to the wide variety of study types included in the systematic review, a random-

effects model was selected *a priori* for all analyses, and heterogeneity statistics were reported

using this model, in accordance with the recommendations from the Cochrane Handbook for

Systematic Reviews of Interventions (version 6.3) [47]. To ensure that the combined numerical

data from the different studies is comparable and can be adequately synthesised, analyses by

sorting studies with the same evaluation measures time frame were performed for each

outcome, and a narrative synthesis approach was adopted when a quantitative combination of

the studies was not feasible.

Three types of meta-analyses were performed. (1) A random-effects model with

Hedges' g [48] and 95% confidence intervals (CI) were calculated for suicidal ideation. Effect

sizes are considered small ($g \geq 0.2$), medium ($g \geq 0.5$), or large ($g \geq 0.8$) [48]. (2) A random-effects

model with the effect estimates weighted by the inverse of the variance was calculated for

suicide attempts. As proposed by Daly [49] and Zhang & Yu [50], the risk ratio (RR) and 95% CI

were analysed. Finally, (3) a random-effects model with the effect estimates weighted by the

Peto $et\ al.$ method [51] was calculated for suicide deaths since the observed frequencies are

very low. The Cochrane's Q test and tau^2 statistic were employed to identify heterogeneity

among studies. The proportion of variability due to heterogeneity was indicated by I^2 value.

Heterogeneity was considered moderate if I^2 was greater than 30%. All analyses were performed

using the Meta-Essentials tool for Microsoft Excel (version 16.65) [52], applying the Knapp-

Hartung adjustment by default in fitting random effects models. The trials with multiple

assessment measurement periods were included as separate studies in the appropriate meta-

analyses [47].

Certainty of evidence

The certainty of evidence was assessed using the Grading of Recommendations Assessment,

Development, and Evaluation (https://gdt.gradepro.org/) [53]. The GRADE guidelines establish

standardised and transparent criteria for grading the certainty of evidence [54]. In classifying

the level of evidence quality, GRADE defines four categories: high, moderate, low, and very low

[55]. The main domains used are risk of bias, inconsistency, indirect evidence, imprecision, and

publication bias [55]. Two reviewers (LC and MPJ) independently evaluated the certainty of the

evidence of the included studies; any discrepancies were resolved through consultation with a

third reviewer (AS). The assessment was carried out for each outcome of suicidal behaviours.

Summary tables of the findings were presented to synthesise the process.

Patient and public involvement

Not involved in the review protocol, design, reporting, or dissemination plans.

RESULTS

Study selection

The initial search retrieved 4930 studies. After removing duplicates, 2802 articles were screened

for title, abstract, and keywords. Of these, 131 studies were selected for full-text review. Finally,

28 publications were included in the systematic review and meta-analyses [56–83]. The article

selection process was described in a PRISMA flowchart [84] (Figure 1). Studies that might appear

to meet the inclusion criteria but were excluded and the reasons for exclusion are presented in

Supplemental File 4: Table S2. Inter-rater agreement was calculated for initial screening using

Delta (δ=0.91, 95% CI 0.88 to 0.94) [85], and for eligibility and extraction process using Cohen's

Kappa (κ=0.82, 95% CI 0.62 to 1.0; κ=0.98, 95% CI 0.97 to 0.99, respectively) [86].

243   [Please insert here Figure 1]

Figure 1 PRISMA flowchart for systematic review and meta-analyses including database

searches and other sources.

Study characteristics

A descriptive summary of the characteristics of the included studies is presented in Table 1 and

Supplemental File 5: Table S3. The current systematic review included 28 studies, of which 19

were RCT (67.9%) [56–58, 61–63, 66, 69, 71, 72, 74–77, 79–83]. The sample size ranges from 10

to 3068 in the intervention group (IG) and from 10 to 3694 in the control group (CG). The mean

age ranged from 23 to 45 years in the IG and from 23 to 47 years in the CG. The proportion of

women in most of the studies was above 50% in both IG and CG, except in three studies [63, 79,

82]. The eligible articles were from 12 countries: France [64, 65, 72, 74, 78, 89, 81], Spain [59,

60, 67, 68, 70], United States of America [63, 73, 79, 82], Iran [69, 71, 75, 76], Brazil, India, Sri

Lanka [57, 66], French Polynesia [56], Canada [58], Sweden [61], Australia [62], United Kingdom

[77], and China [83].

As for outcome variables, 8 studies reported outcomes relating to suicidal ideation [58,

61–63, 77, 79, 82, 83], 26 studies reported the proportion of participants who reattempted

suicide [56–61, 63–65, 67–83], and 13 reported the proportion of suicide deaths [56, 58, 60, 63,

66, 68–71, 73, 74, 80, 81]. There was high heterogeneity in the diagnostic methods used to

identify suicide behaviours: European Parasuicide Study Interview Schedule (EPSIS), Scale for

Suicide Ideation (SSI), General Health Questionnaire (GHQ-28), Columbia Suicide Severity Rating

Scale (CSSRS), Beck Scale for Suicidal Ideation (BSS), Suicide Attempt Self-Injury Count (SASI-

Count), Beck's Suicide Intent Scale (SIS), Multisite Intervention Study on Suicidal Behaviours

(SUPRE-MISS) questionnaire, forensic records of suicide deaths, data from electronic medical

records, etc. The most frequently reported psychiatric diagnoses were affective disorders (n=13;

46.4%) [56, 58, 61, 64, 65, 68, 70, 72, 73, 74, 77, 79, 81], followed by anxiety disorders (n=10;

35.7%) [56, 58, 61, 64, 65, 72, 73, 77, 79, 81], personality disorders (n=5; 17.9%) [56, 58, 65, 68,

74], substance use disorders (n=5; 17.9%) [56, 58, 64, 72, 77], trauma-related disorders (n=5;

17.9%) [58, 64, 74, 79, 81], psychotic disorders (n=4; 14.3%) [56, 65, 68, 73], and eating disorders

(n=3; 10.7%) [65, 72, 81] according to the Diagnostic and Statistical Manual of Mental Disorders

(DSM-III-R, DSM-IV-TR) or the International Classification of Diseases (ICD-10).

The interventions were categorised as (a) telephone follow-up intervention (*n*=16;

57.1%) [59–61, 64, 65, 67, 68, 70–72, 75, 76, 78, 80, 81, 83], (b) Brief Intervention and Contact

(BIC; *n*=5; 17.9%) [56, 57, 66, 69, 79], (c) web-based therapist-delivered Eye Movement

Desensitization and Reprocessing (EMDR; *n*=1; 3.6%) [58], (d) 24-hour call-in services (*n*=1; 3.6%)

[62], (e) instant text messages (*n*=1; 3.6%) [63], (f) Emergency Department Safety Assessment

and Follow-up Evaluation (ED-SAFE intervention; *n*=1; 3.6%) [73], (g) postal cards, 24-hour

telephone, and telephone follow-up (*n*=1; 3.6%) [74], (h) Safety Planning and telephone follow-

up intervention (SAFETEL intervention; *n*=1; 3.6%) [77], and (i) Dialectical Behaviour Therapy

Skills-Based Intervention (DBT-BSI) and telephone follow-up (*n*=1; 3.6%) [82]. On the other hand,

the control group in most studies was TAU (*n*=26; 92.9%) [56–75, 77–81, 83], except for two

studies, which was a face-to-face follow-up [76] and a relaxation training condition [82].

In relation to setting, the included studies were conducted in emergency departments

(*n*=19; 67.9%) [56, 57, 59–61, 65, 66, 69–76, 78, 80, 81, 83], mental health services (*n*=4; 14.3%)

[58, 64, 68, 82], general hospitals (*n*=2; 7.1%) [67, 77], military installations (*n*=2; 7.1%) [63, 79],

or 24-hour call-in services (*n*=1; 3.6%) [62]. Interventions were performed by nurses (*n*=7; 25%)

[59–61, 65, 67, 68, 70], psychologists (*n*=7; 25%) [63, 64, 71, 72, 77, 79, 81], psychiatrists (*n*=6;

21.4%) [58, 74–76, 78, 80], physicians (*n*=1; 3.6%) [73], and by a combination of health

professionals– i.e., medical doctor, psychiatric nurse, psychologist, or social worker (*n*=4; 14.3%)

[56, 57, 66, 69]– although, few interventions were undertaken by non-health professionals (*n*=2;

7.1%) [62, 83] or not reported (*n*=1; 3.6%) [82]. The onset of the intervention– i.e., initial

contact– could be categorised in a time interval of days (*n*=5; 17.9%) [58, 63, 64, 74, 77], weeks

(*n*=19; 67.9%) [56, 57, 59, 60, 62, 65–72, 75, 76, 78, 81–83], months (*n*=2; 7.1%) [61, 80], or not

reported (*n*=2; 7.1%) [73, 79]. The total number of contacts ranged from one [64, 72, 80, 81] to

14 [71], with a minimum duration of five minutes [59, 60, 66] and a maximum of 90 minutes

[58]. In addition, the follow-up period ranged from two months [65] to five years [60], with 12

months being the most prevalent period range (*n*=10; 35.7%) [59, 61–63, 67, 68, 70, 71, 73, 74].

Table 1

Overview of participants, interventions, comparisons, outcomes, and study design (PICOS) for

studies evaluating suicide ideation, attempted suicide, and suicide death.

Table 1. Overview of participants, interventions, comparisons, outcomes, and study design (PICOS) for studies evaluating suicide ideation, attempted suicide, and suicide death

Study (authors, date of publication, country)	Study design	Setting, sample size (N)	Details of participants	Intervention (n) and comparison (n)	Follow-up	Outcome measure (Instrument)
Amadéo et al., 2015 [56]. French Polynesia	RCT	Psychiatric ED N=190	Patients with suicidal behaviour; mean age IG 33, CG 31.48; 64.2% women	BIC (n=90) vs. TAU (n=100). 1h psycho-educational session combined with telephone contacts at 1, 2, 4, 7, 11 weeks, 4, 6, 12, and 18 months by psychologists and psychiatrists.	18 months	Suicide attempt (EPSIS) [87], Suicide death (Forensic records of suicide deaths)
Bertolote et al., 2010 [57]. Brazil, India, Sri Lanka, Iran, China	RCT	ED N=1867	Patients with suicide attempt; mean age IG 26.3, CG 25.5; 58.2% women	BIC (n=922) vs. TAU (n=945). 1h brief educational intervention combined with telephone contacts or visits at 1, 2, 4, 7, 11 weeks, 4, 6, 12, 18 months by medical doctor, psychiatric nurse, psychologist, or social worker.	18 months	Suicide attempt (EPSIS) [87]
Burback et al., 2024 [58]. Canada	RCT	Mental health services. N=42	Patients with suicidal ideation in the past week; age IG 36.0, CG 36.0; 79.0% women	Web-based therapist-delivered EMDR (n=20) vs. TAU (n=22). 12 sessions of web-based EMDR (90 min.) delivered twice weekly via videoconferencing by psychiatrists.	4 months	Suicidal ideation (BSS) [88], Suicide attempt (CSSRS) [89], Suicide death (CSSRS) [89]
Cebrià et al., 2013 [59]. Spain	nRCT; case-control	2 Hospital ED N=514	Patients discharged from ED after a suicide attempt; age IG 41.92, CG 40.73; 66.5% women	Telephone follow-up (n=296) vs. TAU (n=218). Telephone contacts (5-10 min.) at 1 week, 1, 3, 6, 9, and 12 months by a psychiatric nurse.	12 months	Suicide attempt (Interview + medical record)
Cebrià et al., 2015 [60]. Spain	nRCT; retrospective cohort	2 Hospital ED N=485	Patients discharged from ED after a suicide attempt; age IG 38.78, CG 41.92; 64.5% women	Telephone follow-up (n=287) vs. TAU (n=198). Telephone contacts (5-10 min.) at 1 week, 1, 3, 6, 9, and 12 months by a psychiatric nurse.	60 months	Suicide attempt, Suicide death (Interview + Medical records)
Cedereke et al., 2002 [61]. Sweden	RCT	Hospital, Emergency Unit N=216	Patients treated after a suicide attempt; mean age IG 40, CG 42; 66% women	Telephone follow-up (n=107) vs. TAU (n=109). Telephone contacts (20-25 min.) at 4 and 8 months after a suicide attempt, by a psychiatric nurse or social counsellor.	12 months	Suicidal ideation (SSI) [90], Suicide attempt (Semi-structured interview)
Christensen et al., 2013 [62]. Australia	RCT	Australia's Lifeline. N=155	Callers to a national 24h-helpline service with moderate-to-high psychological distress; mean age 41.49; 81.9% women	Web-based CBT (n=38) vs. Web-based CBT + telephone follow-up (n=45) vs. telephone follow-up (n=37) vs. TAU (n=35). CBT delivered across five modules. Telephone contacts (10 min.), weekly, by a Lifeline telephone counsellor.	12 months	Suicidal ideation (GHQ-28) [91]
Comtois et al., 2019 [63]. USA	RCT	3 military installations N=657	Soldiers and marines identified as being at risk of suicide; mean age IG 25.6, CG 24.8; 17.9% women	Caring contacts (n=329) vs. TAU (n=328). Text messages delivered on day 1, at week 1, at months 1, 2, 3, 4, 6, 8, 10, and 12, and on participants' birthdays, by mental health clinicians. Text message replies from participants were monitored 24 hours a day. Replies indicating suicide risk or distress were handled by a few text messages exchanged or a single telephone call.	12 months	Suicidal ideation (SSI) [90], Suicide attempt (Interview + SASI-Count) [92], Suicide death (Interview)
Demesmaeker et al., 2024 [64]. France	nRCT; prospective cohort	General medical or psychiatric services N=733	Patients discharged from the hospital after a suicide attempt (non-first-time attempters); mean age IG 40.2, CG 39.4; 61.5% women	Telephone contact (n=236) vs. Telephone contact and postcards (n=411) vs. Postcards (n=34) vs. TAU (n=52). Telephone contact between the 10th and 21st days after discharge for suicide attempts by psychologists. Personalised postcards are sent at months 2–5 to those unavailable for a phone call or facing a suicidal crisis.	6 months	Suicide attempt (Standardised telephone interview)
Exbrayat et al., 2017 [65]. France	nRCT; non-randomised controlled	Emergency Psychiatry Dept. N=827	Patients admitted to the ED for suicide attempt; mean age IG 40.2, CG 39.7; 69.5% women	Telephone follow-up (n=436) vs. TAU (n=387). Telephone contacts at 8, 30, 60 days after suicide attempt by nurses.	2 months	Suicide attempt (medical records)
Fleischmann et al., 2008 [66]. Brazil, India, Sri Lanka, Iran, China	RCT	ED N=1867	Patients admitted to the ED for suicide attempt; mean age IG 23, CG 23; 58.2% women	BIC (n=922) vs. TAU (n=945). 1h brief educational intervention combined with telephone contacts (5 min.) or visits at 1, 2, 4, 7, 11 weeks, 4, 6, 12, and 18 months by a medical doctor, psychiatric nurse, psychologist, or social worker.	18 months	Suicide death (EPSIS) [87]
Gabilondo et al., 2020 [67]. Spain	nRCT; non-randomised controlled	3 General Hospitals N=586	Patients discharged from a General Hospital after a suicide attempt; mean age IG 41.2, CG 45.2; 58.5% women	Telephone follow-up (n=123) vs. TAU (n=463). 6 months telephone contacts (10-15 min.) after 1, 2 weeks, 1, 3, 6 months by general nurses.	12 months	Suicide attempt (medical records)
Goñi-Sarriés et al., 2022 [68]. Spain	nRCT; prospective cohort	2 Psychiatric Units N=191	Patients admitted to the ED after attempted suicide; mean age IG 44.96, CG 46.91; 52.9% women	Telephone follow-up (n=101) vs. TAU (n=90). Telephone contacts (15-20 min.) the day after discharge, at 15 days, and at 2, 4, 8, and 12 months by a specialist nurse.	12 months	Suicide attempts, Suicide death (Interview)
Hassanzadeh et al., 2010 [69]. Iran	RCT	5 General Hospitals ED N=632	Patients with suicide attempt; mean age IG 24 (SD=8.3), mean age CG 25; 62% women	BIC (n=319) vs. TAU (n=310). 1-hour psycho-educational session combined with telephone contacts at 1, 2, 4, 7, 11 weeks, 4, and 6 months by psychologists, nurses, consultants, and psychiatric residents.	6 months	Suicide attempts, Suicide death (SUPRE-MISS questionnaire) [93]

Study	Design	Setting	Participants	Intervention	Follow-up	Outcome (measure)
López-Goñi & Goñi-Sarriés 2021 [70]. Spain	nRCT; prospective cohort	Psychiatric ED *N*=410	Patients discharged from the ED for a suicide attempt; mean age IG 41.7, CG 44.6; 63.7% women	Telephone follow-up (*n*=203) vs. TAU (*n*=207). Telephone follow-up at 15 days, 2, 4, 8, and 12 months by mental health nurses.	12 months	Suicide attempt, Suicide death (medical records)
Malakouti *et al.*, 2022 [71]. Iran	RCT	Hospital ED *N*=305	Patients admitted to the ED after attempted suicide; 66.9% women	Telephone contact programme (*n*=153) vs. TAU (*n*=152). Telephone contact calls (15-20 min.) at 1, 2, and 4 weeks, and monthly for up to 12 months by clinical psychologists with a master's degree.	12 months	Suicide attempts, Suicide death (SUPRE-MISS questionnaire) [93]
Messiah *et al.*, 2019 [72]. France	RCT	23 Hospital ED and psychiatric crisis centres *N*=1040	Patients discharged from the ED for a suicide attempt; 65.9% women	AlgoS: Telephone follow-up for multi-attempters (*n*=317) or crisis card for first attempters (*n*=203) vs. TAU (*n*=520). Telephone follow-up between the 10th and 21st by a psychologist. The patient could not be contacted after 3 attempts, on 3 different days, at 3 different times of day: postcards were sent at 2, 3, 4, and 5 months.	14 months	Suicide attempt (Semi-structured interview)
Miller *et al.*, 2017 [73]. USA	nRCT; non-randomised controlled	8 Hospital ED *N*=1376	Patients with suicide attempt or ideation presented to hospital ED; mean age IG 36, CG 37; 56% women	Universal screening and ED-SAFE intervention (*n*=502) vs. Universal screening (*n*=377) vs. TAU (*n*=497). Risk screening by the ED physician, discharge resource, and 7 post-ED telephone calls (10-20 min.).	12 months	Suicide attempt, Suicide death (CSSRS) [89]
Mouaffak *et al.*, 2015 [74]. France	RCT	Psychiatric ED *N*=320	Patients admitted to the ED for suicide attempt; mean age IG 39, CG 38.6; 70% women	Postal cards, 24-hour telephone and telephone follow-up (*n*=160) vs. TAU (*n*=160). Postal cards, few days, 1, 6, and 11 months providing 24-hour crisis telephone consultation with a psychiatrist, and telephone calls at 2 weeks, 1- and 3 months PD.	12 months	Suicide attempt (SIS) [94], Suicide death (Semi-structured interview)
Mousavi *et al.*, 2014 [75]. Iran	RCT	Hospital ED *N*=139	Patients admitted to the ED for suicide attempt in 2010-2011; 63.3% women	Telephone follow-up (*n*=69) vs. TAU (*n*=70). Telephone contacts (30 min.) at 2, 4 weeks, 2, 3, 4, 5, and 6 months by psychiatry.	6 months	Suicide attempt (*ad hoc* questionnaire)
Mousavi *et al.*, 2016 [76]. Iran	RCT	Hospital ED *N*=55	Patients admitted to ED for suicide attempt; mean age IG 27.07, CG 29.69; 87.3% women	Telephone follow-up (*n*=29) vs. face-to-face (*n*=26). Telephone contacts (20 min.) at 2, 4 weeks, 2, 3, 4, 5, 6, and 8 months by an assistant of psychiatry.	8 months	Suicide attempt (Interview)
O'Connor *et al.*, 2022 [77]. UK	RCT	4 general hospitals *N*=120	Patients with a history of a suicide attempt; mean age IG 36.1, CG 37.0 (*SD*=14.8); 62.5% women	SAFETEL intervention (*n*=80) vs. TAU (*n*=40). Safety planning intervention session and 5 optional telephone calls at 72h of patient discharge from the hospital and weekly thereafter depending on participants' availability by clinical psychologists.	6 months	Suicide Ideation, Suicide attempt (CSSRS) [89]
Plancke *et al.*, 2020 [78]. France	nRCT; retrospective cohort	Hospital ED *N*=6762	Patients discharged from the hospital after a suicide attempt; mean age IG 37.9, mean age CG 39.4; 61.4% women	Telephone follow-up (*n*=3068) vs. TAU (*n*=3694). Telephone contacts at 10, 21 days, and 6 months by psychiatrists. The patient could not be reached by telephone: post-cards were sent at 2, 3, 4, and 5 months.	6 months	Suicide attempt (Interview)
Riblet *et al.*, 2022 [79]. USA	RCT	Veterans Medical Center *N*=20	Patients at risk for suicide; mean age IG 54.9, CG 48.1; 35% women	BIC (*n*=10) vs. TAU (*n*=10). 6 contacts by phone or video over 3 months, educational materials mailed out to the patient, and 1-hour educational intervention on suicide prevention either over the phone or video by a clinical psychologist.	3 months	Suicide Ideation (BSS) [88] + Suicide attempt (CSSRS) [89]
Vaiva *et al.*, 2006 [80]. France	RCT	13 ED *N*=605	Patients discharged from an ED after attempted suicide by deliberate self-poisoning; mean age IG 37, CG 35; 72.9% women	Telephone contact 1 month (*n*=147) or 3 months (*n*=146) vs. TAU (*n*=312). Telephone contact (20-25 min.) at 1 or 3 months after discharge from an ED for attempted suicide by psychiatrists.	13 months	Suicide attempts, Suicide death (medical records)
Vaiva *et al.*, 2018 [81]. France	RCT	23 ED and psychiatry crisis centers *N*=987	Patients discharged from the ED for a suicide attempt; mean age IG 38.4 (*SD*=13.4), mean age CG 38.1 (*SD*=13.1); 63.4% women	AlgoS algorithm (*n*=493) vs. TAU (*n*=494). Telephone contacts between the 10th and 21st day after a suicide attempt by a psychologist. The patient could not be contacted after 3 attempts, on 3 different days, at 3 different times of day, or decline follow-up care from other services, or identified being under stress/experiencing a suicidal crisis: programmed mailing post-cards at months 2, 3, 4, and 5.	13 months	Suicide attempts, Suicide death (Interview)
Ward-Ciesielski *et al.*, 2017 [82]. USA	RCT	University Outpatient Clinic. *N*=93	Participants with suicidal ideation; mean age IG 38.6 (*SD*=15.0), mean age CG 41.8 (*SD*=15.3); 37% women	DBT-BSI and telephone follow-up (*n*=46) vs. RT control condition (*n*=47). Single-session DBT-BSI, presenting pre-selected skills (mindfulness, opposite-to-emotion action, distraction, and changing your body chemistry) and 3 months telephone contact (30-45 min.) at 1, 4, and 12 weeks.	3 months	Suicidal ideation (SSI) [90], Suicide attempts (Interview)
Wei *et al.*, 2013 [83]. China	RCT	4 General Hospitals ED *N*=239	Patients admitted to the ED after attempted suicide; mean age IG 34.06 (*SD*=15.84), mean age CG 32.12 (*SD*=13.87); 77.1% women	Cognitive Therapy group (*n*=82) vs. Telephone intervention group (*n*=80), (3) vs. TAU (*n*=77). 12 weekly phone calls (20-40 min.) within a 3-month period by professors.	12 months	Suicide Ideation (SSI) [90], Suicide attempt (Structured Interview)

Risk of bias in studies

The results of the RoB 2 [45] and ROBINS-I [46] evaluations are presented in Figure 2. Inter-rater

agreement was calculated using Cohen's Kappa (κ=0.94, 95% CI 0.87 to 1.0) [86]. The overall RoB

ratings were classified as low, moderate, or high. A total of 8 (28.6%) studies met the criteria to

be classified as low risk of bias; in 13 (46.4%) studies the risk of bias was moderate; while in 7

(25%) studies the risk of bias was high for the outcome of suicidal behaviours.

320   [Please insert here Figure 2]

Figure 2 RoB assessment. (**A**) RoB 2 evaluation; (**B**) ROBINS-I evaluation.

Results of individual studies

Suicidal ideation

The results of the meta-analyses of the effects of synchronous remote-based intervention on

suicidal ideation are shown in Figure 3. Among the 28 trials, four studies reported the effects of

the intervention on suicidal ideation in short-term (≤ 3 months) [58, 62, 79, 82] and four studies

in medium-term (6–12 months) [61–63, 77]. There was a very small size, non-statistically

significant effect of IG vs. CG on suicidal ideation at short-term (Hedges' g -0.17, 95% CI -0.62 to

0.28 at 1 month; Hedges' g -0.07, 95% CI -0.34 to 0.20 at 3 months). Also, there was a small size,

non-statistically significant effect at medium-term (Hedges' g -0.28, 95% CI -0.89 to 0.34 at 6

months; Hedges' g -0.27, 95% CI -0.56 to 0.03 at 12 months). Heterogeneity was substantial for

nearly all the measurement moments (I^2=59.4% at 1 month, I^2=69.3% at 6 months, I^2=51.9% at

12 months) except for the measurement at 3 months (I^2=0.0%).

Suicide attempts

The results of the meta-analyses of the effects of synchronous remote-based intervention on

suicide reattempts are shown in Figure 4. Among the 28 trials, seven studies reported the effects

of the intervention on suicide attempts in short-term (≤ 3 months) [58, 65, 75, 76, 79, 82, 83],

17 studies in medium-term (6–12 months) [59, 63, 64, 67–78, 81, 83], and six studies in long-

term (>12 months) [56, 57, 61, 72, 80, 81]. There was a statistically significant effect (RR 0.73,

95% CI 0.62 to 0.85) on suicide attempts at 1 month favourable to the intervention group, with

a 27% reduction of the risk of making at least one suicide attempt, relative to control group.

There was a statistically significant effect (RR 0.56, 95% CI 0.34 to 0.95) on suicide attempts at 6

months, with a 44% lower risk of making at least one suicide attempt in the intervention group

compared to the control group, despite considerable heterogeneity (I^2=85.4%). In addition, a

statistically significant effect (RR 0.68, 95% CI 0.49 to 0.96) on suicide attempts at 12 months

was observed, with a 32% lower risk of making at least one suicide attempt in the intervention

group compared to the control group, although there was considerable heterogeneity

(I^2=87.6%). For the remaining measurement points, the heterogeneity was non-obvious (I^2=0.0%

at 1 month, 3 months, 13 months, and 18 months).

Suicide deaths

The results of the meta-analyses of the effects of synchronous remote-based intervention on

suicide deaths are shown in Figure 5. Among the 28 trials, seven studies reported the effects of

the intervention on suicide deaths in medium-term (6–12 months) [63, 68–70, 73, 74, 81], and

four studies in long-term (>12 months) [56, 66, 80, 81]. There was a statistically significant effect

on suicide deaths at 18 months (Peto OR 0.18, 95% CI 0.08 to 0.44), with the likelihood of dying

by suicide being 0.18 times lower in the intervention group compared to the control group. The

heterogeneity was low for all measurement points (I^2=17.9% at 6 months, I^2=0.0% at 12 months,

13 months, and 18 months).

362   [Please insert here Figures 3, 4, and 5]

Figure 3 Forest plot comparing the effects of synchronous – remote interventions on suicidal

intention.

Figure 4 Forest plot comparing the effects of synchronous – remote interventions on suicide

attempts.

Figure 5 Forest plot comparing the effects of synchronous – remote interventions on suicide

deaths.

In order to simplify interpretation, the results can be grouped into three timepoints: short- (≤ 3

months), medium- (6–12 months), and long-term (>12 months) (Table 2).

Table 2

Summary of intervention effects grouped by follow-up period (short-, medium-, and long-term).

Outcome	Term	Month	Effect (95% CI)	Indicators of heterogeneity	Interpretation
Suicidal ideation	Short-term	1 month	Hedges' g -0.17 (-0.62 to 0.28)	I^2=59.4%, Q=7.38, tau^2=0.13	Non-statistically significant
		3 months	Hedges' g -0.07 (-0.34 to 0.20)	I^2=0.0%, Q=1.16, tau^2=0.00	
	Medium-term	6 months	Hedges' g -0.28 (-0.89 to 0.34)	I^2=69.3%, Q=3.26, tau^2=0.14	Non-statistically significant
		12 months	Hedges' g -0.27 (-0.56 to 0.03)	I^2=51.9%, Q=4.16, tau^2=0.03	
Suicide attempts	Short-term	1 month	RR 0.73 (0.62 to 0.85)	I^2=0.0%, Q=0.70, tau^2=0.00	Statistically significant
		3 months	RR 0.50 (0.21 to 1.16)	I^2=0.0%, Q=1.32, tau^2=0.00	Non-statistically significant
	Medium-term	6 months	RR 0.56 (0.34 to 0.95)	I^2=85.4%, Q=54.92, tau^2=0.36	Statistically significant

		12 months	RR 0.68 (0.49 to 0.96)	I^2=87.6%, Q=72.63, tau^2=0.27	
	Long-term	13 months	RR 0.84 (0.66 to 1.06)	I^2=0.0%, Q=3.89, tau^2=0.00	Non-statistically significant
		18 months	RR 1.09 (0.30 to 3.91)	I^2=0.0%, Q=0.49, tau^2=0.00	
Suicide deaths	Medium-term	6 months	Peto OR 0.40 (0.00 to 789.93)	I^2=17.9%, Q=1.22, tau^2=0.15	Non-statistically significant
		12 months	Peto OR 0.87 (0.24 to 3.15)	I^2=0.0%, Q=3.63, tau^2=0.00	
	Long-term	13 months	Peto OR 0.42 (0.11 to 1.65)	I^2=0.0%, Q=0.77, tau^2=0.00	Non-statistically significant
		18 months	Peto OR 0.18 (0.08 to 0.44)	I^2=0.0%, Q=0.03, tau^2=0.00	Statistically significant

CI: confidence interval; OR: odds ratio; RR: risk ratio.

Results of narrative synthesis

Effects of Telephone Contact Interventions on Suicidal Behaviours

A total of 15 articles examined the effectiveness of telephone follow-up interventions in the

prevention of suicidal ideation, suicide attempts, and suicide deaths [59–61, 64, 65, 67, 68, 70,

71, 75, 76, 78, 80, 81, 83]. The studies by Cebrià et al. [59], Exbrayat et al. [65], Malakouti et al.

[71], Plancke et al. [78], and Vaiva et al. [80] reported differences in suicide reattempts in the

intervention group compared to the control group, and a significant effect of the intervention

group in reducing the rate of reattempts. Cebrià et al. [60], after a 60-month follow-up, indicated

that the significant differences found at 12 months were not maintained in either group in terms

of suicide reattempts (31.4% vs. 34.4%, p=0.40).

387   Demesmaeker et al. [64] examined the impact of the Vigilans programme by comparing

interventions of a telephone call only between the 10th and 21st days after discharge, postcards

at months 2–5 (if unreachable), a combination of both (if in suicidal crisis), and TAU. The Kaplan-

Meier method, illustrating the time to the first suicide reattempt within 6 months, showed the

highest survival probability for individuals receiving only a telephone contact and the lowest for

those receiving TAU.

Cedereke *et al*. [61] and Mousavi *et al*. [76] found significant reductions in suicidal

ideation, regardless of the intervention condition. On the other hand, Wei *et al*. [83] compared

a telephone intervention with a cognitive therapy group and a control group, showing

reductions in the number of suicide reattempts between the intervention and control groups

(1.3% vs. 6.5%, *p*=0.08), although the differences were marginally significant. No statistically

significant differences in suicidal ideation were shown between the three groups at any

assessment point (baseline: *p*=0.43, 3-month: *p*=0.72, 6-month: *p*=0.75, 12-month: *p*=0.67).

Gabilondo *et al*. [67], Goñi-Sarriés *et al*. [68], López-Goñi & Goñi-Sarriés [70], Mousavi

et al. [75], Mousavi *et al*. [76], and Vaiva *et al*. [81] reported no significant differences between

intervention and control groups for suicide reattempts. In addition, Goñi-Sarriés *et al*. [68]

informed that there were no significant differences in suicide deaths. More detailed information

is in Table 1.

Effects of Text Message Interventions on Suicidal Behaviours

Comtois *et al*. [63] conducted a follow-up text-messaging intervention with soldiers and marines

considered at risk of suicide. Eleven contacts were made by mental health clinicians, with

participants' text message responses monitored 24 hours a day. The authors reported no

significant effect of the intervention on the presence or severity of current suicidal ideation;

furthermore, the intervention did not significantly decrease the odds of a suicide risk incident

during the 12-month follow-up period.

Effects of Multi-component Interventions on Suicidal Behaviours

A total of 12 articles examined the effectiveness of a multi-component synchronous

telepsychiatry strategies– e.g., 1-hour psychoeducational session plus telephone contacts– in

the prevention of suicidal ideation, suicide attempts, and suicide deaths [56–58, 62, 66, 69, 72–

74, 77, 79, 82]. Miller *et al.* [73] evaluated the ED-SAFE intervention, which consisted of a risk

screening and seven post-ED telephone calls. No significant differences in risk reduction were

found between the TAU and screening phases (22.9% vs. 21.5%, *p*=0.99). Mouaffak *et al.* [74]

delivered postcards, in combination with the possibility of calling a 24-hour telephone, and with

telephone follow-up at 2 weeks, 1, and 3 months after discharge from the ED. No significant

differences were found between the intervention and control groups for suicide reattempts

(14.5% vs. 14%, *p*=0.89). Messiah *et al.* [72] evaluated the effectiveness of the AlgoS Algorithm,

which consisted of a telephone follow-up at 10 and 21 days, for multi-attempters, or a crisis

card, for first-attempts, versus TAU. There was no significant difference for multi-attempters in

the intervention group compared to the control group for suicide reattempts at 6 (RR 0.84, 95%

CI 0.57 to 1.25) and 14 months (RR 1.00, 95% CI 0.73 to 1.37).

Burback *et al.* [58] investigated the impact of 12 web-based, therapist-administered

EMDR sessions targeting experiences associated with suicidal ideation. Both EMDR and TAU

group participants reported significant differences (*p*=0.009) in suicide ideation scores between

baseline and 4-month follow-up (EMDR group baseline: *M*=20, *SD*=7.6; EMDR group 4-month:

M=13, *SD*=9.6; TAU group baseline: *M*=20, *SD*=7.6; TAU group 4-month: *M*=13, *SD*=9.6).

Additionally, EMDR group participants reported fewer serious adverse events compared to TAU

group participants.

Amadéo *et al.* [56], Bertolote *et al.* [57], Fleischmann *et al.* [66], and Hassanzadeh *et al.*

[69] investigated the efficacy of BIC using a brief 1-hour educational intervention combined with

telephone contacts at 1, 2, 4, 7, 11 weeks, 4, 6, 12, and 18 months delivered by physicians,

psychiatric nurses, psychologists, psychiatrists, or social workers. No significant reduction was

found in the intervention group compared to the control group for suicide reattempts, although,

a significant reduction in suicide deaths in the intervention group compared with the control

group (0.2% vs. 2.2%, *p*<0.001) at 18 months was reported in the study by Fleischmann *et al.*

[66]. In a similar intervention protocol, Riblet *et al*. [79] contacted participants by telephone or

video over 3 months, mailed educational materials, and performed a 1-hour educational

intervention on suicide prevention also by telephone or video. A reduction in suicidal ideation

was found in the intervention group compared with the control group at 1 and 3 months.

Christensen *et al*. [62] conducted a web-based CBT intervention combined with

telephone follow-up compared with a CBT-only intervention and a TAU group. CBT was delivered

in five modules, and telephone contacts were provided weekly by a 24-hour telephone

counselling service. Significant reductions in suicidal ideation, in the telephone call condition, at

1-month post-intervention (p=0.003), 6-month follow-up (p=0.03), and 12-month follow-up

(p=0.01) were found. Similarly, Ward-Ciesielski *et al.* [82] conducted a single-session Dialectical

Behaviour Therapy Skills-Based Intervention (DBT-BSI) and telephone follow-up at 1, 4, and 12

weeks, compared with a control group. A significant reduction in levels of suicidal ideation was

reported in both conditions, although there were no significant differences between conditions

for suicidal ideation and the number of suicide attempts. O'Connor *et al*. [77] evaluated the

effects of the SAFETEL intervention against TAU. A safety planning intervention (SPI) session and

five weekly optional telephone calls were conducted. No significant differences were found

between the intervention group and the control group for suicide attempts (Incidence Rate

Ratio; IRR 1.10, 95% CI 0.43 to 2.79, p=0.85).

Effects of Synchronous Remote-Based Interventions on Treatment Adherence

Nineteen studies reported on outcomes relating to treatment attendance [56–61, 64, 66–72,

77, 80–83]. Concerning the moderating effect of adherence on the effectiveness of the

intervention: Gabilondo *et al*. [67] observed that the intervention group showed significantly

higher adherence to outpatient follow-up (p<0.001); in addition, Hassanzadeh *et al*. [69]

revealed an increase in patients' perceived need for support (p<0.001) and positively influenced

attitudes towards seeking support from services and family. Moreover, Vaiva *et al*. [80] reported

that participants in the intervention group discussed their suicide attempts with their primary

care physician more frequently than controls.

According to Cebrià *et al*. [60], suicide attempters tend to be ambivalent to treatment,

do not attend treatment, or drop out prematurely; complementarily, López-Goñi & Goñi-Sarriés

[70] showed in their study that completion of the Telephone Follow-up Programme (TFP)

correlated with a reduction in suicidal behaviours. Along the same lines, Demesmaeker *et al.*

[64] highlight the importance of addressing the specific challenges faced by people who are

unwilling to enter treatment in the context of suicide prevention interventions. Interventions

may not uniformly benefit all patients, possibly due to differences in baseline suicidality or

programme adherence.

Ward-Ciesielski *et al*. [82], and Wei *et al*. [83] also mention that the 75% retention rate

underscores the acceptability and effectiveness of the study in engaging a historically hard-to-

reach and difficult-to-retain population. As mentioned by Messiah *et al*. [72] telephone calls

could complement a multimodal strategy, providing a surveillance tool to drive more aggressive

interventions. Nevertheless, technical challenges, lack of access to computers, or lack of privacy

could prevent the enrollment of some individuals, as noted by Burback *et al.* [58].

Fleischmann *et al*. [66], Gabilondo *et al*. [67], and Demesmaeker *et al.* [64] mention that

BIC operates similarly to psychosocial counselling, providing temporary social support and

fostering feelings of connectedness through systematic follow-up contacts. BCI may help to

reduce social isolation and facilitate the progressive engagement of patients who may initially

be reluctant to participate in structured psychotherapies.

Concerning the predictor or moderator variables of adherence: Amadéo *et al*. [56],

López-Goñi & Goñi-Sarriés [70], O'Connor *et al*. [77], Vaiva *et al*. [80], and Vaiva *et al*. [81]

concurred that patients lost to follow-up were more likely to be male. Gabilondo *et al*. [67],

Goñi-Sarriés *et al*. [68], López-Goñi & Goñi-Sarriés [70], Vaiva *et al*. [80], and Vaiva *et al*. [81]

concluded that the intervention could be more effective for higher-risk populations requiring

additional support; in these studies, patients who completed PTF demonstrated a higher

severity of suicide attempts, greater psychopathological severity, used more violent methods,

and experienced predominantly affective disorders. In contrast, Ward-Ciesielski *et al*. [82] found

that participants lost to follow-up reported higher levels of depression and anxiety. In addition,

Demesmaeker *et al.* [64] reported that post-traumatic stress disorder and generalised anxiety

disorder appear to be predictors of non-adherence to the programme. On the other hand, four

studies [66, 71, 77, 83] discussed stigma around suicide as a factor that may affect patients'

attitudes towards seeking support from mental health services and family. More detailed

information is in Supplemental File 6: Table S4.

Causes of heterogeneity among study results

The investigations on the possible causes of heterogeneity between the outcomes of suicidal

ideation (I^2=59.4% at 1 month, I^2=69.3% at 6 months, I^2=51.9% at 12 months) and suicide

attempts (I^2=85.4% at 6 months, I^2=87.6% at 12 months) of the selected studies raise the

pertinence of qualitatively synthesising the results due to heterogeneity in: a) the evaluation

measures time frame of the outcomes (see Supplemental File 7. Table S5), b) the protocol of

intervention design and the characteristics of the intervention and control group– i.e., target

effect, measures of effect, method of implementation, administrator-participant interaction,

periodicity, number and duration of the intervention, professional delivering the intervention,

setting, participants, etc.– (see Supplemental File 5. Table S3), and c) assessment of RoB and

study design.

Subgroup analysis

Subgroup analysis was conducted based on length of intervention, type of treatment, research

design, and RoB assessment, if feasible [95]. Analysis revealed, for 1-month suicidal ideation

outcomes, that adjustment for confounding variables altered the direction of the effect, with

adjusted analysis indicating a negative effect (Hedges' *g* -0.41, 95% CI -1.12 to 0.31) and

unadjusted analysis showing a small positive effect (Hedges' *g* 0.17, 95% CI 0.00 to 0.33). On the

other hand, the results suggest that telephone contact interventions tend to show greater

effectiveness in reducing suicide attempts at the 6-month follow-up (RR 0.37, 95% CI 0.22 to

0.62) compared to multi-component interventions (RR 0.99, 95% CI 0.59 to 1.65). Additionally,

nRCTs demonstrated a stronger effect (RR 0.31, 95% CI 0.18 to 0.53) compared to RCTs (RR 0.84,

95% CI 0.60 to 1.18) at 6 months follow-up. More detailed information is in Supplemental File

8: Table S6, S7 and S8.

Publication bias

Publication bias was examined using both funnel plots and Egger's test. The visual inspection of

the funnel plots suggested no evidence of publication bias (see Supplemental File 9: Figure S1,

S2, and S3); supported by the results of Egger's test, which indicated absence of evidence for

publication bias in the included primary studies for suicidal ideation (p=0.92), suicide attempts

(p=0.24), and suicide deaths (p=0.12).

Certainty of evidence

The overall certainty of the evidence, according to the Grading of Recommendations

Assessment, Development, and Evaluation (GRADE) [53], was moderate for RCTs reporting

suicidal ideation and suicide death, and low for RCTs and nRCTs reporting suicide attempts. The

main reasons for downgrading were risk of bias, inconsistency, and imprecision in the results

(Supplemental File 10. Table S9).

DISCUSSION

The aim of the study was to assess the effectiveness of synchronous remote interventions in relation to three indicators related to suicidal behaviour: suicidal ideation, suicide attempt, and death by suicide. The results of our systematic review and meta-analyses are largely consistent with previous findings on the effectiveness of telematic interventions in preventing suicidal behaviour. In line with the Shoib *et al*. study [36], our conclusions reinforce the value of telepsychiatry in reducing suicide reattempt rates, although with heterogeneity in effects. Nevertheless, our research represents a significant advancement, offering more recent findings, a greater number of studies, additional variables, a wider range of telepsychiatry interventions, and an analysis of the temporal evolution of the intervention effects, providing a more nuanced understanding of their impact. Furthermore, we specifically analysed suicidal ideation as a possible mediator of the effects of these interventions, providing new insights into the mechanisms that might explain their effectiveness.

Meta-analyses: Effectiveness of synchronous remote-based interventions

The evidence suggests that, although synchronous remote-based interventions appear to reduce suicide attempts and, to some extent, deaths by suicide, their impact on suicidal ideation remains limited. The results seem to indicate that interventions have a clear effect in reducing suicide attempts, with an inverted U-shaped evolution over time, showing a peak of effectiveness at 6 months (-44%, 95% CI -66% to -5% in the intervention group compared to the control group) which subsequently declines. On the contrary, the results of the study do not show statistically significant effects on suicidal ideation, which exhibit a small effect size throughout the four evaluation moments. From the set of studies analysed in relation to deaths by suicide, it is observed that the confidence intervals cover very large ranges of odds ratios.

This is probably because they refer to a very low-frequency phenomenon, which exceeds the

sampling power of conventional empirical clinical trial studies. The juxtaposition of several of

these studies in a meta-analysis also does not solve the problem of lack of sampling power for

this rare occurrence. Likewise, the difficulty of obtaining reliable information on suicide deaths

is highlighted, which adds another challenge to reaching consistent conclusions. Moreover, in

some of our meta-analyses (e.g., 6-month time point in Figure 5), the use of the Knapp-Hartung

adjustment in random-effects models led to unusually wide confidence intervals when a few

studies were included (i.e., n=2). In situations like the one highlighted above, a high level of

uncertainty is generated, and the results clearly do not reflect the current weight of the evidence

in an intuitive sense.

In addition to the marked differences in the effect sizes of synchronous remote-based

interventions on the different indicators, their evolution over time differs, which can be

interpreted as a probable consequence of variations in the studies used for the evaluation of

the three indicators across the time points, compromising comparability. But it could also be

revealing differential effects of the interventions on each of the outcomes. In particular, the

marked difference in the effect sizes observed between suicidal behaviour and suicidal ideation,

which appears to be consistent across the assessment moments, along with the differing

evolution of these outcomes over time, suggests that the clear behavioural effect of the

intervention is not cognitively mediated. Most theoretical models of suicidal behaviour [96]

assume that it is constituted by a continuum of processes ranging from passive ideation and

planning to death through suicide attempt. Considering this sequencing, the apparent effect of

synchronous remote-based interventions on suicide attempts— but not on suicidal ideation—

suggests the probable involvement of an inhibitory mechanism. This mechanism may act directly

on behaviour, temporarily counteracting the influence of suicidal ideation on suicidal actions,

while the ideation itself probably persists as a latent risk factor. In this sense, it should be

considered that a key aspect of synchronous remote-based interventions is that they provide

the person at risk of death by suicide with regular, high-quality social interaction and support,

which is related to the concept of 'connectivity' discussed further. The discrepancy may also be

explained by the nature of the interventions, which often emphasise external support,

monitoring, and engagement rather than targeting the cognitive and emotional processes

underlying suicidal thoughts. While such strategies may enhance safety and prevent behavioural

outcomes like suicide attempts, they may not sufficiently address internal experiences. A recent

meta-analysis of safety planning-type interventions for suicide prevention [97] also finds

differences in reducing suicidal attempts but not suicidal ideation. The mechanism behind the

effectiveness of these interventions lies in risk management, preventing the transition from

suicidal ideation to suicidal action, even though no significant reduction in suicidal thoughts was

observed.

Despite differences in the trends over time of the three indicators evaluated, some

commonality exists since they generally tend to improve over time, probably reflecting a

cumulative effect of the intervention that plateaus around 12-13 months. However, this finding

should be taken with caution as the studies included in the different evaluation moments are

not always the same, which may partially explain apparent changes in effect size over time.

Quality, comparability, and integrability of the studies included in the meta-analyses

There are various aspects of the nature of the studies analysed, as well as the

methodology followed in the meta-analyses, that condition the interpretation of the results.

Firstly, of the 28 studies analysed, only eight (29%) had a low RoB, including seven randomised

controlled trials (37%) and one non-randomised trial (11%). Additionally, unexplained statistical

heterogeneity across studies warrants further investigation to determine effects.

Secondly, although the selected studies share key commonalities– namely, all being

clinical trials and involving synchronous, remote-based interventions– there is substantial

variability in their action protocols and evaluation points. Significant variations were found in

the procedure of the intervention, and comparison group– i.e., periodicity, number of contacts,

moment of assessment, content, practitioner, setting, type of usual treatment, etc.–, as well as

in the methodology– i.e., study design and RoB– of the reviewed studies. Heterogeneity of

results could be due to differences in the length of follow-up, the methods used in the different

programmes, variability in the number of sessions, or different patient inclusion criteria. This

heterogeneity complicates the integration of findings into a meta-analysis. However, it has been

partially addressed by conducting specific meta-analyses based on temporal groupings, which

enhance comparability while also assessing the dynamic evolution of the intervention's

therapeutic effects. Replication of these interventions according to a standardised protocol

would facilitate more consistent results [98].

Narrative synthesis

Firstly, the neologism 'connectivity' seems to serve as a central concept in the majority

of studies [35]. In addition, the importance of a high frequency of follow-up contacts, earlier

intervention, and involvement in the most critical period– i.e., 6 months post-discharged from

ED– is also underlined [10, 13, 35, 80]. Reducing loss to follow-up is a critical challenge in suicide

prevention, especially during the post-discharge period [13, 99], given historical problems with

treatment completion among people involved in suicidal behaviour [100–103]. According to

Gabilondo *et al*. [67] BIC reduces loneliness and improves adherence to routine treatment. Along

the same lines, Ward-Ciesielski *et al*. [82] reported that a notable percentage of participants

engaged with mental health services, or initiated psychotherapy, or pharmacotherapy during

the synchronous remote intervention follow-up, suggesting a significant benefit of these

interventions for a treatment-unengaged population.

Recommendations, implications for clinical practice, and future research

Following the synthesis of information and the articles collected on the topic of the

review, the authors would like to provide a series of recommendations to standardise the

implementation and development of these interventions. Firstly, the authors of future studies

should consider describing in detail a minimum set of component characteristics of the

intervention and comparison groups– particularly for the TAU group– as there is considerable

heterogeneity between studies. A comprehensive quality checklist is presented in Supplemental

File 11. Table S10.

Secondly, as concluded in the systematic review by Noh *et al*. [33] more structured and

theory-based interventions for the prevention of suicidal behaviours should be conducted and

examined, emphasising the importance of a conceptual and theoretical framework in the

development of interventions, and increasing measurable impacts on key outcomes in complex

therapies [104]. A universal intervention design protocol should be adopted according to the

characteristics of the studies in which an effect has been observed and the recommendations

of the authors of previous studies on the topic [59, 62, 65, 66, 71, 78, 80, 81]. In summary, the

preferred approach is a RCT design that delivers early interventions within a maximum of 7 days

post-ED discharge with long-term follow-up of approximately 12 months, often as concomitant

therapy to TAU– i.e., emergency medical care and follow-up based on the severity of suicidal

behaviour; psychopharmacology, or psychotherapy– conducted by a mental health professional

to enhance suicide risk assessments, clinician-patient connection, and adherence to routine

treatment.

Overall, the findings emphasise the importance of standardising interventions to

enhance both research comparability and clinical application. Moreover, the results open up

several avenues for further investigation into effectiveness, underscoring the need to

standardise the outcomes to be assessed, the timing of evaluations, and the measurement

instruments. Addressing the impact of this intervention on suicide deaths will require much

larger cohorts, which in turn implies improving the protocols for recording this indicator to

ensure data reliability. Finally, the suspicion that synchronous remote-based interventions could

have an effectiveness on the reduction of suicide attempts not mediated by a reduction in

suicidal ideation makes it necessary to focus the future of research on the processes that

mediate the effectiveness of interventions, which may be crucial for improving their efficacy and

cost-effectiveness. In particular, future research could benefit from integrating components

designed to promote cognitive change and emotional regulation in order to more effectively

influence ideation.

Strengths and limitations

The primary limitation is the significant heterogeneity observed in the overall meta-

analyses and subgroup analyses, which may be attributed to the small number of studies

included in some analyses or the substantial variability between their results. In addition, using

ED data records or the authorities' forensic register may lead to the possibility of under-

reporting the actual number of non-fatal suicidal behaviours [105]. Furthermore, suicide is a

rare event, making the design of studies with high statistical power particularly difficult. Finally,

the included trials used intervention and comparison groups that received TAU; these

treatments are often different across studies. Relatedly, although study characteristics were

described, the lack of standardised reporting across research limited further stratification.

The comprehensive literature review and the identification of key characteristics for

improving future study designs are fundamental strengths. Other strengths of the study are the

number of databases explored, the reporting according to PRISMA guidelines, the validation of

the boolean searches according to PRESS guidelines, and the inter-reviewer consensus in the

different phases of screening, data extraction, and RoB assessment.

CONCLUSIONS

Innovation in the prevention and management of suicidal behaviour is a therapeutic

challenge [106]. The current research outlines emerging evidence on suicide prevention through

telemedicine in reducing suicidal behaviours. The main conclusion of the study is that the results

seem to indicate that synchronous remote-based interventions demonstrate a potential effect

in reducing suicide attempts and suicide deaths, although their influence on suicidal ideation

appears limited. The mechanisms underlying these effects remain unclear but may be related to

the promotion of social bonding or 'connectedness'. The difficulties in studying the effectiveness

of such preventive interventions are reflected by the great heterogeneity of studies to date and

the problems in obtaining accurate information on suicidal behaviours in the absence of reliable

registries [107]. Finally, an indirect positive effect on the treatment of patients at risk of death

by suicide seems to derive from the increase in adherence fostered by remote synchronous

interventions.

ETHICS AND DISSEMINATION

Ethics approval is not needed, as systematic reviews are based on published studies. The results

will be disseminated through peer-reviewed publications.

ETHICS STATEMENTS

Patient consent for publication

Not applicable.

AUTHOR CONTRIBUTIONS

AS is the guarantor. LC, JML, DP, AC, and AS: Writing - Original Draft. LC, AS, MPJ, JPS, and CM:

Software. LC, JML, DP and AS: Project administration, Supervision. All authors:

Conceptualization, Methodology, Writing - Review & Editing. JML, AS, JPS, and CM provided

statistical expertise. DP and AC provided expertise on suicidal behaviours. All authors approved

the final manuscript.

AI USE

Google Translate was used to check (and modify where necessary) the linguistic quality and

suitability of the manuscript. No other AI tools were used in either the study or the writing of

the manuscript.

ACKNOWLEDGEMENTS

Authors' thanks to Guillem Cebrián, director of the Library at Parc Taulí University Hospital and

head of the Unitat de Gestió del Coneixement de l'Institut d'Investigació i Innovació Parc Taulí

(I3PT-CERCA), for his support in the refinement of the search strategies. DP thanks the support

of Spanish Ministry of Science and Innovation/ISCIII/MICIU/FEDER (PI21/01148); the Secretaria

d'Universitats i Recerca del Departament d'Economia i Coneixement of the Generalitat de

Catalunya (e-MH-PEMN – 2021 SGR 01431); the CERCA programme / Generalitat de Catalunya;

the Instituto de Salud Carlos III; and the CIBER of Mental Health (CIBERSAM CB 19/09/00029).

The research has been previously presented at a conference and has been published as a

conference abstract [108].

FUNDING

This research was funded by the Instituto de Salud Carlos III, Subdirección General de Evaluación

y Fomento de la Investigación (ISCIII), Ministerio de Ciencia, Innovación y Universidades (MICIU)

co-funded by the European Union, Fondo Europeo de Desarrollo Regional (FEDER) (grant

number PI21/01148 — COGNISUI) and MCIN/AEI/10.13039/501100011033 'ERDF A way of

making Europe' (grant number PID2022-141403NB-I00). Fundació Institut d'Investigació i

Innovació Parc Taulí (I3PT-CERCA), SENDO PI21/01148 (Fondo de Investigación Sanitaria-ISCIII/

FEDER), Instituto de Salud Carlos III. The funders had no role in the design of the study; in the

collection, analysis, or interpretation of data; in the writing of the manuscript; or in the decision

to publish the results. The Department of Mental Health at the Parc Taulí University Hospital

and the Institut d'Investigació i Innovació Parc Taulí (I3PT-CERCA) are the sponsors.

COMPETING INTERESTS

DP has received grants and served as a consultant or advisor for Rovi, Johnson & Johnson, and

Lundbeck. The other authors declare no conflicts of interest.

PATIENT AND PUBLIC INVOLVEMENT

Patients and/or the public were not involved in the design, conduct, reporting, or dissemination

plans of this research.

PATIENT CONSENT FOR PUBLICATION

Not applicable.

PROVENANCE AND PEER REVIEW

Not commissioned; externally peer reviewed.

DATA AVAILABILITY STATEMENT

Data are available in a public, open access repository. The dataset is available in [Dataset]

Comendador L; Jiménez-Villamizar MP; Sanabria-Mazo, JP, *et al*. Data from: Effect of

synchronous remote-based interventions on suicidal behaviours: systematic review and meta-

analysis. CORA Repository, February 9, 2025. https://doi.org/10.34810/data1310 [109]

SUPPLEMENTAL MATERIAL

Supplemental File 1. PRISMA 2020 Checklist (DOCX 32 KB).

Supplemental File 2. Amendments made to the original protocol (DOCX 11 KB).

Supplementary File 3. Search strategy for databases, registers, and websites (DOCX 13 KB).

Supplemental File 4. List of articles excluded after full-text screening (DOCX 23 KB).

Supplemental File 5. Information on the components and characteristics of the intervention and

comparison groups (DOCX 20 KB).

Supplemental File 6. Information on therapeutic adherence (DOCX 15 KB).

Supplemental File 7. Evaluation measures the time frame of the selected studies (DOCX 14 KB).

Supplemental File 8. Subgroup analysis (DOCX 27 KB).

Supplemental File 9. Publication bias (DOCX 130 KB).

Supplemental File 10. GRADE (DOCX 689 KB).

Supplemental File 11. Report of the minimum set of component characteristics (DOCX 11 KB).

OPEN ACCESS

This is an open-access article distributed by the Creative Commons Attribution Non-Commercial

(CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work

non-commercially, and license their derivative works on different terms, provided the original

ORCID iDs

Laura Comendador Vázquez https://orcid.org/0000-0002-5221-4794

María P Jiménez-Villamizar https://orcid.org/0000-0003-2264-7422

Josep-Maria Losilla https://orcid.org/0000-0002-5140-5847

Juan P Sanabria-Mazo https://orcid.org/0000-0003-1688-435X

Corel Mateo-Canedo https://orcid.org/0000-0002-0620-9257

Antoni Sanz Ruíz https://orcid.org/0000-0002-7952-4477

Ana-Isabel Cebrià Meca https://orcid.org/0000-0002-2632-8130

Diego Palao Vidal https://orcid.org/0000-0002-3323-6568

REFERENCES

1. World Health Organisation. Suicide [online]. 2023. https://www.who.int/news-room/fact-

sheets/detail/suicide (accessed 06 April 2024).

2. World   Health   Organisation.   Suicide   worldwide   in   2019   [online].   2021.

https://www.who.int/publications-detail-redirect/9789240026643 (accessed 30 Jan 2024).

3. World Health Organisation. Suicide in the world: Global health estimates [online]. 2019.

https://apps.who.int/iris/bitstream/handle/10665/326948/WHO-MSD-MER-19.3-

eng.pdf?sequence=1&isAllowed=y (accessed 30 Jan 2024).

4. Artieda-Urrutia P, Parra-Uribe I, Garcia-Pares G, *et al*. Management of suicidal behaviour: Is

the   world   upside   down?   *Aust   N   Z   J   Psychiatry*   2014;48:399–401.

https://doi.org/10.1177/0004867414525847

5.   Mokdad AH, Forouzanfar MH, Daoud F, *et al*. Global burden of diseases, injuries, and risk

factors for young people's health during 1990–2013: A systematic analysis for the Global

Burden of Disease Study 2013. *Lancet* 2016;387:2383–401. https://doi.org/10.1016/S0140-

6736(16)00648-6

6.   Parra-Uribe I, Blasco-Fontecilla H, García-Parés G, *et al*. Attempted and completed suicide:

Not   what   we   expected?   *J   Affect   Disord*   2013;150:840–6.

https://doi.org/10.1016/j.jad.2013.03.013

7.   World Health Organisation. Suicide: key facts. Geneva: World Health Organisation [online].

2021.   https://www.who.int/news-room/fact-sheets/detail/suicide   (accessed   06   April

2024).

8.   Olfson M, Wall M, Wang S, *et al*. Suicide following deliberate self-harm. *Am J Psychiat*

2017;174:765–74. https://doi.org/10.1176/appi.ajp.2017.16111288

9.   Shand F, Vogl L, Robinson J. Improving patient care after a suicide attempt. *Australas*

Psychiatry 2018;26:145–8. https://doi.org/10.1177/1039856218758560

10. Parra-Uribe I, Blasco-Fontecilla H, Garcia-Parés G, *et al*. Risk of re-attempts and suicide death

after   a   suicide   attempt: A   survival   analysis.   *BMC   Psychiatry*   2017;17:163.

https://doi.org/10.1186/s12888-017-1317-z

11. Bostwick JM, Pabbati C, Geske JR, McKean AJ. Suicide Attempt as a Risk Factor for Completed

Suicide: Even More Lethal Than We Knew. *Am J Psychiatry*. 2016;173(11):1094-1100.

https://doi.org/10.1176/appi.ajp.2016.15070854

12. Zalsman G, Hawton K, Wasserman D, *et al*. Evidence-based national suicide prevention

taskforce in Europe: A consensus position paper. *Eur Neuropsychopharmacol* 2017;27:418–

21. https://doi.org/10.1016/j.euroneuro.2017.01.012

13. Inagaki M, Kawashima Y, Yonemoto N, *et al*. Active contact and follow-up interventions to

prevent repeat suicide attempts during high-risk periods among patients admitted to

emergency departments for suicidal behavior: A systematic review and meta-analysis. *BMC*

Psychiatry 2019;19:44. https://doi.org/10.1186/s12888-019-2017-7

14. Ganz D, Braquehais MD, Sher L. Secondary Prevention of Suicide. *PLoS Med*

2010;7:e1000271. https://doi.org/10.1371/journal.pmed.1000271

15. World Health Organisation. The European Mental Health Action Plan 2013–2020 [online].

2013.   https://www.euro.who.int/__data/assets/pdf_file/0020/280604/WHO-Europe-

Mental-Health-Acion-Plan-2013-2020.pdf (accessed 15 Feb 2024).

16. World Health Organisation. Preventing suicide: A global imperative [online]. 2014.

https://www.who.int/mental_health/suicide-prevention/exe_summary_english.pdf?ua=1

(accessed 15 Feb 2024).

17. World Health Organisation. Comprehensive mental health action plan 2013–2030 [online].

2021. https://apps.who.int/iris/rest/bitstreams/1371507/retrieve (accessed 15 Feb 2024).

18. Inagaki M, Kawashima Y, Kawanishi C, *et al*. Interventions to prevent repeat suicidal

behavior in patients admitted to an emergency department for a suicide attempt: A meta-

analysis. *J Affect Disord* 2015;175:66–78. https://doi.org/10.1016/j.jad.2014.12.048

19. Zalsman G, Hawton K, Wasserman D, *et al*. Suicide prevention strategies revisited: 10-year

systematic review. *Lancet Psychiatry* 2016;3:646–59. https://doi.org/10.1016/S2215-

0366(16)30030-X

20. Andersson G, Titov N, Dear BF, Rozental A, Carlbring P. Internet-delivered psychological

treatments: from innovation to implementation. *World Psychiatry*. 2019;18(1):20-28.

https://doi.org/10.1002/wps.20610

21. Carlbring P, Hadjistavropoulos H, Kleiboer A, Andersson G. A new era in Internet

interventions: The advent of Chat-GPT and AI-assisted therapist guidance. *Internet Interv.*

2023;32:100621. https://doi.org/10.1016/j.invent.2023.100621

22. Laranjeira C, Carvalho D, Valentim O, *et al*. Therapeutic Adherence of People with Mental

Disorders: An Evolutionary Concept Analysis. *Int J Environ Res Public Health.*

2023;20(5):3869. https://doi.org/10.3390/ijerph20053869

23. Deng M, Zhai S, Ouyang X, Liu Z, Ross B. Factors influencing medication adherence among

patients with severe mental disorders from the perspective of mental health professionals.

BMC Psychiatry. 2022;22(1):22. https://doi.org/10.1186/s12888-021-03681-6

24. Kauppi K, Hätönen H, Adams CE, Välimäki M. Perceptions of treatment adherence among

people with mental health problems and health care professionals. *J Adv Nurs*.

2015;71(4):777-788. https://doi.org/10.1111/jan.12567

25. Gilat I, Shahar G. Emotional first aid for a suicide crisis: Comparison between telephonic

hotline and Internet. Psychiatry-Interpers *Biol Process* 2007;70:12–8.

https://doi.org/10.1521/psyc.2007.70.1.12

26. Seong JM, Cho Y, Cho GC, *et al*. Effects of mobile messenger counseling on case management

success for individuals engaging in self-harm or suicide attempts who were discharged from

emergency departments. *Clin Exp Emerg Med* 2021;8:48–54.

https://doi.org/10.15441/CEEM.20.133

27. Vijayakumar L, Umamaheswari C, Shujaath Ali Z, *et al*. Intervention for suicide attempters:

A randomized controlled study. *Indian J Psychiatry* 2011;53:244-8.

https://doi.org/10.4103/0019-5545.86817

28. Martínez-Alés G, Cruz Rodríguez JB, Lázaro P, *et al*. Cost-effectiveness of a Contact

Intervention and a Psychotherapeutic Program for Post-discharge Suicide Prevention. *Can J*

Psychiatry. 2021;66(8):737-746 https://doi.org/10.1177/0706743720980135

29. De Leo D, Buono MD, Dwyer J. Suicide among the elderly: The long-term impact of a

telephone support and assessment intervention in northern Italy. *Br J Psychiatry*

2002;181:226–9. https://doi.org/10.1192/bjp.181.3.226

30. Gould MS, Munfakh JLH, Kleinman M, *et al*. National Suicide Prevention Lifeline: Enhancing

mental health care for suicidal individuals and other people in crisis. *Suicide Life-Threat*

Behav 2012;42:22–35. https://doi.org/10.1111/j.1943-278X.2011.00068.x

31. Aiello LM, Dadashzadeh S, Lynn JM, *et al*. Using telemedicine to reduce suicide ideation and

behavior: a   systematic   literature   review.   *THMT*   2021;6:1.

https://doi.org/10.30953/tmt.v6.226

32. Milner AJ, Carter G, Pirkis J, *et al*. Letters, green cards, telephone calls and postcards:

Systematic and meta-analytic review of brief contact interventions for reducing self-harm,

suicide   attempts   and   suicide.   *Br   J   Psychiatry*   2015;206:184–90.

https://doi.org/10.1192/bjp.bp.114.147819

33. Noh D, Park YS, Oh EG. Effectiveness of telephone-delivered interventions following suicide

attempts: A   systematic   review.   *Arch   Psychiatr   Nurs*   2016;30:114–9.

https://doi.org/10.1016/j.apnu.2015.10.012

34. Shin HD, Durocher K, Sequeira L, *et al*. Information and communication technology-based

interventions for suicide prevention implemented in clinical settings: a scoping review. *BMC*

Health Serv Res. 2023;23:281. https://doi.org/10.1186/s12913-023-09254-5

35. Luxton DD, June JD, Comtois KA. Can postdischarge follow-up contacts prevent suicide and

suicidal   behavior?   A   Review   of   the   Evidence.   *Crisis*   2013;34:32–41.

https://doi.org/10.1027/0227-5910/a000158

36. Shoib S, Shaheen N, Anwar A, *et al.* The effectiveness of telehealth interventions in suicide

prevention: A systematic review and meta-analysis. *Int J Soc Psychiatry* 2024;70:415–423.

https://doi.org/10.1177/00207640231206059

37. Page MJ, Moher D, Bossuyt PM, *et al*. PRISMA 2020 explanation and elaboration: updated

guidance and exemplars for reporting systematic reviews. *BMJ*. 2021;372:n160.

https://doi.org/10.1136/bmj.n160

38. Comendador L, Jiménez-Villamizar MP, Losilla JM, *et al*. Effect of synchronous remote-based

interventions on suicidal behaviours: protocol for a systematic review and meta-analysis.

BMJ Open. 2023;13(12):e075116. https://doi.org/10.1136/bmjopen-2023-075116

39. World Health Organisation. ICD-11 for mortality and morbidity statistics [online]. 2021.

https://icd.who.int/browse11/l-m/en#/http://id.who.int/icd/entity/778734771 (accessed

15 Feb 2024).

40. Gómez A, Silva H, Amon R. El Suicidio. Teoría, Clínica y Manejo. Barcelona, ES: Editorial

Mediterráneo 2018.

41. Savović J, Whiting PF, Dekkers OM. Systematic reviews of nonrandomized studies of

interventions. In: Egger M, Higgins JPT, Davey Smith G, editors. *Systematic reviews in health*

research.   2nd   ed.   Hoboken,   NJ: Wiley   2022.   Chapter   15.

https://doi.org/10.1002/9781119099369.ch15

42. Saldanha IJ, Skelly AC, Ley KV, *et al*. Inclusion of nonrandomized studies of interventions in

systematic reviews of intervention effectiveness: an update [online]. Rockville (MD): Agency

for   Healthcare   Research   and   Quality   (US);   2022.

https://www.ncbi.nlm.nih.gov/books/NBK584468/table/ch4.tab1/ (accessed 21 Jun 2025).

43. McGowan J, Sampson M, Salzwedel DM, *et al*. PRESS Peer Review of Electronic Search

Strategies: 2015   Guideline   Statement.   *J   Clin   Epidemiol*   2016;75:40–6.

https://doi.org/10.1016/j.jclinepi.2016.01.021

44. Ouzzani M, Hammady H, Fedorowicz Z, *et al*. Rayyan—a web and mobile app for systematic

reviews. *Syst Rev* 2016;5(1):1-10. https://doi.org/10.1186/s13643-016-0384-4

45. Sterne JA, Savović J, Page MJ, *et al*. RoB 2: A revised tool for assessing risk of bias in

randomised trials. *BMJ* 2019;366:l4898. https://doi.org/10.1136/bmj.l4898

46. Sterne JA, Hernán MA, Reeves BC, *et al*. ROBINS-I: A tool for assessing risk of bias in non-

randomised   studies   of   interventions.   *BMJ*   2016;366:i4919.

https://doi.org/10.1136/bmj.i4919

47. Higgins JPT, Thomas J, Chandler J, *et al*. Cochrane Handbook for Systematic Reviews of

Interventions version 6.3. [online]. 2022. https://training.cochrane.org/handbook (accessed

20 Feb 2025).

48. Hedges LV. Distribution Theory for Glass's Estimator of Effect Size and Related Estimators. *J*

Stat Educ 1981;6:107-128. https://doi.org/10.2307/1164588

49. Daly LE. Confidence limits made easy: interval estimation using a substitution method. *Am J*

Epidemiol. 1998;147(8):783-790. https://doi.org/10.1093/oxfordjournals.aje.a009523

50. Zhang J, Yu KF. What's the relative risk? A method of correcting the odds ratio in cohort

studies   of   common   outcomes.   *JAMA*.   1998;280(19):1690-1691.

https://doi.org/10.1001/jama.280.19.1690

51. Peto R, Pike MC, Armitage P, *et al*. Design and analysis of randomized clinical trials requiring

prolonged   observation   of   each   patient.   I.   Introduction   and   design.   *Br J Cancer*.

1976;34(6):585-612. https://doi.org/10.1038/bjc.1976.220

52. Suurmond R, van Rhee H, Hak T. Introduction, comparison, and validation of Meta-

Essentials: A free and simple tool for meta-analysis. *Res Synth Methods*. 2017;8(4):537-553.

https://doi.org/10.1002/jrsm.1260

53. Schünemann H, Brożek J, Guyatt G, *et al*. GRADE Handbook for grading quality of evidence

and   strength   of   recommendations   [online].   2013.   Available:

https://gdt.gradepro.org/app/handbook/handbook.html [Accessed 30 Jun 2024].

54. Balshem H, Helfand M, Schünemann HJ, *et al*. GRADE guidelines: 3. Rating the quality of

evidence. *J Clin Epidemiol* 2011;64:401–6. https://doi.org/10.1016/j.jclinepi.2010.07.015

55. Guyatt G, Oxman AD, Akl EA, *et al*. GRADE guidelines: 1. Introduction-GRADE evidence

profiles and summary of findings tables*. J Clin Epidemiol* 2011;64:383–94.

https://doi.org/10.1016/j.jclinepi.2010.04.026

56. Amadéo S, Rereao M, Malogne A, *et al*. Testing Brief Intervention and Phone Contact among

Subjects with Suicidal Behavior: A Randomized Controlled Trial in French Polynesia in the

Frames of the World Health Organization/Suicide Trends in At-Risk Territories Study. *Ment*

Illn. 2015;7(2):5818. https://doi.org/10.4081/mi.2015.5818

57. Bertolote JM, Fleischmann A, De Leo D, *et al*. Repetition of suicide attempts: data from

emergency care settings in five culturally different low- and middle-income countries

participating in the WHO SUPRE-MISS Study. *Crisis*. 2010;31(4):194-201.

https://doi.org/10.1027/0027-5910/a000052

58. Burback L, Yap S, Purdon SE, *et al*. Randomized controlled trial investigating web-based,

therapist delivered eye movement desensitization and reprocessing for adults with suicidal

ideation. *Front Psychiatry.* 2024;15(1):1361086.

https://doi.org/10.3389/fpsyt.2024.1361086

59. Cebrià AI, Parra I, Pàmias M, *et al*. Effectiveness of a telephone management programme

for patients discharged from an emergency department after a suicide attempt: controlled

study in a Spanish population. *J Affect Disord*. 2013;147(1-3):269-276.

https://doi.org/10.1016/j.jad.2012.11.016

60. Cebrià AI, Pérez-Bonaventura I, Cuijpers P, *et al*. Telephone Management Program for

Patients Discharged From an Emergency Department After a Suicide Attempt: A 5-Year

Follow-Up Study in a Spanish Population. *Crisis*. 2015;36(5):345-352.

https://doi.org/10.1027/0227-5910/a000331

61. Cedereke M, Monti K, Ojehagen A. Telephone contact with patients in the year after a

suicide attempt: does it affect treatment attendance and outcome? A randomised

controlled study. *Eur Psychiatry*. 2002;17(2):82-91. https://doi.org/10.1016/s0924-
9338(02)00632-6

62. Christensen H, Farrer L, Batterham PJ, Mackinnon A, Griffiths KM, Donker T. The effect of a
web-based depression intervention on suicide ideation: secondary outcome from a
randomised controlled trial in a helpline. *BMJ Open*. 2013;3(6):e002886.
https://doi.org/10.1136/bmjopen-2013-002886

63. Comtois KA, Kerbrat AH, DeCou CR, *et al*. Effect of Augmenting Standard Care for Military
Personnel With Brief Caring Text Messages for Suicide Prevention: A Randomized Clinical
Trial. *JAMA Psychiatry*. 2019;76(5):474-483.
https://doi.org/10.1001/jamapsychiatry.2018.4530

64. Demesmaeker A, Amad A, Blekic W, *et al*. Increased risk of suicide reattempt according to
the type of brief contact interventions in the VigilanS program: The critical role of PTSD and
anxiety disorders. *J Psychiatr Res*. 2024;181(1):29-35.
https://doi.org/10.1016/j.jpsychires.2024.11.031

65. Exbrayat S, Coudrot C, Gourdon X, *et al*. Effect of telephone follow-up on repeated suicide
attempt in patients discharged from an emergency psychiatry department: a controlled
study. *BMC Psychiatry*. 2017;17(1):96. https://doi.org/10.1186/s12888-017-1258-6

66. Fleischmann A, Bertolote JM, Wasserman D, *et al*. Effectiveness of brief intervention and
contact for suicide attempters: a randomized controlled trial in five countries. *Bull World
Health Organ*. 2008;86(9):703-709. https://doi.org/10.2471/blt.07.046995

67. Gabilondo A, Aristegi E, Gonzalez-Pinto A, *et al*. Prevention of Suicidal Behavior with
Telemedicine in Patients with a Recent Suicide Attempt: Is a 6-month Intervention Long
Enough?. *Suicide Life Threat Behav*. 2020;50(1):211-219.
https://doi.org/10.1111/sltb.12576

68. Goñi-Sarriés A, Yárnoz-Goñi N, López-Goñi JJ. Psychiatric Hospitalization for Attempted

Suicide and Reattempt at the One-Year Follow-Up. *Psicothema*. 2022;34(3):375-382.

https://doi.org/10.7334/psicothema2021.455

69. Hassanzadeh M, khajeddin N, Nojomi M, Fleischmann A, Eshrati T. Brief Intervention and

Contact after Deliberate Self-Harm: An Iranian Randomized Controlled Trial. *Iran J Psychiatry*

Behav Sci. 2010;4(2):5-12.

70. López-Goñi JJ, Goñi-Sarriés A. Effectiveness of a telephone prevention programme on the

recurrence of suicidal behaviour. One-year follow-up. *Psychiatry Res*. 2021;302:114029.

https://doi.org/10.1016/j.psychres.2021.114029

71. Malakouti SK, Nojomi M, Ghanbari B, Rasouli N, Khaleghparast S, Farahani IG. Aftercare and

Suicide   Reattempt   Prevention   in   Tehran,   Iran.   *Crisis*.   2022;43(1):18-27.

https://doi.org/10.1027/0227-5910/a000754

72. Messiah A, Notredame CE, Demarty AL, Duhem S, Vaiva G; AlgoS investigators. Combining

green cards, telephone calls and postcards into an intervention algorithm to reduce suicide

reattempt (AlgoS): P-hoc analyses of an inconclusive randomized controlled trial. *PLoS One*.

2019;14(2):e0210778. https://doi.org/10.1371/journal.pone.0210778

73. Miller IW, Camargo CA Jr, Arias SA, *et al*. Suicide Prevention in an Emergency Department

Population: The   ED-SAFE   Study.   *JAMA   Psychiatry*.   2017;74(6):563-570.

https://doi.org/10.1001/jamapsychiatry.2017.0678

74. Mouaffak F, Marchand A, Castaigne E, Arnoux A, Hardy P. OSTA program: A French follow

up intervention program for suicide prevention. *Psychiatry Res*. 2015;230(3):913-918.

https://doi.org/10.1016/j.psychres.2015.11.024

75. Mousavi SG, Zohreh R, Maracy MR, Ebrahimi A, Sharbafchi MR. The efficacy of telephonic

follow up in prevention of suicidal reattempt in patients with suicide attempt history. *Adv*

Biomed Res. 2014;3:198. https://doi.org/10.4103/2277-9175.142043

76. Mousavi SG, Amini M, Mahaki B, Bagherian-Sararoudi R. Effect of phone call versus face-to-

face follow-up on recurrent suicide attempts prevention in individuals with a history of

multiple suicide attempts. *Adv Biomed Res*. 2016;5:184. https://doi.org/10.4103/2277-

9175.190990

77. O'Connor RC, Smillie S, McClelland H, *et al*. SAFETEL: a pilot randomised controlled trial to

assess the feasibility and acceptability of a safety planning and telephone follow-up

intervention to reduce suicidal behaviour. *Pilot Feasibility Stud*. 2022;8(1):156.

https://doi.org/10.1186/s40814-022-01081-5

78. Plancke L, Amariei A, Danel T, *et al*. Effectiveness of a French Program to Prevent Suicide

Reattempt (VigilanS). *Arch Suicide Res*. 2021;25(3):570-581.

https://doi.org/10.1080/13811118.2020.1735596

79. Riblet NB, Kenneally L, Stevens S, *et al*. A virtual, pilot randomized trial of a brief intervention

to prevent suicide in an integrated healthcare setting. *Gen Hosp Psychiatry*. 2022;75:68-74.

https://doi.org/10.1016/j.genhosppsych.2022.02.002

80. Vaiva G, Vaiva G, Ducrocq F, *et al*. Effect of telephone contact on further suicide attempts in

patients discharged from an emergency department: randomised controlled study. *BMJ*.

2006;332(7552):1241-1245. https://doi.org/10.1136/bmj.332.7552.1241

81. Vaiva G, Berrouiguet S, Walter M, *et al*. Combining Postcards, Crisis Cards, and Telephone

Contact Into a Decision-Making Algorithm to Reduce Suicide Reattempt: A Randomized

Clinical Trial of a Personalized Brief Contact Intervention. *J Clin Psychiatry*.

2018;79(6):17m11631. https://doi.org/10.4088/JCP.17m11631

82. Ward-Ciesielski EF, Tidik JA, Edwards AJ, Linehan MM. Comparing brief interventions for

suicidal individuals not engaged in treatment: A randomized clinical trial. *J Affect Disord*.

2017;222:153-161. https://doi.org/10.1016/j.jad.2017.07.011

83. Wei S, Liu L, Bi B, *et al*. An intervention and follow-up study following a suicide attempt in

the emergency departments of four general hospitals in Shenyang, China. *Crisis*.

2013;34(2):107-115. https://doi.org/10.1027/0227-5910/a000181

84. Page MJ, McKenzie JE, Bossuyt PM, *et al*. The PRISMA 2020 statement: an updated guideline

for reporting systematic reviews. *BMJ*. 2021;372:n71. https://doi.org/10.1136/bmj.n71

85. Andrés AM, Marzo PF. Delta: a new measure of agreement between two raters. *Br J Math*

Stat Psychol. 2004;57:1-19. https://doi.org/10.1348/000711004849268

86. Cohen J. A coefficient of agreement for nominal scales. *Educ PsycholMeas*. 1960; 20: 37-46.

87. De Leo D, Scocco P, Marietta P, *et al*. Physical illness and parasuicide: evidence from the

European Parasuicide Study Interview Schedule (EPSIS/WHO-EURO). *Int J Psychiatry Med*.

1999;29(2):149-163. https://doi.org/10.2190/E87K-FG03-CHEE-UJD3

88. Beck AT, Steer RA. *Beck scale for suicide ideation: Manual*. San Antonio, TX: Psychological

Corporation 1993.

89. Posner K, Brown GK, Stanley B, *et al*. The Columbia-Suicide Severity Rating Scale: initial

validity and internal consistency findings from three multisite studies with adolescents and

adults. *Am J Psychiatry*. 2011;168(12):1266-1277.

https://doi.org/10.1176/appi.ajp.2011.10111704

90. Beck AT, Kovacs M, Weissman A. Assessment of suicidal intention: the Scale for Suicide

Ideation*. J Consult Clin Psychol*. 1979;47(2):343-352. https://doi.org/10.1037//0022-

006x.47.2.343

91. Gibbons P, de Arévalo HF, Mónico M. Assessment of the factor structure and reliability of

the 28-item version of the General Health Questionnaire (GHQ-28) in El Salvador. *Int J Clin*

Health Psychol. 2004;4(2):389-398.

92. Linehan M, Comtois KA. Suicide Attempt and Self-Injury Count (SASI-Count) [online]. 1996.

http://depts.washington.edu/uwbrtc/wp-content/uploads/L-SASI-Count.pdf (accessed 28

April 2024)

93. Fleischmann A, Bertolote JM, De Leo D, Wasserman D. Instruments used in SUPRE-MISS: The

WHO Multisite Intervention Study on Suicidal Behaviours. In: Wasserman D, Wasserman C,

eds. *Oxford Textbook of Suicidology and Suicide Prevention*. 1th ed. Oxford, OED: Oxford

Textbooks; 2009. https://doi.org/10.1093/med/9780198570059.003.0042

94. Beck AT, Schuyler D, Herman I. Development of suicidal intent scales. In: Beck AT, Resnik

HLP, Lettieri DJ, eds. *The prediction of suicide*. Bowie, MD: Charles Press; 1974:45–56.

95. Borenstein M, Hedges LV, Higgins JP, Rothstein HR. *Introduction to meta-analysis.*

Chichester, UK: John Wiley & Sons, 2021.

96. Díaz-Oliván I, Porras-Segovia A, Barrigón ML, Jiménez-Muñoz L. E. Baca-García. Theoretical

models of suicidal behaviour: A systematic review and narrative synthesis. *Eur J Psychiatry*.

2021;35:181-192. https://doi.org/10.1016/j.ejpsy.2021.02.002

97. Nuij C, van Ballegooijen W, de Beurs D, *et al*. Safety planning-type interventions for suicide

prevention: meta-analysis.   *Br   J   Psychiatry.*   2021;219(2):419-426.

https://doi.org/10.1192/bjp.2021.50

98. Turecki G, Brent DA, Gunnell D, *et al*. Suicide and suicide risk. *Nat Rev Dis Primers*. 2019;5(1).

http://dx.doi.org/10.1038/s41572-019-0121-0

99. Hunt IM, Kapur N, Webb R, *et al*. Suicide in recently discharged psychiatric patients: a case-

control   study.   *Psychol   Med*.   2009;39(3):443-449.

https://dx.doi.org/10.1017/S0033291708003644

100.   Andersen B, Bille-Brahe U. Aftercare of suicide attempters. In: Bille-Brahe U, Schiödt H,

Eds. 4th European Symposium on Suicidal Behaviour. *The Odense suicide prevention project.*

Intervention and Prevention. Odense: University Press; 1994. p. 177-84.

101.   Borg S. Active intervention and comprehensive services following suicide attempts: a

community-based program. In: Bille-Brahe U, Schiödt H, Eds. 4th European Symposium on

Suicidal Behaviour. *The Odense suicide prevention project. Intervention and prevention*.

Odense: University Press; 1994. p. 159-67.

102.   Motto J, Boström AG. Post-crisis suicide preventing without therapy. *Alcohol Dis*. 1993;

5: 564-7.

103.   O'Brien G, Holton AR, Hurren K, Watt L, Hassanyeh F. Deliberate self-harm and

predictors of out-patient attendance. *Br J Psychiatry*. 1987; 150: 246-7.

104.   Corry M, Clarke M, While AE, Lalor J. Developing complex interventions for nursing: a

critical   review   of   key   guidelines.   *J   Clin   Nurs*.   2013;22(17-18):2366-86.

http://dx.doi.org/10.1111/jocn.12173

105.   Serrano-Gimeno V, Diestre A, Agustin-Alcain M, *et al*. Non-fatal suicide behaviours

across phases in the COVID-19 pandemic: a population-based study in a Catalan cohort.

Lancet Psychiatry. 2024; 11(5):348-358. http://dx.doi.org/10.1016/S2215-0366(24)00065-8

106.   Goñi-Sarriés A, Zandio-Zorrilla M. Suicide: The Challenge of Prevention. *Anales Sis San*

Navarra. 2017;40(3):335–8. https://dx.doi.org/10.23938/assn.0122

107.   Pérez V, Elices M, Prat B, *et al*. The Catalonia Suicide Risk Code: A secondary prevention

program   for   individuals   at   risk   of   suicide.   *J   Affect   Disord*.   2020;268:201-205.

https://doi.org/10.1016/j.jad.2020.03.009

108.   Comendador Vázquez L, Cebrià Meca A, Sanz Ruíz A, et al. Efficacy of synchronous

remote-based interventions for suicide prevention among adolescent and adult patients: a

systematic review and meta-analysis. *Eur Psychiatr* 2022;65:S295–6.

109.   [Dataset] Comendador L; Jiménez-Villamizar MP; Sanabria-Mazo, JP, et al. Data from:

Effect of synchronous remote-based interventions on suicidal behaviours: systematic review

and meta-analysis. *CORA Repository*, February 9, 2025. https://doi.org/10.34810/data1310

A RoB 2 for randomised controlled trials

	D1	D2	D3	D4	D5	Overall bias
Amadéo et al. 2015	+	?	?	+	+	+
Bertolote et al. 2010	+	+	+	+	?	+
Burback et al. 2024	+	+	+	?	+	+
Cedereke et al. 2002	+	?	–	+	?	?
Christensen et al. 2013	+	?	–	+	+	?
Comtois et al. 2019	+	–	?	+	?	?
Fleischmann et al. 2008	+	+	?	+	+	+
Hassanzadeh et al. 2010	+	+	+	+	?	+
Malakouti et al. 2021	+	?	+	+	–	?
Messiah et al. 2019	+	–	+	+	?	?
Mouaffak et al. 2015	+	?	+	+	?	+
Mousavi et al. 2014	+	+	+	?	?	+
Mousavi et al. 2016	+	–	?	–	?	–
O'Connor et al. 2022	+	?	–	+	+	?
Riblet et al. 2022	+	+	+	+	–	?
Vaiva et al. 2006	+	+	–	+	?	–
Vaiva et al. 2018	+	+	–	+	+	?
Ward-Ciesielski et al. 2017	+	+	–	+	+	?
Wei et al. 2012	+	–	–	+	?	–

RoB domains:
D1: Bias arising from the randomisatio
D2: Bias due to deviations from intend
D3: Bias due to missing outcome data.
D4: Bias in measurement of the outcor
D5: Bias in selection of the reported re

Judgement:
+ Low risk of bias.
? Moderate risk of bias.
– High risk of bias.

B ROBINS-I for non-randomised studies

	D1	D2	D3	D4	D5	D6	D7	Overall bias
Cebrià et al. 2013	–	+	+	+	+	?	+	?
Cebrià et al. 2015	–	?	+	+	+	?	+	–
Demesmaeker et al. 2024	?	+	?	+	+	+	?	?
Exbrayat et al. 2017	?	+	+	+	?	?	+	?
Gabilondo et al. 2020	–	?	?	–	–	?	+	–
Goñi-Sarriés et al. 2022	–	+	+	?	?	+	?	–
López-Goñi & Goñi-Sarriés 2021	–	+	+	?	?	?	?	–
Miller et al. 2017	–	+	+	+	+	+	?	?
Plancke et al. 2020	?	+	+	+	+	+	?	+

Study	Hedges' g	CI Lower limit	CI Upper limit	Weight	Q	P_Q	I²	tau ²	-1,50	-1,00	-0,50
Burback et al. 2024	0,23	-0,40	0,86	24,42%							
Christensen et al. 2013	-0,78	-1,31	-0,25	27,64%							
Riblet et al. 2022	0,05	-0,85	0,95	17,58%							
Ward-Ciesielski et al. 2017	-0,06	-0,52	0,40	30,36%							
1 month	**-0,17**	**-0,62**	**0,28**	**100,00%**	**7,38**	**0,06**	**59,35%**	**0,13**			
Burback et al. 2024	0,12	-0,58	0,81	27,12%							
Riblet et al. 2022	0,18	-0,72	1,09	17,15%							
Ward-Ciesielski et al. 2017	-0,24	-0,72	0,24	55,73%							
3 months	**-0,07**	**-0,34**	**0,20**	**100,00%**	**1,16**	**0,56**	**0,00%**	**0,00**			
Christensen et al. 2013	-0,63	-1,21	-0,04	44,05%							
O'Connor et al. 2022	0,00	-0,38	0,38	55,95%							
6 months	**-0,28**	**-0,89**	**0,34**	**100,00%**	**3,26**	**0,07**	**69,28%**	**0,14**			
Cedereke et al. 2002	-0,25	-0,55	0,04	35,33%							
Christensen et al. 2013	-0,74	-1,33	-0,15	15,45%							
Comtois et al. 2019	-0,13	-0,31	0,06	49,22%							
12 months	**-0,27**	**-0,56**	**0,03**	**100,00%**	**4,16**	**0,13**	**51,92%**	**0,03**			

Study	Risk Ratio	CI Lower limit	CI Upper limit	Weight	Q	p_Q	I^2	tau^2	0,01	0,05	0,20
Burback et al. 2024	1,10	0,16	7,51	2,79%							
Exbrayat et al. 2017	0,71	0,51	0,98	90,83%							
Mousavi et al. 2014	0,34	0,01	8,39	0,96%							
Mousavi et al. 2016	0,90	0,02	47,89	0,64%							
Riblet et al. 2022	1,00	0,02	59,73	0,66%							
Ward-Ciesielski et al. 2017	1,06	0,22	5,02	4,12%							
1 month	**0,73**	**0,62**	**0,85**	**100,00%**	**0,70**	**0,98**	**0,00%**	**0,00**			
Burback et al. 2024	0,22	0,03	1,84	25,33%							
Mousavi et al. 2016	0,30	0,01	7,59	10,78%							
Riblet et al. 2022	1,00	0,02	59,73	7,33%							
Ward-Ciesielski et al. 2017	0,84	0,18	4,00	45,94%							
Wei et al. 2012	0,36	0,01	9,02	10,63%							
3 months	**0,50**	**0,21**	**1,16**	**100,00%**	**1,32**	**0,86**	**0,00%**	**0,00**			
Demesmaeker et al. 2024	0,26	0,12	0,55	12,31%							
Hassanzadeh et al. 2010	1,21	0,73	2,03	14,48%							
Messiah et al. 2019	0,84	0,57	1,25	15,54%							
Mousavi et al. 2014	0,25	0,03	2,26	3,93%							
Mousavi et al. 2016	0,21	0,02	1,84	4,08%							
O'Connor et al. 2022	1,11	0,55	2,23	12,84%							
Plancke et al. 2020	0,31	0,25	0,38	16,74%							
Vaiva et al. 2018	0,73	0,53	1,00	16,10%							
Wei et al. 2012	0,24	0,03	2,16	3,98%							
6 months	**0,56**	**0,34**	**0,95**	**100,00%**	**54,92**	**0,00**	**85,43%**	**0,36**			
Cebria et al. 2013	0,38	0,21	0,68	10,00%							
Comtois et al. 2019	0,60	0,36	1,01	10,55%							
Gabilondo et al. 2020	1,14	0,63	2,05	9,91%							
Goñi-Sarriés et al. 2022	0,89	0,58	1,36	11,32%							
López-Goñi & Goñi-Sarriés 2021	1,17	0,81	1,68	11,73%							
Malakouti et al. 2021	0,42	0,23	0,76	9,95%							
Miller et al. 2017	0,80	0,63	1,02	12,53%							
Mouaffak et al. 2015	1,03	0,59	1,80	10,20%							
Plancke et al. 2020	0,36	0,30	0,42	12,96%							
Wei et al. 2012	0,76	0,01	39,95	0,84%							
12 months	**0,68**	**0,49**	**0,96**	**100,00%**	**72,63**	**0,00**	**87,61%**	**0,27**			
Cedereke et al. 2002	1,00	0,51	1,95	6,53%							
Messiah et al. 2019	1,00	0,73	1,37	28,76%							
Vaiva et al. 2006a	0,55	0,31	0,96	9,36%							
Vaiva et al. 2006b	0,76	0,46	1,25	11,63%							
Vaiva et al. 2018	0,82	0,63	1,06	43,72%							
13 months	**0,84**	**0,66**	**1,06**	**100,00%**	**3,89**	**0,42**	**0,00%**	**0,00**			
Amadéo et al. 2015	1,27	0,76	2,13	30,14%							
Bertolote et al. 2010	1,02	0,73	1,43	69,86%							
18 months	**1,09**	**0,30**	**3,91**	**100,00%**	**0,49**	**0,48**	**0,00%**	**0,00**			

Study	Peto Odds Ratio	CI Lower limit	CI Upper limit	Weight	Q	p_Q	I^2	tau^2	0,00	0,001	0,05
Hassanzadeh et al. 2010	0,97	0,14	6,96	30,94%							
Vaiva et al. 2018	0,27	0,08	0,88	69,06%							⊢
6 months	**0,40**	**0,00**	**789,93**	**100,00%**	**1,22**	**0,27**	**17,86%**	**0,15**	⊢———————		
Comtois et al. 2019	1,91	0,20	18,58	17,70%							
Goñi-Sarriés et al. 2022	0,45	0,05	4,49	17,53%							⊢—
López-Goñi & Goñi-Sarriés 2021	0,30	0,05	1,78	29,34%							⊢—
Miller et al. 2017	1,48	0,25	8,59	29,51%							
Mouaffak et al. 2015	7,29	0,14	373,41	5,93%							
12 months	**0,87**	**0,24**	**3,15**	**100,00%**	**3,63**	**0,46**	**0,00%**	**0,00**			
Vaiva et al. 2006a	0,23	0,01	4,51	11,44%						⊢———	
Vaiva et al. 2006b	1,07	0,09	12,27	17,09%							⊢
Vaiva et al. 2018	0,37	0,11	1,23	71,47%							
13 months	**0,42**	**0,11**	**1,65**	**100,00%**	**0,77**	**0,68**	**0,00%**	**0,00**			
Amadéo et al. 2015	0,15	0,01	2,44	9,12%						⊢———	
Fleischmann et al. 2008	0,19	0,08	0,45	90,88%							⊢
18 months	**0,18**	**0,08**	**0,44**	**100,00%**	**0,03**	**0,87**	**0,00%**	**0,00**			⊢