Open source machine learning pipeline automatically flags instances of acute respiratory distress syndrome from electronic health records

Félix L. Morales[1][†], Feihong Xu[2], Hyojun Ada Lee[1], Helio Tejedor Navarro[1,3], Meagan A. Bechel[4,5], Eryn L. Cameron[6], Jesse Kelso[6], Curtis H. Weiss[1,7], Luís A. Nunes Amaral[1,3,8,9]*

[1] Department of Chemical and Biological Engineering, Northwestern University, Evanston, IL

[2] Interdepartmental Biological Sciences Program, Northwestern University, Evanston, IL

[3] Northwestern Institute on Complex Systems, Northwestern University, Evanston, IL

[4] Medical Scientist Training Program, Northwestern University Feinberg School of Medicine, Chicago, IL

[5] Department of Radiology, Emory University, Atlanta, GA

[6] Department of Medicine, Endeavor Health, Evanston, IL

[7] Division of Pulmonary and Critical Care Medicine, Endeavor Health, Evanston, IL

[8] Department of Physics and Astronomy, Northwestern University, Evanston, IL

[9] Department of Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL

[†] Current affiliation: Vizient, Inc. Chicago, IL

* Corresponding Author: Luís A. Nunes Amaral (amaral@northwestern.edu)

Abstract

Physicians could greatly benefit from automated diagnosis and prognosis tools to help address information overload and decision fatigue. Intensive care physicians stand to benefit greatly from such tools as they are at particularly high risk for those factors. Acute Respiratory Distress Syndrome (ARDS) is a life-threatening condition affecting >10% of critical care patients, and has a mortality rate over 40%. However, recognition rates for ARDS have been shown to be low (30-70%) in clinical settings. In this work, we present a reproducible computational pipeline that automatically adjudicates ARDS on retrospective datasets of mechanically ventilated adult patients. This pipeline automates the steps outlined by the Berlin Definition through implementation of natural language processing tools and classification algorithms. We train an XGBoost model on chest imaging reports to detect bilateral infiltrates, and another on a subset of attending physician notes labeled for the most common ARDS risk factor in our data. Both models achieve high performance—a minimum area under the receiver operating characteristic curve (AUROC) of 0.86 for adjudicating chest imaging reports in out-of-bag test sets, and an out-of-bag AUROC of 0.85 for detecting a diagnosis of pneumonia. We validate the entire pipeline on a cohort of MIMIC-III encounters and find a sensitivity of 93.5%—an extraordinary improvement over the 22.6% ARDS recognition rate reported for these encounters—along with a specificity of 73.9%. We conclude that our reproducible, automated diagnostic pipeline exhibits promising accuracy, generalizability, and probability calibration, thus providing a valuable resource for physicians aiming to enhance ARDS diagnosis and treatment strategies. We surmise that proper implementation of the pipeline has the potential to aid clinical practice by facilitating the recognition of ARDS cases at scale.

Introduction

Physicians, including intensivists, process large amounts of dispersed information on many patients. This potential information overload poses serious risks to patient safety. Several studies[1–3] have estimated that 100,000-400,000 fatalities per year may be due to medical errors. While information overload is a challenge for humans, vast amounts of information become advantageous if used as an input for machine learning (ML) approaches. Recent advances in artificial intelligence, ML, and data science are enabling the development of protocols to extract knowledge from large datasets. However, some of those approaches lack interpretability and have been shown to be fragile (e.g., recent re-analysis of attempts to diagnose COVID-19 from chest X-ray images[4]).

In this study, we report the development and validation of a ML pipeline to help physicians adjudicate acute respiratory distress syndrome (ARDS). ARDS, a syndrome of severe acute hypoxemia resulting from inflammatory lung injury[5,6], is an ideal case for the development of a diagnostic aid tool. ARDS recognition requires physicians to synthesize information from multiple distinct data streams and determine whether it fits a standard definition of ARDS. The criteria of the clinically-based Berlin Definition of ARDS include quantitative data ($p_aO_2:F_IO_2 \leq 300$ mm Hg), unstructured data (bilateral opacities on chest imaging), and subjective data (assessing for the presence of ARDS risk factors and cardiac failure)[5]. Despite ARDS' high prevalence, morbidity, and mortality, prior research has shown that many patients with ARDS are not recognized by their treating physicians[6,7]. The poor recognition rate of ARDS is at least partially due to the difficulty in evaluating the Berlin Definition criteria, which requires the physician to access laboratory data, chest images or radiology notes, other physicians' notes, and echocardiographic data or notes, then apply the Berlin criteria to determine whether ARDS is present. Under-recognition of ARDS plays an important role in under-utilization of evidence-based ARDS treatment (e.g., low tidal volume ventilation and prone positioning), even when physicians believe these interventions are warranted[8]. An automated approach to help identify ARDS diagnostic criteria has the potential to be a powerful aid to physician decision-making, leading to improved ARDS recognition and therefore improved ARDS management.

Previous studies have demonstrated some success in automating the recognition of individual ARDS diagnostic components using electronic health record (EHR) screening "sniffers"[9–11]. In addition, a ML algorithm to risk stratify patients for ARDS using structured clinical data derived from the EHR was shown to have good discriminative performance[12]. Regarding automating the entire ARDS diagnostic algorithm, two studies[13,14] have recently reported implementation of keyword search (i.e. rule-based approach) in the EHR with validation conducted for 100 intensive care unit (ICU) admissions from a single time period and from a single institution. A third study recently reported on a computable Berlin Definition which employed a previously developed neural network approach to adjudicate chest imaging reports restricted to patients with a single, known ARDS risk factor (COVID-19), with promising performance (93% sensitivity, 92% specificity)[15]. However, no study has successfully and reproducibly automated the entire sequence of steps required by the Berlin Definition of ARDS, or tested the discriminative performance of the tool on a general population of critically ill patients who received invasive mechanical ventilation. Addressing this gap is the goal of this study.

Data and Methods

Cohort Development and Data Collection

We developed three patient cohorts for this study: Cohort MC1-T1, Cohort MC1-T2, and Cohort MC2-T3, where MC indicates the "Medical Center" and T indicates the "Time Period." In addition, we obtained data from the Medical Information Mart for Intensive Care (MIMIC-III) database [16,17]. MIMIC-III is a large, single-center database of critically ill patients at a tertiary care medical center. It includes all the components necessary to identify ARDS and therefore apply our pipeline, and the data is freely available. For all four cohorts, patients were included if they were at least 18 years old; were admitted to an adult ICU; and had acute hypoxemic respiratory failure requiring invasive mechanical ventilation (at least one recorded $P_aO_2/F_IO_2 \leq 300$ mm Hg while receiving positive-end expiratory pressure ≥ 5cm H_2O)[7]. The study was approved by the Institutional Review Boards of Northwestern University

(STU00208049) and Endeavor Health (EH17-325). Table 1 summarizes the data collected for the four cohorts.

Table 1. Cohort characteristics and data annotation. MC refers to "Medical Center", T refers to "Time Period".

Source	Report type	Number	% bilateral infiltrates	Rater IDs
MC1-T1	Chest imaging reports	5,839 reports (800 patients)	60%	1
MC1-T1	Attending physician notes	12,582 notes (790 patients)		
MC1-T1	Adjudicated Attending physician notes	2,034 notes (400 patients)		1, 5
MC1-T1	Echocardiogram reports	1,006 reports (681 patients)		
MC1-T2	Chest imaging reports	6,040 reports (749 patients)	44%	1
MC2-T3	Chest imaging reports	631 reports (90 patients)	34%	1, 2, 3, 4
MIMIC-III	Chest imaging reports	975 reports (100 patients)	22%	1,6
MIMIC-III	Attending physician notes	887 notes (100 patients)		1,6
MIMIC-III	Echocardiogram reports	89 reports (100 patients)		1,6

Cohort MC1-T1

We previously characterized a cohort of 943 patients, which we denote here as MC1-T1, who met the above inclusion criteria at a single academic medical center between June and December, 2013. We collected the following data: all P_aO_2/F_IO_2 ratios; the unstructured text of all radiologist reports for chest imaging (radiographs and CT scans), critical care attending physician notes, and echocardiogram reports;

and B-type natriuretic peptide (BNP) values obtained from hospital admission to the earliest of extubation, death, or discharge. Data were reviewed by study personnel to determine whether each individual Berlin Definition criterion was present, and whether all criteria taken together were consistent with a diagnosis of ARDS [7].

We collected 5,839 chest imaging reports from 800 Cohort MC1-T1 patients. Study personnel adjudicated 57% of these chest imaging reports as describing bilateral infiltrates consistent with the Berlin Definition[5]. We developed our machine learning (ML) approach to bilateral infiltrate adjudication using these Cohort MC1-T1 reports.

For 790 of the 800 Cohort MC1-T1 patients with a chest imaging report, we also had at least one attending physician note. We collected 12,582 attending physician notes for these patients, of which 2,034 notes from a subset of 400 patients were annotated by study personnel for the presence of ARDS risk factors (e.g., pneumonia, sepsis, aspiration, etc.)[5]. We used this annotated subset of 2,034 notes to develop our ML and regular expression (regex) approach for finding ARDS risk factors and cardiac failure language in attending physician notes.

We collected 1,006 echocardiogram (echo) reports from 681 Cohort MC1-T1 patients. Study personnel from a prior analysis[7] text-matched and adjudicated each echo report for the presence or absence of left ventricular ejection fraction < 40%, cardiopulmonary bypass at time of echo, left ventricular hypertrophy, left atrial dimension > 4cm or left atrial volume index > 28 mL/m^2, and Grade II or III diastolic dysfunction. Separately, 35 BNP values were included for 32 patients in Cohort MC1-T1. We used echo reports and BNP values to develop our objective cardiac failure rule-out approach.[7]

Since identification of Berlin Definition-consistent bilateral infiltrates is the most challenging task[11,18] in our computational pipeline, we analyzed the chest imaging reports from two additional cohorts of patients to test our ML algorithm, Cohort MC1-T2 and Cohort MC2-T3.

Cohort MC1-T2

Cohort MC1-T2 comprises 749 patients admitted during 2016 at the same medical center as Cohort MC1-T1, and meeting the same inclusion criteria. We collected 6,040 chest imaging reports for

these patients. Study personnel adjudicated 44% of Cohort MC1-T2 reports as describing bilateral infiltrates. We used this cohort only to train a second ML algorithm for bilateral infiltrate adjudication.

Cohort MC2-T3

Cohort MC2-T3 comprises 90 patients admitted to a different medical center in 2017-2018, and meeting the same inclusion criteria as Cohorts MC1-T1 and MC1-T2. We collected 631 chest imaging reports for these 90 patients. Study personnel adjudicated 34% of these chest imaging reports as describing bilateral infiltrates. We used these reports from Cohort MC2-T3 only to test ML algorithms for bilateral infiltrate adjudication.

MIMIC-III

We identified the set of patients in the MIMIC-III dataset who satisfied the inclusion criteria used to develop cohort MC1-T1. This resulted in a set comprising 3,712 encounters. We then used our automated pipeline to adjudicate the presence or absence of ARDS for all those encounters, and randomly selected a balanced cohort comprising 100 encounters, which we denote as the MIMIC-III cohort. Each of the encounters in the MIMIC-III cohort was adjudicated by one critical care physician and one internal medicine physician for whether each individual Berlin Definition criterion was present, and whether all criteria taken together were consistent with a diagnosis of ARDS. This cohort, with physician adjudications, is publicly available at Northwestern's ARCH database.

The records of these 100 MIMIC-III patients included 975 chest imaging reports, 887 attending physician notes, and 89 echocardiogram (echo) reports. The critical care physician adjudicated 22.3% of these chest imaging reports as describing bilateral infiltrates consistent with the Berlin Definition[5]. The same individual also annotated 887 attending physician notes for the presence of ARDS risk factors[5] and cardiac failure language, and 89 echo reports for the presence or absence of: left ventricular ejection fraction < 40%, cardiopulmonary bypass at time of echo, left ventricular hypertrophy, left atrial dimension > 4cm or left atrial volume index > 28 mL/m^2, and Grade II or III diastolic dysfunction.[7] We used these adjudicated datasets to evaluate the performance of our ML algorithm.

Analysis

Adjudication of bilateral infiltrates from chest imaging reports

We preprocessed chest imaging reports to remove patient information, non-informative sections (e.g. technique, indication, history, etc.), and non-informative words. We then tokenized the remaining sections (i.e., separated the text into sets of unigrams and bigrams) and prepared the data for use of a "bag of words" approach (i.e, we vectorized these tokens according to their counts in the imaging reports). When training a ML model on a given corpus, we used the 200 most frequently appearing tokens across the imaging reports in the respective corpus as model features.

We trained four different ML models (decision trees, logistic regression, random forest classifiers, and extreme gradient boosting 'XGBoost'[19]) on chest imaging reports from Cohort MC1-T1, which was also used to perform hyperparameter tuning for the four models. We performed hyperparameter tuning using Bayesian optimization, which is available through the *hyperopt* package (v.0.2.7)[20] for Python (v.3.10.12). For each model, we performed 5-fold cross-validation to obtain the mean Area under the Receiver Operating Characteristic (AUROC) curve for each hyperparameter combination considered. We then selected the optimal combination of hyperparameters as the one yielding the highest 5-fold cross-validation mean AUROC after at least 100 iterations. We also derived another set of optimal hyperparameters for XGBoost trained with chest imaging reports from Cohort MC1-T2 in the same fashion.

Unless otherwise noted, all cross-validation strategies used healthcare encounters (a.k.a, patient admissions), not individual reports. We split the reports this way to avoid having chest imaging reports from the same encounter found in both the training and validation data (a problem known as "data leakage"). Thus, we ensured all reports from a given encounter can only be found on either the training or validation data (but not both). We also used nested cross-validation to prevent data leakage, as this avoids tuning hyperparameters on validation data.

For comparing the performance of the four models, we used nested cross-validation by doing 5-fold cross-validation to obtain a mean AUROC across five different folds. Furthermore, each fold's

training set was used to tune that model's hyperparameters as described above (i.e., five separate hyperparameter tuning exercises). However, we note that we used a 3-fold cross-validation strategy for this tuning due to the computational cost of running nested cross-validation. We repeated this nested cross validation 10 times, each time with a different resampling with replacement of the data (i.e. a bootstrap) to yield a distribution of mean AUROCs on test sets.

We used 95% confidence intervals to compare ROC curves and AUROCs across different models. For obtaining feature/token importance during training, we employed the default "importance" method that version 1.1.3 of the scikit-learn package implements for decision tree, logistic regression, and random forest algorithms, and version 1.7.4 of *xgboost* package for XGBoost. For decision tree and random forest, feature importance corresponds to the mean decrease in Gini impurity; for logistic regression, importances correspond to the mean value of coefficients in the fitted linear equation; and for XGBoost, the importance corresponds to the mean gain in predictive performance obtained by including a particular feature in the trees.

We also evaluated the inter-rater disagreement rate for chest imaging reports from MIMIC III. For this purpose, we obtained two independent adjudications (one critical care physician and one internal medicine physician) for 975 reports available, and split imaging reports into three groups according to XGBoost output probabilities. For each group, we then calculated the fraction of imaging reports for which these independent raters disagreed on their adjudications.

Finally, to assess how our XGBoost implementation for chest imaging reports generalizes to other datasets, we tuned hyperparameters and then trained XGBoost models on all chest imaging reports from Cohort MC1-T1 and MC1-T2, separately. We then tested each of the two models on the two other chest imaging corpora the model had not yet seen by comparing the AUROC values. We used 100 bootstrapped samples to gather 95% confidence intervals for the mean AUROC values.

Adjudication of risk factors from physician notes

The Berlin Definition of ARDS requires the presence of at least one risk factor—e.g., pneumonia, sepsis, shock, inhalation, pulmonary contusion, vasculitis, drowning, drug overdose—within seven days

of non-cardiogenic acute respiratory failure. We preprocessed attending physician notes from Cohort MC1-T1 to remove identifiable information from the text of these notes. We then used regular expressions (regex v2022.10.31) to match keywords related to risk factor and heart failure language (see SI: Regular expression list 1, for a complete list of risk factors). In order to validate this strategy, we ensured that this regex approach matched 100% of the notes that had a positive adjudication for a particular risk factor (or close to 100% as possible). We also corrected common spelling errors on important keywords, such as 'pneunonia', 'spetic' or 'cardigenic'. To prevent data leakage during ML development, we again split the adjudicated notes into train and test sets by encounter, not note.

Adjudicating the presence of a risk factor is not as simple as finding a particular keyword in a physician note. For example, a note stating "patient is unlikely to have pneumonia" should not be classified as evidence of pneumonia. To account for such possibilities, we implemented a strategy in which, after matching a particular keyword, we extract a text string from the note starting 100 characters prior to the occurrence of the keyword and extending 100 characters post the keyword. Subsequently, we tokenized and vectorized the strings as described in the previous subsection.

Using the vectorized tokens, we trained XGBoost models for a select group of risk/cardiac failure factors, employing a similar nested cross-validation strategy as the one pursued for the adjudication of chest imaging reports (except in this case we used 100 resamples instead of 10). Note that not all risk factors were amenable to a ML approach: we chose risk factors that had more than 100 notes annotated, were risk factors for ARDS (or a cardiac failure criterion), and had relatively balanced yes/no proportions after regex-matching (between 33% and 66%, see SI Table 1). This resulted in the use of 1409 adjudicated attending notes from 337 patients for ML development (see SI Table 1 for a breakdown of notes used for each risk factor/cardiac failure criterion).

Objective adjudication of cardiac failure from echocardiogram reports

We preprocessed echo reports from MC1-T1 to remove identifiable information from text. We then developed regex patterns that first matched keywords associated with the following parameters: left ventricular ejection fraction, cardiopulmonary bypass, left atrial diameter, left atrial volume index, left

ventricular hypertrophy, and grade II or III diastolic dysfunction (see SI: Regular expression list 2 for regex patterns). Once these parameters were found in the echo report text, we then extracted numerical values of numerical variables (left ventricular ejection fraction, left atrial diameter, and left atrial volume index), and the matched text otherwise.

Design of ARDS adjudication pipeline: chaining Berlin Definition steps

Each of the steps outlined above automates the adjudication of specific criteria in the Berlin Definition, with the modifications specified previously[7]. We integrate the different criteria into a single pipeline to build an automated ARDS adjudication pipeline.

The ARDS adjudication pipeline first flags encounters with at least one hypoxemia measurement (i.e., one instance of PF ratio $\leq$ 300 mm Hg while PEEP $\geq$ 5 cm H20), and then uses the predictions of an XGBoost model trained on chest imaging reports from MC1-T1 to adjudicate presence of bilateral infiltrate language in chest imaging reports. Upon settling these two criteria, the pipeline flags whether the hypoxemia record and the report consistent with bilateral infiltrates have timestamps within 48 hours of each other (which we term "qualified hypoxemia"). In addition, at this step the pipeline ensures that the hypoxemia record was taken at or after intubation. Next, the pipeline uses the predictions of an XGBoost model trained on attending physician notes that have pneumonia keywords to adjudicate pneumonia on all notes, and uses regex to flag presence of other risk factors, cardiac failure language (e.g., cardiac arrest), and indicators of cardiogenic and noncardiogenic language. Finally, the pipeline flags whether an attending physician note has a timestamp that falls between one day prior to and seven days after the latter timestamp of any of the qualifying hypoxemia-bilateral infiltrates pairs. Once these annotations are integrated, the pipeline proceeds to adjudicate whether an ARDS diagnosis is warranted.

If any risk factor is identified in this time window for an encounter, via XGBoost or regex, the pipeline adjudicates the encounter as an ARDS case. If no risk factors are identified, but cardiac failure language is identified in the notes using regex, the pipeline adjudicates the encounter as a "No ARDS" case.

For all other encounters not meeting the risk factor or cardiac failure language criteria described above, the pipeline flags the case for objective cardiac failure assessment[7]. This assessment is done sequentially instead of by flagging. If any encounter had BNP greater than 100 pg/mL (an indicator of heart failure), the pipeline adjudicates "No ARDS" for such an encounter. The pipeline then considers the remaining encounters for each subsequent criteria, adjudicating "No ARDS" if the encounter had any of the following: left ventricular ejection fraction < 40%, cardiopulmonary bypass found in echocardiogram report, or at least two of the following present in the echocardiogram report: (i) left atrial diameter > 4 cm or left atrial volume index > 28 mL/m2, (ii) left ventricular hypertrophy, or (iii) Grade II or III diastolic dysfunction.

Any encounter that is not ruled out for ARDS after the objective cardiac failure assessment step is adjudicated as an ARDS case. That is, the pipeline adds these encounters to those adjudicated as positive for ARDS via risk factor identification.

Results

Adjudication of bilateral infiltrates

Figure 1a shows the Receiver Operator Characteristic (ROC) curves for the decision tree model applied to chest imaging reports from Cohort MC1-T1. We quantify model predictive performance using the areas under the ROC curves (AUROCs). We observe that once hyperparameters for each model are optimized, all models trained on chest imaging reports from MC1-T1 achieve AUROCs of at least 0.90 on the training set (Decision tree: AUROC = 0.89, 95%CI = [0.89, 0.91]; logistic regression: AUROC = 0.92, 95%CI = [0.91, 0.93]; random forest: AUROC = 0.94, 95%CI = [0.93, 0.95]; XGBoost: AUROC = 0.95, 95%CI = [0.94, 0.95]) (Fig. 1b).

We next calculated the importance that each model assigned to the 200 tokens used as features. Reassuringly, we find that the four models consistently identify tokens such as edema, bilateral, clear, and atelectasis as the most predictive (Fig. 1c). These tokens correspond closely to the inclusion/exclusion

language we developed to address Berlin Definition shortcomings[7], which we also observed when

implementing Shapley-additive explanations (SHAP) values to assess feature importance (SI Fig. 1).

Figure 1. Machine learning (ML) models achieve high-performance in adjudicating the presence of bilateral infiltrates from chest imaging reports. Error bars show 95% confidence intervals for estimates of the mean obtained using bootstrapping. **a)** Receiver operating characteristic (ROC) curve for the decision tree model trained on chest imaging reports from Cohort MC1-T1. **b)** Bootstrapped mean areas under the ROC (AUROCs) show that all four ML approaches yield AUROCs greater or equal to 0.90. **c)** Feature importances for the four different ML approaches considered. Features in bold are highly ranked in importance in at least 3 of the 4 approaches.

We then assessed how calibrated were the outcome probabilities from the models by comparing

model outcome probability after training to actual probability of occurrence in considered encounters

from MC1-T1 chest imaging reports. Figures 2a-c suggest that logistic regression and XGBoost models output probabilities that are well calibrated, which is expected given they both use log loss (a.k.a. cross-entropy) for fitting. In contrast, random forest produces poorly calibrated probabilities, being over-confident when forecasting with confidence levels lower than 50%, and under-confident with confidence levels greater than 50%. We thus select XGBoost as the implemented approach for our pipeline as it offers the highest predictive performance (AUROC = 0.95, 95%CI = [0.94, 0.95]) and well-calibrated forecasts on the training set (Durbin-Watson statistic = 1.55, 95%CI = [1.10-1.87]).

Figure 2. Assessment of ML implementation probabilities. Comparing calibration of MC1-T1 probabilities by **a)** logistic regression, **b)** random forest, and **c)** XGBoost. A perfectly-calibrated model would have a 1:1 relationship between fraction of positive labels and mean probabilities (i.e., it would overlay the diagonal line). The Durbin-Watson statistic, DW, probes for correlations in the residuals, if DW is close to 2, then one can rule out correlations in the residuals, implying good linear behavior. **d)** Comparing inter-rater disagreement rate to the confidence in adjudicating bilateral infiltrates on chest imaging reports from MIMIC III by an XGBoost model trained on chest imaging reports from Cohort MC1-T2.

The good calibration of the prediction of the XGBoost model prompted us to test the hypothesis that the estimated probabilities by the model may contain information regarding the agreement rate of physician adjudications of the same chest imaging reports. To this end, we leveraged MIMIC III report annotations by two independent raters, a critical care physician and an internal medicine physician. We evaluated how the agreement between these two independent raters in adjudicating chest imaging reports from MIMIC III associated with the output probabilities of an XGBoost classifier trained on chest imaging reports from Cohort MC1-T1. Specifically, we binned predictions into three groups: high-confidence 'No' (probability of bilateral infiltrates <10%), high-confidence 'Yes' (probability of bilateral infiltrates >90%), low confidence (all other probabilities of bilateral infiltrates). As seen on Figure 2d, the interrater disagreement was highest (16.6%) for the cases of high-confidence 'Yes' predictions. This suggests that the model could be an effective way to avoid false negatives by a physician's mis-interpretation of chest imaging reports, and can be used to alert that there is a high chance of a report being consistent with bilateral infiltrates. It also suggests that this model is most reliable when it indicates a low probability of a chest imaging report showing bilateral infiltrates.

Figure 3. Evaluation of XGBoost's generalization performance to MC2-T3 and MIMIC-III cohorts. We show AUROCs with bootstrapped 95% confidence intervals as error bars for XGBoost models trained on **a)** chest imaging reports from cohort MC1-T1, and **b)** chest imaging reports from cohort MC1-T2.

To assess how a model developed for a specific cohort generalizes to a different health system dataset, we tested this MC1-T1-trained XGBoost model on chest imaging reports from Cohort MC2-T3 and MIMIC-III (Fig. 3a). The MC1-T1-trained XGBoost model yields AUROCs of 0.88 (95%CI = [0.84, 0.91]) and 0.87 (95%CI = [0.83, 0.90]) when applied to Cohort MC2-T3 and MIMIC-III, respectively. We also trained a second XGBoost model on chest imaging reports from Cohort MC1-T2, and tested this model against chest imaging reports from MC2-T3 and MIMIC-III. We found similar results to the first XGBoost model, with an AUROC of 0.90 (95%CI = [0.87, 0.92]) when applying this MC1-T2-trained model to chest imaging reports from MC2-T3, and an AUROC of 0.86 (95%CI = [0.82, 0.89]) when applying this model to chest imaging reports from MIMIC-III (Fig. 3b).

Figure 4. Confusion matrices for the performance of regex approach to capture risk factors in attending physician notes. 'Sepsis' and 'shock' are the most prevalent risk factors for ARDS after pneumonia. 'Cardiogenic' and 'cardiac arrest' are the most prevalent cardiac failure keywords in attending physician notes. Notice the absence of false negatives, which indicates that regex-matching can capture all instances in which a physician adjudicated the language as being present in the attending physician note.

Extracting ARDS risk factors in attending physician notes

We first developed regex patterns to match keywords for the risk factors. As shown in Fig. 4, the developed regex patterns match 100% of the notes that were annotated 'yes' for a particular risk factor in MC1-T1. When we count the total number of notes annotated as either yes or no, the most prevalent matches were sepsis (744 notes annotated vs. 748 notes regex-matched), pneumonia (636 notes annotated vs. 955 regex-matched), and shock (604 notes annotated vs. 607 regex-matched). For the cardiac failure criteria, the relevant matches were the cardiogenic keyword (176 notes annotated vs. 725 regex-matched) to qualify the matching of shock, and congestive heart failure (254 notes annotated vs. 352 regex-matched). We thus feel confident that the built regex-patterns can match nearly the entirety of the notes annotated as "yes" for specific risk factors.

Next, we trained an XGBoost model on attending physician notes from Cohort MC1-T1 to adjudicate pneumonia, aspiration, congestive heart failure, and sepsis. We chose these risk factors for ML because at least 100 attending physician notes were annotated for them, and their annotations have relatively balanced yes/no proportions after regex-matching (between 33% and 66%, except for sepsis; see Table S1). We did not use all 2,034 records from Cohort MC1-T1 to train each of the models since not every record had an annotation or a keyword for a given risk factor. For instance, only 636 notes in Cohort MC1-T1 included an annotation for pneumonia, whereas we were able to match 955 notes for pneumonia using regex. Therefore, our training dataset for each of the three models consisted of all notes that were regex-matched for that particular risk factor. For notes that were regex-matched but did not have an annotation, we imputed the annotation as 'No', or zero.

We used nested cross-validation for every XGBoost implementation (pneumonia, sepsis, etc.). This involved splitting data into a train and test set, tuning hyperparameters on the train set using 3-fold cross validation, and measuring AUROC on the test set. Since we used 5-fold cross-validation, this process was repeated 5 times per bootstrapped sample, yielding 5 AUROCs. We repeated the above process for a total of 100 bootstrapped data samples to evaluate the mean AUROCs obtained by each of the models. We observed that out of the three XGBoost implementations, the pneumonia model yielded

the best discriminative performance during cross-validation (Pneumonia: AUROC = 0.93, 95%CI = [0.90, 0.95]; CHF: AUROC = 0.82, 95%CI = [0.75, 0.88]; Aspiration: AUROC = 0.73, 95%CI = [0.60, 0.83]; Sepsis: AUROC = 0.76, 95%CI = [0.66, 0.85]; Fig. 5a), the best calibration (DW = 1.70, 95%CI = [1.13-2.22]; Fig. 5b), and the overall better generalizability to attending notes from MIMIC-III (Pneumonia: AUROC = 0.86, 95%CI = [0.81, 0.89]; Aspiration: AUROC = 0.42, 95%CI = [0.31, 0.51]; Sepsis: AUROC = 0.81, 95%CI = [0.70, 0.91]; Fig. 5c). This is in stark contrast to the XGBoost models for congestive heart failure, aspiration, and sepsis: These models have underwhelming performance on cross-validation, generalizability to MIMIC-III, and turn out poorly calibrated on attending notes from MC1-T1 (Fig. 5d). For these reasons, we decided against integrating these three ML models into our pipeline.

Instead, we use regex-matching and a simple heuristic to adjudicate other ARDS risk factor/cardiac failure language in Cohort MC1-T1. This is actually not a real weakness since many other types of ARDS risk factor/cardiac failure language are more predictable in their adjudication. For example, of the 105 attending physician notes matching for 'cardiac arrest', 103 were annotated as 'yes' (Table S1). Thus, our heuristic was to adjudicate a risk factor as 'present' if it was annotated as 'yes' in more than 80% of the matched notes. These included shock, cardiac arrest, pulmonary contusion, vasculitis, drowning, and overdose.

Adjudication of cardiac failure from echocardiogram (echo) reports

The criteria for the objective assessment of cardiac failure rely on the following six factors: left ventricular ejection fraction, cardiopulmonary bypass, left atrial diameter, left atrial volume index, left ventricular hypertrophy, and grade II or III diastolic dysfunction. Because echo reports are highly standardized, it is possible to extract these factors from the reports using regex. Moreover, we had access to echo reports from Cohort MC1-T1 which were previously text-matched, enabling us to validate our regex approach.

Figure 5. XGBoost model performance in adjudicating for presence of risk factors in attending physician notes amenable to ML techniques. **a)** Cross-validated performance of XGBoost models trained to adjudicate pneumonia, congestive heart failure, aspiration, and sepsis on MC1-T1 attending notes. **b)** Training set calibration curve for the pneumonia XGBoost model. **c)** Test set performance of XGBoost models trained to adjudicate pneumonia, aspiration, and sepsis using MC1-T1 attending notes. We did not have labels for CHF available for MIMIC-III, therefore we did not explore the generalizability of this model. **d)** Training set calibration curve for the sepsis XGBoost model.

Using the regex patterns listed in the SI, we analyze Cohort MC1-T1's echo reports for the presence or absence of each of the six factors of interest. Figure 6 demonstrates that not all six factors were present in every echo report. For three of the six factors—'left ventricular ejection fraction', 'left atrial dimension/diameter', and 'left atrial volume index'—we found excellent agreement between regex and text-matching. Two of the other three, 'cardiopulmonary bypass' and 'diastolic function', were not

text-matched, so no comparison can be made. For 'left ventricular hypertrophy', the regex-matching procedure correctly captured the desired language, indicating that the original text-matching procedure failed to identify 13 echo reports. In addition, we validated the numerical values extracted through this regex approach by randomly selecting 10% of echo reports for visual inspection of values, and comparing against values extracted through regex. We found 100% concordance between values extracted and those retrieved manually (SI Table 2).

Figure 6. Confusion matrices comparing the flagging performance of regex-matching against text-matching for 'Left ventricular ejection fraction', 'Left atrial dimension/diameter', 'Left atrial volume index', and 'Left ventricular hypertrophy'. Note the large discrepancy for the annotations of 'left ventricular hypertrophy', which is explained in text.

Adjudication of ARDS for entire MC1-T1 cohort

We are now ready to compare the performance of our complete pipeline against the previously reported ARDS adjudication[7]. For the XGBoost model components of the pipeline (chest imaging reports and pneumonia risk factor adjudications), we use a threshold of 50% to map estimated probabilities into binary yes/no decisions.

Figure 7. Machine learning computational pipeline for adjudication of MC1-T1 cohort yields a small fraction of false negatives and a manageable fraction of false positives. a) Flowchart of ARDS diagnosis by computational pipeline (blue) vs. physician (black). **b)** Confusion matrix comparing physician adjudication from previous publication[7] against ML computational adjudication pipeline.

We conduct the evaluation of our pipeline for the 943 patients in the MC1-T1 ARDS adjudication cohort who were 18 years and older, received invasive mechanical ventilation, and had acute hypoxemic

respiratory failure (Fig. 7). 143 patients had no chest imaging report available and were adjudicated as negative for ARDS. The remaining 800 patients had at least one chest imaging report available. The XGBoost model trained on MC1-T1 adjudicated bilateral infiltrates for 529 patients. Of these 529 patients, 448 had at least one hypoxemic event occurring post-intubation and within 48 h of the positive chest imaging report. Finally, 322 of these patients had a risk factor within 7 days of the qualified hypoxemia event and were adjudicated as being positive for ARDS.

The remaining 126 patients were then evaluated for cardiac failure. For 21 patients, the physician notes indicated cardiac failure and they were adjudicated as negative for ARDS. The last 105 patients were then adjudicated using the objective cardiac failure assessment step; 62 were adjudicated to have cardiac failure and thus negative for ARDS and the remaining 43 were adjudicated as positive for ARDS. In total, the pipeline adjudicated 365 patients as positive for ARDS and 578 as negative for ARDS.

In summary, using a simple 50% probability cutoff for both ML algorithms, our pipeline yields close agreement with the physician adjudication of ARDS for this cohort[7] (Fig. 7a). Specifically, the pipeline yields a sensitivity or true positive rate of 93.4% on this cohort, which compares favorably to the 19% ARDS diagnosis rate we found on this cohort[7]. Importantly, this high sensitivity is achieved while maintaining a very low 12.5% rate of false positives.

Adjudication of ARDS for MIMIC-III labeled subset

We applied our automated ARDS adjudication to the 100 patient encounters in the MIMIC-III cohort. We then compared physician adjudication against the pipeline's (Fig. 8). Reassuringly, and again using a simple 50% probability cutoff for both ML algorithms, we find that the overall performance of our pipeline on the MIMIC-III cohort subset is strikingly similar to its performance on the development cohort[7] (Fig. 8b). Specifically, the pipeline yields a sensitivity or true positive rate of 93.5% on this cohort, which compares favorably to the 22.6% ARDS documentation rate we found in this subset. This high sensitivity is achieved while maintaining a relatively low 26.1% rate of false positives. Moreover, the

false negative rate of pipeline adjudication is lower than that of a less experienced physician, highlighting the pipeline's potential to aid physicians in ARDS diagnosis.

Figure 8. Machine learning computational pipeline for adjudication of MIMIC-III cohort yields a small fraction of false negatives and a manageable fraction of false positives. a) Flowchart of ARDS diagnosis by computational pipeline (blue) vs. physician (black). **b)** Confusion matrix comparing physician adjudication (ground truth) against ML computational adjudication pipeline (left panel), and physician adjudication (ground truth) against a less experienced physician adjudication.

Discussion

We believe that computational pipelines aiming to help physicians with the diagnosis of complex conditions must follow two principles. First, they should act as physicians aids, not physician replacements. That is, they should flag a potential diagnosis for consideration by the responsible physician, rather than mandate a diagnosis as certain. Second, and a consequence of the first, they should

provide interpretable insights. Others have pointed out[21] that machine learning (ML) should only be considered as the final decision maker for problems that can be interpreted as deterministic, such as differentiating a dog from a cat in a photo. However, for tasks where the characteristics of the two classes overlap and the outcome of the decision has important consequences, such as medical decision-making, the ML approaches should be used to provide an estimation of probabilities, not a final determination.

In this study, we construct and validate a ML pipeline for automating the adjudication of ARDS according to the Berlin Definition based on data from EHRs. We constructed high performing decision tree-based models (XGBoost) to adjudicate chest imaging reports for bilateral infiltrate language and attending physician notes for the presence of pneumonia. These tree-based methods estimate probabilities, the first stage of any classification problem, and enable physicians to optimize the false positive vs. false negative tradeoff by adjusting the decision cutoff. In addition, our implementation of XGBoost allows language-level interpretation of estimated probabilities, which can enhance physician trust in ML models.

Supplementing the above XGBoost models with regular expressions to identify other ARDS risk factors and cardiac failure, and additional structured data to objectively rule out cardiac failure,[7] enabled the automated adjudication of the complete set of Berlin Definition criteria. This pipeline demonstrated excellent test characteristics, including false negative and false positive rates of 6.9% and 12.4%, respectively, at the 50% decision cutoff. We then validated the generalizability of the pipeline on a subset of the MIMIC-III dataset, demonstrating a similar – high – level of performance.

To our knowledge, this study is the first to automate the entire Berlin Definition process using ML and rules-based methods in a multi-center, open-source, generalizable manner. Previous attempts at automated ARDS adjudication used single-center EHR data to adjudicate individual Berlin criteria or used non-reproducible methods. Afshar et. al. used text features in chest imaging reports for ARDS identification, achieving a maximum AUROC of 0.80 for that task[22]. However, our work identifies ARDS by considering data beyond chest imaging reports. Sathe et. al. developed EHR-Berlin, evaluating the Berlin Definition using ML and rules-based methods, but their focus was limited to COVID-19 patients[15]; by using a cohort of patients who were already defined as having an ARDS risk factor, they effectively

eschewed the need to identify ARDS risk factors or cardiac failure. In contrast, our study considers any

adult patient placed on mechanical ventilation, which requires evaluating all Berlin Definition

components. Finally, Song and Li developed a fully rules-based tool that automates the entire Berlin

Definition, both achieving identical high performance[13,14]. However, their models were constructed within

a single hospital, which may not be reproducible across different health systems. Conversely, the

generalizability and reproducibility of our ARDS adjudication pipeline have been demonstrated

conclusively.

Machine learning offers a powerful solution for efficiently analyzing large volumes of data that

would otherwise require countless hours of human effort. A recent study that combined NLP techniques

with manual chart abstraction to evaluate clinical trial outcomes confirms this assertion. The time spent

on review decreased from 2,000 laborious hours for manual chart abstraction to just 34.3 hours with

NLP-screened manual chart abstraction[23]. Similarly, our pipeline demonstrates the ability to adjudicate

ARDS for multiple cohorts of hundreds of patients in under five minutes by training XGBoost models at

runtime (with even faster runtimes possible using pre-trained models on inference mode).

A potential limitation is that previous studies have shown significant variability in the diagnosis

of ARDS among critical care physicians, particularly in relation to interpretation of chest imaging[18]. We

attempted to mitigate concerns raised by this challenge by only relying on chest imaging reports written

by radiologists. This allowed us to use previously developed Berlin Definition-based inclusion and

exclusion language as a guideline for critical care physicians reviewing chest imaging reports to minimize

inter-rater disagreements[7]. Consequently, we are also able to leverage this inclusion and exclusion

language for the NLP processing of chest imaging reports needed for our ML approach. We believe this

choice allowed us to increase the signal-to-noise ratio of the data used for ML development; however, we

recognize that choosing chest imaging reports over the images themselves might limit implementation of

this pipeline for real-time use. On the other hand, we explored the relationship between interrater

disagreement and ML model confidence. We found that lower disagreement rates among our raters

correlated with relatively lower model confidence of "yes", indicating that our algorithm can confidently

"discard" cases which are not likely to have bilateral infiltrates. Interestingly, we also observed that our raters exhibited higher levels of disagreement when the model had high model confidence of "yes". We speculate this could be attributed to MIMIC III containing more reports that are not consistent with bilateral infiltrates (88%).

Concerning the adjudication of ARDS risk factors, we faced a significant challenge in implementing machine learning techniques for parsing attending physician notes due to the lack of clearly defined inclusion/exclusion language for adjudicating ARDS risk factors. In addition, we only had 744 attending physician notes labeled for sepsis, the most of any risk factor, compared to more than 12,000 labeled chest imaging reports. Nonetheless, it is striking that even a regex approach for this step yielded high overall pipeline performance and a small fraction of false negatives.

A possible path to improve performance could involve refining the note-parsing process using advanced natural language processing techniques such as Med-BERT, cTAKES, or leveraging the capabilities of open-source large language models. The latter technique has the potential to eliminate laborious pre-processing steps and facilitate the development of general-purpose models instead of task-specific ones[23]. This is especially true since we use regex patterns for attending physician notes and echocardiogram reports, which would very likely need redevelopment for each health system in which our pipeline would get implemented. However, while such advanced approaches offer exciting opportunities, we must remain cautious as it is in the interest of patients and physicians to implement approaches that prioritize interpretability and transparency. Not to mention the cost-effectiveness of developing and deploying pipelines such as ours instead of those relying on large language models (in token consumption and/or computational resources, among other costs).

Our pipeline compellingly answers the specific question being posed: can we automate the identification of ARDS in a way that is clinically relevant? In the clinical realm, minimizing the false negative rate (at the expense of a still manageable but higher false positive rate) means applying ARDS treatment to patients who do not have ARDS, which is likely to be less harmful than not treating patients

who do have ARDS but were not recognized as such [26,27]. The pipeline powerfully addresses this clinical goal.

While our pipeline could aid researchers and quality reviewers during retrospective reviews, its greatest potential impact lies in its integration with clinical decision support systems, enabling timely alerts to critical care physicians about the probability of ARDS in their patients. Future studies should evaluate the pipeline in several ways, such as a physician fully trusting the pipeline when ML exhibits high confidence in its probability estimations, or a physician double-checking a case only if their adjudication differs from the pipeline.

To properly implement this pipeline, decision theory considerations should be considered. The XGBoost models in our pipeline generate probabilities that need to be binarized into "yes" or "no" for each component of the Berlin Definition. In this regard, Youden's J statistic is used by some to identify optimal thresholds that yield a good balance between false negatives and false positives[24]. While the use of Youden's J for cutoff determination is common practice in theoretical studies, it assumes a similar degree of undesirability for false positives and false negatives while implicitly using disease prevalence as a cost ratio[25]. We believe that implementation is more likely to be successful if a health system explicitly considers the particulars of ARDS when deciding on optimal probability cutoffs. Given the poor recognition of ARDS in clinical practice, prioritizing comparatively low false negative rates is crucial for making life-saving decisions, such as implementing low tidal volume ventilation and prone-positioning strategies[28]. This benefit must be balanced with the potential risk of alert fatigue caused by excessive false positives[29–31]. An implementation study that explores the attitudes of critical care physicians towards the balance of false positives and false negatives could provide valuable insights for implementing decision-support tools like our pipeline.

References

1. James, J. T. A New, Evidence-based Estimate of Patient Harms Associated with Hospital Care. *J. Patient Saf.* **9**, 122–128 (2013).

2. Makary, M. A. & Daniel, M. Medical error—the third leading cause of death in the US. *BMJ* i2139 (2016) doi:10.1136/bmj.i2139.

3. Landrigan, C. P. *et al.* Temporal Trends in Rates of Patient Harm Resulting from Medical Care. *N. Engl. J. Med.* **363**, 2124–2134 (2010).

4. DeGrave, A. J., Janizek, J. D. & Lee, S.-I. AI for radiographic COVID-19 detection selects shortcuts over signal. *Nat. Mach. Intell.* **3**, 610–619 (2021).

5. The ARDS Definition Task Force*. Acute Respiratory Distress Syndrome: The Berlin Definition. *JAMA* **307**, 2526–2533 (2012).

6. Bellani, G. *et al.* Epidemiology, Patterns of Care, and Mortality for Patients With Acute Respiratory Distress Syndrome in Intensive Care Units in 50 Countries. *JAMA* **315**, 788–800 (2016).

7. Weiss, C. H. *et al.* Low Tidal Volume Ventilation Use in Acute Respiratory Distress Syndrome*. *Crit. Care Med.* **44**, (2016).

8. Weiss, C. H. *et al.* A Critical Care Clinician Survey Comparing Attitudes and Perceived Barriers to Low Tidal Volume Ventilation with Actual Practice. *Ann. Am. Thorac. Soc.* **14**, 1682–1689 (2017).

9. Koenig, H. C. *et al.* Performance of an automated electronic acute lung injury screening system in intensive care unit patients*. *Crit. Care Med.* **39**, (2011).

10. Herasevich, V., Yilmaz, M., Khan, H., Hubmayr, R. D. & Gajic, O. Validation of an electronic surveillance system for acute lung injury. *Intensive Care Med.* **35**, 1018–1023 (2009).

11. Laffey, J. G., Pham, T. & Bellani, G. Continued under-recognition of acute respiratory distress syndrome after the Berlin definition: what is the solution? *Curr. Opin. Crit. Care* **23**, (2017).

12. Zeiberg, D. *et al.* Machine learning for patient risk stratification for acute respiratory distress syndrome. *PLOS ONE* **14**, e0214465 (2019).

13. Song, X., Weister, T. J., Dong, Y., Kashani, K. B. & Kashyap, R. Derivation and Validation of an

Automated Search Strategy to Retrospectively Identify Acute Respiratory Distress Patients Per Berlin Definition. *Front. Med.* **8**, (2021).

14. Li, H. *et al.* Rule-Based Cohort Definitions for Acute Respiratory Distress Syndrome: A Computable Phenotyping Strategy Based on the Berlin Definition. *Crit. Care Explor.* **3**, e0451 (2021).

15. Sathe, N. A. *et al.* Evaluating construct validity of computable acute respiratory distress syndrome definitions in adults hospitalized with COVID-19: an electronic health records based approach. *BMC Pulm. Med.* **23**, 292 (2023).

16. Johnson, A. E. W. *et al.* MIMIC-III, a freely accessible critical care database. *Sci. Data* **3**, 160035 (2016).

17. Johnson, A., Pollard, T. & Mark, R. MIMIC-III Clinical Database. [object Object] https://doi.org/10.13026/C2XW26 (2015).

18. Sjoding, M. W. *et al.* Interobserver Reliability of the Berlin ARDS Definition and Strategies to Improve the Reliability of ARDS Diagnosis. *Chest* **153**, 361–367 (2018).

19. Chen, T. & Guestrin, C. XGBoost: A Scalable Tree Boosting System. in *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining* 785–794 (Association for Computing Machinery, New York, NY, USA, 2016). doi:10.1145/2939672.2939785.

20. Bergstra, J., Yamins, D. & Cox, D. D. Making a Science of Model Search: Hyperparameter Optimization in Hundreds of Dimensions for Vision Architectures. in *Proceedings of the 30th International Conference on International Conference on Machine Learning - Volume 28* I-115-I–123 (JMLR.org, 2013).

21. van den Goorbergh, R., van Smeden, M., Timmerman, D. & Van Calster, B. The harm of class imbalance corrections for risk prediction models: illustration and simulation using logistic regression. *J. Am. Med. Inform. Assoc.* **29**, 1525–1534 (2022).

22. Afshar, M. *et al.* A Computable Phenotype for Acute Respiratory Distress Syndrome Using Natural Language Processing and Machine Learning. *AMIA Annu. Symp. Proc. AMIA Symp.* **2018**, 157–165 (2018).

23. Lee, R. Y. *et al.* Assessment of Natural Language Processing of Electronic Health Records to Measure Goals-of-Care Discussions as a Clinical Trial Outcome. *JAMA Netw. Open* **6**, e231204–e231204 (2023).

24. Youden, W. J. Index for rating diagnostic tests. *Cancer* **3**, 32–35 (1950).

25. Smits, N. A note on Youden's Jand its cost ratio. *BMC Med. Res. Methodol.* **10**, 89 (2010).

26. Serpa Neto, A. *et al.* Association Between Use of Lung-Protective Ventilation With Lower Tidal Volumes and Clinical Outcomes Among Patients Without Acute Respiratory Distress Syndrome: A Meta-analysis. *JAMA* **308**, 1651–1659 (2012).

27. Determann, R. M. *et al.* Ventilation with lower tidal volumes as compared with conventional tidal volumes for patients without acute lung injury: a preventive randomized controlled trial. *Crit. Care* **14**, R1 (2010).

28. Petrucci, N. & De Feo, C. Lung protective ventilation strategy for the acute respiratory distress syndrome. *Cochrane Database Syst. Rev.* (2013) doi:10.1002/14651858.CD003844.pub4.

29. Ancker, J. S. *et al.* Effects of workload, work complexity, and repeated alerts on alert fatigue in a clinical decision support system. *BMC Med. Inform. Decis. Mak.* **17**, 36 (2017).

30. Lee, E. K., Wu, T.-L., Senior, T. & Jose, J. Medical Alert Management: A Real-Time Adaptive Decision Support Tool to Reduce Alert Fatigue. *AMIA. Annu. Symp. Proc.* **2014**, 845–854 (2014).

31. Cvach Maria. Monitor Alarm Fatigue: An Integrative Review. *Biomed. Instrum. Technol.* **46**, 268–277 (2012).

Acknowledgements

The authors thank Catherine Gao for insightful discussions and suggestions.

Funding: FX was supported in part by the National Institutes of Health Training Grant (T32GM008449) through Northwestern University's Biotechnology Training Program; R.A.K.R. CHW was supported by the National Heart Lung and Blood Institute (R01HL140362 and K23HL118139). LANA was supported by the National Heart Lung and Blood Institute (R01HL140362). LANA and FX are supported by the National Institute of Allergy and Infectious Diseases (U19AI135964).

Author Contributions: **FM -** Methodology, Software, Validation, Data Curation, Writing – Original Draft, Writing – Review & Editing, Visualization. **HAL -** Methodology, Software, Validation, Data Curation, Writing – Review & Editing. **HTN -** Software, Validation, Data Curation. **MB -** Methodology, Validation, Data Curation, Writing – Review & Editing. **FX -** Methodology, Software, Validation, Data Curation, Writing – Review & Editing, Visualization. **JK -** Data Curation. **ELC -** Data Curation, Writing – Review & Editing. **CHW -** Conceptualization, Methodology, Validation, Data Curation, Resources, Writing – Original Draft, Writing – Review & Editing, Visualization, Supervision, Project Administration, Funding Acquisition. **LANA -** Conceptualization, Methodology, Software, Validation, Formal Analysis, Resources, Writing – Original Draft, Writing – Review & Editing, Visualization, Supervision, Project Administration, Funding Acquisition.

Competing Interests: The authors declare that they have no competing interests.

Data Availability. The datasets analyzed in this study will be made available upon publication at ARCH repository hosted by Northwestern University (https://arch.library.northwestern.edu).

Code Availability. The Python code to reproduce the reported results will be made available upon

publication at the Amaral lab GitHub repository (https://github.com/amarallab).