Title: COVID-19 vaccination atlas: an integrative systems vaccinology approach

Authors: Wasim Aluísio Prates-Syed[1,2], Dennyson Leandro Mathias da Fonseca[3], Shahab
Zaki Pour[3], Igor Salerno Filgueiras[2], Lena F Schimke[2], Aline Lira[1,2], Nelson Cortes[1,2],
Jaqueline Dinis Queiroz Silva[1,6], Evelyn Carvalho[1,6], Tania Geraldine Churascari Vinces[5],
Lorena C. S. Chaves[7], Gerhard Wunderlich[5], Ricardo Durães-Carvalho[4,8], Niels O. S.
Câmara[1], Haroldo Dutra Dias[13], Hans D. Ochs[12], Ester Cerdeira Sabino[9,10], José E. Krieger[11],
Helder I Nakaya[6,14], Otávio Cabral-Marques[1,4,6,15,16,.], Gustavo Cabral-Miranda[1,2,6,9,*].

Emails: wasim.syed@usp.br, dennyson@usp.br, shahab.178@gmail.com,
igor.filgueiras@usp.br, lenaschimke@hotmail.com, cv.tania@usp.br, aline.llira@usp.br,
nelson.cortes@usp.br, jaquelinedinis@gmail.com, evelyn.carvalho@unesp.br,
lorenacschaves@gmail.com, gwunder@usp.br, rdcarval@gmail.com, niels@icb.usp.br,
haroldodd@gmail.com, hans.ochs@seattlechildrens.org, aLLgau@u.washington.edu,
sabinoec@usp.br, j.krieger@hc.fm.usp.br, hnakaya@usp.br, otavio.cmarques@usp.br,
gcabral.miranda@usp.br

Affiliations:

1. Department of Immunology, Institute of Biomedical Sciences (ICB), University of
São Paulo (USP), São Paulo, Brazil.
2. The Interunits Graduate Program in Biotechnology of the University of São Paulo, the
Butantan Institute, and the Technological Research Institute of the State of São Paulo,
Brazil.
3. Interunit Postgraduate Program on Bioinformatics, Institute of Mathematics and
Statistics (IME), University of Sao Paulo (USP), Sao Paulo, Brazil.
4. Laboratory of Molecular Evolution and Bioinformatics, Department of Microbiology,
Biomedical Sciences Institute, University of São Paulo, São Paulo, Brazil;
5. Department of Parasitology, Institute of Biomedical Sciences (ICB), University of São
Paulo (USP), São Paulo, Brazil.
6. Department of Clinical and Toxicological Analyses, School of Pharmaceutical
Sciences, University of São Paulo, São Paulo, Brazil.
7. Department of Microbiology and Immunology, School of Medicine, Emory
University, Atlanta, Georgia, United States.
8. São Paulo School of Medicine, Department of Microbiology, Immunology and
Parasitology, and Department of Morphology and Genetics, Federal University of São
Paulo (UNIFESP), São Paulo, SP, Brazil.
9. Institute of Tropical Medicine, Faculty of Medicine of the University of São Paulo,
Brazil
10. Department of Infectious and Parasitic Diseases, Faculty of Medicine, University of
São Paulo, São Paulo, Brazil
11. Heart Institute, Clinical Hospital, Faculty of Medicine, University of São Paulo,
Brazil; Laboratory of Genetics and Molecular Cardiology, Clinical Hospital, Faculty
of Medicine, University of São Paulo, Brazil.
12. Department of Pediatrics, University of Washington School of Medicine, and Seattle
Children's Research Institute, Seattle, WA, USA.
13. Department of Neuroscience, Institute of Biomedical Sciences, Federal University of
Minas Gerais (UFMG), Belo Horizonte, Brazil
14. Hospital Israelita Albert Einstein, São Paulo, SP, Brazil
15. DO'R Institute for research, São Paulo, Brazil

16. Department of Medicine, Division of Molecular Medicine, University of São Paulo
School of Medicine, São Paulo, Brazil.

*Correspondence to: gcabral.miranda@usp.br or gcabral.miranda@gmail.com, Gustavo
Cabral-Miranda, Department of Immunology, Institute of Biomedical Sciences, University of
São Paulo, São Paulo, Brazil.

· Equal contribution as supervisor and senior author.

One-Sentence Summary:

We developed a COVID-19 vaccination atlas using an integrative systems vaccinology
approach, revealing critical markers that distinguish different vaccine types and regimens,
including infections, highlighting how full vaccination regimens limit immune response shifts
during natural infection, and identifying common transcripts shared across COVID-19 and
other pathogen vaccines.

Abstract:

COVID-19 vaccinations have significantly controlled the pandemic. To elucidate their impact
on the immune system, we developed a COVID-19 vaccination atlas through an integrative
systems vaccinology approach. Our atlas includes both healthy individuals and those infected
with or without prior vaccination, and covers the administration of five vaccines in different
regimens: Covilo®, Zifivax®, Vaxzebria® or Covishield®, Spikevax®, and Comirnaty®..
We identified critical markers to discriminate the various types of vaccines and infection, in
which infection was associated with GATA3, ZNF3, KMT2A, ASXL1, SP100, and GZMM,
and vaccine types were marked by ITGAM, ACTG1, LGALS3, and STAT5B. Additionally,
we described the immunological signatures resulting from heterologous vaccination and
infection, and also showed how a full vaccination regimen markedly limited the shift of
immune responses during natural infection, thereby constraining disease progression. Finally,
we described the common transcripts shared across COVID-19 vaccines and vaccines against
other pathogens.

Main text

1. Introduction

Coronavirus Disease 2019 (COVID-19) is a respiratory disease caused by Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), with its first documented outbreak in Wuhan, China, in 2019 *(1)*. The emergence of variants of concern (VOCs) urged adjustments in strategies to effectively control their spread due to the heterogeneity of symptoms, severity, transmission, and decreased efficacy of vaccines against new VOCs, especially the Omicron variant (Chung et al., 2021; Xia et al., 2022).

Remarkably, it is estimated that COVID-19 vaccinations prevented more than 14 million deaths in only one year after global vaccination enrollment*(4)*. This was accomplished by breakthrough innovations in vaccinology, in addition to global efforts to fund worldwide clinical trials, including classical and new-generation vaccines*(5)*. The mRNA vaccines (RNA) Comirnaty® (BNT162b2, BNT) and Spikevax® (mRNA-1273, MO), the viral vectored (VV) vaccines Covishield®/Vaxzebria® (AZD1222, ChAdOx-1, ChAd) and Jcovden® (Janssen), and inactivated vaccines (IN) such as Coronavac® and Covilo® (BBIBP-CORV, BBIBP) were among the first and most used vaccines worldwide until 2023 *(6)*. Particularly, ChAd and BNT were the most used vaccines during the first years of vaccination and 2 and 3-dose heterologous regimens were widely used and assessed *(7, 8)*. The combination of different vaccines and platforms shows how classical and new-generation vaccines can help control epidemics and potentially trigger complementary systemic immune responses that individual vaccine technologies are intrinsically incapable of *(8–11)*. The complete homologous vaccination regimen showed high efficacy in clinical trials against COVID-19, as high as 95.0%, 94.1%, and 90% for new-generation vaccines, such as BNT, MO, and ChAd, respectively *(11–13)*. Classical technologies, such as inactivated virus (IN), and protein subunit (SU) vaccines, have also shown high efficacy and effectiveness. BBIBP and Zifivax® (ZF2001) had 78.1% and 75.7% efficacy against symptomatic infections, respectively *(14, 15)*. Although these vaccines were initially tested and approved in homologous regimens, heterologous vaccination has been tested in clinical trials and real-world scenarios to address ongoing waves of infection and new VOCs *(8, 16)*.

To understand how different vaccination regimens and vaccine technologies affect the immune system in healthy individuals and infected patients during, and after infection by SARS-CoV-2, we integrate RNAseq data from five distinct studies that include vaccinations with Covilo®, Vaxzebria® or Covishield®, Spikevax®, Comirnaty®, and Zifivax®. These studies, however, present data from different time points, populations and regimens, and there is a need to obtain a unified perspective of their immunological signatures. In this work, we present a systems vaccinology atlas that describes the common and individual profiles of each condition, in both healthy and infected individuals. We show that a full vaccination regimen can limit the progression of disease. We also show that heterologous vaccination can diversify the immunological signatures and potentially protect more effectively against the disease. Additionally, we explored the immune responses induced by vaccines against other pathogens and showed that COVID-19 vaccines share a substantial amount of genes related to the innate immune system.

2. Results

Differential expression of immune-related genes is influenced by both vaccine type and regimen, as well as the underlying infection status.

We analyzed 562 samples derived from peripheral blood mononuclear cell (PBMCs) from 245 participants across five separate studies (**Table 1, Fig. 1A**). These participants were either vaccinated (V) with one or more of the four different types of vaccines (IN, SU, RNA, and VV), which comprised both homologous (HO) and heterologous (HE) vaccination regimens and were administered in one (V1), two (V2), or three doses (V3), or were infected patients (I) with or without prior vaccination (**Fig. 1B-C, Fig. S1-2**). We found many issues with the GSE206023 count matrix, in which most of the genes were annotated in GenBank IDs with multiple identifier versions for the same genes. Additionally, we found that only 10% of these identifiers could be annotated, comprising less than 2% of the genes in the ImmuneGO dataset, in contrast to more than 90% present in the other datasets. Given the importance of this dataset regarding the diversity of vaccine technologies, we opted to reprocess the read files in FASTQ and align it with the reference transcriptome using pseudoalignment in the Kallisto package *(17)*. Although other datasets used HISAT2, we opted for pseudoalignment because it is faster, more robust, and requires fewer computational resources than HISAT2 *(18)*.

The temporal dynamics of differentially expressed genes (DEGs) was influenced by vaccine type, number of doses, and administration regimen. This happened across convalescent and non-convalescent individuals, as well as infected patients with previous vaccination (**Fig. 2A, Fig. S3**). Notably, the BNT vaccine exhibited a significant increase in total DEGs across the three doses, with three DEGs six days after the first dose (BNT V1), followed by 613 and 96 DEGs 1 day after the second (BNT V2) and third doses (BNT V3), respectively, with a more substantial increase with two doses of BNT followed by a single dose of MO (BNT-MO V3) (**Fig. 2A**). In contrast, we observed the ChAd vaccine exhibited a divergent pattern compared to BNT in non-convalescent individuals, with decreased up- and downregulated DEGs over the first week after two homologous doses (ChAd V2), and a higher number of DEGs in the heterologous vaccinations (ChAd-BNT V2 and ChAd-BNT V3) (**Fig. 2A, Fig. S3**). A similar trend was observed in homologous and heterologous vaccinations with the BBIBP vaccine after a third dose (**Fig. 2A, Fig. S3**). It was evident that homologous vaccination with an inactivated virus vaccine does not generate significant DEGs in a third dose (BBIBP V3) and thus requires heterologous vaccination with a different technology, such as a SU vaccine (ZF2001 V3) (**Fig. 2A, Fig. S3**).

Furthermore, infections in vaccinated and unvaccinated patients notably affected the number of downregulated DEGs (**Fig. 2A, Fig. S3**). Among unvaccinated infected individuals, DEGs were higher in severe disease cases than in moderate cases over 51 days. Conversely, in infected patients with prior complete vaccination, primarily mild and moderate disease, this number was at least 10 times lower. However, infected hospitalized patients with moderate and severe disease who were vaccinated with a single dose of BNT 10-11 days before hospitalization, the number of DEGs was similar to that of infected patients on days 10 and 26. This pattern could also be influenced by the age range (71-90 years) of these patients, which is a major risk factor for COVID-19*(19)*. Of the four infected patients, three recovered, and one died (**Fig. S3**). The surviving patients continued dexamethasone treatment and received a second dose of the vaccine 40 days post-infection*(20)*. Samples were collected 15-20 days post-vaccination, exhibiting more DEGs than at a later time post-vaccination in non-convalescent individuals with BNT or ChAd (**Fig. 2A**). The gene expression patterns in

unvaccinated and single-dose vaccinated patients during the second infection were similar,
but vaccinated individuals exhibited a notable, threefold reduction in DEGs compared to with
unvaccinated counterparts. Only in the total DEGs analysis, without discriminating by
ontology (immune and non-immune), was an apparent relationship between vaccine type,
disease severity, and time.

When we filtered immune-related DEGs, it was evident that the fraction of immune-related
genes decreased over time in vaccinees and infected patients (**Fig. 2A, Fig. S3**). We
investigated only the immune-related genes excluding genes from the BCR and TCR
repertoire [immunoglobulin chains (IG@) and T-cell receptors (TR@)]. Particularly in
infected patients, the first days showed a greater fraction of immune genes, which decreased
over time, whereas the number of other non-immune-related genes increased (**Fig. 2A, Fig.
S3A**). In addition, infected patients with prior complete immunization showed fewer DEGs
and a smaller fraction of immune genes. Nevertheless, vaccination results in the activation
substantially more immune genes, which follow proportionally to other genes, either
increasing or decreasing, depending on the vaccine type. This contrast indicates that the
immune response induced by vaccines is quantitatively more balanced in terms of the
expression of immune genes than infection. To identify patterns among these conditions,
immune-related and non-immune DEGs were compared distinctively (q-value < 0.10) (**Fig.
2B, Fig. S4A**). We found that most of the shared genes among the different vaccine types
were non-immune genes, such as ANKRD22, GBP1P1, GPRB4, and HES4 (**Fig. S4A-B).**
Moreover, the most shared genes, such as LRRN3, ADAMTS2, and FAM209B, were more
present among infected patients (**Fig. 5B**). This shows the role of other processes in
vaccination and, to a greater extent, infection.

To understand these non-immune processes during vaccination, we analyzed the non-immune
genes. We found that most of them were related to ubiquitous cellular and metabolic
processes, and the regulation of biological processes (**Fig. 2A-B, Fig. S4**). However, we
focused on immune genes in the further analysis. The shared immune DEGs exhibited one
central cluster associated with the groups infected for the first time (vaccinated and
unvaccinated) and another cluster comprising the first day of vaccination with homologous or
heterologous second and third doses of ChAd, BNT, and MO (**Fig. 2B**). Notably, the number
of shared genes among infected groups was 2- to 10-fold higher than non-convalescent-
vaccinated groups. In addition, we observed a decrease in shared DEGs between infected
patients with prior vaccination (BNT-I) and infected patients with prior infection and prior
vaccination (I-BNT-I), in a similar proportion to non-convalescent vaccinated groups, which
was not evident in the non-immune DEGs context (**Fig. 3B, Fig. S4A**). This finding was also
associated with vaccine type, viral vector vaccines (ChAd and ChAd-BNT), and subunit
vaccines (ZF2001), which were more evident with non-immune DEGs (**Fig. S4A**).

In contrast, we observed a decreasing number of genes on subsequent days, particularly on
day 7, suggesting the initiation of homeostatic processes over time. This reduction in DEGs
becomes more pronounced on subsequent days. However, this decrease contrasts with the
trend observed with increasing doses, particularly notable in homologous RNA vaccines and
heterologous ChAd with subsequent vaccination with BNT. This indicates a broader
activation of diverse genes with a higher number of doses. Particularly, vaccination with BNT
following one or two doses of ChAd led to a significantly greater number of total DEGs
compared to the two-dose homologous regimen of either vaccine (**Fig. 2A-B, Fig. S4**).
Specifically, when compared to the two doses of ChAd on day 3, there was a three-fold
increase in upregulated DEGs, which reduced by half on day 7. Conversely, downregulated

DEGs were 13-fold higher in the ChAd-BNT regimen but increased two-fold on day 7.
Remarkably, even though the assessment was conducted only one day post-vaccination,
upregulated DEGs exhibited in the ChAd-ChAd-BNT regimen were approximately two-fold
higher than those with two doses of ChAd and 1.2-fold higher than those with BNT-MO.

In contrast, the number of up- and downregulated genes was nearly equivalent in infected
individuals, regardless of prior vaccination status (**Fig. 2A, Fig. S4A**). Additionally, genes
from unvaccinated infected individuals were largely constrained within their group and had a
limited overlap with vaccinated individuals.

To describe the immune transcriptome of individual vaccination regimens, we compared all
the genes present in the datasets with the immune system related genes (**Fig. 4D**).
Subsequently, we analyzed the enrichment of gene sets associated with the immune response
(**Fig. 4C**).

Limited Immunological Profile Observed with Three Homologous doses of BBIBP

The BBIBP exhibited enrichment of processes associated with the innate immune system,
including the antiviral response mediated by interferons, neutrophils, eosinophils, and
monocytes, which were consistently found to be negatively regulated (**Fig. 4C**). Further
analysis of gene expression within these processes and cells revealed specific genes of interest
(**Fig. S4-7**). On day 14, we identified genes associated with inflammation (NLRP9,
upregulated), immune regulation (IDO2), macrophage-mediated antiviral response (CCL3,
downregulated), neutrophil activation (CXCL8), and lymphocyte-macrophage signaling
(WNT4, upregulated)*(21)*. On day 28, a transcription factor (NFKBIA, downregulated)
present in ARPP and B cells, which regulate immune system processes, and CCL2
(downregulated), a monocyte chemokine, were observed.

ZF2001 as Heterologous Booster Elicits Distinct Immune Response Profile

A heterologous regimen with the ZF2001 vaccine in individuals who had previously received
two doses of the BBIBP vaccine resulted in a distinct gene expression profile compared to the
homologous vaccine. Although most DEGs were upregulated, a small fraction of genes related
to the immune system were downregulated (**Fig. 2A**). On the seventh day post-vaccination,
there was a suppression of genes associated with inflammatory processes (IL1RN, FOSL2,
and CD58), antiviral response and interferons (CXCL2, CXCL3, CCL3, CCL4, CCL20,
BCL3, ADM, LGALS3, FAU), as well as genes related to antigen recognition and
presentation (TNFAIP3, FOSL1, RAB3B, and ICAM). This suppression also extended to
genes expressed in eosinophils (VAMP2), NK cells (RNF19B), dendritic cells (NFKBIA,
FOSL1), T cells (IL2), B cells, and immunoglobulin-mediated responses (BCL3, BCL6) (**Fig.
2C**, **Fig. S6-9**).

On the other hand, a predominance of upregulated genes was observed on the seventh day,
particularly associated with other chemokines (CX3CL1, CXCL8, CXCL14, CCL8,
CX3CL1), mast cells (CPLX2, CLNK, LGALS3), ARPP (ITGB8, TRIL), APCs (ADGRF5,
FLT1, IL33, WNT4), adaptive response processes and cells (CACNB4, EOMES, SPNS2,
BCAR1), as well as the complement system (**Fig. S6-9**). Specifically, within the complement
system, genes from the classical pathway (C4B and C4BPB) and lectins (MASP1 and
MASP2) were overexpressed (**Fig. S6-9**). In the adaptive immune system, there were genes
related to T cell activation (IL13, EPHB1, TNFAIP8L2, TARM1), specifically Th2 response

(IL33), Th17 (CCR2), TCD8 (EMP2), B cell and Ig-mediated response (WNT3A) (**Fig. S6-**
9).

Viral Vectored Vaccine ChAd Induces Early Gene Expression of Complement and
Adaptive Immune Systems

Genes related to T cells and the cellular response to the viral vector were expressed as early
as the third-day post-vaccination with the first dose. Notably, genes such as CCNB2 and
CAV1, which play roles in the cell cycle and T cell activation, respectively, exhibited
sustained expression levels up to the seventh day. Additionally, the innate immune system
processes are enriched, including interferon-mediated antiviral responses, acute phase
response proteins (ARPP), and NK cell, related proteins along with upregulated genes from
the complement system associated with the alternative pathway (CFB and CFD) (**Fig. S7-9**).

It is also essential to highlight the distinct expression of genes related to humoral responses
on the third and seventh days (**Fig. S8**). These expression patterns were also observed in B-
cell genes, with overexpression between days 3, 6, and 7, returning to the initial profile by the
sixth day. Conversely, except for genes associated to cellular response, B cells processes, and
the humoral immune response were enriched on the first day after the second dose. However,
gene-level analysis still revealed diverse and high gene expression at this stage, especially for
the cytokine CCL2 and immunoregulatory enzyme IDO1. When comparing genes related to
the humoral response between doses, it was noted that these genes were less expressed and
less diverse at the second dose. On the third day, only two DEGs associated with the humoral
response were identified (SYK and KIT). Particularly, on day 7, CD27, a gene involved in
immunoglobulin synthesis, *(22)* was increased after the first dose and suppressed after the
second dose.

Distinct Gene Expression Patterns Elicited by mRNA Vaccines BNT and MO

The mRNA vaccines BNT and MO demonstrate distinct gene expression patterns. Six days
after the first dose of BNT, only 3 DEGs were identified, with IFI27 being the only immune-
related gene (**Fig. 2A, Fig. S4-7**). Subsequent second and third BNT doses revealed elevated
activation of antiviral and interferon responses, ARPP, and the function of neutrophils,
eosinophils, monocytes and dendritic cells (**Fig. 4C**). These sets were similarly enriched
following the third homologous vaccination, except for dendritic cells. A third heterologous
vaccination with MO induced dendritic cell function, inflammation, and the activation of the
cellular adaptive immune system. Notably, IFI27 expression was exclusive to all BNT doses
except for the third dose of MO (**Fig. S4-7**). Neutrophils shared genes in various vaccination
regimens, including CCL8, CXCL9, and CCL2 (**Fig. S4-7**) *(23)*. In addition, eosinophils
exhibited pronounced CCL3L1 expression following MO vaccination, which plays a pivotal
role in virus clearance from the lungs *(24)*.

Immune Signatures in Infected and Convalescent Patients Before and After Vaccination
with BNT

To understand the impact of vaccination before and after infection, we analyzed infected and
convalescent patients. We found that innate immune cells and processes were highly activated
and persisted for several days, especially interferons 1 and 2, neutrophils, APCs, and
bacterium-directed PRRs (TLR4 and NOD) (**Fig. 4C, Fig. S4-7**). Additionally, B-cell
processes, including chemotaxis and homeostasis, are activated (**Fig. 4C, Fig. S4-7**). There
was also a Th2 response in BNT-I (D26, mild), although T cell processes were predominantly

downregulated under other infection conditions. In a gene-level analysis, many genes related
to the adaptive immune system were upregulated and downregulated. Likewise, it was evident
that infection induced different subsets of genes compared to vaccination in every immune
process and cell type (**Fig. S4-7**). Considering the innate immune system and classical
pathway complement system genes (such as CR1 and C1R), high and persistent gene
expression, especially those associated with inflammation (e.g., IL10, HMGB2, ELANE, and
NLRP6), activation of neutrophils (e.g., CD177, ANXA3, and IL18RAP) and mast cells (e.g.,
S100A12 and SLC15A4), may be associated with the initiation of cytokine storms, allergic
reactions, and thrombosis *(25, 26)*. Although most genes participating in adaptive immune
system processes were highly downregulated, there were also many highly expressed genes
in the humoral response, including genes related to Ig-mediated responses, ARPP, and other
B-cell processes, as well as in cellular response and T- cell processes. This pattern varied
among convalescent and infected patients who had received prior vaccination, regardless of
whether they had a completed or incomplete vaccine regimen (**Fig. S4**).

Stratification of COVID-19 vaccination and infection using multivariate analyses

To describe the similarities and differences among all conditions, we conducted a Principal
components analysis (PCA) with all immune DEGs (**Fig. 3A-D**). Principal component 1
(PC1) accounted for 66.8% of the variance, whereas PC2 accounted for 15.7%, totaling
81.9%. PC1 was predominantly defined by genes associated with interleukin (IL) signaling
pathways (IL18RAP, SOS2, MAP2K6), apoptosis (HMGB2), inflammation-related processes
(RC3H1, ALOX5, PPARG, and FCER1A), as well as processes linked to lymphocytes
(ETS1), particularly to T cells (VNN1, CD6, TESPA1, RPL22, PCLG1, TCF7, TRAT1, and
CD3G) (**Fig. 3C**). On the other hand, PC2 was defined by genes related to antiviral and
interferon responses (IFI35, IFITM3, STAT1-2, IDO1, ADAR, BST2, GBP1, and PML), as
well as APCs (BATF2, LILRB1) and antigen receptors, processing and presentation (ARPP)
(TAP2, PDIA3, TRIM26, and CTSS) (**Fig. 3D).**

The results revealed the presence of three primary clusters (**Fig. 3A**). The first significant
cluster indicated patients with ongoing infection, characterized by genes associated with IL-
signaling pathways, apoptosis, and inflammation-related processes in cluster 1 (**C1; Fig.
3A,C).** While this cluster included specific individuals with previous vaccination, we did not
anticipate a substantial influence of vaccination, particularly considering that it was
administered 10 days before the onset of COVID-19 symptoms. Nevertheless, it is evident
that patients who received a complete vaccination regimen and were subsequently infected
(BNT-I), as well as a patient who received a second dose of BNT 10 days after recovery
(BNT-I-BNT, D51), showed overlap with Cluster 2 (C2). The C2 cluster comprises
homologous vaccinations for both convalescent patients and non-convalescent individuals.
Additionally, healthy-vaccinated individuals were part of this cluster, but it also included
patients with two infections (without severe outcomes) and one patient initially experiencing
severe disease 51 days after the onset of symptoms, who were hospitalized and treated with
steroids. Given that this cluster occupies the central position in the plot and represents
comparatively healthy conditions, the proximity of eigenvalues from the first cluster to the
second cluster suggests an ongoing, time-dependent homeostatic process. In contrast, the third
cluster (C3) consisted of homologous and heterologous vaccinations involving BNT, ChAd,
and MO, specifically within the initial three days post-vaccination. This cluster is
characterized by genes in PC2 and shares a minimum number of certain genes with infected
patients in C1. Nevertheless, the expression of the top 20 contributing genes in PC2 was
significantly higher in C3 than in C1 conditions (**Fig. 3B,D**). These differences were marked

by genes such as IDO1, IL31RA, and LILRB1, which were not expressed during infection
and are associated with antimicrobial response, chemotaxis, and MHC1 presentation,
respectively.

Machine learning classification of COVID-19 vaccine types and infection based on immune genes

Using the PCA results, we filtered the genes in PC1 and PC2 (cos2 > 0.5) and trained a random
forest model for classification based on the variable "type", including infection and the
different vaccine types (**Fig. 4A**). This model demonstrated remarkable performance, with an
accuracy of 91.2%, an area under the receiver-operator curve (ROC) of 0.990, and an out-of-
bag (OOB) error of 0.086. The high expression of the transcriptional regulation associated
genes GATA3, ZNF3, KMT2A, and ASXL1, an antiviral response gene (SP100), and the
granzyme GZMM, were key markers for infection, along with the suppression of genes that
play roles in ARPP (ITGAM, ACTG1, LGALS3), and T-cell response (STAT5B) (**Fig. 4B**).
SP100 was also notably expressed in vaccination, in the first 3 days after the second and third
doses of both homologous and heterologous regimens. LGALS3 was also markedly expressed
in the first 3 days of vaccination with ChAd. This observation aligns with the fact that viral
vectored vaccines, like ChAd, act through certain infection associated mechanisms to deliver
genetic material into the nucleus*(27)*.

Comparative Analysis of Vaccines Against Other Pathogens

Understanding how COVID-19 vaccines relate to classical and consolidated vaccines used in
routine immunization is essential. Therefore, we compared all immune-related DEGs
associated with various vaccines, including COVID-19, using MSigDB Vax collection *(28)*.
We manually annotated the dataset and selected all up- and downregulated genes from
vaccination-only sets (**see Methods, Fig. 1A**). This approach originated from several gene
sets, including those correlated with antibody titers, T-cell responses, and training sets. To
understand the extent of gene overlap, we performed a gene-level analysis of these shared
genes (**Fig. 5A-C, Fig. S8 and 9**).

Our analysis revealed a significant sharing of genes, both up- and downregulated, between
vaccines targeting viruses and bacteria based on inactivated (IN) and live-attenuated (LA)
pathogens and COVID-19 vaccines (**Fig. 5A-C**). This overlap was most prominent in the
initial two weeks and following the second and third doses. Specifically, downregulated DEGs
were predominantly associated with infection, particularly with LA vaccines (**Fig. 5A**).
Conversely, upregulated DEGs were common among vaccinated individuals, comprising both
non-convalescent and convalescent individuals. This overlap extended to include infected
patients who had undergone prior one-dose vaccination.

Our observations revealed two principal clusters of genes, predominantly representing the
innate and adaptive immune systems (**Fig. 6**). While the second cluster encompasses genes
present in the adaptive immune system, these genes also exhibit a lower degree of overlap
with the innate immune system. Notably, the second cluster appeared more heterogeneous and
demonstrated significantly fewer shared genes across vaccines than the first cluster. In this
first cluster, we observed that COVID-19 vaccines and others shared most innate genes that
also play a role in the adaptive immune system (**Fig. 5D**).

Some of the most common genes between vaccines included chemokines and receptors
(CCL2, CCL3, CCL8, CCL20, CCR1, and CXCL8), interferons and antiviral responses
(IRF1, IFIT1, STAT1, and STAT2), ARPP (TAP1, TAP2, HLA-DMB, and HLA-DMA), and
some adaptive immune system genes (RIGI, FAS, CD83, BCL6, and FCER1G) (**Fig. 5B**). On
the other hand, some of the genes present mainly in COVID-19 vaccine conditions are
chemokines, such as CCL2, which play a diverse role in the innate, adaptive, and complement
systems (**Fig. 5C**). CCL2 is also shared with the monovalent influenza vaccine adjuvanted
with AS03 and the LA vaccine Fluzone® *(29)*. Although most non-COVID-19 vaccines
sharing genes with COVID-19 vaccines are targeted against the influenza virus, Stamaril®, a
yellow fever LA vaccine, shares many innate immune system genes with COVID-19 vaccines
(30). Indeed, an overrepresentation analysis (ORA) of immune genes showed that general
leukocyte processes and antiviral and interferon processes were enriched in most vaccines and
vaccine types (**Fig. 5E**). VV vaccines against COVID-19 and tuberculosis (MVA85A) were
significantly (qvalue < 0.10) enriched with genes related to eosinophils, monocytes, and
neutrophils (**Fig. 5E**).

Discussion

This COVID-19 vaccination atlas comprehensively analyzes the dynamic and diverse immune
responses generated by different vaccination strategies against COVID-19. These strategies
include homologous and heterologous vaccination regimens, diverse vaccine types, and
consideration of prior vaccination status at the time of infection, alongside the time elapsed
since infection (**Fig. 1A**).

Our analysis reveals that immune-related genes constitute a smaller proportion of the total
DEGs than other biological processes, including cellular, metabolic, and regulation of
biological processes (**Fig. 2A, Fig. S4**). A limitation of this analysis by gene ontology of
biological process is that some genes that are not annotated as immune play ubiquitous roles
in cellular and metabolic processes present in immune cells. This can also be attributed to the
inoculation of immunogens through syringe injection, which leads to local and limited tissue
damage at the injection site. Indeed, intramuscular injections are prone to human errors, and
questions remain regarding which injection site and procedure are optimal and safest *(31)*.
This vaccine administration issue and the molecular characteristics of different vaccine
platforms might explain reactogenic behavior in some individuals. In the context of infection,
this pattern illustrates the regulation of numerous genes influenced by viral infection cycle
processes and the host response *(32, 33)*. This variability is primarily attributed to diverse
response mechanisms among individuals, impacting disease progression or immunity*(34)*.
However, this aspect will not be discussed in more details since it diverges from the primary
objective of our study, which focuses on describing the transcriptome of the immune system.
Nonetheless, it is noteworthy as it raises questions regarding non-immunological factors and
their role in vaccination and disease development.

Moreover, we found that these DEGs are highly shared between the studied conditions,
encompassing both infection and vaccination conditions. In addition, **a** higher representation
of immune-related genes was observed in the vaccination group compared to infection (**Fig.
2A**). Specifically, genes associated with inflammation, the complement system, neutrophil
function, antiviral responses, and interferon pathways exhibit prolonged activation during
infection, while adaptive immune-related genes are significantly downregulated (**Fig. S4-7**).
In contrast, vaccination triggers the activation of chemokines and adaptive immune-related
genes.

The surge in up- and downregulated DEGs in the early days after the second and third doses
of BNT or MO indicates the ability to activate the innate immune response. Considering the
rapid mRNA expression, post-inoculation might also trigger antigen-presenting cells to
initiate an adaptive immune response *(35)*. Notably, despite having the same mRNA platform
as BNT and encoding the same spike amino acid sequence, the MO vaccine differs in
nucleotide sequence due to codon optimization and the absence of pseudouridine
modification*(36, 37)*. However, whether this substantial increase is attributable to an immune
response to the third dose, the technology, or other factors, such as LNP composition, remains
unclear.

The heterologous vaccination regimen was also explored with different types of vaccines,
namely ChAd and BNT, and BBIBP and ZF2001. We found that it can contribute to the
diversification of immune responses, address some of the technological limitations of the
vaccine platforms, and consequently enhance protection against COVID-19.

The ChAd vaccine contains a chimpanzee adenovirus viral vector that shares epitopes with
prevalent human adenoviruses, which can trigger an adaptive immune response against the
vector*(38)*. It is therefore possible that the administration of multiple doses of the vaccine may
generates antibodies targeting the vector, potentially limiting the delivery of genetic material,
unlike mRNA vaccines *(39)*. Hence, heterologous vaccination is a clinical and
immunologically important option, allowing individuals who receive the complete
homologous ChAd regimen to obtain better protection through vaccination with vaccines that
use different technologies, particularly those based on mRNA *(16, 40)*. Nevertheless, the
discussion regarding the continued efficacy of the ChAd viral vector in these individuals
compared to other antigen vaccine formulations is crucial.

Heterologous mRNA vaccination regimens, especially those following the ChAd vaccine,
demonstrate similarity to the second and third mRNA homologous and heterologous
vaccinations. This can be advantageous for a vaccination regimen that is initiated by a VV
vaccine and boosted by mRNA vaccines, potentially mitigating the impact of antibodies
directed against the viral vector. However, the use of BNT as a third dose after two ChAd
doses is mainly associated with antiviral and interferon responses as well as neutrophils.
Additionally, as the samples were limited to the first day after vaccination in this condition,
we could not assess the impact of these responses on the adaptive immune system.

The Covilo®/BBIBP vaccine is an inactivated virus vaccine formulated with alum adjuvant
and evaluated in clinical trials with a homologous two-dose regimen(Wang et al., 2020; Al
Kaabi et al., 2021; Zhang et al., 2022). The regimen of three homologous doses of BBIBP,
specifically when given on days 7, 14, and 28, seems to present a limited immunological
profile owing to the nature of the immune response to inactivated viruses*(43)*. In a study
assessing the effectiveness of two doses of an inactivated influenza vaccine, employing a
similar strategy to the development of the BBIBP vaccine, significant changes in gene
expression were not observed one week after the second dose. Additionally, no considerable
antibody titers were recorded, corroborating the BBIBP vaccine data*(44)*. However, in another
BBIBP vaccine study, this limited immunological profile was not observed, in which a second
boost induced more robust humoral and cellular immune responses detected after two
weeks*(45)*. Despite the limited data regarding the efficacy of a third dose, it is evident that
employing a homologous vaccination strategy with three doses of the inactivated BBIBP
vaccine may offer limited immunological benefits. Nevertheless, the third dose demonstrated
significant efficacy against the disease *(15)*. It is important to emphasize the need to evaluate
the immune transcriptome after the first two doses in samples from infected patients who have

received two or three vaccinations*(42)*. Furthermore, investigating the early stages of
vaccination, regardless of the number of doses administered, could provide valuable insights
into identifying the genes linked to the innate immune response.

The Zifivax®/ZF2001 vaccine was not initially evaluated at the onset of the pandemic when
the population had not yet been vaccinated. Instead, it was assessed as a booster dose in
individuals who had previously received two doses of the BBIBP vaccine *(15)*. As expected,
in a heterologous regimen, vaccination resulted in a distinct gene expression profile when
compared to the homologous vaccine *(11)*. These findings suggest that immunological
memory plays a pivotal role in the first-week post-vaccination, and antibodies generated by
the inactivated vaccine recognize, opsonize, and trigger a response to the ZF2001 vaccine
protein through the classical pathway of the complement system. Additionally, the original
study revealed a significantly higher antibody titer than the homologous regimen, supporting
the importance of heterologous vaccination *(15)*.

Importantly, our findings show the significant contribution of complete vaccination regimens
compared to natural infection and incomplete vaccination. Considering that infected patients
with incomplete vaccination were hospitalized 10-11 days after vaccination with a single dose
of BNT, infection must have occurred some days before hospitalization, and we anticipated
that vaccination would not have protected these patients. In the PCA, individuals with prior-
vaccination overlapped between the infected cluster and the vaccination cluster C2, which
encompassed homologous vaccinations for both convalescent and non-convalescent
individuals, along with healthy-vaccinated individuals (**Fig. 3A**). This overlapping was
notable among individuals who were infected after a complete vaccination regimen with BNT.
Conversely, an incomplete vaccination regimen was associated with infection, suggesting a
limited protection of vaccination administered right before infection. We found that in
infected patients with no prior vaccination or with incomplete vaccination the genes related
to innate immune cells and processes were highly activated and persisted for several days,
especially IFN1 and IFN2, neutrophils, and APCs. Interestingly, bacterium-directed PRRs
(TLR4 and NOD) were also highly expressed in these conditions, suggesting that these
patients were also coinfected by bacteria, which is a risk factor for hospitalization*(46)*.

To provide a model of key genes that describe infection and the different types of COVID-19
vaccines, we trained a classification random forest model with 244 genes selected from the
PCA. Our model identified the 10 genes that are most important for classification, with high
accuracy (91.2%) and sensitivity (ROC AUC of 99.0%), and low OOB ratio (8.6%). The
genes that classified infection were associated with transcriptional regulation (GATA3, ZNF3,
KMT2A, and ASXL1), antiviral response (SP100), and granzymes (GZMM); vaccine types
were marked by ARPP (ITGAM, ACTG1, LGALS3) and T-cell response (STAT5B) (**Fig.
4B**).

In addition, we compare COVID-19 vaccines with other vaccines targeting different
pathogens, as outlined in the Vax MSigDB dataset (**Fig. 5C**). This comparison is crucial for
understanding how new-generation vaccines, such as mRNA-based and viral-vectored
vaccines, relate to classical and consolidated vaccine technologies. Given the pivotal role of
the innate immune response in the early stages of both vaccination and infection, the
overlapping genes identified likely contribute to the innate immune system *(47)*. In particular,
bias in FCER1G towards non-COVID-19 vaccines shows a predominance of Th2 response,
presumably due to the widespread use of alum adjuvants, a known Th2 response activator that
is used in the formulation of BBIBP and ZF2001 *(48)*. This comparison was notable with
influenza IN vaccines and LA vaccines against yellow fever and smallpox, which share

common DEGs with COVID-19 vaccines in the initial weeks, especially those related to the innate immune system. Additionally, LA vaccines serve as a valuable reference for a robust immune response, mimicking the wild-type virus infection. The observed overlap of LA vaccines with COVID-19 vaccination conditions adds an intriguing dimension to our findings. The diversity in the immunome in vaccines sharing the same technology may be attributed to the specific and varied nature of the adaptive immune system and other technical limitations of the MSigDB Vax dataset. It primarily relies on transcriptomic data from microarrays, which predates the widespread adoption of high-throughput RNA sequencing. In addition, this dataset lacked gene expression fold-change values, highlighting a constraint in our analysis.

Although this study provides valuable insights, there are notable limitations regarding the datasets and transcriptomic analyses utilized. Specifically, the use of only bulk RNA-seq data restricts our understanding of the dynamic and heterogeneous behavior of individual immune cells. Furthermore, collecting vaccination and infection samples at different time points presents a limitation in describing the chronological dynamics of the immune response. Specifically, this difference restricts the characterization of the adaptive immune response in vaccination, predominantly assessed within a one-week period. There are additional challenges in analyzing third-dose vaccinations with BNT, MO, and ChAd, which were assessed on the first-day post vaccination, primarily constraining the analysis to the innate and complement immune systems. These restrictions remain significant despite their interconnectedness with the adaptive immune responses elicited by previous doses. In addition, the analysis of BBIBP and ZF2001 vaccines is restricted by the limited time points on the days during the first week, crucial for the innate immune response, and also did not allow us to compare with the preceding doses. Moreover, the observed gene expression pattern was in agreement with the immunological finding of a study, but was contradictory to another(Aydillo et al., 2022; Ying Chen et al., 2023). Exploring how different vaccine technologies contribute to protecting infected vaccinated individuals, as observed in BNT-vaccinated individuals, would provide further insight.

To improve analysis and interpretation by incorporating supplementary data sources, such as antibody titers, neutralizing antibodies, T cell counts, and clinical trial data, would greatly enhance our understanding. Although sex, age, and VOCs data were present, we did not adjust the contribution of these variables to the immunological signatures described. Adopting a multidimensional approach that integrates various datasets and immunological methodologies would offer a more comprehensive understanding of the immunological landscape after vaccination. Furthermore, introducing innovative tools such as VaxGO, described in this study, can promote the exploration and comparison of immunological processes across different vaccines, thereby facilitating more informed vaccine designs and evaluation strategies.

3. Methods

Data curation and processing

The selected datasets were first curated using available metadata from Gene Expression Omnibus (GEO) and supplemented with information obtained from pertinent publications (**Fig. S1-3, Table 1**). The inclusion criteria were as follows: 1) studies related to SARS-CoV-2 immunization; 2) detailed and precise information about methodology; 3) convalescent and non-convalescent patients with or without prior vaccination; and 4) bulk or single-cell RNA-seq PBMC-derived samples. Exclusion criteria: 1) absence of description of methods or references, especially on sample collection time points between doses; 2) TCR and BCR repertoire sequencing; 3) inclusion of patients with distinct medical conditions, such as autoimmune diseases and cancer; and 4) datasets without standardized count matrices.

The count matrices for each dataset were obtained from the Gene Expression Omnibus (GEO) *(49)* and their metadata from the R package GEOquery *(50)*. To standardize the gene identifiers as HGNC symbols, we used the R package biomaRT *(51)*. For reprocessing of reads in the GSE206023 study, FASTQ files were downloaded directly from GEO under the accession number GSE206023. **The details of the dataset and its accession numbers can be found in Supplementary Table ST1.** The average number of reads per sample was approximately 36 million after trimming, which can also be found in the supplementary material. Quality control of raw and trimmed reads was performed using FastQC v.0.11.8 (Andrews, 2010). Trimming of the adapter content and quality trimming was performed using Trimmomatics v0.36, with the following settings: LEADING 20 TRAILING 20 SLIDINGWINDOW 4:25 MINLEN 31, and the Kallisto index was built with reference transcriptome GRCh38 (Ensembl) with a k-mer length of 31 using the Kallisto v.46.0 programs, and the abundance of the transcripts was quantified using the Kallisto pseudo-alignment, which provides estimates of transcript levels *(17, 52, 53)*. Subsequently, we used the tximport R package to summarize count estimates at the gene level *(54)*.

Differential gene expression analysis

Differential gene expression analysis was performed independently for each dataset using the R package DESeq2 *(55)*. We used the Wald test with the interaction terms vaccine and time point. DEGs were filtered based on the Benjamini-Hochberg adjusted p-values (padj <0.05). We classified up and downregulated genes by log2 fold change ($-1 <$ L2FC > 1, respectively).

Gene set enrichment analysis

The ontology of biological processes within the immune system, specifically the "immune system process" (GO:0002376), was chosen to analyze immune-related DEGs. This dataset was obtained from the GO.db package *(56)*. To facilitate subsequent analyses, biological processes associated with the immune system were categorized into three distinct groups: "Immune system," "Immune subsystem," and, when related to specific cells, classified under the "Immune cell" category. The VAX MSigDB database was obtained via the MSigDB portal and converted into a wide-table format. Subsequently, gene sets were enriched with various attributes through manual annotation derived from the available information on the descriptive pages of the dataset. We used only peripheral blood mononuclear cell-derived samples.

The enrichment analysis of DESeq2 results was conducted with ImmuneGO using the Enricher and Clusterprofiler packages for over-representation analysis (ORA) and gene set enrichment analysis (GSEA) *(57)*. ORA was conducted with COVID-19-related samples and MSigDB Vax collection, using up- and downregulated genes filtered (log2-fold change $\neq 0$, p-adj < 0.05). For GSEA, we used all DEG-enriched gene sets with an adjusted p-value (FDR, Benjamini-Hochberg) of <0.25. Heatmaps were visualized using the ComplexHeatmap package in R and Morpheus (https://software.broadinstitute.org/morpheus) *(58, 59)*.

Principal component analysis and machine learning classification

PCA was performed using L2FC values of immune-related DEGs, through the factoextra R package *(60)*. For cluster analysis, we used the k-nearest neighbors (KNN) algorithm. To optimize the efficiency of our random forest classification, we initially filtered genes based on their cosine squared values in the first two principal components, setting a threshold above 0.5. We then constructed our random forest classifier by leveraging the tidymodels package. Our dataset was partitioned into training (70%) and testing (30%) sets, with stratification for the "type" variable and upsampling to address class imbalance. Employing cross-validation, we fine-tuned the model's hyperparameters using a grid search method within the tidymodels framework (Kuhn & Wickam, 2020). Performance evaluation metrics, including AUC, ROC, OOB, and accuracy, guided the optimization of hyperparameters, including mtry and min_n. Finally, the optimized random forest model was trained on the entire training dataset and subsequently assessed on the testing dataset to evaluate its predictive performance.

Data and code availability

Raw and processed data, as well as codes used are available on the project Github repository (github.com/wapsyed/COVIDvax_atlas).

Author Contributions:

WAPS, DLMF — Conceptualization and manuscript writing;

LV, AL, NC, EC, JSD — Manuscript contributions;

WAPS, DLMF, SZP — Data analysis;

OC, HIN, LFS, ISF, GW, RDC, NOSC, HDD, HDO, ECS, JEK – Review and editing;

GCM, OCM, HIN – Supervision, editing, and approval of the final version.

Funding: We thank the São Paulo Research Foundation (FAPESP grants 2019/14526-0 and 2020/05146-7 to GCM; 2018/18886-9 to OCM; 2023/07806-2 to ISF; 2020/16246-2 and 2023/13356-0 to DLMF;2019/01255-9 and 2021/03684-4 to RDC) for financial support. We acknowledge the National Council for Scientific and Technological Development (CNPq) Brazil (grants: 309482/2022-4 to OCM and 102430/2022-5 to LFS).

Conflicts of Interest: The authors declare that they have no competing interests.

638 **Tables**

639
640 <mark>(This table is also displayed in the main figures file, because it gives an error on formatting when included in this document.)</mark>

Vaccination								
Vaccine	Commercial name	Regimen	Timepoints (days)	Doses	Technology	Adjuvant	GSE ID	Participants
BNT162b2 (BNT)	Comirnarty	HO	0, 1, 6	3x BNT	RNA	No	GSE199750	25
		HO	0, 1, 6	2x BNT	RNA	No	GSE199750	46
		HO	0, 6	1x BNT	RNA	No	GSE199750	67
		HE	0, 1, 3, 7	1x ChAd + 1x BNT	VV + RNA	No	GSE201530	16
		HE	1	2x ChAd + 1x BNT	VV + RNA	No	GSE199750	5
		BI	10, 26, 51	2x BNT + 1x I	RNA + I	No	GSE201530	13
		BI	10, 26, 51	1x BNT (10 dpi) + 1x I	RNA + I	No	GSE189039	5
		AI	10	1x BNT + 1x I + 1x BNT	RNA + I	No	GSE189039	2
		AI	0, 10, 26, 51	1x I + RNA + 1x I	I + RNA + I	No	GSE201530	1
AZD1222 (ChAd)	Covishield, Vaxzebria	HO	0, 1, 3, 6, 7	2x ChAd	VV	No	GSE199750, GSE201533	12

Vaccine	Brand		Timepoints (days)	Doses	Type	Adjuvant	GSE ID	Participants
		HO	0, 3, 6	1x ChAd	VV	No	GSE199750, GSE201533	53
		HE	0, 1, 3, 7	1x ChAd + 1x BNT	VV + RNA	No	GSE201530	17
		HE	1	2x ChAd + 1x BNT	VV + RNA	No	GSE199750	5
mRNA-1273 (MO)	Spikevax	HE	1	2x BNT + 1x MO	RNA	No	GSE199750	10
BBIBP-CorV (BBIBP)	Covilo	HO	0, 7, 14, 28	3 (3rd dose)	IN	Alum	GSE206023	6
ZF2001	Zifivax	HE	0, 7, 14, 28	1 (3rd dose)	SU	Alum	GSE206023	6

Infection						
GSE ID	Prior infection	Prior vaccination	Timepoints (days)	Variant	Severity	Participants
GSE189039	No	Yes	0, 10, 26, 51	Beta	MI (1), S (3)	4
	No	No	0, 10, 26, 51	Beta	MO (2), S (3)	5
GSE201530	No	No	0, 1, 3, 5	Omicron	MI	1
	Yes	No	0, 1, 3, 5	Omicron	MI	6
	No	Yes	0, 1, 3, 5	Omicron	MI	38

643   **Table 1.** Vaccines and infection conditions selected in this study. Legend: HO: Homologous;
644   HE: Heterologous; BI: Before infection; AI: After infection; I: Infection; RNA: mRNA; VV:
645   Viral vector; IN: Inactivated; SU: Subunit. MI: Mild; MO: Moderate; S: Severe; A:
646   Asymptomatic.

Figures

Fig. 1. (A) Methods and description of datasets and study population. We manually curated the GEO datasets and performed differential gene expression analysis. Next, we determined the overlapping genes between all the conditions. Finally, genes from immune system biological process (GO:) and the VAX collection from MSigDB were used in GSEA and ORA, and individual gene expression analysis was analyzed across the studied conditions. Created with Biorender. Number of participants (B) and samples (C) were categorized by vaccine and infection conditions. Legend: BBIBP: BBIBP-CORV; BNT: BNT162b2, ChAd: ChAdOx-1; MO: mRNA-1273; I: Infected; V: Vaccinated.

Fig. 2. A. Differentially expressed genes (DEGs) were categorized as up- (left panel) and down-regulated (right panel) with the absolute and relative number of genes. The conditions were then plotted with the total number of genes and immune genes. Labels correspond to the total number of immune genes and the percentage over total DEGs. B. Shared DEGs between infection and vaccination. The upper-left cluster include individuals infected for the first time, and the bottom-right cluster comprise the first week of vaccination by RNA, VV, and SU vaccines. C. Twenty most shared immune-related genes. Legend: BBIBP: BBIBP-CORV; BNT: BNT162b2, ChAd: ChAdOx-1; MO: mRNA-1273; I: Infected; V: Vaccinated; IN: Inactivated; RNA: mRNA vaccines; SU: Subunit; VV: Viral vector. B. Shared DEGs in vaccination and infection.

Fig. 3. Principal component analysis. A. Two principal components were employed to plot all conditions, capturing 86.2% of the total variance. K-nearest neighbors (KNN) clustering identified three distinct clusters. Cluster 1 (C1) includes infected patients and overlaps with Cluster 2 (C2). C2 is predominantly characterized by vaccination in healthy individuals,

encompassing convalescent patients and non-convalescent individuals. Cluster 3 (C3)
comprises the initial three days post-vaccination with mRNA vaccines and ChAd. B. The
loading plot displays cos2 values for the top 20 genes in PC1 and PC2. The top 20 genes
contributing to PC1 (C) and PC2 (D), along with the corresponding Log2 Fold Change
(L2FC). Genes were displayed with correspondent log2-fold change ranging from -2 (blue) to
2 (red). The left annotation on the heatmap depicts the contribution to different immune
processes and cells, with colors corresponding to processes and cells related to the
complement system (pink), the innate (greens) and adaptive immune systems (purples), and
general processes and cells (leukocytes, black). Legend: BBIBP: BBIBP-CORV; BNT:
BNT162b2, ChAd: ChAdOx-1; MO: mRNA-1273; I: Infected; V: Vaccinated.

Fig. 4. DEGs with cos2 > 0.5 were used to train a random forest model to predict the types of
vaccines and infection, and the ten most important genes were selected (A). The final model
was selected after an iterative process of hyperparameter tuning. The best model was defined
with 2000 trees in the forest, where mtry was set to 6 and min_n to 6. The most important
genes were used to plot the log2-fold change (L2FC) (B). C. Gene set enrichment analysis of
COVID-19 vaccines and infection. Major immunological gene sets were enriched with GSEA.
Normalized enrichment scores (NES) for enriched gene sets, with circle size corresponding
to -log(qvalue). D. Representation of upregulated (red) and downregulated (blue) genes in
COVID-19 conditions associated with various immune biological processes. Visualization IN

stacked hierarchical layout was generated using Cytoscape *(61)*. Node size is determined by
degree centrality, and node colors are determined by type and infection status. Illustration
made in **BioRender.** Legend: BBIBP: BBIBP-CORV; BNT: BNT162b2, ChAd: ChAdOx-1;
MO: mRNA-1273; I: Infected; V: Vaccinated.

Fig. 5. Gene-level comparison of COVID-19 vaccines and the MSigDB Vax collection.
Overlapping immune-related (A) and other (B) genes between conditions. C. Genes with the
most sharing between COVID-19 vaccination and other vaccines. Visualization in stacked
hierarchical layout was generated using Cytoscape *(61)*. C. All immune-related genes shared

at least once between conditions. D. Genes included in the first cluster (see complete heatmap
in **Fig. S9**) were set with L2FC as 1 (upregulated) and -1 (downregulated). These genes were
annotated by their corresponding contribution to subprocesses and cells among the innate,
adaptive and complement immune systems (left). E. Over-representation analysis of vaccines
against COVID-19 and other pathogens with immune cells and processes. Legend: BBIBP:
BBIBP-CORV; BNT: BNT162b2, ChAd: ChAdOx-1; MO: mRNA-1273; I: Infected; V:
Vaccinated.

Supplementary Material

Supplementary Figures

Fig. S1. Data curation. Datasets were obtained from NCBI BioProject and curated. All datasets, including the selected and non-selected, were classified by vaccine technology and vaccination regimen, and by sample source.

Fig. S2. Description of the selected datasets. The datasets include healthy vaccinees without
prior infection, infected patients without vaccination, and patients with prior full or
incomplete vaccination. Sample collection timelines are shown as black lines, with red lines
indicating the first sample collection during hospitalization and gray lines representing events
without sample collection. Days post-infection or vaccination are depicted, with blue lines
representing hospitalization periods and purple lines indicating dexamethasone treatment.

Fig. S3. DEGs were categorized as up- (A) and down-regulated (B) with the absolute and
relative number of genes, and grouped based on infection (red) and vaccination (blue)

conditions. The conditions were then plotted with the relative number of genes corresponding
to their roles in essential biological processes, namely immune system, cellular processes,
metabolism, and regulation of biological processes, respectively. Shapes correspond to type,
colors to disease or infection, and size to total number of DEGs. I: Infection; IN: Inactivated;
RNA: mRNA vaccines; SU: Subunit; VV: Viral vector.

Fig. S4. A. Shared non-immune-related DEGs between infection and vaccination. B. Twenty
most shared immune-related genes. BBIBP: BBIBP-CORV; BNT: BNT162b2, ChAd:
ChAdOx-1; MO: mRNA-1273; I: Infected; V: Vaccinated; IN: Inactivated; RNA: mRNA
vaccines; SU: Subunit; VV: Viral vector.

Fig. S5. Genes related to complement system, antimicrobial response, and innate immune
system processes and cells. I: Infection; IN: Inactivated; RNA: mRNA vaccines; SU: Subunit;
VV: Viral vector.

Fig. S6. Genes related to humoral adaptive immune system, including B cell processes and
immunoglobulin-mediated response. I: Infection; IN: Inactivated; RNA: mRNA vaccines;
SU: Subunit; VV: Viral vector.

747 **Fig. S7.** Genes related to the cellular adaptive immune system, including cellular response
748 and T- and T helper cell processes. I: Infection; IN: Inactivated; RNA: mRNA vaccines; SU:
749 Subunit; VV: Viral vector.

 Fig. S8. General leukocyte processes. I: Infection; IN: Inactivated; RNA: mRNA vaccines;
752 SU: Subunit; VV: Viral vector.

754 **Fig. S9.** Gene expression in vaccination against COVID-19 and other pathogens. I: Infection;
755 IN: Inactivated; RNA: mRNA vaccines; SU: Subunit; VV: Viral vector; CONJ: Conjugated;
756 LA: Live-attenuated; VLP: Virus-like particle.

References

References

1. C. Wang, P. W. Horby, F. G. Hayden, G. F. Gao, A novel coronavirus outbreak of global
health concern. *The Lancet* **395**, 470–473 (2020).

2. H. W. Chung, C. Apio, T. Goo, G. Heo, K. Han, T. Kim, H. Kim, Y. Ko, D. Lee, J. Lim, S.
Lee, T. Park, Effects of government policies on the spread of COVID-19 worldwide. *Sci Rep*
11, 20495 (2021).

3. S. Xia, L. Wang, Y. Zhu, L. Lu, S. Jiang, Origin, virological features, immune evasion and
intervention of SARS-CoV-2 Omicron sublineages. *Signal Transduct Target Ther* **7**, 241
(2022).

4. O. J. Watson, G. Barnsley, J. Toor, A. B. Hogan, P. Winskill, A. C. Ghani, Global impact
of the first year of COVID-19 vaccination: a mathematical modelling study. *Lancet Infect Dis*
22, 1293–1302 (2022).

5. Y. Li, R. Tenchov, J. Smoot, C. Liu, S. Watkins, Q. Zhou, A Comprehensive Review of the
Global Efforts on COVID-19 Vaccine Development. *ACS Cent Sci* **7**, 512–533 (2021).

6. World Health Organization (WHO), Status of COVID-19 Vaccines within WHO EUL/PQ
evaluation
process*https://extranet.who.int/prequal/sites/default/files/document_files/Status_COVID_VA*
X_08AUgust2023.pdf (2023).

7. E. Mathieu, H. Ritchie, E. Ortiz-Ospina, M. Roser, J. Hasell, C. Appel, C. Giattino, L.
Rodés-Guirao, A global database of COVID-19 vaccinations. *Nat Hum Behav* **5**, 947–953
(2021).

8. R. L. Atmar, K. E. Lyke, M. E. Deming, L. A. Jackson, A. R. Branche, H. M. El Sahly, C.
A. Rostad, J. M. Martin, C. Johnston, R. E. Rupp, M. J. Mulligan, R. C. Brady, R. W. Frenck,
M. Bäcker, A. C. Kottkamp, T. M. Babu, K. Rajakumar, S. Edupuganti, D. Dobrzynski, R. N.
Coler, C. M. Posavad, J. I. Archer, S. Crandon, S. U. Nayak, D. Szydlo, J. A. Zemanek, C. P.
Dominguez Islas, E. R. Brown, M. S. Suthar, M. J. McElrath, A. B. McDermott, S. E.
O'Connell, D. C. Montefiori, A. Eaton, K. M. Neuzil, D. S. Stephens, P. C. Roberts, J. H.
Beigel, Homologous and Heterologous Covid-19 Booster Vaccinations. *New England Journal*
of Medicine **386**, 1046–1057 (2022).

9. Z. S. Vasconcelos, H. A. Salem, S. P. Veiga, F. E. B. de Lima, C. R. da S. Gonçalves, E.
C. dos Santos, A. R. J. Brandão, K. N. Couceiro, J. A. de O. Guerra, M. das G. V. B. Guerra,
Immunogenicity Characterization of COVID-19 Vaccines: A Systematic Review and Meta-
analysis. *Rev Soc Bras Med Trop* **56** (2023), doi:10.1590/0037-8682-0661-2022.

10. A. J. Pollard, E. M. Bijker, A guide to vaccinology: from basic principles to new
developments. *Nat Rev Immunol* **21**, 83–100 (2021).

11. S. A. Costa Clemens, L. Weckx, R. Clemens, A. V. Almeida Mendes, A. Ramos Souza,
M. B. V Silveira, S. N. F. da Guarda, M. M. de Nobrega, M. I. de Moraes Pinto, I. G. S.
Gonzalez, N. Salvador, M. M. Franco, R. N. de Avila Mendonça, I. S. Queiroz Oliveira, B. S.
de Freitas Souza, M. Fraga, P. Aley, S. Bibi, L. Cantrell, W. Dejnirattisai, X. Liu, J.
Mongkolsapaya, P. Supasa, G. R. Screaton, T. Lambe, M. Voysey, A. J. Pollard, M. Bittaye,
D. Woods, S. Davies, H. Smith, M. Ulaszewska, H. Sanders, R. Mabette, S. Vernon, Z. Valliji,

G. Mead, C. Tejpal, J. Park, A. Beveridge, A. Eldawi, S. Felle, M. Fraga, T. Muniz Martins,
C. L. Martins Medrado, L. J. de Arruda Cordeiro Matos, Heterologous versus homologous
COVID-19 booster vaccination in previous recipients of two doses of CoronaVac COVID-19
vaccine in Brazil (RHH-001): a phase 4, non-inferiority, single blind, randomised study. *The*
Lancet **399**, 521–529 (2022).

12. L. R. Baden, H. M. El Sahly, B. Essink, K. Kotloff, S. Frey, R. Novak, D. Diemert, S. A.
Spector, N. Rouphael, C. B. Creech, J. McGettigan, S. Khetan, N. Segall, J. Solis, A. Brosz,
C. Fierro, H. Schwartz, K. Neuzil, L. Corey, P. Gilbert, H. Janes, D. Follmann, M. Marovich,
J. Mascola, L. Polakowski, J. Ledgerwood, B. S. Graham, H. Bennett, R. Pajon, C. Knightly,
B. Leav, W. Deng, H. Zhou, S. Han, M. Ivarsson, J. Miller, T. Zaks, Efficacy and Safety of
the mRNA-1273 SARS-CoV-2 Vaccine. *New England Journal of Medicine* **384**, 403–416
(2021).

13. M. Voysey, S. A. C. Clemens, S. A. Madhi, L. Y. Weckx, P. M. Folegatti, P. K. Aley, B.
Angus, V. L. Baillie, S. L. Barnabas, Q. E. Bhorat, S. Bibi, C. Briner, P. Cicconi, A. M.
Collins, R. Colin-Jones, C. L. Cutland, T. C. Darton, K. Dheda, C. J. A. Duncan, K. R. W.
Emary, K. J. Ewer, L. Fairlie, S. N. Faust, S. Feng, D. M. Ferreira, A. Finn, A. L. Goodman,
C. M. Green, C. A. Green, P. T. Heath, C. Hill, H. Hill, I. Hirsch, S. H. C. Hodgson, A. Izu,
S. Jackson, D. Jenkin, C. C. D. Joe, S. Kerridge, A. Koen, G. Kwatra, R. Lazarus, A. M.
Lawrie, A. Lelliott, V. Libri, P. J. Lillie, R. Mallory, A. V. A. Mendes, E. P. Milan, A. M.
Minassian, A. McGregor, H. Morrison, Y. F. Mujadidi, A. Nana, P. J. O'Reilly, S. D.
Padayachee, A. Pittella, E. Plested, K. M. Pollock, M. N. Ramasamy, S. Rhead, A. V
Schwarzbold, N. Singh, A. Smith, R. Song, M. D. Snape, E. Sprinz, R. K. Sutherland, R.
Tarrant, E. C. Thomson, M. E. Török, M. Toshner, D. P. J. Turner, J. Vekemans, T. L.
Villafana, M. E. E. Watson, C. J. Williams, A. D. Douglas, A. V. S. Hill, T. Lambe, S. C.
Gilbert, A. J. Pollard, M. Aban, F. Abayomi, K. Abeyskera, J. Aboagye, M. Adam, K. Adams,
J. Adamson, Y. A. Adelaja, G. Adewetan, S. Adlou, K. Ahmed, Y. Akhalwaya, S. Akhalwaya,
A. Alcock, A. Ali, E. R. Allen, L. Allen, T. C. D. S. C. Almeida, M. P. S. Alves, F. Amorim,
F. Andritsou, R. Anslow, M. Appleby, E. H. Arbe-Barnes, M. P. Ariaans, B. Arns, L. Arruda,
P. Azi, L. Azi, G. Babbage, C. Bailey, K. F. Baker, M. Baker, N. Baker, P. Baker, L. Baldwin,
I. Baleanu, D. Bandeira, A. Bara, M. A. S. Barbosa, D. Barker, G. D. Barlow, E. Barnes, A.
S. Barr, J. R. Barrett, J. Barrett, L. Bates, A. Batten, K. Beadon, E. Beales, R. Beckley, S.
Belij-Rammerstorfer, J. Bell, D. Bellamy, N. Bellei, S. Belton, A. Berg, L. Bermejo, E. Berrie,
L. Berry, D. Berzenyi, A. Beveridge, K. R. Bewley, H. Bexhell, S. Bhikha, A. E. Bhorat, Z.
E. Bhorat, E. Bijker, G. Birch, S. Birch, A. Bird, O. Bird, K. Bisnauthsing, M. Bittaye, K.
Blackstone, L. Blackwell, H. Bletchly, C. L. Blundell, S. R. Blundell, P. Bodalia, B. C.
Boettger, E. Bolam, E. Boland, D. Bormans, N. Borthwick, F. Bowring, A. Boyd, P. Bradley,
T. Brenner, P. Brown, C. Brown, C. Brown-O'Sullivan, S. Bruce, E. Brunt, R. Buchan, W.
Budd, Y. A. Bulbulia, M. Bull, J. Burbage, H. Burhan, A. Burn, K. R. Buttigieg, N. Byard, I.
Cabera Puig, G. Calderon, A. Calvert, S. Camara, M. Cao, F. Cappuccini, J. R. Cardoso, M.
Carr, M. W. Carroll, A. Carson-Stevens, Y. de M. Carvalho, J. A. M. Carvalho, H. R. Casey,
P. Cashen, T. Castro, L. C. Castro, K. Cathie, A. Cavey, J. Cerbino-Neto, J. Chadwick, D.
Chapman, S. Charlton, I. Chelysheva, O. Chester, S. Chita, J.-S. Cho, L. Cifuentes, E. Clark,
M. Clark, A. Clarke, E. A. Clutterbuck, S. L. K. Collins, C. P. Conlon, S. Connarty, N.
Coombes, C. Cooper, R. Cooper, L. Cornelissen, T. Corrah, C. Cosgrove, T. Cox, W. E. M.
Crocker, S. Crosbie, L. Cullen, D. Cullen, D. R. M. F. Cunha, C. Cunningham, F. C.
Cuthbertson, S. N. F. Da Guarda, L. P. da Silva, B. E. Damratoski, Z. Danos, M. T. D. C.
Dantas, P. Darroch, M. S. Datoo, C. Datta, M. Davids, S. L. Davies, H. Davies, E. Davis, J.
Davis, J. Davis, M. M. D. De Nobrega, L. M. De Oliveira Kalid, D. Dearlove, T. Demissie,

A. Desai, S. Di Marco, C. Di Maso, M. I. S. Dinelli, T. Dinesh, C. Docksey, C. Dold, T. Dong,
F. R. Donnellan, T. Dos Santos, T. G. dos Santos, E. P. Dos Santos, N. Douglas, C. Downing,
J. Drake, R. Drake-Brockman, K. Driver, R. Drury, S. J. Dunachie, B. S. Durham, L. Dutra,
N. J. W. Easom, S. van Eck, M. Edwards, N. J. Edwards, O. M. El Muhanna, S. C. Elias, M.
Elmore, M. English, A. Esmail, Y. M. Essack, E. Farmer, M. Farooq, M. Farrar, L. Farrugia,
B. Faulkner, S. Fedosyuk, S. Felle, S. Feng, C. Ferreira Da Silva, S. Field, R. Fisher, A.
Flaxman, J. Fletcher, H. Fofie, H. Fok, K. J. Ford, J. Fowler, P. H. A. Fraiman, E. Francis, M.
M. Franco, J. Frater, M. S. M. Freire, S. H. Fry, S. Fudge, J. Furze, M. Fuskova, P. Galian-
Rubio, E. Galiza, H. Garlant, M. Gavrila, A. Geddes, K. A. Gibbons, C. Gilbride, H. Gill, S.
Glynn, K. Godwin, K. Gokani, U. C. Goldoni, M. Goncalves, I. G. S. Gonzalez, J. Goodwin,
A. Goondiwala, K. Gordon-Quayle, G. Gorini, J. Grab, L. Gracie, M. Greenland, N.
Greenwood, J. Greffrath, M. M. Groenewald, L. Grossi, G. Gupta, M. Hackett, B. Hallis, M.
Hamaluba, E. Hamilton, J. Hamlyn, D. Hammersley, A. T. Hanrath, B. Hanumunthadu, S. A.
Harris, C. Harris, T. Harris, T. D. Harrison, D. Harrison, T. C. Hart, B. Hartnell, S. Hassan, J.
Haughney, S. Hawkins, J. Hay, I. Head, J. Henry, M. Hermosin Herrera, D. B. Hettle, J. Hill,
G. Hodges, E. Horne, M. M. Hou, C. Houlihan, E. Howe, N. Howell, J. Humphreys, H. E.
Humphries, K. Hurley, C. Huson, A. Hyder-Wright, C. Hyams, S. Ikram, A. Ishwarbhai, M.
Ivan, P. Iveson, V. Iyer, F. Jackson, J. De Jager, S. Jaumdally, H. Jeffers, N. Jesudason, B.
Jones, K. Jones, E. Jones, C. Jones, M. R. Jorge, A. Jose, A. Joshi, E. A. M. S. Júnior, J.
Kadziola, R. Kailath, F. Kana, K. Karampatsas, M. Kasanyinga, J. Keen, E. J. Kelly, D. M.
Kelly, D. Kelly, S. Kelly, D. Kerr, R. de Á. Kfouri, L. Khan, B. Khozoee, S. Kidd, A. Killen,
J. Kinch, P. Kinch, L. D. W. King, T. B. King, L. Kingham, P. Klenerman, F. Knapper, J. C.
Knight, D. Knott, S. Koleva, M. Lang, G. Lang, C. W. Larkworthy, J. P. J. Larwood, R. Law,
E. M. Lazarus, A. Leach, E. A. Lees, N.-M. Lemm, A. Lessa, S. Leung, Y. Li, A. M. Lias, K.
Liatsikos, A. Linder, S. Lipworth, S. Liu, X. Liu, A. Lloyd, S. Lloyd, L. Loew, R. Lopez
Ramon, L. Lora, V. Lowthorpe, K. Luz, J. C. MacDonald, G. MacGregor, M. Madhavan, D.
O. Mainwaring, E. Makambwa, R. Makinson, M. Malahleha, R. Malamatsho, G. Mallett, K.
Mansatta, T. Maoko, K. Mapetla, N. G. Marchevsky, S. Marinou, E. Marlow, G. N. Marques,
P. Marriott, R. P. Marshall, J. L. Marshall, F. J. Martins, M. Masenya, M. Masilela, S. K.
Masters, M. Mathew, H. Matlebjane, K. Matshidiso, O. Mazur, A. Mazzella, H. McCaughan,
J. McEwan, J. McGlashan, L. McInroy, Z. McIntyre, D. McLenaghan, N. McRobert, S.
McSwiggan, C. Megson, S. Mehdipour, W. Meijs, R. N. Á. Mendonça, A. J. Mentzer, N.
Mirtorabi, C. Mitton, S. Mnyakeni, F. Moghaddas, K. Molapo, M. Moloi, M. Moore, M. I.
Moraes-Pinto, M. Moran, E. Morey, R. Morgans, S. Morris, S. Morris, H. C. Morris, F.
Morselli, G. Morshead, R. Morter, L. Mottal, A. Moultrie, N. Moya, M. Mpelembue, S.
Msomi, Y. Mugodi, E. Mukhopadhyay, J. Muller, A. Munro, C. Munro, S. Murphy, P. Mweu,
C. H. Myasaki, G. Naik, K. Naker, E. Nastouli, A. Nazir, B. Ndlovu, F. Neffa, C. Njenga, H.
Noal, A. Noé, G. Novaes, F. L. Nugent, G. Nunes, K. O'Brien, D. O'Connor, M. Odam, S.
Oelofse, B. Oguti, V. Olchawski, N. J. Oldfield, M. G. Oliveira, C. Oliveira, A. Oosthuizen,
P. O'Reilly, P. Osborne, D. R. J. Owen, L. Owen, D. Owens, N. Owino, M. Pacurar, B. V. B.
Paiva, E. M. F. Palhares, S. Palmer, S. Parkinson, H. M. R. T. Parracho, K. Parsons, D. Patel,
B. Patel, F. Patel, K. Patel, M. Patrick-Smith, R. O. Payne, Y. Peng, E. J. Penn, A. Pennington,
M. P. Peralta Alvarez, J. Perring, N. Perry, R. Perumal, S. Petkar, T. Philip, D. J. Phillips, J.
Phillips, M. K. Phohu, L. Pickup, S. Pieterse, J. Piper, D. Pipini, M. Plank, J. Du Plessis, S.
Pollard, J. Pooley, A. Pooran, I. Poulton, C. Powers, F. B. Presa, D. A. Price, V. Price, M.
Primeira, P. C. Proud, S. Provstgaard-Morys, S. Pueschel, D. Pulido, S. Quaid, R. Rabara, A.
Radford, K. Radia, D. Rajapaska, T. Rajeswaran, A. S. F. Ramos, F. Ramos Lopez, T.
Rampling, J. Rand, H. Ratcliffe, T. Rawlinson, D. Rea, B. Rees, J. Reiné, M. Resuello-Dauti,
E. Reyes Pabon, C. M. Ribiero, M. Ricamara, A. Richter, N. Ritchie, A. J. Ritchie, A. J.
Robbins, H. Roberts, R. E. Robinson, H. Robinson, T. T. Rocchetti, B. P. Rocha, S. Roche,

C. Rollier, L. Rose, A. L. Ross Russell, L. Rossouw, S. Royal, I. Rudiansyah, S. Ruiz, S. Saich, C. Sala, J. Sale, A. M. Salman, N. Salvador, S. Salvador, M. Sampaio, A. D. Samson, A. Sanchez-Gonzalez, H. Sanders, K. Sanders, E. Santos, M. F. S. Santos Guerra, I. Satti, J. E. Saunders, C. Saunders, A. Sayed, I. Schim van der Loeff, A. B. Schmid, E. Schofield, G. Screaton, S. Seddiqi, R. R. Segireddy, R. Senger, S. Serrano, R. Shah, I. Shaik, H. E. Sharpe, K. Sharrocks, R. Shaw, A. Shea, A. Shepherd, J. G. Shepherd, F. Shiham, E. Sidhom, S. E. Silk, A. C. da Silva Moraes, G. Silva-Junior, L. Silva-Reyes, A. D. Silveira, M. B. V. Silveira, J. Sinha, D. T. Skelly, D. C. Smith, N. Smith, H. E. Smith, D. J. Smith, C. C. Smith, A. Soares, T. Soares, C. Solórzano, G. L. Sorio, K. Sorley, T. Sosa-Rodriguez, C. M. C. D. L. Souza, B. S. D. F. Souza, A. R. Souza, A. J. Spencer, F. Spina, L. Spoors, L. Stafford, I. Stamford, I. Starinskij, R. Stein, J. Steven, L. Stockdale, L. V. Stockwell, L. H. Strickland, A. C. Stuart, A. Sturdy, N. Sutton, A. Szigeti, A. Tahiri-Alaoui, R. Tanner, C. Taoushanis, A. W. Tarr, K. Taylor, U. Taylor, I. J. Taylor, J. Taylor, R. te Water Naude, Y. Themistocleous, A. Themistocleous, M. Thomas, K. Thomas, T. M. Thomas, A. Thombrayil, F. Thompson, A. Thompson, K. Thompson, A. Thompson, J. Thomson, V. Thornton-Jones, P. J. Tighe, L. A. Tinoco, G. Tiongson, B. Tladinyane, M. Tomasicchio, A. Tomic, S. Tonks, J. Towner, N. Tran, J. Tree, G. Trillana, C. Trinham, R. Trivett, A. Truby, B. L. Tsheko, A. Turabi, R. Turner, C. Turner, M. Ulaszewska, B. R. Underwood, R. Varughese, D. Verbart, M. Verheul, I. Vichos, T. Vieira, C. S. Waddington, L. Walker, E. Wallis, M. Wand, D. Warbick, T. Wardell, G. Warimwe, S. C. Warren, B. Watkins, E. Watson, S. Webb, A. Webb-Bridges, A. Webster, J. Welch, J. Wells, A. West, C. White, R. White, P. Williams, R. L. Williams, R. Winslow, M. Woodyer, A. T. Worth, D. Wright, M. Wroblewska, A. Yao, R. Zimmer, D. Zizi, P. Zuidewind, Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. *The Lancet* **397**, 99–111 (2021).

14. L. Dai, L. Gao, L. Tao, S. R. Hadinegoro, M. Erkin, Z. Ying, P. He, R. T. Girsang, H. Vergara, J. Akram, H. I. Satari, T. Khaliq, U. Sughra, A. P. Celi, F. Li, Y. Li, Z. Jiang, D. Dalimova, J. Tuychiev, S. Turdikulova, A. Ikram, N. Flores Lastra, F. Ding, M. Suhardono, E. Fadlyana, J. Yan, Z. Hu, C. Li, I. Y. Abdurakhmonov, G. F. Gao, Efficacy and Safety of the RBD-Dimer–Based Covid-19 Vaccine ZF2001 in Adults. *New England Journal of Medicine* **386**, 2097–2111 (2022).

15. N. Al Kaabi, Y. Zhang, S. Xia, Y. Yang, M. M. Al Qahtani, N. Abdulrazzaq, M. Al Nusair, M. Hassany, J. S. Jawad, J. Abdalla, S. E. Hussein, S. K. Al Mazrouei, M. Al Karam, X. Li, X. Yang, W. Wang, B. Lai, W. Chen, S. Huang, Q. Wang, T. Yang, Y. Liu, R. Ma, Z. M. Hussain, T. Khan, M. Saifuddin Fasihuddin, W. You, Z. Xie, Y. Zhao, Z. Jiang, G. Zhao, Y. Zhang, S. Mahmoud, I. ElTantawy, P. Xiao, A. Koshy, W. A. Zaher, H. Wang, K. Duan, A. Pan, X. Yang, Effect of 2 Inactivated SARS-CoV-2 Vaccines on Symptomatic COVID-19 Infection in Adults. *JAMA* **326**, 35 (2021).

16. N. Wu, K. Joyal-Desmarais, P. A. B. Ribeiro, A. M. Vieira, J. Stojanovic, C. Sanuade, D. Yip, S. L. Bacon, Long-term effectiveness of COVID-19 vaccines against infections, hospitalisations, and mortality in adults: findings from a rapid living systematic evidence synthesis and meta-analysis up to December, 2022. *Lancet Respir Med* **11**, 439–452 (2023).

17. N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* **34**, 525–527 (2016).

18. D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907–915 (2019).

19. K. Romero Starke, D. Reissig, G. Petereit-Haack, S. Schmauder, A. Nienhaus, A. Seidler,
The isolated effect of age on the risk of COVID-19 severe outcomes: a systematic review with
meta-analysis. *BMJ Glob Health* **6**, e006434 (2021).

20. L. Knabl, H. K. Lee, M. Wieser, A. Mur, A. Zabernigg, L. Knabl, S. Rauch, M. Bock, J.
Schumacher, N. Kaiser, P. A. Furth, L. Hennighausen, BNT162b2 vaccination enhances
interferon-JAK-STAT-regulated antiviral programs in COVID-19 patients infected with the
SARS-CoV-2 Beta variant. *Communications Medicine* **2**, 17 (2022).

21. Haseeb, Pirzada, Ain, Choi, Wnt Signaling in the Regulation of Immune Cell and Cancer
Therapeutics. *Cells* **8**, 1380 (2019).

22. Y. Xiao, V. Peperzak, A. M. Keller, J. Borst, CD27 Instructs CD4+ T Cells to Provide
Help for the Memory CD8+ T Cell Response after Protein Immunization. *The Journal of
Immunology* **181**, 1071–1082 (2008).

23. T. Mohan, W. Zhu, Y. Wang, B.-Z. Wang, Applications of chemokines as adjuvants for
vaccine immunotherapy. *Immunobiology* **223**, 477–485 (2024).

24. A. B. Adewoye, N. Shrine, L. Odenthal-Hesse, S. Welsh, A. Malarstig, S. Jelinsky, I.
Kilty, M. D. Tobin, E. J. Hollox, L. V Wain, Human CCL3L1 copy number variation, gene
expression, and the role of the CCL3L1-CCR5 axis in lung function. *Wellcome Open Res* **3**,
13 (2024).

25. M. Merad, C. A. Blish, F. Sallusto, A. Iwasaki, The immunology and immunopathology
of COVID-19. *Science (1979)* **375**, 1122–1127 (2022).

26. B. Afzali, M. Noris, B. N. Lambrecht, C. Kemper, The state of complement in COVID-
19. *Nat Rev Immunol* **22**, 77–84 (2022).

27. A. Almuqrin, A. D. Davidson, M. K. Williamson, P. A. Lewis, K. J. Heesom, S. Morris,
S. C. Gilbert, D. A. Matthews, SARS-CoV-2 vaccine ChAdOx1 nCoV-19 infection of human
cell lines reveals low levels of viral backbone gene transcription alongside very high levels of
SARS-CoV-2 S glycoprotein gene transcription. *Genome Med* **13**, 43 (2021).

28. A. Liberzon, A. Subramanian, R. Pinchback, H. Thorvaldsdóttir, P. Tamayo, J. P. Mesirov,
Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**, 1739–1740 (2011).

29. C. A. DiazGranados, A. J. Dunning, M. Kimmel, D. Kirby, J. Treanor, A. Collins, R.
Pollak, J. Christoff, J. Earl, V. Landolfi, E. Martin, S. Gurunathan, R. Nathan, D. P.
Greenberg, N. G. Tornieporth, M. D. Decker, H. K. Talbot, Efficacy of High-Dose versus
Standard-Dose Influenza Vaccine in Older Adults. *New England Journal of Medicine* **371**,
635–645 (2014).

30. F. Piras-Douce, F. Raynal, A. Raquin, Y. Girerd-Chambaz, S. Gautheron, M. E. N.
Sanchez, M. Vangelisti, N. Mantel, Next generation live-attenuated yellow fever vaccine
candidate: Safety and immuno-efficacy in small animal models. *Vaccine* **39**, 1846–1856
(2021).

31. R. H. Behrens, V. Patel, Avoiding shoulder injury from intramuscular vaccines. *The
Lancet* **397**, 471 (2021).

32. M. Salgado-Albarrán, E. I. Navarro-Delgado, A. Del Moral-Morales, N. Alcaraz, J.
Baumbach, R. González-Barrios, E. Soto-Reyes, Comparative transcriptome analysis reveals
key epigenetic targets in SARS-CoV-2 infection. *NPJ Syst Biol Appl* **7**, 21 (2021).

33. T. M. Delorey, C. G. K. Ziegler, G. Heimberg, R. Normand, Y. Yang, Å. Segerstolpe, D.
Abbondanza, S. J. Fleming, A. Subramanian, D. T. Montoro, K. A. Jagadeesh, K. K. Dey, P.
Sen, M. Slyper, Y. H. Pita-Juárez, D. Phillips, J. Biermann, Z. Bloom-Ackermann, N. Barkas,
A. Ganna, J. Gomez, J. C. Melms, I. Katsyv, E. Normandin, P. Naderi, Y. V. Popov, S. S.
Raju, S. Niezen, L. T.-Y. Tsai, K. J. Siddle, M. Sud, V. M. Tran, S. K. Vellarikkal, Y. Wang,
L. Amir-Zilberstein, D. S. Atri, J. Beechem, O. R. Brook, J. Chen, P. Divakar, P. Dorceus, J.
M. Engreitz, A. Essene, D. M. Fitzgerald, R. Fropf, S. Gazal, J. Gould, J. Grzyb, T. Harvey,
J. Hecht, T. Hether, J. Jané-Valbuena, M. Leney-Greene, H. Ma, C. McCabe, D. E.
McLoughlin, E. M. Miller, C. Muus, M. Niemi, R. Padera, L. Pan, D. Pant, C. Pe'er, J.
Pfiffner-Borges, C. J. Pinto, J. Plaisted, J. Reeves, M. Ross, M. Rudy, E. H. Rueckert, M.
Siciliano, A. Sturm, E. Todres, A. Waghray, S. Warren, S. Zhang, D. R. Zollinger, L. Cosimi,
R. M. Gupta, N. Hacohen, H. Hibshoosh, W. Hide, A. L. Price, J. Rajagopal, P. R. Tata, S.
Riedel, G. Szabo, T. L. Tickle, P. T. Ellinor, D. Hung, P. C. Sabeti, R. Novak, R. Rogers, D.
E. Ingber, Z. G. Jiang, D. Juric, M. Babadi, S. L. Farhi, B. Izar, J. R. Stone, I. S. Vlachos, I.
H. Solomon, O. Ashenberg, C. B. M. Porter, B. Li, A. K. Shalek, A.-C. Villani, O. Rozenblatt-
Rosen, A. Regev, COVID-19 tissue atlases reveal SARS-CoV-2 pathology and cellular
targets. *Nature* **595**, 107–113 (2021).

34. N. L. Pereira, F. Ahmad, M. Byku, N. W. Cummins, A. A. Morris, A. Owens, S. Tuteja,
S. Cresci, COVID-19: Understanding Inter-Individual Variability and Implications for
Precision Medicine. *Mayo Clin Proc* **96**, 446–463 (2021).

35. F. Liang, G. Lindgren, A. Lin, E. A. Thompson, S. Ols, J. Röhss, S. John, K. Hassett, O.
Yuzhakov, K. Bahl, L. A. Brito, H. Salter, G. Ciaramella, K. Loré, Efficient Targeting and
Activation of Antigen-Presenting Cells In Vivo after Modified mRNA Vaccine
Administration in Rhesus Macaques. *Molecular Therapy* **25**, 2635–2647 (2017).

36. K. Karikó, H. Muramatsu, F. A. Welsh, J. Ludwig, H. Kato, S. Akira, D. Weissman,
Incorporation of Pseudouridine Into mRNA Yields Superior Nonimmunogenic Vector With
Increased Translational Capacity and Biological Stability. *Molecular Therapy* **16**, 1833–1840
(2008).

37. X. Xia, Detailed Dissection and Critical Evaluation of the Pfizer/BioNTech and Moderna
mRNA Vaccines. *Vaccines (Basel)* **9**, 734 (2021).

38. M. D. J. Dicks, A. J. Spencer, N. J. Edwards, G. Wadell, K. Bojang, S. C. Gilbert, A. V.
S. Hill, M. G. Cottingham, A Novel Chimpanzee Adenovirus Vector with Low Human
Seroprevalence: Improved Systems for Vector Derivation and Comparative Immunogenicity.
PLoS One **7**, e40385 (2012).

39. T. Travieso, J. Li, S. Mahesh, J. D. F. R. E. Mello, M. Blasi, The use of viral vectors in
vaccine development. *NPJ Vaccines* **7**, 75 (2022).

40. P. Nordström, M. Ballin, A. Nordström, Effectiveness of heterologous ChAdOx1 nCoV-
19 and mRNA prime-boost vaccination against symptomatic Covid-19 infection in Sweden:
A nationwide cohort study. *The Lancet Regional Health - Europe* **11**, 100249 (2021).

41. H. Wang, Y. Zhang, B. Huang, W. Deng, Y. Quan, W. Wang, W. Xu, Y. Zhao, N. Li, J.
Zhang, H. Liang, L. Bao, Y. Xu, L. Ding, W. Zhou, H. Gao, J. Liu, P. Niu, L. Zhao, W. Zhen,
H. Fu, S. Yu, Z. Zhang, G. Xu, C. Li, Z. Lou, M. Xu, C. Qin, G. Wu, G. F. Gao, W. Tan, X.
Yang, Development of an Inactivated Vaccine Candidate, BBIBP-CorV, with Potent
Protection against SARS-CoV-2. *Cell* **182**, 713-721.e9 (2020).

42. Y. Zhang, J. Belayachi, Y. Yang, Q. Fu, L. Rodewald, H. Li, B. Yan, Y. Wang, Y. Shen,
Q. Yang, W. Mu, R. Tang, C. Su, T. Xu, M. Obtel, A. Mhayi, R. Razine, R. Abouqal, Y.
Zhang, X. Yang, Real-world study of the effectiveness of BBIBP-CorV (Sinopharm) COVID-
19 vaccine in the Kingdom of Morocco. *BMC Public Health* **22**, 1584 (2022).

43. B. Sanders, M. Koldijk, H. Schuitemaker, in *Vaccine Analysis: Strategies, Principles, and*
Control, (Springer Berlin Heidelberg, 2015), pp. 45–80.

44. T. Aydillo, A. S. Gonzalez-Reiche, D. Stadlbauer, M. A. Amper, V. D. Nair, C. Mariottini,
S. C. Sealfon, H. van Bakel, P. Palese, F. Krammer, A. García-Sastre, Transcriptome
signatures preceding the induction of anti-stalk antibodies elicited after universal influenza
vaccination. *NPJ Vaccines* **7**, 160 (2022).

45. Y. Chen, C. Hu, Z. Wang, J. Su, S. Wang, B. Li, X. Liu, Z. Yuan, D. Li, H. Wang, B. Zhu,
Y. Shao, Immunity Induced by Inactivated SARS-CoV-2 Vaccine: Breadth, Durability,
Potency, and Specificity in a Healthcare Worker Cohort. *Pathogens* **12**, 1254 (2023).

46. M. J. Patton, C. J. Orihuela, K. S. Harrod, M. A. N. Bhuiyan, P. Dominic, C. G. Kevil, D.
Fort, V. X. Liu, M. Farhat, J. L. Koff, C. V. Lal, A. Gaggar, R. P. Richter, N. Erdmann, M.
Might, A. Gaggar, COVID-19 bacteremic co-infection is a major risk factor for mortality,
ICU admission, and mechanical ventilation. *Crit Care* **27**, 34 (2023).

47. A. Sette, S. Crotty, Immunological memory to <scp>SARS-CoV</scp> -2 infection and
<scp>COVID</scp> -19 vaccines. *Immunol Rev* **310**, 27–46 (2022).

48. B. Pulendran, P. S. Arunachalam, D. T. O'Hagan, Emerging concepts in the science of
vaccine adjuvants. *Nat Rev Drug Discov* **20**, 454–475 (2024).

49. R. Edgar, Gene Expression Omnibus: NCBI gene expression and hybridization array data
repository. *Nucleic Acids Res* **30**, 207–210 (2024).

50. S. Davis, P. S. Meltzer, GEOquery: a bridge between the Gene Expression Omnibus
(GEO) and BioConductor. *Bioinformatics* **23**, 1846–1847 (2007).

51. S. Durinck, Y. Moreau, A. Kasprzyk, S. Davis, B. De Moor, A. Brazma, W. Huber,
BioMart and Bioconductor: a powerful link between biological databases and microarray data
analysis. *Bioinformatics* **21**, 3439–3440 (2024).

52. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

53. S. E. Hunt, W. McLaren, L. Gil, A. Thormann, H. Schuilenburg, D. Sheppard, A. Parton,
I. M. Armean, S. J. Trevanion, P. Flicek, F. Cunningham, Ensembl variation resources.
Database **2018** (2018), doi:10.1093/database/bay119.

54. C. Soneson, M. I. Love, M. D. Robinson, Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences. *F1000Res* **4**, 1521 (2016).

55. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

56. M. Carlson, GO.db: A set of annotation maps describing the entire Gene Ontology.*R*
package version 3.8.2. .

57. T. Wu, E. Hu, S. Xu, M. Chen, P. Guo, Z. Dai, T. Feng, L. Zhou, W. Tang, L. Zhan, X.
Fu, S. Liu, X. Bo, G. Yu, clusterProfiler 4.0: A universal enrichment tool for interpreting
omics data. *The Innovation* **2**, 100141 (2021).

58. Z. Gu, Complex heatmap visualization. *iMeta* **1** (2022), doi:10.1002/imt2.43.

59. Z. Gu, R. Eils, M. Schlesner, Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

60. S. Lê, J. Josse, F. Husson, **FactoMineR** : An *R* Package for Multivariate Analysis. *J Stat*
Softw **25** (2008), doi:10.18637/jss.v025.i01.

61. P. Shannon, A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, D. Ramage, N. Amin, B.
Schwikowski, T. Ideker, Cytoscape: A Software Environment for Integrated Models of
Biomolecular Interaction Networks. *Genome Res* **13**, 2498–2504 (2003).

1079

Vaccination								
Vaccine	Commercial name	Regimen	Timepoints (days)	Doses	Technology	Adjuvant	GSE ID	Participants
BNT162b2 (BNT)	Comirnarty	HO	0, 1, 6	3x BNT	RNA	No	GSE199750	25
		HO	0, 1, 6	2x BNT	RNA	No	GSE199750	46
		HO	0, 6	1x BNT	RNA	No	GSE199750	67
		HE	0, 1, 3, 7	1x ChAd + 1x BNT	VV + RNA	No	GSE201530	16
		HE	1	2x ChAd + 1x BNT	VV + RNA	No	GSE199750	5
		BI	10, 26, 51	2x BNT + 1x I	RNA + I	No	GSE201530	13
		BI	10, 26, 51	1x BNT (10 dpi) + 1x I	RNA + I	No	GSE189039	5
		AI	10	1x BNT + 1x I + 1x BNT	RNA + I	No	GSE189039	2
		AI	0, 10, 26, 51	1x I + RNA + 1x I	I + RNA + I	No	GSE201530	1
AZD1222 (ChAd)	Covishield, Vaxzebria	HO	0, 1, 3, 6, 7	2x ChAd	VV	No	GSE199750, GSE201533	12
		HO	0, 3, 6	1x ChAd	VV	No	GSE199750, GSE201533	53
		HE	0, 1, 3, 7	1x ChAd + 1x BNT	VV + RNA	No	GSE201530	17
		HE	1	2x ChAd + 1x BNT	VV + RNA	No	GSE199750	5
mRNA-1273 (MO)	Spikevax	HE	1	2x BNT + 1x MO	RNA	No	GSE199750	10
BBIBP-CorV (BBIBP)	Covilo	HO	0, 7, 14, 28	3 (3rd dose)	IN	Alum	GSE206023	6
ZF2001	Zifivax	HE	0, 7, 14, 28	1 (3rd dose)	SU	Alum	GSE206023	6

Infection						
GSE ID	Prior infection	Prior vaccination	Timepoints (days)	Variant	Severity	Participants
GSE189039	No	Yes	0, 10, 26, 51	Beta	MI (1), S (3)	4
	No	No	0, 10, 26, 51	Beta	MO (2), S (3)	5
GSE201530	No	No	0, 1, 3, 5	Omicron	MI	1
	Yes	No	0, 1, 3, 5	Omicron	MI	6
	No	Yes	0, 1, 3, 5	Omicron	MI	38

Table 1. Vaccines and infection conditions selected in this study. Legend: HO: Homologous; HE: Heterologous; BI: Before infection; AI: After infection; I: Infection; VV: Viral vector; IN: Inactivated; SU: Subunit. MI: Mild; MO: Moderate; S: Severe; A: Asymptomatic.

A

Data curation

562 PBMC samples
245 participants
4 Bulk-RNAseq datasets
5 SARS-CoV-2 vaccines
2 Studies with infected patients

BNT
ChAd
MO
ZF2001
BBIBP

1x or 2x BNT

1x BNT

Gene sets annotation

GENEONTOLOGY
Unifying Biology

Immune system
biological process

Immune process

Antimicrobial response | Complement system | Innate immune system | Adaptive immune system
Processes | Cells | Processes | Cells

MSigDB
Molecular Signatures
Database

VAX collection

Differential gene expression

DESeq2

p-adj < 0.05
-1 < L2FC > 1
Wald test

DEGs overlap

Fisher's exact test
qvalue < 0.10

GSEA and ML classification

Random forest classifier

Immune cells
and processes

Gene markers for
vaccine type and
infection

Other
vaccines

B

Participants by vaccine

1x BNT	1x ChAd	2x BNT	1x Inf	3x BNT	1x ChAd + 1x BNT	2x BNT + 1x Inf	2x ChAd	2x BNT + 1x MO			
67	53	46	38	25	17	13	12	10	6	6	6

Type
- BBIBP
- BNT
- ChAd
- Infection
- MO
- ZF2001

2x Inf
3x BBIBP
2x BBIBP + 1x ZF2001
1x BNT + 1x Inf (5)
1x Inf (5)
2x ChAd + 1x BNT (5)
1x BNT + 1x Inf (4)
1x BNT + 1x Inf + 1x BNT (3)
1x Inf + 1x BNT + 1x Inf (1)

C

Samples

75, 64, 53, 46, 37, 36, 25, 17, 16, 15, 13, 12, 11, 10, 8, 8, 6, 6, 6, 6, 6, 6, 6, 6, 6, 6, 5, 5, 5, 4, 3, 3, 3, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1

BNT (V1, D0), BNT (V1, D6), ChAd (V1, D0), BNT (V2, D1), ChAd (V1, D7), ChAd (V1, D3), BNT (V3, D1), ChAd-BNT (V2, D3), ChAd-BNT (V2, D0), ChAd (V1, D6), BNT-I (D0), ChAd-BNT (V2, D2), BNT-MO (V3, D7), ChAd (V2, D1), H (D0), BBIBP (V3, D0), BBIBP (V3, D07), BBIBP (V3, D14), BBIBP (V3, D28), ZF2001 (V3, D0), ZF2001 (V3, D07), ZF2001 (V3, D14), ZF2001 (V3, D28), BNT (V1, D7), BNT-I (D1), BNT-I (V2, D3), ChAd-BNT (V2, D3), I-(D10, severe), I-(D10, moderate), I-(D26, D10), BNT (V1, D28, severe), BNT-I (D26, severe), BNT (V1, D4), BNT-MO (V1, D0), BNT-MO (V1, D6), I-(D10, severe), I-(D26, severe), I-(D51, moderate), I-(D51, severe), BNT-I (D5), BNT-I (V1, D11), BNT-I (D10, mild), BNT-I (D26, mild), BNT-I (D51, severe), BNT-I-BNT (D51, mild), BNT-I-BNT (D51, severe), ChAd (V2, D7), I-(D1), I-BNT-I (D5), I-I (D1), I-I (D2), I-I (D3)

A

B

C

Figure 3

Figure 5