Development and Validation of a Diagnostic Prediction Rule for Osteopenia

Thammabhorn Janwittayanuchit[1]

Naritsaret Kaewboonlert[2,*]

Pornthep Tangkanjanavelukul[2]

Kitirat Phattaramarut[2]

Pattama Thongdee[2]

1 Suranaree University of Technology Hospital, Nakhon Ratchasima 30000, Thailand

2 Institute of Medicine, Suranaree University of Technology, Nakhon Ratchasima 30000, Thailand

* Corresponding author: School of Surgery, Institute of Medicine, Suranaree University of Technology, Nakhon Ratchasima 30000, Thailand. Tel: +66 4422 3956, E-mail: naritcvt@gmail.com

Abstract

Objectives

To triage patients with a high likelihood of osteopenia before referring them for a standard bone mass density test for diagnosis.

Introduction

Osteopenia defined by low bone mineral density, is a precursor for osteoporosis and is primarily associated with aging-linked natural bone loss in adulthood. The model and findings can be used to adopt an inclusive screening and swift treatment model that can work in most settings where resources are limited.

Methods

We developed a diagnostic prediction rule based on clinical characteristics. A retrospective cohort of 798 patients who were going to be diagnosed with osteopenia or osteoporosis, within January-September 2022. The multivariable logistic regression to assess potential predictors. The logistic coefficients were transformed as a risk-based scoring system. The internally validation was performed using a bootstrapping procedure.

Results

The model initially included seven predictors: sex, age, height, weight, body mass index, diabetes mellitus, and estimated glomerular filtration rate. However, after using backward elimination for model reduction, only three predictors—sex, age, and weight—were retained in the final model. The discrimination performance was assessed with the area under the receiver operating characteristic curve (AuROC); it was 0.779 (95%CI 0.74-0.82), and the calibration plot showed

good calibration. For internal validation, bootstrap resampling was utilized, yielding an AuROC of

0.768 (95% CI 0.73-0.81), indicating robust performance of the model.

Conclusions

This study developed and internally validated the Osteopenia Simple Scoring System. This clinical

risk score could be one of the important tools for diagnosing osteopenia and allocating resources

in resource-limited settings.

Keywords

Osteopenia, Osteoporosis, Risk prediction, Diagnostic prediction rule, Clinical prediction

Introduction

Osteopenia is a decrease in bone mineral density (BMD) below normal values for their

ages, is the initial stage of bone loss, which may progress to a more severe condition i.e.

osteoporosis. However, that doesn't always lead to osteoporosis depending on many factors. The

primary cause of osteopenia is the natural bone loss that occurs gradually during adulthood.

Secondary causes supposed to accelerate bone loss include lifestyle factors[1] such as smoking,

certain underlying diseases, steroid usage, early menopausal woman, rheumatoid arthritis, and

some medications as well. Osteopenia is often a precursor to osteoporosis, which are now

diagnosed by measuring bone mineral density using dual energy X-ray absorptiometry bone

scans.[2] The osteopenia, as defined by the World Health Organization (WHO) is a t-score

between -1 to -2.5, while values less than -2.5 are diagnostic for osteoporosis.[3, 4] Osteopenia

is not considered a disease while osteoporosis is. In the other hand, osteopenia is considered a

marker for risk of fractures.[5]

The potential predictors of bone mass density in patients with fractures treated in hospitals

were found that factors such as age, sex, smoking, history of adult wrist fractures, spinal

deformities[6, 7], history of adult hip fractures, and osteoarthritis of the spine[8] significantly

differed statistically between groups with normal bone mass density and those with bone mass

density below -1 standard deviation. Low body mass index, low vitamin D level[9] and diabetes[10,

11], chronic kidney disease[12] are also associated with osteoporosis.

There has been increasing attention to the clinical predictive models of diagnostic

screening models for the prediction of fracture risk in patients diagnosed with osteoporosis.[13-

16] Clinical predictive models are commonly used in clinics for the purpose of disease diagnosis,

outcome prediction, and evaluation of the clinical response.[17, 18] We used multivariable logistic

regression to develop predictive models for possible use in the facilitation of early treatment and

screening for osteopenia.

In countries with limited resources, access to a test for bone mass density would be far-

fetched. This research aims to help triage patients at high risk of having osteopenia before they

are referred for a standard BMD test for diagnosis. This work could be extended to programs

aimed to osteopenia or osteoporosis screening in the community.

Methods

Study design and setting

This diagnostic prediction research, utilizing a retrospective cohort design, was conducted

at Suranaree University of Technology Hospital in Nakhon Ratchasima, located in the lower

northeastern region of Thailand. Our university hospital conducts more than 2,000 bone mass

density tests annually.

This study gathered demographic and laboratory data, including age, weight, height, body

mass index, smoking status, early menopause, rheumatoid arthritis, diabetes mellitus, chronic

kidney disease, serum creatinine, and estimated glomerular filtration rate levels from the

electronic medical records of Suranaree University Technology Hospital. The data collection

period spanned from January to September 2022. BMD was used in the analysis. Participants

previously diagnosed with osteopenia or osteoporosis were excluded from the study.

Confirmation of cases

102   All patients included in this study underwent a BMD test on a single machine at the

103   hospital's checkup center. In this study, osteopenia is defined as a T-score of less than -1, and

104   osteoporosis is defined as a T-score of less than -2.5.

Statistical analysis and sample size calculations

Continuous variables were assessed for normality and presented as means and standard

deviations if normally distributed, or medians and interquartile ranges if not. Mean differences of

the variables between the two groups were compared by using an independent t-test, or rank-

sum test based on the distribution of the data. Categorical data were expressed as frequencies

and percentages of the total in each group and compared between groups using either the exact

probability test or chi-square test where applicable. We assessed diagnostic performance and

potential prediction by univariable logistic regression, using crude odds ratios (OR) and their

corresponding area under the receiver operating characteristic curves (AuROC). Statistically

significant two-sided p-values less than 0.05 were considered in applicable cases. All analyses

were done using Stata statistical software version 17. In the development of clinical prediction

rules based on methods described by Riley et al.[19], it was estimated that the minimum sample

size for a multivariable prediction model with a binary outcome was required. This was estimated

from a model c-statistic of 0.8, six candidate predictors, and an assumed prevalence of

osteopenia from a preliminary study, standing at 46%, to get the minimum sample size of 382

cases with 176 events.

Model development and validation

All potential predictor variables necessary for diagnostic prediction of osteopenia in our

routine practice were extracted from the hospital's electronic medical records. These included

age, sex, height, weight, smoking status, serum creatinine levels, diabetes mellitus (with or

without insulin use), hypertension, early menopause, chronic kidney disease, steroid use, and

rheumatoid arthritis.

We identified potential predictors based on prior knowledge; about the biological process,

a review of the literature, and available prediction models. Subsequently, the exploratory analysis

of significant predictors was done using a univariable logistic regression. We assessed the

significance of the predictors through the diagnostic odds ratio and the corresponding p-value.

Additionally, we assessed the area under the receiver operating characteristic curve (AuROC) for

each univariable logistic model. Any predictor variable showing an odds ratio >1.00, significant p-

value of <0.05, and higher AuROC than others was included in the model. Continuous potential

variables were categorized into ordinal following the preceding model and review of literature.

Therefore, in this respect, understanding the nature of the relationship between the dependent

variable and the outcome to determines the cut-off point.

The model to be used for the study is derived from the multivariable logistic regression

with a binary outcome. The factors that not contributing to the outcome were removed using the

backward elimination method. A total of four predictors got pruned from the model: height, body

mass index, diabetes mellitus, and estimated glomerular filtration rate. Diagnostic performance of

the developed model was assessed using the reduced multivariable model by means of

calibration and discrimination. Calibration was assessed by the Hosmer-Lemeshow goodness-of-

fit test, and a plot was applied to show both the model-estimated disease probabilities and

observed disease data. The discriminative ability of the model was graphically tested through a

distributional plot. It was reported with the area under the curve of the receiver operating

characteristic. The internal validation was conducted using the bootstrapping procedure of 1000

replications.

Simplified score derivation

Each predictor in the multivariable model was assigned a specific score based on the

logistic regression coefficients. The coefficient of each predictor was divided by the smallest

coefficient and then rounded up to the nearest whole number. Utilizing a population-analogue

approach, the positive predictive value (PPV) was calculated to demonstrate the predictive

performance. Calibration and discrimination measurements were also conducted using the score-

based multivariable logistic model.

Ethical considerations

This research was conducted based on ethical standards of clinical research. According

to the Helsinki Declaration and began its activity only after it received approval and permission

from the Institutional Review Board of Suranaree University of Technology regarding the review

of the research protocol. Retrospective data were extracted through data record forms. The

patients were treated by the routine hospital staff and were not affected by any research protocols,

the informed consent was waived. The study adhered to the reporting guidelines outlined in the

Transparent Reporting of a multivariable prediction model for Individual Prognosis Or Diagnosis

(TRIPOD) statement. The study protocol received approval from the Institutional Review Board

of Suranaree University of Technology, with approval number COA No.32/2567.

Results

Participants

A total of 798 participants were evaluated for osteopenia at the Suranaree University of

Technology check-up center from January to September 2022. After excluding 242 patients

previously diagnosed with osteopenia or osteoporosis, the remaining 556 patients were divided

into two groups: 230 in the osteopenia group and 326 in the non-osteopenia group. The

prevalence of osteopenia and osteoporosis was 41.4% and 5.4%, respectively. Of these, 198

patients in the osteopenia group and 188 in the non-osteopenia group were female. The mean

age for the osteopenia group was 65.07±10.34 years, compared to 59.03±9.14 years for the non-

osteopenia group. The average weight was 57.40±9.58 kg for osteopenia cases and 57.40±12.71

kg for non-osteopenia cases. Average heights were 155.22±7.50 cm for the osteopenia group

and 160.53±8.00 cm for the non-osteopenia group. The mean BMI was 24.68±14.44 kg/m^2 for

osteopenia cases and 25.64±4.02 kg/m^2 for non-osteopenia cases (Table 1). There were

significant differences in groups in terms of female gender, age, weight, height, BMI, and current

underlying diseases, including diabetes mellitus, as well as in laboratory factors such as estimated

glomerular filtration rate (eGFR) (p < 0.001). There were no significant differences in the serum

creatinine, early menopause, rheumatoid arthritis, and smoking status.

Model development and internal validation

The following continuous variables were converted to ordinal variables: eGFR, body mass

index, weight, age, and height. The cut points were determined based on information gathered

from the literature and prior models. The model was developed using multivariable logistic

regression, which demonstrated the relevant characteristics included in the regression for

predicting osteopenia were female gender, age, height, weight, BMI, diabetes mellitus, and eGFR.

These factors remained significant in the multivariable logistic regression analysis and were

integrated into the full risk prediction model (Table 2).

The reduced model employed a backward elimination strategy. Multivariable logistic

regression indicated that female gender, age, and weight, which were included in the final model,

were found to be statistically significant. The scoring ranged from 1 to a maximum of 5 points,

accumulating to a total of 13 points, summing up to 133. Ages over 70 years were assigned the

highest scores (Table 3).

The calibration plot of the estimated risk of osteopenia compared to the actual risk showed

acceptable calibration, with observed probabilities closely matching the expected probabilities

and exhibiting minimum variation from the ideal (Figure 2). The Hosmer-Lemeshow goodness-of-

fit statistic yielded a non-significant result for the outcome (p = 0.614), suggesting that the

statistical fitness of the model was satisfactory, given that a p-value larger than 0.1 was regarded

indicative of a good fit. Area under the receiver operating characteristic curve (AuROC) was

0.7792 (95% CI 0.74-0.82), showing a good performance in model discrimation (Figure 3).

The process of internal validation was conducted by utilizing a bootstrap resampling

technique with 1000 repeats. Following the adjustment for optimism in discrimination, the

bootstrap analysis produced an area under the receiver operating characteristic curve (AuROC)

of 0.768 (95% CI 0.73-0.81), indicates a good ability and a significant level of agreement between

the estimated and observed probabilities of risk (Supplementary material).

Simple score cut point identification

A very high sensitivity of 94.8% (95% CI 91.1-97.3) was observed at a cut point of ≥4,

while the specificity was 28.5% (95% CI 23.7-33.8). At a score of ≥10, the specificity was high at

91.7% (95% CI 88.2-94.5), but the sensitivity was lower at 44.3% (95% CI 37.8-51.0). The best

cut point was determined by achieving a balance between sensitivity and specificity. For a score

of 6 or above, the sensitivity was 82.2% (95% CI 76.6-86.9), and the specificity was 55.2% (95%

CI 49.6-60.7), with a likelihood ratio of positive result of 1.83 (95% CI 1.60-2.10). The positive

predictive value was 56.4% (95% CI 50.9-61.8), whereas the negative predictive value was 81.4%

(95% CI 75.7-86.3) (Table 4).

Clinical utility

In the context of triaging patients for osteopenia, the newly developed diagnostic

prediction rule was applied. 32.37% (180 patients out of 556) were true negatives, this model can

reduce the unnecessary BMD tests. Among the 211 patients who scored negatively (simple score

below 6), 41 patients (18.55%) were false negatives for osteopenia, and only 2 patients (0.36%

of all patients) were diagnosed with osteoporosis. On the other hand, among those with a positive

score test (simple score above 6), 146 patients had normal BMD and were recommended for

follow-up, while 189 patients were confirmed to have osteopenia and received medical treatment

according to the treatment guidelines.

234 Discussion

Despite the limitations, the current study managed to establish a diagnostic rule for

predicting patients with osteopenia, which could strengthen an early diagnosis and the treatment

of patients in osteopenia. This finding is highly relevant because our hospital faces a significant

burden of bone density testing and a high prevalence of osteopenia. Moreover, the application of

a predictive diagnosis logistic regression model, as conducted in the current study, also supports

the efforts of other studies to implement more predictive analytics in clinical settings[20].

Our results emphasize the importance of a multi-factor approach in novel predictive model

formulation. Specifically, our model uses the same age, sex, and weight as the known risk factors,

established to predict osteopenia and osteoporosis, proven to be significant in other studies.[21]

Similarly, the use of backward elimination for the purpose of precise identification of the factors

allowed reducing the insignificant height and BMI from the model, as these variables have

negligible impact on the model's predictability of outcomes in our patients.[22]

The calibration and discrimination results of our model are satisfactory and being

confirmed. The results of the Hosmer-Lemeshow test and AuROC testify to the model's effective

prediction of true osteopenia cases. This is confirmed by the literature on the appropriateness of

calculating these metrics to check how well a diagnostic tool performs in clinical epidemiology

and diagnostics.[23]

Furthermore, the use of this model demonstrated its significant clinical use. The

prioritization of resources based on a simple score, which identifies high and low-risk patients,

may also lower the risk of untreated osteopenia transitioning to osteoporosis and adhere to WHO,

who suggests that more tests should be conducted on populations at higher risks.[24, 25]

Although the FRAX score is currently used in predicting the 10-year probability of hip

fracture[26, 27], a study by Teeratakulpisarn et al[28]. reports that even though there is

concordance between the 10-year probability of hip fractures for FRAX scores with and without

BMD, this concordance declines in elderly and osteoporotic participants, and in those with FRAX

scores without BMD. Therefore, to achieve higher accuracy, it is advisable to undergo BMD

testing.

In 2001, Koh LK et al.[29] designed a simple tool to categorize postmenopausal Asian

women (OSTA score). They utilized a questionnaire to identify those in the cohort with

osteoporosis, defined as BMD T-scores ≤ -2.5 and use multivariable logistic regression analysis.

The tool had a good performance with an area under the ROC curve of 0.79. Subsequently, it

showed sensitivity of 91%, specificity of 45%, among others.[29] Additionally, upon a validation in

Thai population, OSTA score presented sensitivity of 51.7% and specificity of 77.4% with a false

negative rate of ~20%.[30-33] The OSTA risk classification system showed that high and medium-

risk patients were significantly more likely to sustain injuries in falls and have different femoral

bone fractures patterns compared to low-risk patients. Machine learning models particularly

artificial neural networks offer another opportunity to predict low BMD. Comparison of both ANN

models to logistic regression models to predict low BMD had no significantly different in

performance for either the femoral neck or lumbar spine.[34] Although the OSTA score performs

well within the Thai population and particularly among postmenopausal women, however, it can

be limited use to the general population.

Our model's stability and reliability were internal validation via bootstrapping, process

accounting for the potential optimism that can compromise prediction models developed in narrow

or specific populations. All of these methods make our research more reliable and can be used

to apply to similar settings with limited health care resources.

The study also has its limitations. In particular, the specificity of the model at some cut

points was insufficient for real-life applications, resulting in overdiagnosis and overtreatment. This

compromise between sensitivity and specificity is common in the development of diagnostic

instruments, and it needs to be adjusted depending on the costs and risks of the disease. For

future research, it is possible not only to include new, more prognostic factors but also to progress

statistical instruments, such as machine learning. Further external validation is also required

before adopting this model in other settings.

288   Conclusion

289   The development of a diagnostic prediction rule for osteopenia in a resource-limited

290   context is a major progress in the field of bone health management. This instrument is likely to

291   enhance patient prognosis and maximize the use of available healthcare resources by detecting

292   and offering timely therapeutic treatment to those at risk.

294   Acknowledgement

295   This paper was supported by Suranaree University of Technology.

Funding

None

Conflict of interest

None

Author contribution statement

NK and TJ contributed to all parts of the research.PT, KP, and PT focused on discussing, reviewing, and editing the manuscript.

Figure legends

Figure 1: Study flow.

Figure 2: Calibration plot for model predicted risk for osteopenia versus actual risk.

Figure 3: Discrimination performance of the newly developed model, using clinical characteristics to classify patients with normal and low bone mass density.

Figure 4: Clinical utility.

Tables

315   **Table 1:** Baseline patient characteristics, underlying diseases, and laboratory investigations of

316   the derivation cohort, along with a comparison of osteopenia cases and normal bone mineral

317   density tests (n = 556).

Patient characteristics	Osteopenia cases (n = 230)		Non-osteopenia cases (n = 326)		Univariable OR (95%CI)	p-value	AuROC
	n	(%)	n	(%)			
Female gender	198	(86.09)	188	(57.67)	4.54 (2.94-7.00)	< 0.001	0.64 (0.61-0.68)
Age, years (mean ± SD)	65.07	± 10.34	59.03	± 9.14	1.07 (1.05-1.09)	< 0.001	0.68 (0.64-0.73)
Weight, kg (mean ± SD)	57.40	± 9.58	66.30	± 12.71	0.93 (0.91-0.95)	< 0.001	0.29 (0.25-0.34)
Height, cm (mean ± SD)	155.22	± 7.50	160.53	± 8.03	0.92 (0.89-0.94)	< 0.001	0.32 (0.28-0.37)
BMI, km/m^2 (mean ± SD)	24.68	± 14.44	25.64	± 4.02	0.88 (0.84-0.93)	< 0.001	0.37 (0.33-0.42)
Current smoking	1	(0.43)	5	(1.54)	0.28 (0.03-2.40)	0.245	0.49 (0.49-0.50)
Early menopause	5	(2.17)	10	(3.09)	0.70 (0.24-2.07)	0.516	0.50 (0.48-0.51)
Rheumatoid arthritis	1	(0.43)	2	(0.62)	0.70 (0.06-7.80)	0.774	0.50 (0.49-0.51)
Diabetes mellitus	15	(6.52)	45	(13.89)	0.43 (0.23-0.80)	0.007	0.46 (0.44-0.49)
Creatinine, g/dL (mean ± SD)	0.87	± 0.72	0.87	± 0.37	1.01 (0.72-1.42)	0.967	0.42 (0.37-0.48)
eGFR, ml/min/1.73 m^2 (mean ± SD)	68.59	± 24.13	83.71	± 25.39	0.97 (0.97-0.98)	< 0.001	0.32 (0.28-0.37)
Chronic kidney disease	25	(10.92)	37	(11.46)	0.95 (0.55-1.62)	0.844	0.50 (0.47-0.52)

318   Abbreviations: eGFR, estimated glomerular filtration rate; OR, odds ratio; CI, confidence interval

320   **Table 2:** Full model multivariable logistic regression analysis.

Predictors	Full model		
	mOR	95% CI	P-value
Female	5.33	3.03 - 9.38	0.000
Age, year			
≤ 59	1		
60 – 69	3.59	2.25 - 5.74	< 0.001
≥ 70	6.91	3.62 - 13.20	< 0.001
Height, cm			
≥ 155	1		
< 155	0.97	0.57 - 1.64	0.903
Weight, kg			

≥ 70	1		
60 – 69	1.15	0.61 - 2.17	0.656
50 – 59	1.66	0.72 - 3.82	0.233
≤ 49	1.72	0.54 - 5.54	0.360
Body mass index, kg/m^2			
≥ 25.0	1		
23.0 – 24.9	1.19	0.64 - 2.22	0.577
18.5 – 22.9	1.71	0.79 - 3.69	0.169
< 18.5	3.94	0.87 - 17.90	0.076
Diabetes mellitus	0.35	0.17 - 0.71	0.004
Estimated GFR			
≥ 60	1		
30 – 59	0.84	0.45 - 1.54	0.565
15 – 29	1.89	0.22 - 16.05	0.558
≤ 14	1	Empty	
Model intercept	.062		

321 Abbreviation: mOR, multivariable odds ratio; BMI, body mass index; GFR, glomerular filtration

322 rate

330 **Table 3:** Reduced model with logit coefficients.

Predictors	mOR	95% CI	P-value	β Coefficient	Score
Female	4.09	2.51 - 6.68	0.000	1.41	4
Age, year					
≤ 59	1				
60 – 69	3.29	2.13 - 5.08	0.000	1.19	3.5
≥ 70	5.52	3.21 - 9.50	0.000	1.71	5
Weight, kg					
≥ 70	1				
60 – 69	1.40	0.80 - 2.45	0.240	0.34	1
50 – 59	2.60	1.51 - 4.50	0.001	0.96	3
≤ 49	3.96	2.02 - 7.76	0.000	1.38	4

Model intercept	.065				

331 Abbreviation: mOR, multivariable odds ratio; BMI, body mass index

333 **Table 4:** Selection of score cut-off point with sensitivity, specificity, LHR+, PPV, NPV, and along

334 with 95% confidence interval.

Score cut point	Sensitivity (%)	Specificity (%)	LHR+ (%)	PPV (%)	NPV (%)
≥ 4	94.8 (91.1-97.3)	28.5 (23.7-33.8)	1.33 (1.23-1.43)	48.3 (43.6-53.1)	88.6 (80.9-94.0)
≥ 6	82.2 (76.6-86.9)	55.2 (49.6-60.7)	1.83 (1.60-2.10)	56.4 (50.9-61.8)	81.4 (75.7-86.3)
≥ 10	44.3 (37.8-51.0)	91.7 (88.2-94.5)	5.35 (3.63-7.90)	79.1 (71.0-85.7)	70.0 (65.4-74.3)

335 Abbreviation: LHR+, likelihood ratio for positive test; PPV, positive predictive value; NPV,

336 negative predictive value

References

1.   Rychter AM, Ratajczak AE, Szymczak-Tomczak A, Michalak M, Eder P, Dobrowolska A,
et al. Associations of Lifestyle Factors with Osteopenia and Osteoporosis in Polish Patients with
Inflammatory Bowel Disease. Nutrients. 2021;13(6). Epub 2021/06/03. doi:
10.3390/nu13061863. PubMed PMID: 34070791; PubMed Central PMCID: PMCPMC8227497.
2.   Blake GM, Fogelman I. The role of DXA bone density scans in the diagnosis and
treatment of osteoporosis. Postgrad Med J. 2007;83(982):509-17. Epub 2007/08/07. doi:
10.1136/pgmj.2007.057505. PubMed PMID: 17675543; PubMed Central PMCID:
PMCPMC2600106.
3.   Sözen T, Özışık L, Başaran N. An overview and management of osteoporosis. Eur J
Rheumatol. 2017;4(1):46-56. Epub 2017/03/16. doi: 10.5152/eurjrheum.2016.048. PubMed
PMID: 28293453; PubMed Central PMCID: PMCPMC5335887.

4. Johnell O, Kanis J. Epidemiology of osteoporotic fractures. Osteoporos Int. 2005;16
Suppl 2:S3-7. Epub 2004/09/15. doi: 10.1007/s00198-004-1702-6. PubMed PMID: 15365697.
5. Tran T, Bliuc D, Pham HM, van Geel T, Adachi JD, Berger C, et al. A Risk Assessment
Tool for Predicting Fragility Fractures and Mortality in the Elderly. J Bone Miner Res.
2020;35(10):1923-34. Epub 2020/05/28. doi: 10.1002/jbmr.4100. PubMed PMID: 32460361.
6. Rozental TD, Shah J, Chacko AT, Zurakowski D. Prevalence and predictors of
osteoporosis risk in orthopaedic patients. Clin Orthop Relat Res. 2010;468(7):1765-72. Epub
2009/11/17. doi: 10.1007/s11999-009-1162-6. PubMed PMID: 19911243; PubMed Central
PMCID: PMCPMC2881983.
7. Lane NE. Epidemiology, etiology, and diagnosis of osteoporosis. Am J Obstet Gynecol.
2006;194(2 Suppl):S3-11. Epub 2006/02/02. doi: 10.1016/j.ajog.2005.08.047. PubMed PMID:
16448873.
8. Grams AE, Rehwald R, Bartsch A, Honold S, Freyschlag CF, Knoflach M, et al.
Correlation between degenerative spine disease and bone marrow density: a retrospective
investigation. BMC Med Imaging. 2016;16:17. Epub 2016/02/26. doi: 10.1186/s12880-016-
0123-2. PubMed PMID: 26911278; PubMed Central PMCID: PMCPMC4765052.
9. Tang G, Feng L, Pei Y, Gu Z, Chen T, Feng Z. Low BMI, blood calcium and vitamin D,
kyphosis time, and outdoor activity time are independent risk factors for osteoporosis in
postmenopausal women. Front Endocrinol (Lausanne). 2023;14:1154927. Epub 2023/11/08.
doi: 10.3389/fendo.2023.1154927. PubMed PMID: 37937050; PubMed Central PMCID:
PMCPMC10627178.
10. Tang Y, Zhang L, Ye D, Zhao A, Liu Y, Zhang M. Causal relationship between Type 1
diabetes and osteoporosis and fracture occurrence: a two-sample Mendelian randomization
analysis. Osteoporos Int. 2023;34(6):1111-7. Epub 2023/04/04. doi: 10.1007/s00198-023-
06734-6. PubMed PMID: 37012460.
11. Ali D, Tencerova M, Figeac F, Kassem M, Jafari A. The pathophysiology of osteoporosis
in obesity and type 2 diabetes in aging women and men: The mechanisms and roles of
increased bone marrow adiposity. Front Endocrinol (Lausanne). 2022;13:981487. Epub
2022/10/04. doi: 10.3389/fendo.2022.981487. PubMed PMID: 36187112; PubMed Central
PMCID: PMCPMC9520254.
12. Hampson G, Elder GJ, Cohen-Solal M, Abrahamsen B. A review and perspective on the
assessment, management and prevention of fragility fractures in patients with osteoporosis and
chronic kidney disease. Endocrine. 2021;73(3):509-29. Epub 2021/05/12. doi: 10.1007/s12020-
021-02735-9. PubMed PMID: 33974225; PubMed Central PMCID: PMCPMC8325650.
13. Tan J, Zhang Z, He Y, Xu X, Yang Y, Xu Q, et al. Development and validation of a risk
prediction model for osteoporosis in elderly patients with type 2 diabetes mellitus: a
retrospective and multicenter study. BMC Geriatrics. 2023;23(1):698. doi: 10.1186/s12877-023-
04306-1.
14. Mauck KF, Cuddihy MT, Atkinson EJ, Melton LJ, 3rd. Use of clinical prediction rules in
detecting osteoporosis in a population-based sample of postmenopausal women. Arch Intern
Med. 2005;165(5):530-6. Epub 2005/03/16. doi: 10.1001/archinte.165.5.530. PubMed PMID:
15767529.
15. Leeyaphan J, Rojjananukulpong K, Intarasompun P, Peerakul Y. Development and
Validation of a New Clinical Diagnostic Screening Model for Osteoporosis in Postmenopausal
Women. J Bone Metab. 2023;30(2):179-88. Epub 2023/07/14. doi:
10.11005/jbm.2023.30.2.179. PubMed PMID: 37449350; PubMed Central PMCID:
PMCPMC10346005.
16. Wang J, Kong C, Pan F, Lu S. Construction and Validation of a Nomogram Clinical
Prediction Model for Predicting Osteoporosis in an Asymptomatic Elderly Population in Beijing. J
Clin Med. 2023;12(4). Epub 2023/02/26. doi: 10.3390/jcm12041292. PubMed PMID: 36835828;
PubMed Central PMCID: PMCPMC9967366.

17.   Aibar-Almazán A, Voltes-Martínez A, Castellote-Caballero Y, Afanador-Restrepo DF,
Carcelén-Fraile MDC, López-Ruiz E. Current Status of the Diagnosis and Management of
Osteoporosis. Int J Mol Sci. 2022;23(16). Epub 2022/08/27. doi: 10.3390/ijms23169465.
PubMed PMID: 36012730; PubMed Central PMCID: PMCPMC9408932.
18.   Sun X, Chen Y, Gao Y, Zhang Z, Qin L, Song J, et al. Prediction Models for Osteoporotic
Fractures Risk: A Systematic Review and Critical Appraisal. Aging Dis. 2022;13(4):1215-38.
Epub 2022/07/21. doi: 10.14336/ad.2021.1206. PubMed PMID: 35855348; PubMed Central
PMCID: PMCPMC9286920.
19.   Riley RD, Snell KI, Ensor J, Burke DL, Harrell FE, Jr., Moons KG, et al. Minimum sample
size for developing a multivariable prediction model: PART II - binary and time-to-event
outcomes. Stat Med. 2019;38(7):1276-96. Epub 2018/10/26. doi: 10.1002/sim.7992. PubMed
PMID: 30357870; PubMed Central PMCID: PMCPMC6519266.
20.   Riley RD, Pate A, Dhiman P, Archer L, Martin GP, Collins GS. Clinical prediction models
and the multiverse of madness. BMC Med. 2023;21(1):502. Epub 2023/12/19. doi:
10.1186/s12916-023-03212-y. PubMed PMID: 38110939; PubMed Central PMCID:
PMCPMC10729337.
21.   Pouresmaeili F, Kamalidehghan B, Kamarehei M, Goh YM. A comprehensive overview
on osteoporosis and its risk factors. Ther Clin Risk Manag. 2018;14:2029-49. Epub 2018/11/23.
doi: 10.2147/tcrm.S138000. PubMed PMID: 30464484; PubMed Central PMCID:
PMCPMC6225907.
22.   Chowdhury MZI, Turin TC. Variable selection strategies and its importance in clinical
prediction modelling. Fam Med Community Health. 2020;8(1):e000262. Epub 2020/03/10. doi:
10.1136/fmch-2019-000262. PubMed PMID: 32148735; PubMed Central PMCID:
PMCPMC7032893.
23.   Alba AC, Agoritsas T, Walsh M, Hanna S, Iorio A, Devereaux PJ, et al. Discrimination
and Calibration of Clinical Prediction Models: Users' Guides to the Medical Literature. Jama.
2017;318(14):1377-84. Epub 2017/10/20. doi: 10.1001/jama.2017.12126. PubMed PMID:
29049590.
24.   Kanis JA. Assessment of fracture risk and its application to screening for
postmenopausal osteoporosis: synopsis of a WHO report. WHO Study Group. Osteoporos Int.
1994;4(6):368-81. Epub 1994/11/01. doi: 10.1007/bf01622200. PubMed PMID: 7696835.
25.   Rubin KH, Holmberg T, Rothmann MJ, Høiberg M, Barkmann R, Gram J, et al. The risk-
stratified osteoporosis strategy evaluation study (ROSE): a randomized prospective population-
based study. Design and baseline characteristics. Calcif Tissue Int. 2015;96(2):167-79. Epub
2015/01/13. doi: 10.1007/s00223-014-9950-8. PubMed PMID: 25578146.
26.   Kanis JA, Johnell O, Oden A, Dawson A, De Laet C, Jonsson B. Ten year probabilities
of osteoporotic fractures according to BMD and diagnostic thresholds. Osteoporos Int.
2001;12(12):989-95. Epub 2002/02/16. doi: 10.1007/s001980170006. PubMed PMID:
11846333.
27.   Kanis JA, Johnell O, De Laet C, Jonsson B, Oden A, Ogelsby AK. International
Variations in Hip Fracture Probabilities: Implications for Risk Assessment. Journal of Bone and
Mineral Research. 2002;17(7):1237-44. doi: https://doi.org/10.1359/jbmr.2002.17.7.1237.
28.   Teeratakulpisarn N, Charoensri S, Theerakulpisut D, Pongchaiyakul C. FRAX score with
and without bone mineral density: a comparison and factors affecting the discordance in
osteoporosis treatment in Thais. Arch Osteoporos. 2021;16(1):44. Epub 2021/02/27. doi:
10.1007/s11657-021-00911-y. PubMed PMID: 33635451.
29.   Koh LK, Sedrine WB, Torralba TP, Kung A, Fujiwara S, Chan SP, et al. A simple tool to
identify asian women at increased risk of osteoporosis. Osteoporos Int. 2001;12(8):699-705.
Epub 2001/10/03. doi: 10.1007/s001980170070. PubMed PMID: 11580084.
30.   Chaovisitsaree S, Namwongprom SN, Morakote N, Suntornlimsiri N, Piyamongkol W.
Comparison of osteoporosis self assessment tool for Asian (OSTA) and standard assessment in

Menopause Clinic, Chiang Mai. J Med Assoc Thai. 2007;90(3):420-5. Epub 2007/04/13.
PubMed PMID: 17427514.
31.   Geater S, Leelawattana R, Geater A. Validation of the OSTA index for discriminating
between high and low probability of femoral neck and lumbar spine osteoporosis among Thai
postmenopausal women. J Med Assoc Thai. 2004;87(11):1286-92. Epub 2005/04/14. PubMed
PMID: 15825701.
32.   Chen CC, Rau CS, Wu SC, Kuo PJ, Chen YC, Hsieh HY, et al. Association of
Osteoporosis Self-Assessment Tool for Asians (OSTA) Score with Clinical Presentation and
Expenditure in Hospitalized Trauma Patients with Femoral Fractures. Int J Environ Res Public
Health. 2016;13(10). Epub 2016/10/14. doi: 10.3390/ijerph13100995. PubMed PMID:
27735874; PubMed Central PMCID: PMCPMC5086734.
33.   Panichyawat N, Tanmahasamut P. Comparison of OSTA index and KKOS scoring
system for prediction of osteoporosis in postmenopausal women who attended Siriraj
Menopause Clinic. J Med Assoc Thai. 2012;95(11):1365-71. Epub 2012/12/21. PubMed PMID:
23252200.
34.   Ongphiphadhanakul B, Rajatanavin R, Chailurkit L, Piaseu N, Teerarungsikul K, Sirisriro
R, et al. Prediction of low bone mineral density in postmenopausal women by artificial neural
network model compared to logistic regression model. J Med Assoc Thai. 1997;80(8):508-15.
Epub 1997/08/01. PubMed PMID: 9277083.

Figure 1

Figure 2

Area under ROC curve = 0.7792

Figure 3

Figure 4