Effects of trauma quality improvement programme implementation on mortality: A nonrandomised controlled trial

Date: 02 September, 2025

Cover page

Corresponding author

Johanna Berg, MD Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden Department of Emergency and Internal Medicine, Skåne University Hospital, Malmö, Sweden
ORCID: https://orcid.org/0000-0001-7553-7337
Email: johanna.berg@ki.se
Telephone: +46730659092 Address: Department of Global Public Health, Karolinska Institutet, SE-171 77, Stockholm, Sweden

Author list

Johanna Berg, MD Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden Department of Emergency and Internal Medicine, Skåne University Hospital, Malmö, Sweden. ORCID: https://orcid.org/0000-0001-7553-7337

Siddarth David, PhD Doctors For You, India Department of Global Public Health, Karolinska Institutet, Sweden. ORCID: https://orcid.org/0000-0002-4848-2358

Girish D. Bakhshi, MS, MRCSEd.Uk, FACS, Grant Government Medical College & Sir J.J. Group of Hospitals, Mumbai-400008, Maharashtra, India. ORCID: https://orcid.org/0000-0001-9542-4428

Debojit Basak, SSKM Hospital/IPGME&R

Shamita Chatterjee, MS, FMAS, FAIS, Department of Surgery, IPGME&R-SSKM Hospital, Kolkata, India. ORCID https://orcid.org/0000-0002-9460-108X

Kapil Dev Soni, MD, Department of Critical and Intensive Care, JPN Apex Trauma Centre, All India Institute of Medical Sciences, New Delhi, India. ORCID: https://orcid.org/0000-0003-1214-4119

Ulf Ekelund, MD, PhD, Emergency Medicine, Department of Clinical Sciences Lund, Lund University, Lund, Sweden. Department of Emergency Medicine, Skåne University Hospital, Lund, Sweden. ORCID: https://orcid.org/0000-0002-0951-1582

Li Felländer-Tsai, MD, PhD Division of Orthopaedics and Biotechnology, Department of Clinical Science, Intervention and Technology, Karolinska Institutet, Stockholm, Sweden. Reconstructive Orthopaedics, Karolinska University Hospital, Stockholm, Sweden. ORCID: https://orcid.org/0000-0003-0693-6080

Manjul Joshipura, MD Academy of Traumatology (India), Ahmedabad, India. ORCID: https://orcid.org/0000-0001-5891-8234

Tamal Khan, All India Institute of Medical Sciences, Patna- 801507, India

Monty Khajanchi, DNB, Department of General Surgery, Seth G. S Medical College & K.E. M Hospital, Mumbai, India ORCID: https://orcid.org/0000-0002-0898-6391

Mohan L N, MBBS, MS, FRCS, Vydehi Institute of Medical Sciences and Research Centre, Bangalore. India ORCID: https://orcid.org/0000-0002-3503-832X

Anurag Mishra, MBBS, MS, DNB, MNAMS, FACS, FAIS Department of Surgery, Maulana Azad Medical College, New Delhi, India

Max Petzold, PhD ORCID: 0000-0003-4908-2169 School of Public Health and Community Medicine, Institute of Medicine, University of Gothenburg, Gothenburg, Sweden

Sendhil Rajan, MBBS MS MCh FEBS FRCS Department of Surgery, Norfolk & Norwich University Hospital, Colney Ln, Colney, Norwich NR4 7UY, United Kingdom. ORCID: https://orcid.org/0000-0003-3161-5553

Nobhojit Roy, MS, PhD, MPH The George Institute for Global Health, India Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden

Rajdeep Singh, MS, DNB, FACS, FRCS Department of Surgery, Maulana Azad Medical College, Delhi, India. ORCID: https://orcid.org/0000-0001-6593-2624

Martin Gerdin Wärnberg, MD, PhD Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden Function Perioperative Medicine and Intensive Care, Karolinska University Hospital, Solna, Sweden. ORCID: https://orcid.org/0000-0001-6069-4794

Abstract

Objectives: To determine whether implementing a trauma quality improvement programme using audit filters improves outcomes for adult trauma patients.

Design: Prospective controlled interrupted time series study using a generalised additive model applied to monthly aggregated data, with a secondary analysis using difference-in-differences.

Setting: Multicentre study at four tertiary care hospitals in urban India from 2017 to 2022.

Participants: A total of 10,143 adult trauma patients admitted to hospital with a history of trauma, defined by external causes listed in block V01–Y36, chapter 20 of the ICD-10. Median age was 35 years (IQR 26 to 50), and 83% were men.

Interventions: Two hospitals implemented a trauma quality improvement programme using audit filters after a one-year observation period (intervention arm), while two hospitals continued usual care without intervention (control arm).

Main outcome measures: Primary outcome was all-cause in-hospital mortality. 30-day mortality, originally designated as the primary outcome, was analysed as a secondary outcome due to lower-than-expected prospective inclusion rates, which limited statistical power.

Results: In the intervention arm, in-hospital mortality decreased by an estimated -11.2% (95% CI –16.0% to –5.5%, p<0.001), with no significant change in the control arm -0.5% (95% CI –4.0% to 5.4%, p=0.835). No significant seasonal effects or autocorrelation were observed. Secondary difference-in-differences analysis showed significant reductions in in-hospital mortality –12% (95% CI –16 to –9%; p<0.001) and 30-day mortality –15% (95% CI –19 to –11%; p<0.001) in the intervention arm compared to controls. However, external factors, including the opening of a dedicated trauma centre and the COVID-19 pandemic, may have influenced these results.

Conclusions: Implementing trauma quality improvement programmes using audit filters may reduce mortality in adult trauma patients. Further research is needed to confirm these findings, clarify underlying mechanisms, and ensure sustainability over time.

Trial registration: Trauma Audit Filter Trial, ClinicalTrials.gov ID NCT03235388, https://clinicaltrials.gov/study/NCT03235388

Summary boxes

What is already known on this topic

• Trauma is a leading cause of mortality and morbidity worldwide, with 90% of the burden in low- and middle-income countries. Here, it causes the largest number of quality-related deaths, with more than half of trauma fatalities attributed to poor quality of care.

• Trauma quality improvement programmes have been recommended to address inequalities in care, but despite their widespread use in high-income countries there is no high-level evidence that these programmes improve patient outcomes.

What this study adds

• We present the first prospective quasi-experimental trial investigating if implementing trauma quality improvement programmes improves mortality.

• Our results suggest that these programmes can improve mortality-rates in adult trauma patients, particularly in settings with high numbers of preventable deaths.

• However, the mechanisms behind their effectiveness remain poorly understood.

Introduction

Quality improvement initiatives are frequently proposed to enhance patient safety and improve care.[1] However, evidence for their effectiveness in improving patient outcomes remains limited with several studies report conflicting results, highlighting the need for systematic, evidence-based approaches and rigorous evaluation of these initiatives.[1–3] Additionally, many interventions lack a clear theoretical foundation to explain their mechanisms of action.[1] Despite these uncertainties, improving quality of care remains essential, as poor quality has been identified as a greater cause of preventable deaths worldwide than lack of access to services.[4]

Trauma, defined as injury due to external factors and the bodys associated response, results in more than four million deaths annually and is the leading cause of death among individuals aged 10-29 years.[5] Over two million lives could be saved each year if mortality rates in low- and middle-income countries (LMICs) were the same as those in high-income countries (HICs). Trauma accounts for the largest number of quality related deaths in LMICs, with more than half of trauma deaths estimated to be due to poor quality among those who reach care. Strengthening the quality of trauma services is therefore essential to reduce preventable trauma deaths and global inequities in survival.[4]

The development of trauma systems integrating prevention, pre- and intrahospital care, and rehabilitation has been essential for the improvement in care in high-income countries over the last decades.[6–9] As one part of trauma systems, trauma quality improvement programmes have evolved. These programmes encompass an ongoing

process that includes identifying deviations from care standards, analysing these findings, evaluating factors for improvement, and implementing corrective action plans to improve patient care.[10–12] Identifying cases with deviations from care traditionally uses audit filters, in some areas referred to as process quality indicators. They are predefined statements that represent ideal care standards, for example, "Patients with a Glasgow Coma Scale (GCS) score of less than 8 should receive a definitive airway".[11,13] These filters are used to flag individual patient cases so that the care can be reviewed, potential opportunities for improvement can be identified, and corrective strategies can be implemented. This process is resource-intensive because it requires continuous data collection and audit filter tracking.[11,14]

The American College of Surgeons released the first trauma audit filters as part of the guidelines on trauma care in 1987.[15] Since then, the use of audit filters and trauma quality improvement programmes has been widely applied in HICs, and in 2009 they were recommended by the WHO as one way to mitigate the inequalities of trauma care globally—despite there being no high-level evidence for their effectiveness in improving patient outcomes.[14] Our aim was therefore to assess whether implementing a trauma quality improvement programme using audit filters improved patient outcomes.

Methods

Study design and setting

We conducted a controlled interrupted time series study across four hospitals in urban India (ClinicalTrials.gov identifier NCT03235388[16]). All tertiary care hospitals have approximately 1500 beds and in-house clinical specialities to care for trauma patients. None of the hospitals used any structured trauma quality improvement process prior to this study. The study had three phases: an observation phase lasting 14 months in all four hospitals to establish baseline outcomes, an implementation phase lasting six months, during which two hospitals were randomized to implement a trauma quality improvement programme with audit filters, and an intervention phase lasting 41 months. The intervention phase was extended by 18 months due to the COVID-19 pandemic. We performed an interim analysis 15 months after the initiation of the intervention to asses the quality of the data and identify unexpected changes in outcomes. We followed the TREND checklist for reporting of non-randomised clinical trials.[17]

Intervention and control

For the intervention centres, a multidisciplinary review group was established at each centre, including representatives from all departments involved in trauma care: emergency/casualty officers, general surgery, anaesthesia, burns and plastics, orthopaedics, radiology, and neurosurgery. The local principal investigator at each centre recruited senior healthcare staff from each department. Within this group, an anonymous online Delphi process was conducted to identify which audit filters would be most useful to implement at each hospital [18]. Further details on the Delphi process, including participant composition and procedures, are available in the cited publication. A list of implemented audit filters is included as supplementary material.

A project officer prospectively included patients and identified specific audit filter violations through direct observation and data collection. Structured data were collected on violations, process outcomes, treatments, interventions, and patient outcomes during the hospital stay. These cases were compiled into automated structured reports every 4–6 weeks, which served as the basis for discussions at the multidisciplinary board review meetings. During each meeting, cases were reviewed and improvement actions identified. Progress on these actions was followed up in subsequent meetings. During the implementation phase, members of the core research team with experience in facilitating review discussions attended each meeting to guide and support the discussions, after this, only the local review board attended.

In the control centres, no intervention was implemented. They continued treating trauma patients using standard care, without a quality improvement process. The centre-specific data collected for this study were available to both intervention and control centres but were not used in any structured way in the control centres to inform or direct care. Weekly calls were held between the core research team and project officers at each hospital to support data collection and address operational issues. In intervention centres, multidisciplinary review meetings were conducted independently by the local

teams, with no involvement from the research team beyond facilitating the underlying reports based on collected data.

Participants

Inclusion criteria and enrollment

We included adults aged 18 years and older admitted for in-hospital care with a history of trauma, defined by International Classification of Diseases, tenth revision (ICD-10) chapter 20, codes V01-Y36 for external causes of morbidity and mortality, as the reason for admission. Project officers were trained to record vital signs with standard equipment for patients across rotating shifts (day, evening, and night). Each shift comprised six hours for enrolling new patients in the emergency department and two hours for follow-up of previously included patients.

Retrospective inclusion

Due to lower than expected prospective inclusion rates, particularly during the COVID-19 pandemic, we retrospectively included patients to achieve the required sample size for the primary statistical analysis. Using hospital records, we randomly selected trauma patients admitted each month. As informed consent could not be obtained, only basic, reliably recorded data were collected: date of admission, age, sex, mechanism of injury, and in-hospital mortality. Data were extracted from ward books and paper charts, as no electronic health record was available.

Data collection and quality control

Initial data collection was performed using paper forms, and the data were then periodically entered into a digital data collection tool and uploaded to secure servers. Patient identifiers were not uploaded, and identification was only possible through the original paper records, which were stored at each hospital in compliance with local regulations. Basic data validation was performed at the time of entry, and double entry of all variables was performed after all the data had been collected to minimize the risk of transfer errors. Quality control visits were conducted each quarter with project officers from different hospitals evaluating the data collection processes based on predefined criteria.

Outcomes and covariates

Our primary outcome was all-cause in-hospital mortality. Information on 30-day and 90-day mortality was collected from medical records if the patient died during the hospital stay, otherwise, it was obtained through telephone follow-up. 30-day mortality, originally designated as the primary outcome, was analysed as a secondary outcome due to lower-than-expected prospective inclusion rates. We also collected data on additional secondary outcomes, including length of stay and intensive care unit (ICU) admissions. Data on several covariates, such as age, sex, vital signs, injury severity, and diagnostic procedures, were collected to describe and compare cohorts and, if needed, to adjust for case-mix and differences between hospitals.

Study execution

One principal investigator and one co-principal investigator were recruited for each participating hospital. These were all experienced trauma clinicians with extensive local knowledge and research interest.

Observation phase (Month 1-14)

The participating hospitals started data collection at the same time to establish baseline outcomes. Weekly meetings were held with all project officers and the core research team to identify and address any issues related to data collection.

Implementation phase (Month 15-20)

The participating hospitals were paired so that the two hospitals with the highest and lowest volumes formed one pair, and the two remaining hospitals formed the second pair. One pair of hospitals was then randomly selected to implement the trauma quality improvement programme, becoming intervention hospitals, while the other two were control hospitals. The process of data collection remained the same at all sites.

At the intervention hospitals, a two-day session on the background and rationale of trauma audit filters was held by representatives from the core research team. This session was attended by a multidisciplinary team of surgeons/physicians, anaesthetists, administrators and nurses involved in trauma care. The local review board and audit filters selected as described previously. During the implementation phase, participants from the core research team attended audit filter review meetings to facilitate the discussion and formulate corrective strategies accordingly.

Intervention phase (Months 21-42)

The intervention hospitals continued the data collection, both for base data and for audit filter deviations. The trauma audit review board continued to analyse deviations, implementing corrective strategies. The meetings were held without participation from the core research team. Three months after the intervention phase started, the largest intervention hospital opened a dedicated trauma centre. The control hospitals continued with base data collection.

Statistical analysis

As we collected both prospective and retrospective data, with the prospective cohort being a random subset from the combined prospective and retrospective data, some pre-defined analysis is only possible to do for the prospective cohort.

Primary analysis

Our primary analysis was a segmented interrupted time series model using a generalized additive model (GAM) to estimate the effect of the intervention on in-hospital mortality. To assess robustness, we conducted two secondary analyses: a

difference-in-differences analysis using individual-level data, and an unadjusted pre-post analysis.

For our primary analysis, we applied a segmented Generalized Additive Model (GAM) to estimate the effect of the intervention on mortality. [19] Data were aggregated by study month across intervention and control arms, with each observation representing the average outcome for all patients enrolled during that month. The intervention was modelled as a step change, represented by a binary indicator. Time was included as a smooth term to flexibly capture potential non-linear trends. Calendar month was entered as a cyclic cubic spline with 12 knots to adjust for seasonality. The model used a beta regression distribution with a logit link, suitable for modelling bounded continuous outcomes between 0 and 1. Estimation was performed using restricted maximum likelihood (REML). Autocorrelation was assessed using the Box-Ljung test (lag = 12, degrees of freedom = 0), applied separately to response residuals from the intervention and control arms.

GAM model specification:

$$\text{logit}(\mathbb{E}[Y]) = \beta_0 + \beta_1 \cdot \text{post\_intervention} + f_1(\text{study\_month}) + f_2(\text{month})$$

We developed a counterfactual model to compare predicted outcomes under scenarios without the intervention.

Sample size considerations

The sample size requirements for interrupted time series analyses depend on several factors, including model complexity, data variance and the spread of observations. For our primary analysis, we chose to adhere to published guidelines stating that at least twelve observations are needed during the observation phase and twelve observations are needed during the intervention phase and that each observation should be an aggregate of at least 100 patients. [20] Our use of a large GAM introduced extra complexity, and we estimated that 100 patients/arm/month and an extended intervention phase were needed to detect the potential impact of the intervention. This sample size allowed us to detect a reduction in mortality from 20% to 15% (power 0.8, alpha 0.05) using a pre-post design.

Secondary and subgroup analysis

For the secondary analysis, we conducted a difference-in-differences (DiD), using linear regression on individual-level data for prospectively included patients. We adjusted for confounders selected based on clinical relevance and observed baseline differences between arms. To assess the parallel trends assumption, we plotted pre-intervention trends in in-hospital and 30-day mortality stratified by study arm, and visually inspected the trajectories. In addition, we tested for interaction between time (study month) and group (intervention vs control) using pre-intervention prospective data. All analyses were conducted in R [21], using the `mgcv` package for GAM modelling.

We conducted two pre-specified subgroup analyses for major trauma and potentially salvageable trauma patients. We defined a major trauma patient as any patient who

was in the hospital for more than three days with an Injury Severity Score (ISS) >15 or who died within three days of arrival, and a potentially salvageable trauma patient as any patient with 15<ISS<24. Due to limited statistical power to conduct stratified interrupted time series analyses, these subgroup analyses were conducted using the secondary difference-in-differences approach. To account for multiple comparisons, we applied the Holm correction method [22].

Sensitivity analyses

We conducted two pre-specified sensitivity analyses: an unadjusted pre-post analysis using a two-sample Z-test for proportions, and an interrupted time series analysis excluding the implementation phase. Three post hoc analyses were also conducted: excluding the period after a trauma centre opened at one intervention hospital, excluding the COVID-19 phase, and performing unadjusted and adjusted interrupted time series analyses in the prospectively included cohort to evaluate whether observed effects on mortality were confounded by changes in case mix over time.

Missing data and data processing

We added an analysis of missing data to the analysis plan to assess whether complete case analysis would be appropriate. We examined associations between missingness and key variables using the R package finalfit [23], which applies chi-squared tests for categorical variables and univariate logistic regression for continuous variables, with missingness as the outcome. If no significant associations were found, indicating data were Missing Completely at Random (MCAR), we planned to proceed with complete case analysis. If significant associations were found, suggesting data were Missing at Random (MAR), we planned to use multiple imputation by chained equations (MICE), provided that the proportion of missing data was >5% and that the available data supported stable estimation of the imputation model. If data were deemed likely to be Missing Not at Random (MNAR), complete case analysis would be applied with caution, and sensitivity analyses would be considered. The Abbreviated Injury Scale (AIS) and Injury Severity Score (ISS) were calculated from ICD-10 codes using the R package icdpicr [24].

External events during the study period

During the study period, two external events occurred that may have influenced patient outcomes. First, three months after the intervention phase began, the largest intervention hospital opened a dedicated trauma centre. Although all surgical and radiology services for trauma were previously available, they were distributed across separate buildings and wards, with no dedicated units for trauma patients. With the opening of the trauma centre, these services were consolidated into a single 180-bed facility with in-house CT, operating theatres, and both ICU and general wards dedicated to trauma care. Nurses also received trauma-specific training prior to the opening.

Second, the COVID-19 pandemic vastly disrupted hospital operations at all sites. Prospective patient inclusion was halted between April and December 2020, and review meetings were paused until mid-2021. The second intervention hospital was converted

into a COVID-19 facility for much of 2020 and 2021, effectively eliminating trauma admissions during this period. In addition to straining hospital capacity, the pandemic reduced population mobility and led to an overall decline in trauma incidence.

Patient and public involvement

We have not had any patient or public involvement during the planning or execution of this study.

Deviations from study protocol

The intended primary outcome was 30-day mortality, but this became unfeasible due to low prospective inclusion. We therefore defined in-hospital mortality as the primary outcome, for which the study was adequately powered. 30-day mortality was analyzed as a secondary outcome. We also added two non-prespecified analyses: an analysis of missing data and a difference-in-differences analysis. These were introduced because we lacked sufficient power for a time-series analysis of 30-day mortality and observed an unexpected baseline difference in mortality between study arms. The difference-in-differences approach allowed formal comparison between arms with adjustment for confounders. Not all protocol outcomes are reported here; notably, quality of life results will be published separately.

Transparency declaration

The guarantor affirms that this manuscript is an honest, accurate, and transparent account of the study being reported. No important aspects of the study have been omitted. Discrepancies from the study protocol have been reported and explained.

Ethical considerations

The need for informed consent to participate in the intervention was not deemed applicable, and we were granted waivers of informed consent for recording vital signs, demographic parameters and in-hospital outcomes. The project officers obtained written consent for telephone follow-up. Ethical approval was granted by the Swedish Ethical Review Authority (approved 2017-06-07 2017/930-31/2), as well as by all local ethical review boards at each participating hospital (Maulana Azad Medical College (MAMC) - approved 2017-07-19 F.1/IEC/MAMC/(57/02/2017/No 113. SSKM/IPGME&R, Kolkata – approved 2017-08-21, IPGME&R/IEC/2017/396. JJ Hospital, Mumbai – approved 2017-08-22, No. IEC/Pharm/CT/111/A/2017. St. Johns, Bangalore – approved 2017-08-24, 160/2017).

Data sharing

The code for this publication and an anonymised dataset from the clinical trial are available on Zenodo and GitHub under a Creative Commons Attribution 4.0 International License (CC BY 4.0)[25]. The dataset has been de-identified through categorisation and removal of indirect identifiers to minimise the risk of re-identification.

Results

We included 10143 patients between October 2017 and October 2022. Of these patients, 4126 were prospectively included (Figure 1). We included an average of 166 (SD 61) patients per month, 45 out of 61 months did not reach the target number of inclusions, due to low inclusion rates and lower number of patients during COVID-19.

Patient characteristics and missing data counts are presented in Table 1 (all patients) and Table 2 (prospectively included patients). For the primary analysis, missing data were minimal for in-hospital mortality (0.6%) and absent for study month. No significant associations were found between missingness and examined variables, except for one control arm month, which was deemed unlikely to affect results. In the secondary analysis using the adjusted DiD model, missingness was associated with GCS, ISS, and shock for 30-day mortality. The proportion of missing data was low for GCS (1.6%) and shock (2.2%), but higher for ISS (12.5%). Given the exploratory nature of the secondary analysis and the challenge of correctly imputing ISS, all statistical models used complete case analysis.

In the intervention arm, we prospectively screened and collected structured data on 1454 additional patients for audit filter violations to inform discussions in the multidisciplinary review meetings. A list of audit filters with the most violations is available in the supplementary material. Descriptively, in the intervention arm, intubations rose from 2.6% to 24%, ultrasound increased from 2.4% to 9% and computed tomography (CT) scanning from 31% to 42% indicating changes in clinical practice aligned with key audit filters following implementation of the trauma quality improvement programme (Table 1).

Time series analysis for in-hospital mortality

Primary analysis

In the interrupted time series analysis, including 10,086 patients, the intervention phase was associated with a significant reduction in the odds of in-hospital mortality in the intervention arm *(OR: 0.56, 95% CI 0.4 to 0.77, p<0.001)*, corresponding to an estimated absolute difference of -11.2% *(95% CI -16% to -5.5%, p<0.001)*. No significant change was observed in the control arm *(OR: 0.94, 95% CI 0.5 to 1.75, p=0.83)*, absolute difference -0.5% *(95% CI -4% to 5.4%, p=0.835)*(Figure 2).

The study time variable was not statistically significant *(p=0.14 for intervention; p=0.14 for control)*, suggesting that mortality remained otherwise stable over time, with no indication of other time-dependent trends influencing outcomes. We found no significant seasonal effects *(p=0.26 for intervention; p=0.48 for control)* and no evidence of autocorrelation in either study group *(p = 0.51 for intervention; p = 0.10 for control)*.

Secondary and subgroup analysis

Formally comparing the control and intervention arms in the difference-in-differences analysis, including 3317 prospectively included patients, we found that the intervention

was associated with a -12% difference in the in-hospital mortality rate *(-0.12, 95% CI -0.16 to -0.09, p<0.001)* and a -15% difference in the 30-day mortality rate *(-0.15, 95% CI -0.19 to -0.11, p<0.001)* in the intervention arm compared to the control arm after adjusting for sex, age, GCS score, Injury Severity Score (ISS) and shock. For major trauma and potentially salvageable trauma patients, the reduction in mortality was more pronounced. No significant mortality change was observed in the control arm for either group (Table 3).

Sensitivity analysis

In the unadjusted pre-post analysis of all patients, the intervention was associated with an 8.0% absolute reduction in in-hospital mortality *(95% CI -11% to -5.3%, p<0.001)* and in the prospectively included cohort a 8.5% reduction in in-hospital mortality *(95% CI -12% to -4.8%, p<0.001)* and a 14% reduction in 30-day mortality *(95% CI -18% to -9.5%, p<0.001)*. No statistically significant change was observed in the control arm. Excluding the implementation phase and the COVID-19 period showed results consistent with the main analysis, with a statistically significant reduction in in-hospital mortality in the intervention arm. Excluding the trauma centre period resulted in a non-significant reduction; however, this analysis was limited to nine months of post-intervention data, including the six-month implementation phase. The adjusted analysis in the prospectively included cohort showed a stati stically significant reduction in both in-hospital and 30-day mortality after adjusting for age, sex, ISS, GCS, and shock. Full results are available in the supplementary material.

Discussion

Our results indicate that implementing a trauma quality improvement programme using audit filters was associated with a substantial, significant reduction in-hospital mortality of -11.2% *(95% CI -16.0% to –5.5%, p<0.001). We also observed increased use of ultrasound, intubations and CT scans, reflecting changes in care processes. However, the largest hospital in the intervention arm opened a trauma centre equipped with in-house CT scanner, multiple operating rooms, improved resuscitation capabilities and an increased number of ICU beds nine months after the beginning of the implementation phase. Although our sensitivity analysis revealed a statistical non-significant decrease in mortality before the trauma centre opened, it remains challenging to separate the impacts of the trauma centre and the quality improvement programme or determine whether their effects were synergistic.

Health systems are increasingly being recognised as complex adaptive systems, composed of interdependent agents and processes whose interactions give rise to emergent and often unpredictable outcomes. [26,27] In such systems, effects are non-linear, and identifying direct causal chains between intervention and outcome is inherently difficult. We found that mortality declined after the programme was introduced, but the precise mechanisms responsible for this effect are challenging to disentangle. Although we did not formulate an explicit programme theory in advance, we believe that enabling health care staff to follow the trajectory of individual patients

and their outcomes, and to review the processes of care, helped them to identify opportunities for improvement in their own context.

Several studies have reported improved mortality, particularly in severely injured patients, after trauma quality improvement implementation.[28–32] However, definitions of trauma quality improvement vary, with heterogeneous interventions reported. Two studies conducted over 20 years ago in Germany and Thailand used trauma audit filters with a similar approach as our study, and reported reduced mortality and improved care processes.[33,34] All were cohort studies using pre-post design. Development of audit filters for trauma has been done in more recent years, one study from Ghana and one from Cameroon.[35,36]. These filters differ substantially from those initially developed by ASCOT and later suggested in the WHO guidelines. In our previous study, the usefulness of the filters developed in Ghana and Cameroon was deemed high in India, compared filters developed in HICs.[18]

In HIC trauma quality improvement programmes and audit filters have been criticized for being inefficient in detecting opportunities for improvement and being costly.[37] In obstetric care, audit processes have shown to improve outcomes while also emphasizing the importance of adaptation[38], highlighting that standards of care need to be developed considering local priorities and knowledge to ensure that improvement efforts are directed to areas in greatest need. This is particularity important in settings with less developed trauma systems.

The maturity level of a trauma system is linked to reduced mortality and preventable deaths.[8,39] Therefore, the effectiveness of these programmes in reducing mortality likely varies with system maturity. In a similar urban Indian setting, over 50% of trauma deaths were estimated to be preventable.[40] Our results can likely be generalized to settings with similarly mature trauma systems, though local adaptations and contextual factors greatly influence their success. The core process of data collection and multidisciplinary case review can be utilized to empower providers to identify areas for improvement, learning, and education to develop the care provided. This can be conducted for trauma systems at all levels, even though data quality is a known barrier to this in LMICs.[31,41,42]

It should also be recognised that while this intervention, which is a scaffold for data collection and peer review, is introduced into a complex system. Regognizing health care as a complex adaptive system is important as effects are non-linear, What we can show ere is that that when this scafflod was introduced, mortality did appear to decline. Exactly by which mechanisms or the causla relationship between the intervention and the outcome is per definition in a complex system, challenging to identify. We did not have an explicit pre-porgram theory, we do belive that the ability for health care staff to understand the patient outcomes, follow the process of care for individual patients, helped them understand how to imporve care in the context where they work.

Our study has several limitations. First, the introduction of a new trauma centre at the largest intervention hospital may have influenced outcomes directly or acted synergistically with the trauma quality improvement programme. This hospital also accounted for a larger proportion of patients in the intervention arm, thus contributing

more to the aggregate outcome estimates than the smaller intervention hospital. Second, there was an unexpected difference in baseline mortality between the study arms. However, in the primary analysis, each arm was modeled independently, and the control arm was included to account for background trends over time and external events, rather than for direct comparison. In the secondary analysis, we used a difference-in-differences model, which assumes parallel pre-intervention trends rather than equivalent baseline levels. This allowed us to compare outcome changes over time between arms despite baseline differences in mortality. Third, 45 of 61 months fell below the target inclusion for the primary analysis; however, all months contributed data, and the number of time points before and after the intervention exceeded recommended thresholds for interrupted time series analysis. While smaller monthly samples increased uncertainty and variability in monthly estimates, the observed mortality reduction remained robust across analyses.

Fourth, the COVID-19 pandemic significantly impacted the participating centres. The smaller intervention centre was converted to a COVID-19 hospital, reducing trauma patient intake. Quarantine restrictions also decreased the overall number of trauma patients. Despite extending the study by 18 months and adjusting for the pandemic's effects in the sensitivity analysis, its long-term impacts were likely beyond our ability to fully account for. The pandemic led to the suspension of review meetings, necessitating a restart post-pandemic. Last, both intervention hospitals receive a high number of transferred patients at varying intervals after the time of injury, potentially introducing bias, as patients may have died before arriving at the study hospital.

The intervention was designed to provide a structure containing training in trauma quality improvement processes, data collection and multidisciplinary review boards. The ongoing review process and implementation of changes were led entirely by the local teams, without involvement from the core research team. Finally, although we tracked the number of cases reviewed and meetings conducted, and held weekly calls with all project officers to support operations, ensure data collection, and facilitate the generation of reports for review meetings, we did not apply a formal framework to evaluate implementation fidelity. This limits our insight into how consistently the intervention was sustained and engaged with over time.

Despite these limitations, this is the first quasi-experimental study attempting to assess the impact of quality improvement programmes using audit filters on patient outcomes. We conducted a broad and comprehensive statistical analysis that supports our main findings, which are also in alignment with previous research. We believe that there is knowledge on how to select and adapt audit filters and implement a review process, but our understanding of how this translates into changes in care and outcomes is lacking. We also need better understanding of the implementation and sustainability of these programmes in complex health care systems, including how to best tailor them to local needs and detect potential negative effects, especially for other patient populations also competing for care resources.

In conclusion, our results suggest that system-level changes and data-driven quality improvement programmes using audit filters may reduce mortality in trauma patients.

However, the challenge to understand what makes these programmes effective, useful and sustainable in terms of improving outcomes remains.

Acknowledgement

We would like to thank Johan von Schreeb, professor at the department of Global Public Health at the Karolinska Institutet and Charles Mock, Professor Emeritus of Surgery and Epidemiology, University of Washington for their valuable input during this project. A warm thank you to Manjula Ranagatti and all project officers that have dedicated their time and expertise to this project and ensuring the quality of data collection.

JB and MGW had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Author contributions

MGW conceived the study and has been the PI. MGW, KS, MK, NR, SD, LFT, and MP contributed to the design of the study. SD was the project manager during study execution. MGW, KS, MK, NR, SD, JB, MJ, SR, MLN, RS, AM, SC, GB, DB, TK contributed to the execution of the study. JB prepared the data and conducted the statistical analysis. MGW and MP reviewed the results of the statistical analysis. JB drafted the first version of the article. All authors contributed to interpretation of data and critical revisions of the work for important intellectual content. MGW is the guarantor.

Funding

This work was supported by the Swedish Research Council (2016-02041).

Competing interests

None declared.

Tables and figures

Figure 1: Flow chart

Figure 2: Interrupted time-series analysis

Table 1: All included patients, n = 10143

Characteristic	Control arm		Intervention arm	
	Observation phase N = 832[1]	Intervention phase N = 3,565[1]	Observation phase N = 1,436[1]	Intervention phase N = 4,310[1]
Age	32 (25, 46)	35 (25, 48)	36 (25, 50)	36 (26, 51)
Missing	1	19	0	6
Sex male	667 (80)	2,939 (83)	1,148 (80)	3,612 (84)
Missing	2	10	1	10
Transfers	466 (71)	1,196 (74)	1,250 (90)	2,448 (80)
Missing	175	1,957	42	1,255
Type of injury				
Assault	70 (8.4)	299 (8.7)	100 (7.0)	215 (5.0)
Burns	27 (3.2)	149 (4.4)	51 (3.6)	88 (2.1)
Fall	197 (24)	753 (22)	339 (24)	840 (20)
Intentional self-harm	6 (0.7)	35 (1.0)	17 (1.2)	12 (0.3)
Other accident, not violence	109 (13)	306 (8.9)	79 (5.5)	111 (2.6)
Transport accident	405 (49)	1,815 (53)	840 (59)	3,019 (70)
Undetermined intent	17 (2.0)	65 (1.9)	7 (0.5)	6 (0.1)
Missing	1	143	3	19
Died in hospital	74 (8.9)	409 (12)	455 (32)	1,016 (24)
Missing	4	41	2	10

[1]Median (Q1, Q3); n (%)

Table 2: All prospectively included patients, n = 4126

Characteristic	Control arm		Intervention arm	
	Observation phase N = 583[1]	Intervention phase N = 888[1]	Observation phase N = 938[1]	Intervention phase N = 1,717[1]
Age	31 (24, 45)	34 (25, 46)	40 (25, 55)	37 (27, 52)
Missing	1	2		
Sex male	470 (81)	720 (81)	729 (78)	1,438 (84)
Glasgow coma scale	15 (15, 15)	15 (15, 15)	15 (8, 15)	12 (9, 15)
Missing	17	38	5	6
Shock	5 (0.9)	18 (2.1)	54 (5.9)	25 (1.5)
Missing	22	34	20	14
Injury Severity Score (ISS)	6 (8)	7 (8)	12 (8)	13 (7)
Missing	50	134	164	166
Major trauma	103 (19)	155 (21)	486 (65)	1,164 (75)
Missing	53	142	186	174
CT done				
Before arrival	11 (2.4)	32 (4.6)	506 (54)	833 (51)
No	219 (48)	302 (43)	137 (15)	101 (6.2)
Yes	227 (50)	363 (52)	293 (31)	688 (42)
Missing	126	191	2	95
Ultrasound done				
Before arrival	0 (0)	1 (0.2)	2 (0.2)	31 (1.9)
No	258 (58)	341 (52)	908 (97)	1,455 (89)
Yes	187 (42)	312 (48)	22 (2.4)	147 (9.0)
Missing	138	234	6	84
Intubated				
Before arrival	8 (2.0)	6 (1.1)	2 (0.2)	2 (0.1)
No	350 (89)	496 (90)	907 (97)	1,203 (76)
Yes	34 (8.7)	50 (9.1)	24 (2.6)	377 (24)
Missing	191	336	5	135
Admitted to ICU	62 (16)	109 (20)	13 (1.4)	77 (4.9)
Missing	197	340	12	148
Dead at 24 hours after arrival	14 (2.4)	17 (1.9)	99 (11)	109 (6.3)
Missing	4	7	1	0
Died in hospital	42 (7.3)	79 (9.1)	300 (32)	405 (24)
Missing	4	18	1	0
Dead at 30 days after arrival	46 (8.5)	85 (11)	328 (39)	435 (26)
Missing	43	94	102	20

Characteristic	Control arm		Intervention arm	
	Observation phase N = 583[1]	Intervention phase N = 888[1]	Observation phase N = 938[1]	Intervention phase N = 1,717[1]

[1]Median (Q1, Q3); n (%); Mean (SD), Shock is defined as systolic blood pressure < 95 mmHg with heart rate > 100 beats per minute at arrival. Major trauma defined as ISS > 15 and admitted for more than three days, or, died within three days

Table 3: Difference-in-differences analysis n=3317

Characteristic	All patients(n=3317)			Major trauma patients(n=1714)				Potentially salvageable trauma patients(n=1505)			
	Beta	95% CI	p-value	Beta	95% CI	p-value	Adj. p-value	Beta	95% CI	p-value	Adj. p-value
Outcome: In-Hospital Mortality											
Unadjusted Intervention Effect	-0.129	-0.186 to -0.072	<0.001	-0.360	-0.505 to -0.215	<0.001	<0.001	-0.312	-0.481 to -0.144	<0.001	<0.001
Intervention Effect	-0.124	-0.163 to -0.085	<0.001	-0.239	-0.330 to -0.147	<0.001	<0.001	-0.249	-0.357 to -0.140	<0.001	<0.001
Sex	-0.003	-0.026 to 0.020	0.8	-0.019	-0.056 to 0.017	0.3		0.002	-0.035 to 0.039	>0.9	
Age	0.001	0.001 to 0.002	<0.001	0.002	0.001 to 0.002	<0.001		0.002	0.001 to 0.002	<0.001	
Injury Severity Score (ISS)	0.001	0.000 to 0.002	0.2	-0.007	-0.010 to -0.004	<0.001		0.007	-0.008 to 0.022	0.4	
Glasgow coma scale	-0.083	-0.086 to -0.080	<0.001	-0.092	-0.096 to -0.088	<0.001		-0.090	-0.094 to -0.086	<0.001	
Shock	0.072	0.012 to 0.132	0.018	0.020	-0.055 to 0.096	0.6		0.084	-0.002 to 0.169	0.055	
Outcome: 30 Day Mortality											
Unadjusted Intervention Effect	-0.155	-0.213 to -0.096	<0.001	-0.362	-0.508 to -0.215	<0.001	<0.001	-0.324	-0.496 to -0.151	<0.001	<0.001
Intervention Effect	-0.149	-0.189 to -0.110	<0.001	-0.239	-0.332 to -0.146	<0.001	<0.001	-0.259	-0.368 to -0.149	<0.001	<0.001
Sex	0.000	-0.023 to 0.024	>0.9	-0.013	-0.050 to 0.025	0.5		0.002	-0.036 to 0.039	>0.9	
Age	0.001	0.001 to 0.002	<0.001	0.002	0.001 to 0.002	<0.001		0.002	0.001 to 0.002	<0.001	
Injury Severity Score (ISS)	0.001	0.000 to 0.002	0.13	-0.006	-0.009 to -0.003	<0.001		0.005	-0.010 to 0.021	0.5	
Glasgow coma scale	-0.085	-0.088 to -0.082	<0.001	-0.093	-0.097 to -0.089	<0.001		-0.093	-0.097 to -0.089	<0.001	

Characteristic	All patients (n=3317)			Major trauma patients (n=1714)				Potentially salvageable trauma patients (n=1505)			
	Beta	95% CI	p-value	Beta	95% CI	p-value	Adj. p-value	Beta	95% CI	p-value	Adj. p-value
Shock	0.075	0.014 to 0.136	0.017	0.017	-0.060 to 0.093	0.7		0.082	-0.004 to 0.168	0.063	

Abbreviation: CI = Confidence Interval, Shock is defined as systolic blood pressure < 95 mmHg with heart rate > 100 beats per minute at arrival. Major trauma defined as ISS > 15 and admitted for more than three days, or, died within three days

References

Dixon-Woods M, Martin GP. Does quality improvement improve quality? *Future hospital journal*. 2016;3:191–4. doi: 10.7861/futurehosp.3-3-191

Marshall M, Pronovost P, Dixon-Woods M. Promotion of improvement as a science. *The Lancet*. 2013;381:419–21. doi: 10.1016/S0140-6736(12)61850-9

Hill JE, Stephani A-M, Sapple P, *et al.* The effectiveness of continuous quality improvement for developing professional practice and improving health care outcomes: A systematic review. *Implementation science: IS*. 2020;15:23. doi: 10.1186/s13012-020-0975-2

National Academies of Sciences, Engineering, and Medicine, Health and Medicine Division, Board on Health Care Services, *et al. Crossing the global quality chasm: Improving health care worldwide*. Washington, D.C., DC: National Academies Press 2019.

Global Burden of Disease Collaborative Network. Global burden of disease study 2019 (GBD 2019) results. 2020.

SHACKFORD SR, HOLLINGWORTH-FRIDLUND P, EASTMAN AB. The Effect of Regionalization upon the Quality of Trauma Care as Assessed by Concurrent Audit before and after Institution of a Trauma System: A Preliminary Report. *J Trauma*;26:812. doi: 10.1097/00005373-198609000-00006

Davenport RA, Tai N, West A, *et al.* A major trauma centre is a specialty hospital not a hospital of specialties. *BRITISH JOURNAL OF SURGERY*. 2010;97:109–17. doi: 10.1002/bjs.6806

Alharbi RJ, Shrestha S, Lewis V, *et al.* The effectiveness of trauma care systems at different stages of development in reducing mortality: A systematic review and meta-analysis. *World journal of emergency surgery: WJES*. 2021;16:38. doi: 10.1186/s13017-021-00381-0

Celso B, Tepas J, Langland-Orban B, *et al.* A systematic review and meta-analysis comparing outcome of severely injured patients treated in trauma centers

following the establishment of trauma systems. *The Journal of trauma*. 2006;60:371–8; discussion 378. doi: 10.1097/01.ta.0000197916.99629.eb

Ronald Maier, Michael Rhodes. Trauma Performance Improvement. *Injury control: A guide to research and program evaluation*. Cambridge: Cambridge University Press :236–49.

World Health Organization. *Guidelines for trauma quality improvement programmes*. 2009.

Juillard CJ, Mock C, Goosen J, *et al.* Establishing the Evidence Base for Trauma Quality Improvement: A Collaborative WHO-IATSIC Review. *World journal of surgery*. 2009;33:1075–86. doi: 10.1007/s00268-009-9959-8

Willis CD, Gabbe BJ, Cameron PA. Measuring quality in trauma care. *Injury*. 2007;38:527–37. doi: 10.1016/j.injury.2006.06.018

Evans C, Howes D, Pickett W, *et al.* Audit filters for improving processes of care and clinical outcomes in trauma systems. *Cochrane Database of Systematic Reviews*. 2009;84:782. doi: 10.1002/14651858.cd007590.pub2

American College of Surgeons Committee on Trauma. Hospital and prehospital resources for optimal care of the injured patient. Chicago: American College of Surgeons 1987.

The Trauma Audit Filters Trial (TAFT).

Des Jarlais DC, Lyles C, Crepaz N, *et al.* Improving the reporting quality of nonrandomized evaluations of behavioral and public health interventions: The TREND statement. *American journal of public health*. 2004;94:361–6. doi: 10.2105/ajph.94.3.361

Berg J, Alvesson HM, Roy N, *et al.* Perceived usefulness of trauma audit filters in urban india: A mixed-methods multicentre delphi study comparing filters from the WHO and low and middle-income countries. *BMJ open*. 2022;12:e059948. doi: 10.1136/bmjopen-2021-059948

Shadish WR, Zuur AF, Sullivan KJ. Using generalized additive (mixed) models to analyze single case designs. *Journal of School Psychology*. 2014;52:149–78. doi: 10.1016/j.jsp.2013.11.004

Wagner AK, Soumerai SB, Zhang F, *et al.* Segmented regression analysis of interrupted time series studies in medication use research. *Journal of Clinical Pharmacy and Therapeutics*. 2002;299–309.

R Core Team. R: A language and environment for statistical computing. 2017.

Holm S. A Simple Sequentially Rejective Multiple Test Procedure. *Scandinavian Journal of Statistics*. 1979;6:65–70. doi: 10.2307/4615733

Harrison E, Drake T, Pius R. *Finalfit: Quickly create elegant regression results tables and plots when modelling*. 2023.

Black A, Clark D. *Icdpicr: 'ICD' programs for injury categorization in r*. 2022.

Berg J. Noacs-io/tqip-effect-on-mortality. 2025.

Jeffrey Braithwaite, Kate Churruca, Louise A Ellis, Janet Long, Robyn Clay-Williams, Nikki Damen, Jessica Herkes, Chiara Pomare, Kristiana Ludlow. Complexity science in healthcare – aspirations, approaches, applications and accomplishments: A white paper. 2017.

Institute of Medicine (US) Committee on Quality of Health Care in America. *Crossing the quality chasm: A new health system for the 21st century*. Washington (DC): National Academies Press (US) 2001.

Hashmi ZG, Haider AH, Zafar SN, *et al.* Hospital-based trauma quality improvement initiatives: first step toward improving trauma outcomes in the developing world. *The journal of trauma and acute care surgery*. 2013;75:60–8; discussion 68. doi: 10.1097/TA.0b013e31829880a0

Hemmila MR, Cain-Nielsen AH, Jakubus JL, *et al.* Association of hospital participation in a regional trauma quality improvement collaborative with patient outcomes. *JAMA surgery*. 2018;153:747–56. doi: 10.1001/jamasurg.2018.0985

Juillard CJ, Mock C, Goosen J, *et al.* Establishing the evidence base for trauma quality improvement: A collaborative WHO-IATSIC review. *World Journal of Surgery*. 2009;33:1075–86. doi: 10.1007/s00268-009-9959-8

Reynolds TA, Stewart B, Drewett I, *et al.* The Impact of Trauma Care Systems in Low- and Middle-Income Countries. *Annual review of public health*. 2017;38:507–32. doi: 10.1146/annurev-publhealth-032315-021412

Dinh MM, Bein KJ, Gabbe BJ, *et al.* A trauma quality improvement programme associated with improved patient outcomes: 21 years of experience at an Australian Major Trauma Centre. *Injury*. 2014;45:830–4. doi: 10.1016/j.injury.2013.11.005

Ruchholtz S, Waydhas C, Lewan U, *et al.* A multidisciplinary quality management system for the early treatment of severely injured patients: Implementation and results in two trauma centers. *Intensive Care Med*. 2002;28:1395–404.

Chadbunchachai W, Sriwiwat S, Kulleab S, *et al.* The comparative study for quality of trauma treatment before and after the revision of trauma audit filter, Khon Kaen hospital 1998. *Journal of the Medical Association of Thailand = Chotmaihet thangphaet*. 2001;84:782–90.

Stewart BT, Gyedu A, Quansah R, *et al.* District-level hospital trauma care audit filters: Delphi technique for defining context-appropriate indicators for quality improvement initiative evaluation in developing countries. *Injury*. 2016;47:211–9. doi: 10.1016/j.injury.2015.09.007

Wu N, Carvalho M, Nwanna-Nzewunwa O, *et al.* Developing Trauma Audit Filters for Regional Referral Hospitals in Cameroon: Mixed-Methods Approach. *The Journal of surgical research.* 2018;232:578–86. doi: 10.1016/j.jss.2018.07.048

Willis CD, Gabbe BJ, Cameron PA. Measuring quality in trauma care. *Injury.* 2007;38:527–37. doi: 10.1016/j.injury.2006.06.018

Graham W, Wagaarachchi P, Penney G, *et al.* Criteria for clinical audit of the quality of hospital-based obstetric care in developing countries. *Bull World Health Organ.* 2000;78:614–20.

Teixeira PGR, Inaba K, Salim A, *et al.* Preventable morbidity at a mature trauma center. *Archives of surgery.* 2009;144:536-41; discussion 541-2. doi: 10.1001/archsurg.2009.82

Roy N, Veetil DK, Khajanchi MU, *et al.* Learning from 2523 trauma deaths in India- opportunities to prevent in-hospital deaths. *BMC Health Services Research.* 2017;17:142. doi: 10.1186/s12913-017-2085-7

McIver R, Erdogan M, Parker R, *et al.* Effect of trauma quality improvement initiatives on outcomes and costs at community hospitals: A scoping review. *Injury.* 2024;111492. doi: 10.1016/j.injury.2024.111492

Kapanadze G, Berg J, Sun Y, *et al.* Facilitators and barriers impacting in-hospital trauma quality improvement program (TQIP) implementation across country income levels: A scoping review. *BMJ open.* 2023;13:e068219. doi: 10.1136/bmjopen-2022-068219

Controlled interrupted time-series

Observed and predicted in-hospital mortality over time