Mobility and non-household environments: understanding dengue transmission patterns in urban contexts

Víctor Hugo Peña-García[1, *], Bryson A. Ndenga[2], Francis M. Mutuku[3], Donal Bisanzio[4], A. Desiree LaBeaud[5], Erin A. Mordecai[1]

[1] Department of Biology, Stanford University, Stanford, CA, USA

[2] Kenya Medical Research Institute, Kisumu, Kenya

[3] Department of Environmental and Health Sciences, Technical University of Mombasa, Mombasa, Kenya

[4] RTI International, Washington, DC, USA

[5] School of Medicine, Stanford University, Stanford, CA, USA

[*] Corresponding author: Víctor Hugo Peña-García

email: victorhugopega@gmail.com

Abstract

Households (HH) have been traditionally described as the main environments where people are at risk of dengue and other arbovirus infections. Mounting entomological evidence suggests a larger role for environments other than HH. Recently, an agent-based model (ABM) estimated that over half of infections occur in non-household (NH) environments such as workplaces, markets, and recreational sites. Despite the inferred importance of NH sites, we do not yet know how their urban spatial configurations, and human and vector mobility between them, affects their role in dengue transmission. To address this gap, we expanded an ABM calibrated with field data from Kenya to examine movement of people and vectors under different spatial configurations of buildings. We assessed the number of people traveling between HH and NH and the distances traveled, in three urban configurations: NH distributed randomly (scattered), concentrated in a single center, or clustered in multiple centers. Across simulations, the number of people moving was the most influential variable, with higher movement between HH and NH increasing case numbers. The number of cases was also higher when NH were scattered compared to centered or clustered. Intriguingly, the distance people traveled from HH to NH had little effect on dengue burden but influenced the spatial clustering of infections. These findings underscore the role of NH as major spreaders of infections between HH and NH environments, and the importance of human movement in driving dengue dynamics.

Author summary

Recent evidence describes a major role of non-household (NH) environments in dengue

transmission. This new knowledge implies that spatial distribution of these locations and

the movement of humans among them has implications in dynamics of transmission.

However, these haven't been evaluated neither together nor considering a differential

role in transmission of urban environments. We modified a previously informed agent-

based model to include spatial variables by assessing three different urban

conformations, i.e. when NH are randomly distributed (scattered), centered, or grouped

in differed clusters (clustered). On these, we also evaluated the movement of people

from households (HH) to NH (both distance and number of individuals). Our findings,

which includes a higher burden of dengue when NH are scattered and at higher levels

of human movement, are aligned with the idea that infections are mainly happening in

NH mediated by human mobilization. Infected people reach the households where local

subpopulation of vector spread the virus to remaining inhabitants. This work highlights

the role of NH and human mobility in dengue transmission.

Introduction

Dengue is a vector-borne disease prevalent and on the rise in most of the tropical and

subtropical regions around the globe, with introduced and locally acquired cases being

reported in non-endemic areas like USA and Europe [1, 2]. The main vector, *Aedes*

aegypti, is highly anthropophilic, found very close to human environments, and

impacting the public health of urban environments [3, 4].

Historically, the main strategy to control the disease has been to reduce vector-human

contact by reducing the size of mosquito populations. Conventional wisdom is that for

effective control, these activities should be focused on households (HH) as the main

environment where transmission is happening [5, 6]. However, some studies have

suggested that locations other than households might have an important role because

of a significant presence of mosquitoes [7-13] and infected vectors [14, 15] and the fact

that this vector bites during the day [16-19], when people may be outside the home. In a

recent study, we used an agent-based model (ABM) to quantify the number of infections

in different types of urban spaces based on fieldwork mosquito collections and

seroincidence data and estimated that over half of infections are happening in non-

household (NH) environments, where the main high-risk spaces are workplaces and

markets/shops [20].

These results have implications for dengue epidemiology since the high flux of people

through NH suggests that these spaces can contribute to the spread of infections. In

this way, the total number of infections can be affected by the distribution in space of

NH and the movement of people between HH and NH. However, a key knowledge gap

is how these intra-urban dynamics vary with spatial configuration and mobility levels

[21-24], though Massaro and colleagues used mobile phone data to get estimates for

movement between workplaces [25].

Building on a previous result showing the importance of NH for dengue transmission,

we use an ABM to address this knowledge gap. In particular, what role does spatial

configuration of NH spaces play, along with the extent to which people and mosquitoes

move between spaces, in determining dengue dynamics? To address this, we modified

a previously published ABM [20] to make it spatially explicit by assigning coordinates

that mimic different urban conformations and evaluated different scenarios of movement

of people and vectors. We then assessed how these variables affect the burden of

dengue and the spatial patterns to understand urban-level transmission dynamics.

89   Methods

90   Model overview

To achieve the aims of this study, we modified the ABM previously used to describe the

importance of HH and NH in transmission [20]. The model was developed to quantify

the relative contribution of five different types of NH (workplaces, markets or shops,

recreational, religious, and schools) and HH to dengue burden. The model development

and calibration were based on data from two Kenyan cities: Kisumu in the west and

Ukunda on the coast [20]. Here, we focus on parameters calibrated to Kisumu, although

additional results including dynamics from Ukunda are found in supplementary material.

The model represents the movement of people between HH and two different types of

NH locations. The first are daily-commuting locations where individuals attend daily and

meet with the same individuals, including schools (average number of students per

school set to be 360 [26]) and workplaces (average number of workers per workplace is

19 [27]). The assignment of each individual as student, worker, or neither depends on

their age and the age-specific proportion of individuals in such role according to

governmental information [26-28]. The second type of NH locations are randomly

assigned locations for which both the number and identity of people who visit them is

randomly defined every day, including markets, shops, recreational, and religious

spaces. Movement between HHs (categorized as "visit") is also included in the model

with a daily probability for incoming visits of 0.1, according to information extracted from

a human movement survey conducted in Kenya [20].

Epidemiologically relevant locations where individuals move are those with presence of

mosquitoes. Thus, based on vector surveys conducted over two years of fieldwork

previously published [29], NH and HH environments were assigned to have mosquito

presence or absence based on observed prevalence of mosquitoes. Population

dynamics of vectors were modeled at the building level, whereby sub-population

dynamics are driven by site-specific features—particularly the availability and capacity

of water containers as well as numbers of individuals present—as informed by field

surveillance data. These dynamics as well as infection dynamics of vectors are also

determined by temperature by using functions previously described and widely used

elsewhere [30-32].

Initial baseline prevalence of dengue was set to 0.08%, estimated from previous studies

reporting age-structured seroprevalence [33] with an incidence rate per year estimated

as $IR = -\frac{1}{age}\ln(1 - prevalence)$. Transmission events happen in those locations where

infected vectors bite susceptible humans or vice versa. Mosquitoes bite depending on

both temperature-dependent biting rate and the probability of having a successful

vector-human encounter, which depends on the amount of time that humans spend in

the location. This information was obtained from a human movement survey where the

time and frequency that individuals spend in different types of locations were recorded

[20]. Infection status of mosquitoes can be either susceptible, exposed, or infected while

humans can be either susceptible, exposed, infected, or recovered and (temporarily)

immune. The time that mosquitoes spend as exposed depends on temperature

(extrinsic incubation period) and is determined in the model by equations reported

previously by Mordecai and colleagues [30]. Once infected, mosquitoes remain in this

stage until death, which is evaluated daily following a temperature-dependent death rate

(probabilistic implementation within the model is described in supplementary material).

Humans remain susceptible until they are bitten by infected mosquitoes and moved to a

latent stage where they remain for five days. Then, the individual is moved to the

infectious stage, which lasts seven days, before moving to the recovered stage. Since

the model does not explicitly represent dynamics of different serotypes, waning immune

protection was based on Sabin's classic studies describing the loss of complete

heterotypic protection after roughly three months [34, 35] to set a return to susceptible

after 100 days in recovered status. The number of infections in each location—number

of infectious bites from mosquito to human leading to successful infection within a given

physical location— is recorded daily. Statistics about the total number of infections and

locations are reported weekly. The model simulates transmission dynamics happening

for 731 days (comprising temperature conditions between January 1st of 2020 until

December 31st of 2021) and results are shown as a distribution of the number of

infections over 200 simulations. The model and modifications described in this work

were coded in Julia language (v1.10.0) and simulations were run on Sherlock

computing cluster (Stanford Research Computing Center). A more complete and

detailed description of the model can be found at [20].

Spatial variables

The original model was not spatially explicit and hence the movement of individuals was

assumed to be totally random, with equal probability of visitation for all NH locations

within a given NH type and by any given individual. As such, the resulting infections

arise from complete mixing of individuals among structures, which is not realistic and

does not capture local, intra-urban spatial phenomena. To include mobility-associated

variables and describe such local phenomena, we made the model spatially explicit,

where the transmission dynamics previously described are still happening but in a non-

random, distance-dependent manner. To achieve this, we constructed an artificial

coordinate system and tested different urban configurations using the same spatial

framework. In this way, we can assure that differences are due to the building

designation as HH or NH and not by spatial disparities coming from different sets of

spatial coordinates. The model considers a synthetic population of 20,160 people, so

the total municipality areas were rescaled to fit the total number of structures of the

virtual populations while considering similar densities. Spatial coordinates were

randomly generated to create a synthetic settlement and assigned to each structure of

the population. With the aim of capturing different intra-urban conformations, coordinate

assignments were done based on either "scattered" (randomly distributed), "centered"

(majority of NH concentrated in the center of the city in a single cluster), or "clustered"

(majority of NH concentrated in three clusters) configurations (Fig 1).

Fig 1: Spatial distribution of NH for three different urban conformations to be

tested: scattered (NH randomly distributed in space), centered (majority of NH are

clustered in the center), and clustered (majority of NH are grouped in three clusters).

NH are shown with colors while HH are displayed as light gray points. Religious

(orange) and schools (purple) are represented in the first row; markets/shops (red) and

recreational spaces (blue) in middle row; and the most abundant NH, workplaces

(green), are represented in the lower row.

Movement of people

We included two movement-related variables: the distance from each household (HH) to the nearest non-household (NH) locations and the number of people moving. To control movement distances, we applied three treatments: (1) limiting attendance to the nearest NH locations (assigned as distance zero), (2) allowing attendance to NH locations at least 500 meters away, and (3) allowing attendance to NH locations at least 1000 meters away from each HH. These treatments were applied across the three different urban configurations. The results are entirely based on computational simulations; no experiments involving humans were conducted.

In the clustered or centered configurations, assigning the closest NH locations to HHs would primarily select NH sites at the periphery of each cluster, potentially leading to biased representations. To address this and make it equivalent to scattered configuration, we generated a list of NH locations sorted by proximity for each HH and allocated the closest NHs based on the number of inhabitants in the HH. For example, if a HH had four inhabitants, the four closest NHs were assigned to it. This approach ensured a more representative distribution across all HHs.

The number of people visiting NH was simulated at three levels. First, we included the same levels previously described in the model, categorized as 100% mobility [20]. This treatment includes all students and workers attending their respective school and workplace, and random-attendance locations (religious, markets/shops, and recreational). Unfortunately, no data were available on the number of individuals visiting these types of locations. We anticipated significant variability due to the influence of

several unmodeled factors. To account for this uncertainty, we relied on discussions

with local residents to estimate a range of possible visitor numbers. We then applied a

uniform distribution, with a minimum of 10 and a maximum of 70 individuals, to cover a

broad spectrum of potential scenarios. The number of people visiting a given location is

drawn from this uniform distribution on the interval [10,70] daily. We also evaluated a

medium-mobility scenario by decreasing the number of people moving to school and

workplaces to 50% and remaining NH locations to a uniform distribution with

parameters minimum = 5 and maximum = 35. Finally, we include low-mobility scenario

of 20% for school and workplaces and a uniform distribution with parameters minimum

= 2 and maximum = 14 for random-attendance locations.

215 Movement of mosquitoes

In line with the inclusion of spatial variables and human movement, we included the

movement of vectors. There were no data available to characterize the rate of

movement of mosquitoes into new areas. As a result, we parameterized such

movement by considering two variables—availability of both breeding and blood-feeding

resources—to estimate a baseline migration probability for the number of mosquitoes

moving from given locations.

The model includes a local density-dependent function that allows for the mosquito

population to grow in a location-specific way, representing dynamics previously

described for a fragmented environment [36]. The function, as previous described [20],

depends on temperature (peaking at 29°C) and the density of immatures according to

water availability, as follows:

$$f(D) = \frac{1}{1 + e^{bD-a}} d(T) \#(1)$$

227   Where

$$d(T) = -0.166 + 0.08T - 0.0014T^2 \#(2)$$

In this equation a and b are calibrated parameters, T is environmental temperature in

degrees Celsius, D is the larval density expressed as the ratio of the number of larvae

to liters of water available for breeding in the structure, and $d(T)$ is the term describing

the temperature-dependence of population growth.

By using this function, the growth of the mosquito sub-population (in a given building)

depends on the amount of water resources available in the structure. In this way, when

the mosquito population has grown so the water resources are depleted (reaching the

carrying capacity), the mortality of immatures is higher and the probability for a given

mosquito to migrate out of the building increases (details provided in supplementary

material).

Additionally, we considered for migration the possibility that human blood availability is

not sufficient to support the mosquito subpopulation in a given building. For this purpose,

the number of females that fed in each day was estimated by considering the number of

females biting (N_b) and the probability for those to have a successful feeding encounter

with a human the same day ($P(bit)$). The latter is determined considering the number of

individuals, the time they spend at the building, and the time a visitor spends in each

location (details are provided in supplementary material). The final number of fed

females (N_F) is estimated by

$$N_F \sim Bin\left(N_b, P(bit)\right) \#(3)$$

247 Thus, when the difference between the number of mosquitoes trying to bite and the

248 actual number of mosquitoes that are fed is large, the probability of migration increases.

249 Once a mosquito migrates, a new location is assigned by considering a dispersal kernel

250 [37]. Following previous work [38], we used a lognormal function with the form

$$P_D = \frac{1}{(2\pi)^{\frac{3}{2}} b d^2} e^{-\frac{\log(d/a)^2}{2b^2}} \#(4)$$

251 Where d is distance and both a and b are parameters to be estimated. We fitted a

252 function by assuming a mean dispersal distance of 105.69 meters, as estimated for

253 *Aedes aegypti* by [39].

Spatial autocorrelation

By taking advantage of spatial features introduced in the model, we performed a spatial

autocorrelation analysis to evaluate the level of clustering of dengue cases recorded in

each simulation as a function of urban configuration and human and mosquito mobility.

To do this, the location of each mosquito-human infectious encounter irrespective of the

type—HH or NH—is recorded so a Global Moran's I index [40] could be estimated at

the end of the simulated period. Global Moran's I ranges from -1 to +1, where -1 means

totally dispersed location of cases while 1 represents a spatial distribution that is totally

clustered (total separation between locations with dengue cases and those without

cases). In this sense, the null hypothesis of this analysis is that dengue cases are

randomly distributed in municipalities, represented by Moran's I value of 0 [41]. The

analysis was done for every simulation by using 1,000 permutations for inference in

each of them. Analyses were done using the package SpatialDependence.jl

implemented in Julia language (v 1.10.0). Code is available in the GitHub repository

(https://github.com/vhpenagarcia/ABM_dengue) and have been archived within the

Zenodo repository (https://doi.org/10.5281/zenodo.14036270).

Results

Burden of dengue is strongly affected by number of people visiting NH

We quantified the total number of cases after two years of transmission. When we

simulated the epidemic under different human movement regimes, it was evident that

the number of cases nonlinearly decreased as the number of people moving from HH

decreased. For 100% human movement, irrespective of the urban conformation, we

estimated a median of 4,228 cases (IQR: 3,025 – 4,921), which decreased to a median

of 764 (IQR: 349 – 1,626) and 154 (IQR: 108 – 232) cases for 50% and 20% human

movement, respectively (Fig 2 and Table S1) (all results are derived from the model

calibrated for Kisumu, Kenya; see supplementary material for further results for Ukunda,

Kenya).

Fig 2: Increasing the number of people moving from HH to NH significantly

increased the burden of dengue under three NH spatial distribution scenarios.

Three levels of human movement were assessed (20%, 50%, and 100%) on three

urban conformations (scattered, centered, or clustered). Boxplots show the distribution

of the total number of infections for 200 runs of two-year simulations where median is

the horizontal line, the filled box is the interquartile (IQR) range, the whiskers show the

values above and under the IQR and no more than 1.5·IQR, and dots are representing

values beyond this range.

Additionally, the scenario where NH locations are spatially randomly distributed

produced more cases, though at all movement levels the interquartile ranges for

different spatial configurations overlapped (Fig 2). At 100% movement, scattered

conformation yielded a median of 4,672 (IQR: 3,956 – 5,227) while the centered and

clustered scenarios produced, respectively, medians of 4,432 (IQR: 3,587 – 5,027) and

3,178 (IQR: 1,785 – 4,179) (Table S2). Trends were similar for the 50% and 20%

mobility levels (Fig 2).

The relative role of NH versus HH environments increased with the overall level of

mobility and hence transmission (Fig 3). At lower levels of movement (20%), the

number of infections was slightly higher in HH than NH. However, at higher levels of

movement (50-100%), the number of infections in NH was higher than HH. At 100%

movement, irrespective of urban conformation, NH produced 67% of the cases, but this

proportion decreased to 58.8% and to 42.3% at 50% and 20% of human movement,

respectively (Fig 3 and Table S1). Mosquito movement had the opposite influence on

transmission. Specifically, transmission decreased at higher levels of mosquito

movement due to migration-mediated mortality (see supplementary results and

discussion).

Fig 3: Number of infections are higher in NH than HH at high levels of human movement, but the HH contribution exceeds NH at low mobility and transmission levels. Three levels of human movement were assessed (20%, 50%, and 100%) on three urban conformations (scattered, centered, or clustered). Boxplots show the distribution of total number of infections for 200 runs of two-year simulations, spanning all three urban configurations, where median is the horizontal line, the filled box is the interquartile (IQR) range, the whiskers show the values above and under the IQR and no more than 1.5·IQR, and dots represent values beyond this range. The y-axis is represented in Log_{10} scale.

Distance from HH to NH makes little difference in dengue burden but defines level of spatial structure

Varying the distance between HH and NH had only slight impacts on the total number of

infections, with a slight increase in the number of cases with distance when NH are

clustered (Fig 4 and Table S3). Besides these slight changes, differences among urban

conformations were still evident (Fig 4).

Fig 4: Different distance regimes people travel from HH to NH results in slight

differences in cases and larger differences among configurations. Distance 0

means people only visit the closest NH location from HH. On the three urban

configuration scenarios, three levels of distance from HH to NH were assessed (the

closest [categorized as 0], at least 500 meters, and at least 1000 meters). Boxplots are

showing the total number of infections after two-year simulation for 200 runs.

Intrigued by the apparent lower importance of human movement distance in

transmission, we wanted to explore further by assessing the spatial structure of cases.

Given that the number of people moving affected the number of cases, we evaluated

the spatial structuring when distance traveled is considered as well. In general, we

found that irrespective of urban conformation, when people moved short distances the

level of spatial structure was higher, as expected (Fig 5). Similarly, the spatial

structuring levels was also modified by the number of people moving, where lower

levels of movement decreased Moran's I, thereby making cases more dispersed (Fig

S6).

Fig 5: Spatial structuring decreased with increased distance of movement.

Distribution of Moran's I values for 200 simulations for each of the three distance

regimes is shown. Values that are both significantly (α = 0.05; green) and non-

significantly (orange) different from zero are displayed by color. Three levels of distance

from HH to NH were assessed (the closest [categorized as 0], at least 500 meters, and

at least 1000 meters) on three urban conformations (scattered, centered, or clustered).

Discussion

Building on recent model results showing an important role of NH environments for dengue transmission and control, where over half of infections occur in NH environments, here we addressed how non-random mixing and mobility of humans and vectors affected dengue dynamics [20]. By extending the ABM to consider spatially-explicit urban conformations and movement levels, we showed that human movement is a primary driver of dengue dynamics, irrespective of urban spatial configuration. Further, we found qualitatively similar outcomes to those modeled for Kisumu for the coastal Kenyan city of Ukunda, presented in the supplementary material.

Among urban conformations, when NH spaces are scattered throughout the city it allows for closer connections to HH and therefore increased transmission. In this way, NH spaces serve as spreaders of infection since they are highly visited locations, which increases the chances of having a successful feeding encounter between humans and infected vectors. Once an individual is infected, the chances of infecting mosquitoes inside the household and in turn having another household inhabitant infected increases, generating local household chains of transmission.

For this reason, a lower number of individuals visiting NH locations nonlinearly reduces the burden of dengue; for example, a 50% reduction in mobility from 100% to 50% reduced cases by 82% (Fig 2). In this sense, when the number of people visiting NH decreases, the number of infections happening in these spaces also decreases until becoming roughly even with the number of infections in HH (Fig 3). These results are supported by previous reports showing that when COVID-19 pandemic lockdowns

forced people to stay at home most of the time, the number of dengue cases was

significantly reduced [24, 42, 43]. The idea of human movement affecting dengue

transmission is not new and has been explored previously, especially by Stoddard and

colleagues (2009), who focused their study on the movement of individuals between

houses only [21]. However, here we highlight how human mobility interacts with NH

spaces as drivers of transmission, which then disseminates within households. As a

result, when control is focused on households, it prevents the spread of disease to the

remaining household inhabitants but leaves broader foci of transmission active as long

as NH transmission is not under control.

Distance from HH to NH was not as important as overall levels of human movement for

dengue burden (Fig 4), suggesting that it is not how far people are traveling but the

destination and total amount of movement. This is in line with previous work where a

large, longitudinal study in Iquitos, Peru showed that human infection risk was mainly

driven by individuals visiting locations with presence of infected vectors, irrespective of

the distance [22]. It is important to note that our model does not account for movement

times, which increase with distance (but note that even our largest range of movement,

>1000m, is still very localized within a city's limits, which allowed us to identify patterns

but might be a limitation of such analysis). These NH spaces have been previously

described to have mosquitoes [44] and hence represent some degree of risk for

transmission when people are nearby. Unfortunately, we do not have data about the

time people spend in NH locations. Equally, the limited information collected about the

number of individuals within specific types of NH on a daily basis is a big limitation of

the study. Although our simulations relied on information from observations and

conversation with locals in Kisumu, Kenya, no quantitative activity space data exists in

this setting. Although the results should not be dramatically different, it is likely that true

patterns might be slightly different to what is reported in this work.

The distance that people travel to NH does, however, affect the urban spatial dynamics

of transmission. By increasing people's travel distances we are also increasing the

mixing of individuals. Distance traveled affects the level of clustering of cases, which is

a measure of the level of spatial dependence of cases and hence of how cases are

unevenly distributed in space (Fig 5) [41]. The role of mobility between HH and NH in

driving the clustering of cases is something that has not been explored before and

deserves further exploration to understand its implication for disease control program

design.

Although urban spatial configuration had subtle effects on the number and spatial

structure of dengue infections, human movement between HH and NH had a much

larger impact, with an 82% decline in cases as the number of people moving decreased

from 100% to 50%. Together, these results reflect the importance of NH and human

mobility between NH and HH spaces in dengue epidemiology. This underscores the

importance of vector control in NH spaces, which is not currently implemented in many

dengue endemic regions. Finally, though people's within-city travel distance did not

have a large impact on the number of cases, it is important for shaping spatial patterns,

which can have implications for control activities and for local herd immunity.

Acknowledgements

We would like to acknowledge Dr. Jason R. Andrews for the valuable input in this work.

We would also acknowledge the fieldwork teams (in Kisumu: Joel Mbakaya, Samwel

Ndire and Charles Adipo; in Ukunda: Said Lipi Malumbo, Paul S. Mutuku, Charles M.

Ng'ang'a) who collected the data that made this work possible.

Funding Statement

This research was funded by NIH through the grant R01AI102918 (PI LaBeaud). In

addition, V.H.P.G. is supported by grants R01AI102918 and R35GM133439; E.A.M. is

supported by NIH grants R35GM133439, R01AI102918, and R01AI168097, and NSF

grant DEB-2011147 (with Fogarty International Center); A.D.L. is supported by grants

R01AI102918, R01AI149614, R01AI155959, D43TW011547.

Competing interests

The authors declare that they have no competing interests.

Data availability statement

All the data, codes, and resources to recreate the simulations for the original model and

the modification reported in this work are publicly available at

https://github.com/vhpenagarcia/ABM_dengue and have been archived within the

Zenodo repository (https://doi.org/10.5281/zenodo.14036270).

References

1.   Guzman MG, Harris E. Dengue. Lancet. 2015;385(9966):453-65. doi: 10.1016/S0140-6736(14)60572-9. PubMed PMID: 25230594.

2.   Guzman MG, Gubler DJ, Izquierdo A, Martinez E, Halstead SB. Dengue infection. Nat Rev Dis Primers. 2016;2:16055. Epub 20160818. doi: 10.1038/nrdp.2016.55. PubMed PMID: 27534439.

3.   Scott TW, Amerasinghe PH, Morrison AC, Lorenz LH, Clark GG, Strickman D, et al. Longitudinal studies of *Aedes aegypti* (Diptera: Culicidae) in Thailand and Puerto Rico: blood feeding frequency. J Med Entomol. 2000;37(1):89-101. PubMed PMID: 15218911.

4.   Scott TW, Chow E, Strickman D, Kittayapong P, Wirtz RA, Lorenz LH, et al. Blood-feeding patterns of *Aedes aegypti* (Diptera: Culicidae) collected in a rural Thai village. J Med Entomol. 1993;30(5):922-7. PubMed PMID: 8254642.

5.   Montenegro-Quiñonez CA, Louis VR, Horstick O, Velayudhan R, Dambach P, Runge-Ranzinger S. Interventions against Aedes/dengue at the household level: a systematic review and meta-analysis. EBioMedicine. 2023;93:104660. Epub 20230621. doi: 10.1016/j.ebiom.2023.104660. PubMed PMID: 37352828; PubMed Central PMCID: PMCPMC10333437.

6.   World Health Organization. Dengue guidelines for diagnosis, treatment, prevention and control : new edition. Geneva: World Health Organization; 2009. x, 147 p. p.

7.   Huang CH, Lin CY, Yang CY, Chan TC, Chiang PH, Chen YH. Relationship between the Incidence of Dengue Virus Transmission in Traditional Market and Climatic Conditions in Kaohsiung City. Can J Infect Dis Med Microbiol. 2021;2021:9916642. Epub 20210809. doi: 10.1155/2021/9916642. PubMed PMID: 34422144; PubMed Central PMCID: PMCPMC8371670.

8.   Suwanbamrung C, Promsupa S, Doungsin T, Tongjan S. Risk factors related to dengue infections in primary school students: exploring students' basic knowledge of dengue and examining the larval indices in southern Thailand. J Infect Public Health. 2013;6(5):347-57. Epub 20130621. doi: 10.1016/j.jiph.2013.04.006. PubMed PMID: 23999339.

9.   Barrera R, Acevedo V, Amador M. Role of Abandoned and Vacant Houses on *Aedes aegypti* Productivity. Am J Trop Med Hyg. 2021;104(1):145-50. doi:

10.4269/ajtmh.20-0829. PubMed PMID: 33021195; PubMed Central PMCID:
PMCPMC7790113.

10.   Kampango A, Furu P, Sarath DL, Haji KA, Konradsen F, Schiøler KL, et al. Risk
factors for occurrence and abundance of *Aedes aegypti* and *Aedes bromeliae* at hotel
compounds in Zanzibar. Parasit Vectors. 2021;14(1):544. Epub 20211022. doi:
10.1186/s13071-021-05005-9. PubMed PMID: 34686195; PubMed Central PMCID:
PMCPMC8539800.

11.   Chiuya T, Masiga DK, Falzon LC, Bastos ADS, Fèvre EM, Villinger J. A survey of
mosquito-borne and insect-specific viruses in hospitals and livestock markets in western
Kenya. PLoS One. 2021;16(5):e0252369. Epub 20210528. doi:
10.1371/journal.pone.0252369. PubMed PMID: 34048473; PubMed Central PMCID:
PMCPMC8162702.

12.   Morrison AC, Sihuincha M, Stancil JD, Zamora E, Astete H, Olson JG, et al.
Aedes aegypti (Diptera: Culicidae) production from non-residential sites in the
Amazonian city of Iquitos, Peru. Ann Trop Med Parasitol. 2006;100 Suppl 1:S73-S86.
doi: 10.1179/136485906X105534. PubMed PMID: 16630393.

13.   Pai HH, Lu YL. Seasonal abundance of vectors at outdoor environments in
endemic and nonendemic districts of dengue in Kaohsiung, South Taiwan. J Environ
Health. 2009;71(6):56-60. PubMed PMID: 19192745.

14.   Pérez-Pérez J, Peña-García VH, Calle-Tobón A, Quimbayo-Forero M, Rojo R,
Henao E, et al. Entomovirological Surveillance in Schools: Are They a Source for
Arboviral Diseases Transmission? Int J Environ Res Public Health. 2021;18(11). Epub
20210606. doi: 10.3390/ijerph18116137. PubMed PMID: 34204166; PubMed Central
PMCID: PMCPMC8201003.

15.   García-Rejón JE, Loroño-Pino MA, Farfán-Ale JA, Flores-Flores LF, López-Uribe
MP, Najera-Vazquez MeR, et al. Mosquito infestation and dengue virus infection in
Aedes aegypti females in schools in Merida, Mexico. Am J Trop Med Hyg.
2011;84(3):489-96. doi: 10.4269/ajtmh.2011.10-0654. PubMed PMID: 21363990;
PubMed Central PMCID: PMCPMC3042828.

16.   Christophers SR. *Aedes aegypti* (L.), the yellow fever mosquito; its life history,
bionomics, and structure. Cambridge Eng.: University Press; 1960. xii, 738 p. p.

17.   Trpis M, McClelland GA, Gillett JD, Teesdale C, Rao TR. Diel periodicity in the
landing of *Aedes aegypti* on man. Bull World Health Organ. 1973;48(5):623-9. PubMed
PMID: 4544150; PubMed Central PMCID: PMCPMC2482924.

18.   Ndenga BA, Mutuku FM, Ngugi HN, Mbakaya JO, Aswani P, Musunzaji PS, et al.
Characteristics of *Aedes aegypti* adult mosquitoes in rural and urban areas of western
and coastal Kenya. PLoS One. 2017;12(12):e0189971. Epub 2017/12/19. doi:
10.1371/journal.pone.0189971. PubMed PMID: 29261766; PubMed Central PMCID:
PMCPMC5736227.

19.   Jemberie W, Dugassa S, Animut A. Biting Hour and Host Seeking Behavior
of *Aedes* Species in Urban Settings, Metema District, Northwest Ethiopia. Trop Med
Infect Dis. 2025;10(2). Epub 20250128. doi: 10.3390/tropicalmed10020038. PubMed
PMID: 39998042; PubMed Central PMCID: PMCPMC11860606.

20.   Peña-García VH, LaBeaud AD, Ndenga BA, Mutuku FM, Bisanzio D, Andrews
JR, et al. Non-household environments make a major contribution to dengue
transmission: implications for vector control. R Soc Open Sci. 2025;12(4):241919. Epub
20250416. doi: 10.1098/rsos.241919. PubMed PMID: 40242341; PubMed Central
PMCID: PMCPMC12000688.

21.   Stoddard ST, Morrison AC, Vazquez-Prokopec GM, Paz Soldan V, Kochel TJ,
Kitron U, et al. The role of human movement in the transmission of vector-borne
pathogens. PLoS Negl Trop Dis. 2009;3(7):e481. Epub 20090721. doi:
10.1371/journal.pntd.0000481. PubMed PMID: 19621090; PubMed Central PMCID:
PMCPMC2710008.

22.   Stoddard ST, Forshey BM, Morrison AC, Paz-Soldan VA, Vazquez-Prokopec GM,
Astete H, et al. House-to-house human movement drives dengue virus transmission.
Proc Natl Acad Sci U S A. 2013;110(3):994-9. doi: 10.1073/pnas.1213349110. PubMed
PMID: 23277539; PubMed Central PMCID: PMCPMC3549073.

23.   Wesolowski A, Qureshi T, Boni MF, Sundsøy PR, Johansson MA, Rasheed SB,
et al. Impact of human mobility on the emergence of dengue epidemics in Pakistan.
Proc Natl Acad Sci U S A. 2015;112(38):11887-92. Epub 20150908. doi:
10.1073/pnas.1504964112. PubMed PMID: 26351662; PubMed Central PMCID:
PMCPMC4586847.

24.   Liyanage P, Rocklöv J, Tissera HA. The impact of COVID-19 lockdown on
dengue transmission in Sri Lanka; A natural experiment for understanding the influence
of human mobility. PLoS Negl Trop Dis. 2021;15(6):e0009420. Epub 20210610. doi:
10.1371/journal.pntd.0009420. PubMed PMID: 34111117; PubMed Central PMCID:
PMCPMC8192006.

25.   Massaro E, Kondor D, Ratti C. Assessing the interplay between human mobility
and mosquito borne diseases in urban environments. Sci Rep. 2019;9(1):16911. Epub
20191115. doi: 10.1038/s41598-019-53127-z. PubMed PMID: 31729435; PubMed
Central PMCID: PMCPMC6858332.

26.   Kenya Ministry of Education. Basic Education Statistical Booklet, 2019. Nairobi:
2019.

27.   Kenya National Bureau of Statistics. Quarterly Labour Force Report. Quarter 1.
Nairobi: 2021.

28.   Kenya National Bureau of Statistics. 2019 Kenya Population and Housing
Census. Volume 1: Population by County and Sub-county. Nairobi: 2019.

29.   Peña-García VH, Mutuku FM, Ndenga BA, Mbakaya JO, Ndire SO, Agola GA, et
al. The Importance of Including Non-Household Environments in Dengue Vector Control
Activities. Viruses. 2023;15(7). Epub 20230714. doi: 10.3390/v15071550. PubMed
PMID: 37515236; PubMed Central PMCID: PMCPMC10384488.

30.   Mordecai EA, Cohen JM, Evans MV, Gudapati P, Johnson LR, Lippi CA, et al.
Detecting the impact of temperature on transmission of Zika, dengue, and chikungunya
using mechanistic models. PLoS Negl Trop Dis. 2017;11(4):e0005568. Epub
2017/04/27. doi: 10.1371/journal.pntd.0005568. PubMed PMID: 28448507; PubMed
Central PMCID: PMCPMC5423694.

31.   Huber JH, Childs ML, Caldwell JM, Mordecai EA. Seasonal temperature variation
influences climate suitability for dengue, chikungunya, and Zika transmission. PLoS
Negl Trop Dis. 2018;12(5):e0006451. Epub 20180510. doi:
10.1371/journal.pntd.0006451. PubMed PMID: 29746468; PubMed Central PMCID:
PMCPMC5963813.

32.   Caldwell JM, LaBeaud AD, Lambin EF, Stewart-Ibarra AM, Ndenga BA, Mutuku
FM, et al. Climate predicts geographic and temporal variation in mosquito-borne
disease dynamics on two continents. Nat Commun. 2021;12(1):1233. Epub 20210223.
doi: 10.1038/s41467-021-21496-7. PubMed PMID: 33623008; PubMed Central PMCID:
PMCPMC7902664.

33.   Inziani M, Adungo F, Awando J, Kihoro R, Inoue S, Morita K, et al.
Seroprevalence of yellow fever, dengue, West Nile and chikungunya viruses in children
in Teso South Sub-County, Western Kenya. Int J Infect Dis. 2020;91:104-10. Epub
20191108. doi: 10.1016/j.ijid.2019.11.004. PubMed PMID: 31712089.

34.   Sabin AB. The dengue group of viruses and its family relationships. Bacteriol Rev.
1950;14(3):225-32. PubMed PMID: 14772197.

35.   Sabin AB. Research on dengue during World War II. Am J Trop Med Hyg.
1952;1(1):30-50. doi: 10.4269/ajtmh.1952.1.30. PubMed PMID: 14903434.

36.   McCormack CP, Ghani AC, Ferguson NM. Fine-scale modelling finds that
breeding site fragmentation can reduce mosquito population persistence. Commun Biol.
2019;2:273. Epub 20190725. doi: 10.1038/s42003-019-0525-0. PubMed PMID:
31372512; PubMed Central PMCID: PMCPMC6658551.

37.   Nathan R, Klein E, Robledo-Arnuncio JJ, Revilla E. Dispersal kernels: review. In:
Clobert J, Baguette M, Benton TG, Bullock JM, editors. Dispersal Ecology and
Evolution: Oxford University Press; 2012. p. 0.

38.   Marcantonio M, Reyes T, Barker CM. Quantifying *Aedes aegypti* dispersal in
space and time: a modeling approach. Ecosphere. 2019;10(12):e02977. doi:
https://doi.org/10.1002/ecs2.2977.

39.   Moore TC, Brown HE. Estimating *Aedes aegypti* (Diptera: Culicidae) Flight
Distance: Meta-Data Analysis. J Med Entomol. 2022;59(4):1164-70. doi:
10.1093/jme/tjac070. PubMed PMID: 35640992.

40.   Moran PAP. The Interpretation of Statistical Maps. Journal of the Royal Statistical
Society: Series B (Methodological). 1948;10(2):243-51. doi: 10.1111/j.2517-
6161.1948.tb00012.x.

41.   Pfeiffer DU, Robinson TP, Stevenson M, Stevens KB, Rogers DJ, Clements ACA.
Spatial clustering of disease and global estimates of spatial clustering. Spatial Analysis
in Epidemiology: Oxford University Press; 2008. p. 0.

42.   Wilder-Smith A. Dengue during the COVID-19 pandemic. J Travel Med.
2021;28(8). doi: 10.1093/jtm/taab183. PubMed PMID: 34850050; PubMed Central
PMCID: PMCPMC8690170.

43.   Sasmono RT, Santoso MS. Movement dynamics: reduced dengue cases during
the COVID-19 pandemic. Lancet Infect Dis. 2022;22(5):570-1. Epub 20220302. doi:
10.1016/S1473-3099(22)00062-7. PubMed PMID: 35247322; PubMed Central PMCID:
PMCPMC8890753.

44.   Ngugi HN, Mutuku FM, Ndenga BA, Musunzaji PS, Mbakaya JO, Aswani P, et al.
Characterization and productivity profiles of Aedes aegypti (L.) breeding habitats across
rural and urban landscapes in western and coastal Kenya. Parasit Vectors.
2017;10(1):331. Epub 20170712. doi: 10.1186/s13071-017-2271-9. PubMed PMID:
28701194; PubMed Central PMCID: PMCPMC5508769.

Scattered	Centered	Clustered

Workplace • Market • Religious • Recreational • School