Mobility and non-household environments:

understanding dengue transmission patterns in urban

contexts

Víctor Hugo Peña-García[1, *], Bryson A. Ndenga[2], Francis M. Mutuku[3], Donal Bisanzio[4], A. Desiree LaBeaud[5], Erin A. Mordecai[1]

[1] Department of Biology, Stanford University, Stanford, CA, USA

[2] Kenya Medical Research Institute, Kisumu, Kenya

[3] Department of Environmental and Health Sciences, Technical University of Mombasa, Mombasa, Kenya

[4] RTI International, Washington, DC, USA

[5] School of Medicine, Stanford University, Stanford, CA, USA

[*] Corresponding author: Víctor Hugo Peña-García

email: victorhugopega@gmail.com

Abstract

Households (HH) have been traditionally described as the main environments where people are at risk of dengue and other arbovirus infections. Mounting entomological evidence suggests a larger role for environments other than HH. Recently, an agent-based model (ABM) estimated that over half of infections occur in non-household (NH) environments such as workplaces, markets, and recreational sites. However, the influence of human and vector mobility and urban spatial configurations on the effects of NH on dengue transmission remains understudied. To address this gap, we expanded an ABM calibrated with field data from Kenya to examine movement of people and vectors under different spatial configurations of buildings. We assessed the number of people traveling between HH and NH and the distances traveled, in three urban configurations: NH distributed randomly (scattered), concentrated in a single center, or clustered in multiple centers. Across simulations, the number of people moving was the most influential variable, with higher movement between HH and NH increasing case numbers. The number of cases was also higher when NH were scattered. Intriguingly, the distance people traveled from HH to NH had little effect on dengue burden but influenced the spatial clustering of cases. These findings underscore the role of NH as major spreaders of infections between HH and NH environments, and the importance of human movement in driving dengue dynamics.

39 **Keywords**

40 Dengue transmission, Agent-based model, Human mobility, Urban Configuration, Non-

41 household environments, Spatial Structure

43 Introduction

Dengue is a vector-borne disease prevalent and on the rise in most of the tropical and

subtropical regions around the globe with frequent number of cases being present in

non-endemic areas like USA and Europe.[1, 2] The main vector, *Aedes aegypti,* is highly

anthropophilic, found very close to human environments and impacting the public health

of urban environments.[3, 4]

Historically, the main strategy to control the disease has been to reduce vector-human

contact by reducing the size of mosquito populations. Conventional wisdom is that for

effective control, these activities should be focused on households (HH) as the main

environment where transmission is happening.[5, 6] However, some studies have

suggested that locations other than households might have an important role because of

a significant presence of mosquitoes,[7, 8, 9, 10] and infected vectors.[11] In a recent study, we

used an agent-based model (ABM) to quantify the number of infections in different types

of urban spaces and estimated that over half of infections are happening in non-

household (NH) environments, where the main high-risk spaces are workplaces and

markets/shops.[12]

These results have implications for dengue epidemiology since the high flux of people

through NH suggests that these spaces can contribute to the spread of infections. In this

way, the total number of infections can be affected by the distribution in space of NH

and the movement of people between HH and NH. However, these intra-urban

dynamics have never been described in the context of different roles among HH and

different NH space types,[13, 14, 15, 16] though Massaro and colleagues used mobile phone

data to get estimates for movement between workplaces.[17]

Building on a previous result showing the importance of NH for dengue transmission, we

now evaluate how levels of human and vector mobility as well as different urban spatial

configurations of NH affect dengue transmission. In particular, what role does spatial

configuration of NH spaces play, along with the extent to which people and mosquitoes

move between spaces, in determining dengue dynamics? To address this, we modified

a previously published ABM[12] to make it spatially explicit by assigning coordinates that

mimic different urban conformations and evaluated different scenarios of movement of

people and vectors. We then assessed how these variables affect the burden of dengue

and the spatial patterns to understand urban-level transmission dynamics.

Methods

Model overview

To achieve the aims of this study, we modified the ABM previously used to describe the importance of HH and NH in transmission.[12] The model was developed to quantify the relative contribution of five different types of NH (workplaces, markets or shops, recreational, religious, and schools) and HH to dengue burden. The model development and calibration were based on data from two Kenyan cities: Kisumu in the west and Ukunda on the coast.[12] Here, we focus on parameters calibrated to Kisumu, although additional results including dynamics from Ukunda are found in supplementary material.

The model represents the movement of people between HH and two different types of NH locations: daily-commuting locations where individuals attend daily and meet with the same individuals like schools and workplaces, and randomly assigned locations for which both the number and identity of people who visit them is randomly defined every day. The latter include locations like markets or shops, recreational and religious spaces. Movement between HH (categorized as "visit") is also included in the model for which its frequency is daily determined with probability of 0.1. Based on vector surveys conducted over two years of fieldwork previously published,[18] NH and HH environments were assigned to have mosquito presence or absence based on observed prevalence of mosquitoes. Population dynamics of vectors were modeled at the building level, whereby the sub-population dynamics are determined based on building-level conditions like the presence of water containers and the total amount of water they can

hold, again informed by field vector surveillance data. These dynamics as well as

infection dynamics of vectors are also determined by temperature by using functions

previously described and widely used elsewhere.[19, 20, 21]

Initial baseline prevalence of dengue was set to 0.08%, estimated from previous studies

reporting age-structured seroprevalence[22] with an incidence rate per year estimated as

$IR = -\frac{1}{age}\ln(1 - prevalence)$. Transmission events happen in those locations where

infected vectors contact susceptible humans or vice versa. Mosquitoes bite depending

on both temperature-dependent biting rate and the probability of having a successful

vector-human encounter, which depends on the amount of time that humans spend in

the location (Details can be found in supplementary material). Infection status of

mosquitoes can be either susceptible, exposed, or infected while humans can be either

susceptible, exposed, infected, or recovered and (temporarily) immune. The time that

mosquitoes spend as exposed depends on temperature (extrinsic incubation period)

and is determined in the model by equations reported previously by Mordecai and

colleagues.[19] Once infected, mosquitoes remain in this stage until death, which is

evaluated daily following a temperature-dependent death rate. Humans remain

susceptible until they are bit by infected mosquitoes and moved to latent stage where

they remain for five days. Then, the individual is moved to the infectious stage which

lasts seven days before moving to the recovered stage. Since the model does not

explicitly represent dynamics of different serotypes, waning immune protection was

based on Sabin's classic studies describing the loss of complete heterotypic protection

after roughly three months[23, 24] to set a return to susceptible after 100 days on recovered

status. The number of infections in each location is recorded daily. Statistics about the

total number of infections and locations are provided weekly. The model simulates

transmission dynamics happening for 731 days (comprising temperature conditions

between January 1st of 2020 until December 31st of 2021) and results are shown as a

distribution of the number of infections over 200 simulations. The model and

modifications described in this work were coded in Julia language (v1.10.0) and

simulations were run on Sherlock computational cluster (Stanford Research Computing

Center).

128   Spatial variables

The original model was not spatially explicit and hence the movement of individuals was

assumed to be totally random. As such, the resultant infections arise from complete

mixing of individuals among structures, which is not realistic and does not capture local,

intra-urban spatial phenomena. To include mobility-associated variables and describe

such local phenomena, we made the model spatially explicit. No real spatial coordinates

were used so we could record the outcome when different urban conformations are

tested by using the same set of spatial coordinates. In this way, we can assure that

differences are due to the building designation as HH or NH and not by the specific set

of spatial coordinates. Because the model considers synthetic populations of about

20,000 people, the total municipality areas were rescaled to fit the total number of

structures of the virtual populations while considering similar densities. Spatial

coordinates were randomly generated to create a synthetic settlement and assigned to

each structure of the population. With the aim to capture different real setting

conformations, coordinate assignments were done based on either "Scattered"

(randomly distributed), "Centered" (majority of NH concentrated in the center of the city

in a single cluster), or "Clustered" (majority of NH concentrated in three clusters)

configurations (Figure 1).

Figure 1: Spatial distribution of NH for three different urban conformations to be

tested: scattered (NH randomly distributed in space), centered (majority of NH are

clustered in the center), and clustered (majority of NH are grouped in three clusters).

NH are shown with colors while HH are displayed as light gray points. Religious and

schools are represented in the first row, markets/shops and recreational in middle row

and more frequent NH, workplaces, are represented in the lower row.

[ALT TEXT: Arrangement of plots showing the distribution of non-household (NH)

environments in three patterns: scattered in the left column, centered in a single dense

area in the middle column, and clustered into three major concentrations in the right

column. The top row shows religious and school NH, the middle row shows markets and

recreational NH, and the bottom row shows workplace NH.]

Movement of people

We included two movement-related variables: the distance from each household (HH) to

the nearest non-household (NH) locations and the number of people moving. To control

movement distances, we applied three treatments: (1) limiting attendance to the nearest

NH locations (assigned as distance zero), (2) allowing attendance to NH locations at

least 500 meters away, and (3) allowing attendance to NH locations at least 1000 meters

away from each HH. These treatments were applied across the three different urban

configurations. It is worth noting that no experiments involving humans were conducted,

and the results presented in this work are limited to simulations performed using

synthetic human populations.

In the clustered or centralized configurations, assigning the closest NH locations to HHs

would primarily select NH sites at the periphery of each cluster, potentially leading to

biased representations. To address this, we generated a list of NH locations sorted by

proximity for each HH and allocated the closest NHs based on the number of inhabitants

in the HH. For example, if a HH had four inhabitants, the four closest NHs were assigned

to it. This approach ensured a more representative distribution across all HHs.

The number of people visiting NH was simulated at three levels. First, we included the

same levels previously described in the model, categorized as 100% mobility.[12] This

treatment includes all students and workers attending their respective school and

workplace, and random-attendance locations (religious, markets/shops, and

recreational). Unfortunately, no data were available on the number of individuals visiting

these types of locations. We anticipated significant variability due to the influence of

several unmodeled factors. To account for this uncertainty, we relied on discussions

with local residents to estimate a range of possible visitor numbers. We then applied a

uniform distribution, with a minimum of 10 and a maximum of 70 individuals, to cover a

broad spectrum of potential scenarios. The number of people visiting a given location is

determined daily. We also evaluated the number of cases when decreasing the number

of people moving to school and workplaces to 50% and remaining NH locations to a

uniform distribution with parameters minimum = 5 and maximum = 35. Finally, we

include a level of people movement of 20% for school and workplaces and uniform

distribution with parameters minimum = 2 and maximum = 14 for random-attendance

locations.

Movement of mosquitoes

In line with the inclusion of spatial variables and human movement, we included the movement of vectors. There were no data available to characterize the rate of movement of mosquitoes into new areas. As a result, we parameterized such movement by considering two variables—availability of both breeding and blood-feeding resources—to estimate a baseline migration probability for the number of mosquitoes moving from given locations.

The model includes a local density-dependent function that allows for the mosquito population to grow in a location-specific way, representing dynamics previously described for a fragmented environment.[25] The function is already described elsewhere[12] and depends on temperature (peaking at 29°C) and the density of immatures according to water availability, as follows:

$$f(D) = \frac{1}{1 + e^{bD-a}} d(T) \tag{1}$$

Where

$$d(T) = -0.166 + 0.08T - 0.0014T^2 \tag{2}$$

In this equation a and b are calibrated parameters, D is the larval density expressed as the ratio of the number of larvae to liters of water available for breeding in the structure, and $d(T)$ is the term describing the temperature-dependence of population growth (additional details in supplementary material).

By using this function, the growth of the mosquito sub-population (in a given building)

depends on the amount of water resources available. In this way, when the mosquito

population has grown so the water resources are depleted (reaching the carrying

capacity), the mortality of immatures is higher and the probability for a given mosquito to

migrate out of the building increases (details provided in supplementary material).

Additionally, we considered for migration the possibility that human blood availability is

not satisfying the mosquito subpopulation needs in a given building. For this purpose,

the number of females that fed in each day was estimated by considering the number of

females biting (N_b) and the probability for those to have a successful feeding encounter

with a human the same day ($P(bit)$). The latter is determined considering the number of

inhabitants, the time they spend at home and the time a visitor spends in each location

(details are provided in supplementary material). The final number of fed females (N_F) is

estimated by

$$N_F \sim Bin\big(N_b, P(bit)\big) \tag{3}$$

Thus, when the difference between the number of mosquitoes trying to bite and the

actual number of mosquitoes that are fed is large, the probability of migration increases.

Once a mosquito migrates, a new location is assigned by considering a dispersal

kernel[26]. Following previous work[27], we used a lognormal function with the form

$$P_D = \frac{1}{(2\pi)^{\frac{3}{2}} b d^2} e^{-\frac{\log(d/a)^2}{2b^2}} \tag{4}$$

Where d is distance and both a and b are parameters to be estimated. We fitted a

function by assuming a mean dispersal distance of 105.69 meters, as estimated for

Aedes aegypti by.[28]

Spatial autocorrelation

By taking advantage of spatial features introduced in the model, we performed a spatial

autocorrelation analysis to evaluate the level of clustering of dengue cases recorded in

each simulation as a function of urban configuration and human and mosquito mobility.

To do this, the household location of each infection is recorded so a Global Moran's I

index[29] could be estimated at the end of the simulated period. Global Moran's I ranges

from -1 to +1 where -1 means totally dispersed location of cases while a value of 1

represents a spatial distribution that is totally clustered (total separation between

locations with dengue cases and those without cases). In this sense, the null hypothesis

of this analysis is that dengue cases are randomly distributed in municipalities,

represented by Moran's I value of 0.[30] The analysis was done for every simulation by

using 1,000 permutations for inference in each of them. Analyses were done using the

package SpatialDependence.jl implemented in Julia language (v 1.10.0). Code is

available in the GitHub repository (https://github.com/vhpenagarcia/ABM_dengue) and

have been archived within the Zenodo repository

(https://doi.org/10.5281/zenodo.14036270).

Results

Burden of dengue is strongly affected by number of people visiting NH

We quantified the total number of cases after two years of transmission. When we simulated the epidemic under different human movement regimes, it was evident that the number of cases decreased as the number of people moving from HH decreased. For 100% human movement, irrespective of the urban conformation, we estimated a median of 4,228 cases (IQR: 3,025 – 4,921), which decreased to a median of 764 (IQR: 349 – 1,626) and 154 (IQR: 108 – 232) cases for 50% and 20% human movement, respectively (Figure 2 and S1 Table) (all results are derived from the model calibrated for Kisumu, Kenya; see supplementary material for further results for Ukunda, Kenya).

Figure 2: Increasing the number of people moving from HH to NH significantly

increases the burden of dengue under three NH spatial distribution scenarios.

Three levels of human movement were assessed (20%, 50%, and 100%) on three urban

conformations (scattered, centered, or clustered). Boxplots shows the distribution of the

total number of infections for 200 runs of two-year simulation where median is the

horizontal line, the filled box is the interquartile (IQR) range, the whiskers show the

values above and under the IQR and no more than 1.5·IQR, and dots are representing

values beyond this range.

[ALT TEXT: Box plot showing that the number of cases increases as the percentage of

people moving from HH locations rises. Small variations are also observed among urban

configurations—slightly higher for scattered, followed by centered and clustered—but

differences by movement percentage are more pronounced.]

Additionally, the scenario where NH locations are spatially randomly distributed

produced more cases, though at all movement levels the interquartile ranges for

different spatial configurations overlapped (Figure 2). At 100% movement, scattered

conformation yielded a median of 4,672 (IQR: 3,956 – 5,227) while the centered and

clustered scenarios produced, respectively, medians of 4,432 (IQR: 3,587 – 5,027) and

3,178 (IQR: 1,785 – 4,179) (S2 Table).

When we compared HH and NH environments, the number of infections is slightly

higher in HH at lower levels of movement and hence lower transmission. However, at

higher levels of movement, the number of infections in NH are higher than HH. In this

way, at 100% movement, irrespective of urban conformation, NH produced 67% of the

cases, but this proportion decreased to 58.8% and to 42.3% at 50% and 20% of human

movement, respectively (Figure 3 and S1 Table).

It is worth noting that different levels of mosquito movement were found to influence

transmission. Specifically, transmission decreases at higher levels of movement due to

migration-mediated mortality (see supplementary results and discussion).

Figure 3: Number of infections are higher in NH than HH at high levels of human movement but becomes more even when levels of transmission are low due to lower movement. Three levels of human movement were assessed (20%, 50%, and 100%) on three urban conformations (scattered, centered, or clustered). Boxplots show the distribution of total number of infections for 200 runs of two-year simulation where median is the horizontal line, the filled box is the interquartile (IQR) range, the whiskers show the values above and under the IQR and no more than 1.5·IQR, and dots represent values beyond this range. The y-axis is represented in Log_{10} scale.

303 [ALT TEXT: Box plot comparing HH and NH environments as the percentage of people

304 moving from HH increases. For 20% and 50% movement, case numbers are similar in

305 both environments, while at 100% movement, cases are higher in NH environments.]

307 Distance from HH to NH makes little difference in dengue burden

308 but defines level of spatial structure

309 Varying the distance between HH and NH had only slight impacts on the total number of

310 infections, with a slight increase in the number of cases with distance when NH are

311 clustered (Figure 4 and S3 Table). Besides these slight changes, differences among

312 urban conformations are still evident (Figure 4).

Figure 4: Different distance regimes people travel from HH to NH results in slight differences in cases and larger differences among configurations. Distance 0 means people only visit the closest NH location from HH. On the three urban configuration scenarios, three levels of distance from HH to NH were assessed (the closest [categorized as 0], at least 500 meters, and at least 1000 meters). Boxplots are showing the total number of infections after two-year simulation for 200 runs.

[ALT TEXT: Box plot showing no significant differences in case numbers among the three urban configurations across the three distance categories—closest, 500 meters, and 1000 meters—from household (HH) locations. A slight tendency toward more homogeneous case numbers is observed as distance increases.]

Intrigued by the apparent lower importance of human movement distance in

transmission, we wanted to explore further by assessing the spatial structure of cases.

Given that the number of people moving affects the number of cases, we evaluated the

spatial structuring when distance traveled is considered as well. In general, we found

that irrespective of urban conformation, when people move short distances the level of

spatial structure is higher, as expected (Figure 5). Similarly, the spatial structuring levels

are also modified by the number of people moving, where lower levels of movement

decrease Moran's I, thereby making cases more dispersed (Figure S6).

Figure 5: Spatial structuring decreases with increased distance of movement.

Distribution of Moran's I values for 200 simulations for each of the three distance

regimes is shown. Values that are both significantly (α = 0.05) and non-significantly

different from zero are displayed by color. Three levels of distance from HH to NH were

assessed (the closest [categorized as 0], at least 500 meters, and at least 1000 meters)

on three urban conformations (scattered, centered, or clustered).

[ALT TEXT: Box plot showing that Moran's I values shift toward a higher range as the

distance of people moving from HH locations increases. The category representing zero

distance—people moving to the closest NH location—shows the highest Moran's I

values.]

Discussion

Despite recent results of this model showing an important role of NH environments for

dengue transmission and control, where over half of infections occur in NH

environments, it remained unknown how non-random mixing and mobility of humans

and vectors affected dengue dynamics.[12] By extending the ABM to consider spatially-

explicit urban conformations and movement levels, we showed that human movement is

a primary driver of dengue dynamics, irrespective of urban spatial configuration.

Further, qualitatively similar, outcomes generated by simulating conditions from the

coastal Kenyan city of Ukunda supports these results and can be found in

supplementary material.

Among urban conformations, when NH spaces are scattered throughout the city it

allows for closer connections to HH and therefore increased transmission. In this way,

NH spaces serve as spreaders of infection since they are highly visited locations, which

increases the chances of having a successful feeding encounter between humans and

infected vectors. Once an individual is infected, the chances of infecting mosquitoes

inside the household and in turn having another household inhabitant infected

increases, generating local household chains of transmission.

For this reason, a lower number of individuals visiting NH locations tends to reduce the

burden of dengue (Figure 2). In this sense, when the number of people visiting NH

decreases, the number of infections happening in these spaces also decreases until

becoming roughly even with the number of infections in HH. These results are

supported by previous reports showing that when COVID-19 pandemic lockdowns

forced people to stay at home most of the time, the number of dengue cases was

significantly reduced.[16, 31, 32] The idea of human movement guiding different levels of

transmission is not new and has been explored previously, especially by Stoddard and

colleagues (2009), who focused their study on the movement of individuals between

houses only.[13] However, here we highlight how human mobility interacts with NH spaces

as drivers of transmission, which then disseminates within households. As a result, when

control is focused on households, it prevents the spread of disease to the remaining

household inhabitants but leaves broader foci of transmission active as long as NH

transmission is not under control.

Distance from HH to NH was not as important as overall levels of human movement for

dengue burden (Figure 4). This supports our previous result by suggesting that it is not

how far people are traveling but the destination and total amount of movement. This is in

line with previous work where a large, longitudinal study in Iquitos, Peru showed that

human infection risk is mainly driven by individuals visiting locations with presence of

infected vectors, irrespective of the distance.[14] It is important to note that our model

does not account for movement times, which increase with distance (but note that even

our largest range of movement, >1000m, is still very localized within a city's limits).

These NH spaces have been previously described to have mosquitoes[33] and hence

represent some degree of risk for transmission when people are nearby. Unfortunately,

we do not have data about the time people spend in NH locations. We think that the

number of infections happening while people are moving through NH locations is likely

to be negligible, but this merits further work.

The distance that people travel to NH does, however, affect the urban spatial dynamics

of transmission. By increasing people's traveling distances we are also increasing the

mixing of individuals. Distance traveled affects the level of clustering of cases, which is a

measure of the level of spatial dependence of cases and hence of how cases are

unevenly distributed in space (Figure 5).[30] The clustering of cases considering the major

role in transmission of NH is something that has not been explored before and deserves

further exploration to understand its implication for disease control program design.

Although urban spatial configuration had subtle effects on the number and spatial

structure of dengue infections, human movement between HH and NH had a much

larger impact, with an 82% decline in cases as the number of people moving decreased

from 100% to 50%. Together, these results reflect the importance of NH and human

mobility between NH and HH spaces in dengue epidemiology. This underscores the

importance of vector control in NH spaces, which is not currently implemented in many

dengue endemic regions. Finally, though people's travel distance did not have a large

impact on the number of cases, it is important for shaping spatial patterns, which can

have implications for control activities and for local herd immunity.

Acknowledgements

409   We would like to acknowledge Dr. Jason R. Andrews for the valuable input in this work.

410   We would also acknowledge the fieldwork teams (in Kisumu: Joel Mbakaya, Samwel

411   Ndire and Charles Adipo; in Ukunda: Said Lipi Malumbo, Paul S. Mutuku, Charles M.

412   Ng'ang'a) who collected the data that made this work possible. We also acknowledge

413   NIH for funding through project R01 AI102918 (ADL).

Financial Support

416   This research was funded by NIH through the grant R01AI102918 (PI LaBeaud). In

417   addition, V.H.P.G. is supported by grants R01AI102918 and R35GM133439; E.A.M. is

418   supported by NIH grants R35GM133439, R01AI102918, and R01AI168097, and NSF

419 grant DEB-2011147 (with Fogarty International Center); A.D.L. is supported by grants

420 R01AI102918, R01AI149614, R01AI155959, D43TW011547.

Competing interests

423 The authors declare that they have no competing interests.

Authors' Information:

VHPG: Department of Biology, Stanford University, Stanford, CA, USA.

victorhugopega@gmail.com

BAN: Kenya Medical Research Institute, Kisumu, Kenya, bndenga@yahoo.com

FMM: Department of Environmental and Health Sciences, Technical University of

Mombasa, Mombasa, Kenya, fmutuku73@gmail.com

DB: RTI International, Washington, DC, USA, dbisanzio@rti.org

ADL: School of Medicine, Stanford University, Stanford, CA, USA,

dlabeaud@stanford.edu

EAM: Department of Biology, Stanford University, Stanford, CA, USA,

emordeca@stanford.edu

References

1.   Guzman MG, Harris E, 2015. Dengue. Lancet 385: 453-65.

2.   Guzman MG, Gubler DJ, Izquierdo A, Martinez E, Halstead SB, 2016. Dengue infection.
Nat Rev Dis Primers 2: 16055.

3.   Scott TW, Amerasinghe PH, Morrison AC, Lorenz LH, Clark GG, Strickman D,
Kittayapong P, Edman JD, 2000. Longitudinal studies of *Aedes aegypti* (Diptera:
Culicidae) in Thailand and Puerto Rico: blood feeding frequency. J Med Entomol 37: 89-
101.

4.   Scott TW, Chow E, Strickman D, Kittayapong P, Wirtz RA, Lorenz LH, Edman JD, 1993.
Blood-feeding patterns of *Aedes aegypti* (Diptera: Culicidae) collected in a rural Thai
village. J Med Entomol 30: 922-7.

5.   Montenegro-Quiñonez CA, Louis VR, Horstick O, Velayudhan R, Dambach P, Runge-
Ranzinger S, 2023. Interventions against Aedes/dengue at the household level: a
systematic review and meta-analysis. EBioMedicine 93: 104660.

6.   World Health Organization., 2009. Dengue guidelines for diagnosis, treatment,
prevention and control : new edition. Geneva: World Health Organization.

7.   Huang CH, Lin CY, Yang CY, Chan TC, Chiang PH, Chen YH, 2021. Relationship
between the Incidence of Dengue Virus Transmission in Traditional Market and Climatic
Conditions in Kaohsiung City. Can J Infect Dis Med Microbiol 2021: 9916642.

8.   Suwanbamrung C, Promsupa S, Doungsin T, Tongjan S, 2013. Risk factors related to
dengue infections in primary school students: exploring students' basic knowledge of
dengue and examining the larval indices in southern Thailand. J Infect Public Health 6:
347-57.

9.   Barrera R, Acevedo V, Amador M, 2021. Role of Abandoned and Vacant Houses
on *Aedes aegypti* Productivity. Am J Trop Med Hyg 104: 145-150.

10. Kampango A, Furu P, Sarath DL, Haji KA, Konradsen F, Schiøler KL, Alifrangis M, Saleh
F, Weldon CW, 2021. Risk factors for occurrence and abundance of *Aedes aegypti* and
Aedes bromeliae at hotel compounds in Zanzibar. Parasit Vectors 14: 544.

11.   Pérez-Pérez J, Peña-García VH, Calle-Tobón A, Quimbayo-Forero M, Rojo R, Henao E,
469   Shragai T, Rúa-Uribe G, 2021. Entomovirological Surveillance in Schools: Are They a
470   Source for Arboviral Diseases Transmission? Int J Environ Res Public Health 18.

471   12.   Peña-García VH, LaBeaud AD, Ndenga BA, Mutuku FM, Bisanzio D, Andrews JR,
472   Mordecai EA, 2025. Non-household environments make a major contribution to dengue
473   transmission: implications for vector control. R Soc Open Sci 12: 241919.

474   13.   Stoddard ST, Morrison AC, Vazquez-Prokopec GM, Paz Soldan V, Kochel TJ, Kitron U,
475   Elder JP, Scott TW, 2009. The role of human movement in the transmission of vector-
476   borne pathogens. PLoS Negl Trop Dis 3: e481.

477   14.   Stoddard ST, Forshey BM, Morrison AC, Paz-Soldan VA, Vazquez-Prokopec GM,
478   Astete H, Reiner RC, Vilcarromero S, Elder JP, Halsey ES, Kochel TJ, Kitron U, Scott
479   TW, 2013. House-to-house human movement drives dengue virus transmission. Proc
480   Natl Acad Sci U S A 110: 994-9.

481   15.   Wesolowski A, Qureshi T, Boni MF, Sundsøy PR, Johansson MA, Rasheed SB, Engø-
482   Monsen K, Buckee CO, 2015. Impact of human mobility on the emergence of dengue
483   epidemics in Pakistan. Proc Natl Acad Sci U S A 112: 11887-92.

484   16.   Liyanage P, Rocklöv J, Tissera HA, 2021. The impact of COVID-19 lockdown on dengue
485   transmission in Sri Lanka; A natural experiment for understanding the influence of
486   human mobility. PLoS Negl Trop Dis 15: e0009420.

487   17.   Massaro E, Kondor D, Ratti C, 2019. Assessing the interplay between human mobility
488   and mosquito borne diseases in urban environments. Sci Rep 9: 16911.

489   18.   Peña-García VH, Mutuku FM, Ndenga BA, Mbakaya JO, Ndire SO, Agola GA, Mutuku
490   PS, Malumbo SL, Ng'ang'a CM, Andrews JR, Mordecai EA, LaBeaud AD, 2023. The
491   Importance of Including Non-Household Environments in Dengue Vector Control
492   Activities. Viruses 15.

493   19.   Mordecai EA, Cohen JM, Evans MV, Gudapati P, Johnson LR, Lippi CA, Miazgowicz K,
494   Murdock CC, Rohr JR, Ryan SJ, Savage V, Shocket MS, Stewart Ibarra A, Thomas MB,
495   Weikel DP, 2017. Detecting the impact of temperature on transmission of Zika, dengue,
496   and chikungunya using mechanistic models. PLoS Negl Trop Dis 11: e0005568.

497   20.   Huber JH, Childs ML, Caldwell JM, Mordecai EA, 2018. Seasonal temperature variation
498   influences climate suitability for dengue, chikungunya, and Zika transmission. PLoS Negl
499   Trop Dis 12: e0006451.

500   21.   Caldwell JM, LaBeaud AD, Lambin EF, Stewart-Ibarra AM, Ndenga BA, Mutuku FM,
501   Krystosik AR, Ayala EB, Anyamba A, Borbor-Cordova MJ, Damoah R, Grossi-Soyster
502   EN, Heras FH, Ngugi HN, Ryan SJ, Shah MM, Sippy R, Mordecai EA, 2021. Climate

predicts geographic and temporal variation in mosquito-borne disease dynamics on two
continents. Nat Commun 12: 1233.

22. Inziani M, Adungo F, Awando J, Kihoro R, Inoue S, Morita K, Obimbo E, Onyango F,
Mwau M, 2020. Seroprevalence of yellow fever, dengue, West Nile and chikungunya
viruses in children in Teso South Sub-County, Western Kenya. Int J Infect Dis 91: 104-
110.

23. Sabin AB, 1950. The dengue group of viruses and its family relationships. Bacteriol Rev
14: 225-32.

24. Sabin AB, 1952. Research on dengue during World War II. Am J Trop Med Hyg 1: 30-
50.

25. McCormack CP, Ghani AC, Ferguson NM, 2019. Fine-scale modelling finds that
breeding site fragmentation can reduce mosquito population persistence. Commun Biol
2: 273.

26. Nathan R, Klein E, Robledo-Arnuncio JJ, Revilla E, 2012. Dispersal kernels: review.
Clobert J, Baguette M, Benton TG, Bullock JM, eds. Dispersal Ecology and Evolution:
Oxford University Press, 0.

27. Marcantonio M, Reyes T, Barker CM, 2019. Quantifying *Aedes aegypti* dispersal in
space and time: a modeling approach. Ecosphere 10: e02977.

28. Moore TC, Brown HE, 2022. Estimating *Aedes aegypti* (Diptera: Culicidae) Flight
Distance: Meta-Data Analysis. J Med Entomol 59: 1164-1170.

29. Moran PAP, 1948. The Interpretation of Statistical Maps. Journal of the Royal Statistical
Society: Series B (Methodological) 10: 243-251.

30. Pfeiffer DU, Robinson TP, Stevenson M, Stevens KB, Rogers DJ, Clements ACA, 2008.
Spatial clustering of disease and global estimates of spatial clustering. Spatial Analysis
in Epidemiology: Oxford University Press, 0.

31. Wilder-Smith A, 2021. Dengue during the COVID-19 pandemic. J Travel Med 28.

32. Sasmono RT, Santoso MS, 2022. Movement dynamics: reduced dengue cases during
the COVID-19 pandemic. Lancet Infect Dis 22: 570-571.

33. Ngugi HN, Mutuku FM, Ndenga BA, Musunzaji PS, Mbakaya JO, Aswani P, Irungu LW,
Mukoko D, Vulule J, Kitron U, LaBeaud AD, 2017. Characterization and productivity
profiles of Aedes aegypti (L.) breeding habitats across rural and urban landscapes in
western and coastal Kenya. Parasit Vectors 10: 331.

Scattered	Centered	Clustered

Workplace ● Market ● Religious ● Recreational ● School ●