Risk factors for atopic and non-atopic asthma in school-aged children from high-, and low-and-middle-income countries

Charlotte E Rutter,[1] Harriet Mpairwe,[2] Camila A Figueiredo,[3] Mary Njoroge,[1] Steven Robertson,[4] Hajar Ali,[5] Collin Brooks,[5] Jeroen Douwes,[5] Philip J Cooper,[6,7,8] Martha Chico,[6] Natalia Romero-Sandoval,[7,9] Alvaro A Cruz,[10] Pius Tumwesige,[2] Mauricio L Barreto,[11] Neil Pearce,[1] Lucy Pembrey,[1] and the CAMERA Study Group

Department of Medical Statistics, London School of Hygiene and Tropical Medicine, London, UK
MRC/UVRI and LSHTM Uganda Research Unit, Entebbe, Uganda
Institute of Health Sciences, Federal University of Bahia, Salvador, Brazil
Clinical Trials Unit, London School of Hygiene and Tropical Medicine, London, UK
Centre for Public Health Research, Massey University, Wellington, New Zealand
Fundacion Ecuatoriana Para Investigacion en Salud, Quito, Ecuador
School of Medicine, Universidad Internacional del Ecuador, Quito, Ecuador
Institute of Infection and Immunity, St George's University of London, London, UK
Grups de Recerca d'Amèrica i Àfrica Llatines-GRAAL, Barcelona, Spain
Program for Asthma Control in Bahia (ProAR), Federal University of Bahia, Salvador, Brazil
The Centre for Data and Knowledge Integration for Health (CIDACS), Fiocruz, Bahia, Brazil

Corresponding author:

Dr Charlotte Rutter
Department of Medical Statistics
Faculty of Epidemiology and Population Health
London School of Hygiene and Tropical Medicine
Keppel Street
London
WC1E 7HT
Email: Charlotte.Rutter1@lshtm.ac.uk

Abstract

Background

It is well established that there are different asthma phenotypes, but whereas determinants of atopic asthma are well studied, little is known about non-atopic asthma. We compared risk factors for atopy, atopic asthma (AA) in atopics, and non-atopic asthma (NAA) in non-atopics, in children in a wide variety of countries.

Methods

Using four studies, across 23 countries, we assessed asthma status and atopy (skin prick tests) for children aged 6-17, plus risk factors from housing, heating, pets, family, diet, and air-quality categories. Using mixed effects logistic regression models we assessed 4 pathways to asthma: Pathway 1: comparing NAA with non-atopic non-asthma (risk factors for NAA); Pathway 2: comparing atopic non-asthma with non-atopic non-asthma (risk factors for atopy); Pathway 3: comparing AA with atopic non-asthma (risk factors for AA); Pathway 4: comparing AA with non-atopic non-asthma (risk factors for atopy and AA together).

Results

Our final sample of 32741 children comprised 67% non-atopic non-asthmatics, 15% atopic non-asthmatics, 8% with AA and 10% with NAA. Risk factors were similar for AA in atopic individuals and NAA in non-atopic individuals (Pearson's correlation r=0.81). In contrast, risk factors for atopy in non-asthmatics differed from those for AA in atopic individuals (r=-0.06).

Discussion

These findings indicate that although atopy increases the risk of asthma, the risk factors for subsequently developing asthma are generally the same in atopics and non-atopics. This raises important questions about the role of atopy in asthma, particularly whether it is an inherent part of the aetiological process or is coincidental.

Key messages

1. It is well established that there are different phenotypes of asthma but little is known about risk factors for non-atopic asthma.

2. Using a novel approach, we found that lifestyle and environmental risk factors for developing asthma are generally similar in atopic children and non-atopic children but the risk factors for atopy are quite different.

3. Our findings suggest that atopy and asthma may be coincidental in a large proportion of children who are defined as having atopic asthma. This has important implications for our understanding of the causes and mechanisms of different asthma phenotypes, and therefore prevention and treatment of asthma.

Introduction

Although the importance of intrinsic (or non-allergic or non-atopic) asthma has been recognised since the early 20th century[1], the focus of asthma research in the past 60 years has been predominantly on allergic mechanisms despite compelling evidence that at most half of asthma cases are attributable to atopy.[2,3]

As a consequence, little is known about the causes and pathophysiological mechanisms of non-atopic asthma (NAA), hampering the development of effective prevention and treatment options for a large proportion of asthmatics.

Atopic asthma (AA) is associated with T helper type 2-mediated airways inflammation (Th2), with atopy often identified (with varying results) by skin prick test (SPT) positivity to common allergens or serum allergen-specific IgE measurement (commonly associated with airways eosinophilia).[4] The mechanisms underlying NAA remain unclear but may involve neutrophilic inflammation, neural mechanisms, infection-related responses, or other currently unknown pathophysiological

pathways.[5-7] Different mechanisms may lead to different risk factors associated with disease. To make a valid comparison between NAA and AA we first need to tease apart the risk factors for asthma from the risk factors for atopy.

Figure 1 shows a framework for pathways potentially involved in the development of AA and NAA.[8] These are not often considered separately in studies of risk factors, instead most have done one of the following:[8] (i) compared all asthma cases with all non-asthmatic controls (a mix of all four pathways); (ii) adjusted for atopy (an average of the effects of pathway 1 and pathway 3); (iii) compared AA with all non-asthmatics (a mix of pathways 2-4); (iv) compared NAA with all non-asthmatics (none of the indicated pathways); or a few studies have (v) stratified on atopy (yielding separate estimates for pathways 1 and 3).

A 2014 review of the evidence for NAA risk factors found that, of 30 risk factors evaluated in 43 studies, only three (family history of allergic disease, household dampness/mould, and childhood lower respiratory tract infections) showed consistent associations with NAA. A lack of breastfeeding, or breastfeeding for a limited period, was less consistently associated with NAA. However, this review noted that most of these studies used inappropriate comparison groups.[9] There was also considerable methodological heterogeneity with different definitions of atopy and asthma, and different study designs, hampering valid comparisons.[9]

Here, using data from four studies: the International Study of Asthma and Allergies in Childhood (ISAAC) Phase II, the World Asthma Phenotypes Study (WASP), the Social Change, Asthma and Allergy in Latin America Study (SCAALA), and the Avon Longitudinal Study of Parents and Children (ALSPAC), we aimed to separately assess and compare risk factors for the four pathways (Figure 1), allowing the identification of modifiable factors that may increase/decrease the risk of NAA (pathway 1) and/or AA (pathways 2-4).

Methods

Data sources

The ISAAC Phase II cross-sectional questionnaire survey (with additional objective tests) commenced in 1998, targeting children aged 10-11, and included 30 centres from a mix of high-income and low-middle-income countries. Detailed methods have been published previously.[12]

The WASP cross-sectional case-control study was conducted in five centres in high-income (UK, NZ) and low-middle-income countries (Brazil, Ecuador, Uganda) between 2017 and 2020. Participants in Brazil, Ecuador and New Zealand were recruited from ongoing population-based cohort studies, with additional recruitment through the community (e.g., surveys in schools). In Uganda, participants were recruited from a larger case-control study of asthmatics and non-asthmatics identified through a cross-sectional survey in schools. Full details of the questionnaire and methods have been published previously.[4] The UK centre surveyed young adults (26-27 years) and has therefore been excluded from this analysis.

The SCAALA study involved a cross-sectional survey in Ecuador and a longitudinal study in Brazil, with detailed protocols published previously. In Ecuador, data were collected in 2006 on children aged 5-16 years in the Esmeraldas province.[13] In Brazil, data were collected in 2005 on children aged 4-11 years at baseline in the city of Salvador, Northeastern Brazil.[14]

The ALSPAC birth cohort study recruited pregnant women resident in Avon, UK, with expected delivery dates in 1991-1992. The study included regular surveys and tests throughout childhood, details of which have been reported previously.[15,16] Here we use SPT data at age 7 and surveys collected up to age 7.

From all sources, individuals with missing age or sex data, and those outside the target age group of 6-17 years were excluded from the analysis.

Variable definitions

Asthmatics were defined as having current asthma if they had wheeze symptoms and/or used asthma medications in the past 12 months. Non-asthmatics were defined as never having had asthma or asthma symptoms, determined by negative responses to questions on ever having wheeze symptoms or diagnosed asthma (ISAAC and SCAALA) or by negative responses to questions on recent wheeze symptoms across all surveys in ALSPAC. Participants qualifying for neither status (e.g., those who previously had asthma) were excluded from the analyses.

We defined atopy as SPT positivity (≥3mm wheal in ISAAC, WASP and SCAALA, ≥2mm wheal in ALSPAC) to at least one of a panel of five or more common allergens. Individuals testing positive for saline (negative control), negative for histamine (positive control), or with fewer than five allergens tested, were excluded. Similar SPT methods were used in each study, and allergens appropriate for the local environment of each centre were used (supplementary Table S2).

Study participants were categorised into four outcome groups: non-atopic non-asthmatic (X), atopic non-asthmatic (Y), atopic asthmatic (AA) and non-atopic asthmatic (NAA).

We included potential risk factors if they were available in ISAAC (the largest study) and at least one other study (details in supplementary Table S1). Risk factors included types of bedding, house condition, pets, diet, body mass index (BMI) and early life.

Statistical analyses

Four sets of models were used, representing each pathway in Figure 1.

Pathway 1: comparing NAA with non-atopic non-asthma (risk factors for NAA)

Pathway 2: comparing atopic non-asthma with non-atopic non-asthma (risk factors for atopy)

Pathway 3: comparing AA with atopic non-asthma (risk factors for AA)

Pathway 4: comparing AA with non-atopic non-asthma (risk factors for AA and atopy together)

For the first three pathways we examined which risk factors were associated with NAA (Figure 1, pathway 1), atopy (pathway 2), and AA in atopic children (pathway 3). Pathway 4 was more difficult to address as it is in addition to the two existing pathways, 2 and 3. It was assessed by estimating the overall association of non-atopic non-asthma (X) with AA (not adjusted or stratified on atopy) and comparing it with that predicted assuming no direct path, i.e., the Odds Ratio (OR) for pathway 2 multiplied by the OR for pathway 3. A difference between the two figures was considered to indicate a possible direct path (assuming the difference was not due to misclassification or residual confounding). In addition, we directly compared AA to NAA to see which factors differentiated them and how these compared to the risk factors for atopy alone.

Mixed-effects logistic regression models were fitted on the binary outcome representing each pathway, with a random intercept for centre and adjusting for age, sex, study, and country income level (LMIC/HIC). We first tested for evidence against a linear trend for age using the models with no risk factors by adding a quadratic term; the most appropriate form was then used for the models of each pathway. Individual models were fitted for each risk factor in turn, and results compared between pathways. The overall relationship between pathways was assessed using Pearson's correlation.

Models stratified by country income group were fitted, to test for interactions, and separate models for each study, to assess consistency.

Sensitivity analyses for the main non-stratified models were conducted with data from ISAAC and ALSPAC (due to data availability) to exclude borderline atopics. Two stricter atopy definitions were applied: any wheal of at least 5mm in the core allergens; 2 or more wheals of at least 4mm in the core allergens. Those originally classed as atopic but not meeting the new definition were excluded from that analysis. The definition for non-atopics remained the same. Results were compared to those using the standard atopy definition but limited to the two studies.

Data were analysed using Stata version 17.[17]

Results

From a total sample of 71916 participants (1205 from WASP, 54271 from ISAAC, 8266 from SCAALA and 8174 from ALSPAC), 39175 were excluded due to: missing symptom information (7273); not fitting case/control status (10754); no valid SPT result (20334); either missing demographics or out of the age range (814) (Table 1).

The final sample of 32741 comprised 66% non-atopic non-asthmatics, 15% atopic non-asthmatics, 8% AA and 10% NAA. The WASP sample had a higher proportion of asthmatics (75%) than the others (15-30%) due to its case-control design (Table 2).

Participants in WASP were on average the oldest but with a wide range (9-18 years). Those in ALSPAC were 7-8 years, ISAAC were mainly 10-11 years and SCAALA ranged from 6-16 years (supplementary Figures S1-S3). The WASP dataset was 55% female, SCAALA was 48% and ISAAC and ALSPAC were both 51%. In WASP 64% of the individuals were from LMIC and in ISAAC this was 32%. SCAALA participants were all from LMIC (Brazil and Ecuador) and ALSPAC were all from HIC (UK). Details of the final sample by country are in supplementary Table S3.

The prevalence of each risk factor by study/centre is presented in Table 3. Each risk factor was available for a different subset of individuals, ranging from 14060 for synthetic pillows to 31175 for parental allergic diseases. There was considerable variation between centres for some variables, particularly indoor heating questions as some centres in hot countries do not use heating at all.

Results from the mixed effects logistic regression, with random intercepts for centre and adjusted for study, sex, age and country income level (LMIC/HIC), but without including risk factors, showed evidence that the effect of age was not linear for pathway 1 (*P*=0.03) and for the AA/NAA comparison, but there was no evidence against linearity for pathways 2-4 (*P*=0.18, *P*=0.36, *P*=0.57

respectively). Thus, a quadratic age term was included in all models for pathway 1 and the AA/NAA comparison.

Figure 2 compares the estimated ORs of each risk factor between AA in already atopic individuals (pathway 3), and the risk of NAA in non-atopic individuals (pathway 1). The associations for the included risk factors were generally similar for AA and NAA, with a high positive correlation between the estimated effects (Pearson's correlation r=0.81).

Figure 3 compares risk factors for atopy (pathway 2) with risk factors for AA in atopic individuals (pathway 3) with no correlation between the two (r=-0.06).

We also compared the risk factors for atopy (pathway 2) with the risk factors for AA compared to NAA, as in both of these groups the difference is atopy. We found a moderate positive correlation (r=0.54) between these groups (supplementary Figure S4).

The individual results for each risk factor and outcome are shown in supplementary table S4, along with predictions of pathway 4, based on the results for pathway 2 and 3. The observed and the predicted values are similar so there is likely no direct pathway 4 (non-atopic to AA) over and above the two separate pathways. Results comparing pathways 1-3 by category of risk factor are shown in supplementary Figures S5-S10.

The strongest risk factor for asthma was parental history of allergic disease (asthma, eczema or hay fever) (NAA OR=2.45 [95% CI=2.24-2.69]; AA 2.05 [1.81-2.31]). Breastfeeding (ever) was associated with a lower risk of NAA (0.75 [0.67-0.85]). Premature birth (1.42 [1.19-1.69]) and low birthweight (1.25 [1.05-1.49]) were associated with an increased NAA risk.

Bedding-related risk factors included feather bedding (NAA 0.63 [0.52-0.76]; Atopy 0.87 [0.76-0.99]; AA 0.64 [0.52-0.79]) and feather pillows (NAA 0.80 [0.66-0.96]; Atopy 0.88 [0.77-1.01]; AA 0.64 [0.50-0.81]) which were both associated with a lower risk of asthma and atopy across all pathways.

Conversely synthetic bedding was associated with a higher risk (NAA 1.33 [1.16-1.53]; Atopy 1.18 [1.05-1.31]; AA 1.51 [1.28-1.79]).

Almost all indoor air-related factors including damp (NAA 1.64 [1.41-1.90]; Atopy 0.87 [0.75-1.00]; AA 1.51 [1.22-1.86]), mould (NAA 1.51 [1.29-1.75]; Atopy 0.83 [0.69-1.00]; AA 1.51 [1.20-1.91]) and maternal smoking (current: NAA 1.46 [1.32-1.62]; Atopy 0.84 [0.78-0.92]; AA 1.22 [1.07-1.40]; in first year: NAA 1.56 [1.40-1.74]; Atopy 0.87 [0.78-0.96]; AA 1.37 [1.18-1.60]) were associated with an increased risk of asthma (NAA and AA) and a slight protective effect on atopy. Current use of coal/coke heating was associated with an increased risk of developing AA among atopic individuals (1.75 [1.11-2.78]) but other heating factors did not show a strong association with asthma or atopy.

There was evidence that having pet birds in the first year of life was associated with higher risk of asthma (NAA 1.38 [1.13-1.68]; AA 1.47 [1.15-1.87]), and pet dogs in the first year of life a higher risk of AA 1.41 [1.15-1.72]. Many types of pet (both current and in first year of life) were associated with a lower risk of atopy (any pets 0.87 [0.79-0.95]; cats 0.81 [0.74-0.89]; dogs 0.90 [0.82-0.99]; cats in first year 0.83 [0.72-0.95]; dogs in first year 0.73 [0.63-0.84]). Farm animal contact showed no association with either asthma or atopy.

Obesity (NAA 1.60 [1.28-2.01]; AA 1.55 [1.10-2.16]) and being overweight (NAA 1.31 [1.14-1.50]; AA 1.24 [1.03-1.51]) were associated with higher risk of both forms of asthma but were not related to risk of atopy alone. Regular fruit consumption was associated with a lower risk of asthma and weakly with an increased risk of atopy (NAA 0.80 [0.70-0.93]; Atopy 1.14 [0.99-1.31]; AA 0.78 [0.63-0.97]).

Stratified by country affluence, there was still a positive correlation between risk factors for AA and NAA in both settings but some of the strongest risk factors were quite different by setting (Figure 4). The risk factors for atopy were unrelated to the risk factors for AA in both settings (Figure 5).

Notably, pet birds and dogs were strong risk factors for asthma in LMIC but not in HIC. Coal/coke heating was a risk factor for AA in HIC but not LMIC. The protective effect of feather bedding and

pillows was only found in HIC. Regular fast food was protective for AA (but not NAA) in LMIC only. Detailed results of these stratified models are in supplementary Table S5.

Results for the separate studies (supplementary Tables S6 to S9) showed some differences, though the WASP study alone was relatively small, resulting in wide confidence intervals.

The sensitivity analyses, with different atopy definitions, showed similar outcomes. Using only ISAAC and ALSPAC but with the standard atopy definition (n=26496), the correlation between risk factors for AA in already atopic individuals (pathway 3), and NAA in non-atopic individuals was lower but still quite strongly positive (r=0.68) and the correlation between risk factors for atopy (pathway 2) and AA in atopic individuals (pathway 3) was still very low (r=-0.08). The sample size was 23188 using wheal size >=5mm for any core allergen and 22243 using wheal size >=4mm for 2 or more core allergens. The correlation between risk factors for AA and NAA was slightly less strong (r=0.63 for both definitions). The risk factors for AA and atopy had weak negative correlations (r=-0.13 for wheal>=5mm and r=-0.17 for wheal>=4mm for 2+ allergens) (Supplementary Figures S11-S12 and Table S10).

Discussion

We used a novel approach to assess risk factors for the different pathways to atopic and non-atopic asthma, in a large and diverse population representing both HIC and LMIC.

Our findings from the available variables indicate that lifestyle and environmental risk factors are similar for NAA (in non-atopic children) and AA (in atopic children) but not for atopy. For example, dog contact in the first year of life showed a strong association for AA in atopics (OR=1.41), weaker evidence of effect for NAA in non-atopics (1.10) and a protective association with atopy (0.73); frequent fruit intake showed strong and consistent protective associations for asthma (NAA 0.80; AA 0.78) but a positive association with atopy (1.09); maternal smoking at various times in the child's

life was strongly associated with increased risk of asthma (NAA and AA) but reduced risk of atopy. The exception was parental history of allergic diseases which was a strong risk factor for both forms of asthma (NAA 2.45; AA 2.05) and atopy (1.38).

Although atopy (defined using SPT) is associated with increased risk of asthma, other risk factors are generally the same for AA and NAA. There may be important risk factors that were not included in this study, particularly for NAA which has not been as extensively studied, and the results may have been different if these "other" risk factors (e.g., air pollution, viral infections, stress) had been examined. Nevertheless, we have included a large number of commonly recognised and studied risk factors in our analyses, including most of the risk factors that have been identified as being associated with asthma in major studies such as ISAAC and GAN.[10,18] The fact that we have found these risk factors to have similar associations with both asthma in atopics (AA) and asthma in non-atopics (NAA) raises important questions about the role of atopy in asthma, particularly whether it is an inherent part of the aetiological process, or an exacerbating factor (or a mix of both), or is coincidental. Alternatively, NAA and AA could be separate phenotypes caused by different mechanisms for the initial onset of asthma but with similar exacerbating factors. However, this cannot be established from our cross-sectional studies as it is not possible to differentiate risk factors for initial onset from those for exacerbation of pre-existing asthma.

Risk factors in the animal and air-quality categories showed a particularly striking pattern of negative associations with atopy but positive associations with asthma (both AA and NAA) (Supplementary Figures S7 and S9). This is apparent for risk factors from early life as well as current, so is unlikely to be due to reverse causation or specific to exacerbation of symptoms.

In comparing our method of investigating different pathways to the other methods listed in the introduction, we found, as an example, that fruit intake (which was protective for NAA and AA but a risk factor for atopy) was protective when comparing all asthma to all non-asthma, also when adjusting for atopy, and when comparing NAA to all non-asthmatics. However this effect was not

seen when comparing AA to all non-asthmatics (as the effect was confounded by atopy). If the differences between risk factors for NAA and AA were greater than we observed, consideration of the separate pathways is even more important.

A strength of this study is the inclusion of data from 23 different countries, including both HICs and LMICs, as well as the resulting large sample size. Previous analyses of ISAAC data have generally found the risk factors for asthma to be similar in HICs and LMICs, providing for triangulation of findings,[19] and increasing the likelihood that the associations are causal. However, there are difficulties in ensuring the relevance of questions in varying settings e.g., heating questions in different climates. In addition, some risk factors may be differentially associated with socio-economic position (SEP), which we partially assessed by stratifying the results by country income level, but we do not have the information to assess differences within country.

The strongest risk factor for both types of asthma, and for atopy alone, was parental history of allergic disease, consistent with previously published findings.[9,20] Other strong risk factors for asthma (AA and NAA), but not atopy, were damp and mould in the home, which are already established risk factors for childhood asthma (both aetiology and exacerbation).[9] Obesity was a risk factor for both AA and NAA in HIC, which is consistent with literature,[9] but we did not see this in LMICs; whereas pet birds and dogs were risk factors for AA and NAA only in LMIC. These differences could indicate that associations were due to confounding, or that there were other uncaptured differences between settings e.g., bird contact could mean one pet budgie or a brood of chickens.

Other limitations of these analyses should be considered. The associations reported here will likely be subject to residual confounding as not all asthma risk factors are known or were recorded in all the included studies. Furthermore, the questionnaire was relatively simple, and most questions were binary. For some risk factors, reverse causation is a possibility, e.g., those with asthma (or family members) may have changed to synthetic bedding if feathers exacerbate symptoms. Recall bias could also be an issue, with parents of asthmatics more likely to recall information on known risk

factors than parents of non-asthmatics. There is evidence of a disassociation between SPT and IgE in LMIC where some individuals with negative SPT are IgE-positive to the same allergen.[21] In addition, the European Academy of Allergy and Clinical Immunology, in 2023, proposed a broader classification of allergy, including hypersensitivity reactions far beyond IgE-mediated phenomena (atopy).[22] Both these factors could indicate a higher proportion of asthma cases are in fact allergic (using the above definition) than that which we find here using SPT alone.

Our findings are important as they point towards similar mechanisms in AA and NAA, once the risk factors for atopy itself are removed. With SPT positivity defining atopy, there is no evidence that the risk factors differ between asthma phenotypes, even after using stricter definitions in sensitivity analyses. This suggests that atopy and asthma may be coincidental in a large proportion of children who are defined as having AA. To confirm or refute these findings, further work on identifying different phenotypes of asthma should be undertaken in a variety of populations, including using other markers and data sources (e.g., microbiome and epigenetics). IgE analysis could identify patterns in potential allergens and clinical tests could investigate the role of neurological triggers.

Ethics approval

Ethical approval was obtained from the London School of Hygiene & Tropical Medicine (LSHTM) ethics committee (ref: 26308)

Data availability

ISAAC data are publicly available at the UK Data Service archive http://discover.ukdataservice.ac.uk/catalogue?sn=8131. ALSPAC data can be accessed after application to the ALSPAC Executive Team; application instructions and data use agreements are

available at http://www.bristol.ac.uk/alspac/researchers/access/. The relevant data from the WASP study and SCAALA study will be shared on reasonable request to the corresponding author.

Acknowledgements

We thank all the participants, their parents, and the researchers involved in the WASP, ISAAC, SCAALA and ALSPAC studies for providing the data for this work to be completed.

Author contributions

N.P. conceived of and designed the study with support from C.B., H.M., M.L.B., P.J.C., J.D., C.A.F., A.A.C. and L.P.; C.E.R and S.R. were responsible for data cleaning and management; C.E.R. analysed the data; C.E.R, N.P. and L.P. wrote the first draft of the manuscript; all other authors contributed to revisions of the manuscript. Others also contributed to the planning and conduct of the study, and are listed below as part of the CAMERA Study Group.

Funding

The CAMERA project was supported by the European Research Council under ERC grants 668954, and 101020088. The SCAALA study was supported by the Wellcome Trust under grant 072405/Z/03/Z/WT. Work in Salvador was also supported by: Fundação de Amparo à Pesquisa do Estado da Bahia (FAPESB) under grants PNE0003/2014 and PNX0001/2014; Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Brasil (CAPES) under finance code 001; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) under grant 308521/2019 and CNPq/ CONFAP/FAPESB under grant 40314. Work in Wellington was supported by the Health Research

Council (HRC) of New Zealand and CB was supported by an HRC Sir Charles Hercus Fellowship. Work

in Ecuador was supported by the Universidad Internacional del Ecuador under grant EDM-INV-04-19.

Disclosures

MLB has received grants from the Wellcome Trust and the NIHR. AAC has received grants from NIH,

ERC and NIHR for related work, consulting fees from taking part in advisory boards for AstraZeneca,

GSK, Sanofi and Myralis, and honoraria payments for lectures from AstraZeneca, GSK and Sanofi; JD

has participated on the Board of the New Zealand Environmental Protection Authority, which is

independent and government-funded, and on the Board of the Health Research Council, both of

which are unrelated to this work; PJC has been supported by an NIHR grant; all other authors declare

no competing interests.

References

1. Lötvall J, Akdis CA, Bacharier LB, Bjermer L, Casale TB, Custovic A, Lemanske RF Jr, Wardlaw AJ,
    Wenzel SE, Greenberger PA. Asthma endotypes: a new approach to classification of disease
    entities within the asthma syndrome. J Allergy Clin Immunol. 2011 Feb;127(2):355-60. doi:
    10.1016/j.jaci.2010.11.037. PMID: 21281866.

2. Pembrey L, Brooks C, Mpairwe H, et al. Asthma inflammatory phenotypes on four continents:
    most asthma is non-eosinophilic. Int J Epidemiol. 2022 Aug 30:dyac173. doi:
    10.1093/ije/dyac173.

3. Pearce N, Pekkanen J, Beasley R. How much asthma is really attributable to atopy? Thorax.
    1999;54(3):268-72

4. Pembrey L, Barreto ML, Douwes J, et al. Understanding asthma phenotypes: the World Asthma
    Phenotypes (WASP) international collaboration. ERJ Open Res. 2018 Aug 22;4(3):00013-2018.
    doi: 10.1183/23120541.00013-2018. PMID: 30151371; PMCID: PMC6104297.

5. Gibson PG, Simpson JL, Saltos N. Heterogeneity of airway inflammation in persistent asthma :
    evidence of neutrophilic inflammation and increased sputum interleukin-8. Chest. 2001
    May;119(5):1329-36. doi: 10.1378/chest.119.5.1329. PMID: 11348936.

6. Ali H, Brooks C, Crane J, et al. Enhanced airway sensory nerve reactivity in non-eosinophilic asthma. BMJ Open Respir Res. 2021 Nov;8(1):e000974. doi: 10.1136/bmjresp-2021-000974. PMID: 34728474; PMCID: PMC8565536.

7. Figueiredo CA, Rodrigues LC, Alcantara-Neves NM, et al. Does IFN-γ play a role on the pathogenesis of non-atopic asthma in Latin America children? Allergy Asthma Clin Immunol. 2012 Dec 19;8(1):18. doi: 10.1186/1710-1492-8-18. PMID: 23253516; PMCID: PMC3564733.

8. Barreto ML, Cunha SS, Fiaccone R, Esquivel R, Amorim LD, Alvim S, Prado M, Cruz AA, Cooper PJ, Santos DN, Strina A, Alcantara-Neves N, Rodrigues LC. Poverty, dirt, infections and non-atopic wheezing in children from a Brazilian urban center. Respir Res. 2010 Dec 1;11(1):167. doi: 10.1186/1465-9921-11-167.

9. Strina A, Barreto ML, Cooper PJ, Rodrigues LC. Risk factors for non-atopic asthma/wheeze in children and adolescents: a systematic review. Emerg Themes Epidemiol. 2014;11:5

10. Silverwood RJ, Rutter CE, Mitchell EA, Asher MI, Garcia-Marcos L, Strachan DP, Pearce N; ISAAC Phase Three Study Group. Are environmental risk factors for current wheeze in the International Study of Asthma and Allergies in Childhood (ISAAC) phase three due to reverse causation? Clin Exp Allergy. 2019 Apr;49(4):430-441. doi: 10.1111/cea.13325. Epub 2019 Jan 23. PMID: 30508327; PMCID: PMC6487816.

11. Strachan D, Ait-Khaled N, Foliaki S, et al. Siblings, asthma, rhinoconjunctivitis and eczema: a worldwide perspective from the International Study of Asthma and Allergies in Childhood. Clinical and Experimental Allergy 2015; 45: 126-136.

12. Weiland SK, Björkstén B, Brunekreef B, et al. Phase II of the International Study of Asthma and Allergies in Childhood (ISAAC II): rationale and methods. Eur Respir J. 2004 Sep;24(3):406-12. doi: 10.1183/09031936.04.00090303. PMID: 15358699.

13. Cooper PJ, Chico ME, Vaca MG, et al. Risk factors for asthma and allergy associated with urban migration: background and methodology of a cross-sectional study in Afro-Ecuadorian school children in Northeastern Ecuador (Esmeraldas-SCAALA Study). BMC Pulm Med. 2006 Sep 13;6:24. doi: 10.1186/1471-2466-6-24. PMID: 16970809; PMCID: PMC1578586.Barreto 2006

14. Barreto ML, Cunha SS, Alcântara-Neves N, et al. Risk factors and immunological pathways for asthma and other allergic diseases in children: background and methodology of a longitudinal study in a large urban center in Northeastern Brazil (Salvador-SCAALA study). BMC Pulm Med. 2006 Jun 23;6:15. doi: 10.1186/1471-2466-6-15. PMID: 16796729; PMCID: PMC1559717.

15. Boyd A, Golding J, Macleod J, et al.: Cohort Profile: the 'children of the 90s'--the index offspring of the Avon Longitudinal Study of Parents and Children. Int J Epidemiol. 2013;42(1):111–27. 10.1093/ije/dys064.

16. Fraser A, Macdonald-Wallis C, Tilling K, et al.: Cohort profile: The Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. Int J Epidemiol. 2013;42(1):97–110. 10.1093/ije/dys066.

17. StataCorp. 2021. Stata Statistical Software: Release 17. College Station, TX: StataCorp LLC.

18. Rutter CE, Silverwood RJ, Asher MI, Ellwood P, Pearce N, Garcia-Marcos L, Strachan DP; ISAAC Phase Three Study Group. Comparison of individual-level and population-level risk factors for rhinoconjunctivitis, asthma, and eczema in the International Study of Asthma and Allergies in Childhood (ISAAC) Phase Three. World Allergy Organ J. 2020 Jul 2;13(6):100123. doi: 10.1016/j.waojou.2020.100123. PMID: 32642022; PMCID: PMC7334817.

19. Lawlor DA, Tilling K, Davey Smith G. Triangulation in aetiological epidemiology. International Journal of Epidemiology 2016;45:1866-86.

20. He, P., Moraes, T.J., Dai, D. et al. Early prediction of pediatric asthma in the Canadian Healthy Infant Longitudinal Development (CHILD) birth cohort using machine learning. Pediatr Res (2024). doi: 10.1038/s41390-023-02988-2.

21. Alcantara-Neves NM, Veiga RV, Ponte JC, da Cunha SS, Simões SM, Cruz ÁA, Yazdanbakhsh M, Matos SM, Silva TM, Figueiredo CA, Pontes-de-Carvalho LC, Rodrigues LC, Fiaccone RL, Cooper PJ, Barreto ML. Dissociation between skin test reactivity and anti-aeroallergen IgE: Determinants among urban Brazilian children. PLoS One. 2017 Mar 28;12(3):e0174089. doi: 10.1371/journal.pone.0174089. PMID: 28350867; PMCID: PMC5369757.

22. Jutel et al. Nomenclature of allergic diseases and hypersensitivity reactions: Adapted to modern needs: An EAACI position paper. Allergy. 2023 Nov;78(11):2851-2874.

CAMERA Study Group

Jessica Alchundia, Hospital de Especialidades, Portoviejo, Manabi Province, Ecuador; Hajar Ali, Centre for Public Health Research, Massey University, Wellington, New Zealand; Lucia Alonzo, School of Medicine, Universidad Internacional del Ecuador (UIDE), Quito, Ecuador; Candace Andrade,

Federal University of Bahia (UFBA), Salvador, Brazil; Mauricio L Barreto, The Centre for Data and Knowledge Integration for Health (CIDACS), Fiocruz, Bahia, Brazil; Collin Brooks, Centre for Public Health Research, Massey University, Wellington, New Zealand; Jeroen Burmanje, Centre for Public Health Research, Massey University, Wellington, New Zealand; Martha Chico, Fundacion Ecuatoriana Para Investigacion en Salud (FEPIS), Quito, Ecuador; Philip J Cooper, FEPIS, Quito, Ecuador, School of Medicine, UIDE, Quito, Ecuador, and Institute of Infection and Immunity, St George's University of London, London, UK; Alvaro A Cruz, Program for Asthma Control in Bahia (ProAR), UFBA, Salvador, Brazil; Auxiliadora Damiane Pereira Vieira da Costa, Universidade Federal de Alagoas, Maceió, Brazil; Donna Davoren, Department of Medical Statistics, London School of Hygiene and Tropical Medicine (LSHTM), UK; Jeroen Douwes, Centre for Public Health Research, Massey University, Wellington, New Zealand; Maria Fernanda Arias, School of Medicine, UIDE, Quito, Ecuador; Camila A Figueiredo, Institute of Health Sciences, UFBA, Salvador, Brazil; Luciano Gama da Silva Gomes, UFBA, Salvador, Brazil; Givaneide Lima, UFBA, Salvador, Brazil; Letícia Marques dos Santos, UFBA, Salvador, Brazil; Santiago Mena, School of Medicine, UIDE, Quito, Ecuador; Harriet Mpairwe, MRC/UVRI and LSHTM Uganda Research Unit, Entebbe, Uganda; Irene Nambuya, MRC/UVRI and LSHTM Uganda Research Unit, Entebbe, Uganda; Milly Namutebi, MRC/UVRI and LSHTM Uganda Research Unit, Entebbe, Uganda; Karla Nicole, UFBA, Salvador, Brazil; Mary Njoroge, Department of Medical Statistics, LSHTM, UK; Marble Nnaluwooza, MRC/UVRI and LSHTM Uganda Research Unit, Entebbe, Uganda; Genesis Palma, School of Medicine, UIDE, Quito, Ecuador; Neil Pearce, Department of Medical Statistics, LSHTM, UK; Lucy Pembrey, Department of Medical Statistics, LSHTM, UK; Gabriela Pimentel Pinheiro, Program for Asthma Control in Bahia (ProAR), UFBA, Salvador, Brazil; David E Rebellon Sanchez, Department of Medical Statistics, LSHTM, UK; Steven Robertson, Clinical Trials Unit, LSHTM, UK; Natalia Romero-Sandoval, School of Medicine, UIDE, Quito, Ecuador, and Grups de Recerca d'Amèrica i Àfrica Llatines (GRAAL), Barcelona, Spain; Charlotte E Rutter, Department of Medical Statistics, LSHTM, UK; Bianca Sampaio Dotto Fiuza, UFBA, Salvador, Brazil; Karla Solorzano, Hospital de Especialidades, Portoviejo, Manabi Province, Ecuador; Josephine

Tumusiime, MRC/UVRI and LSHTM Uganda Research Unit, Entebbe, Uganda; Pius Tumwesige,

MRC/UVRI and LSHTM Uganda Research Unit, Entebbe, Uganda; Cinthia Vila Nova Santana, Program

for Asthma Control in Bahia (ProAR), UFBA, Salvador, Brazil;

Figures and Tables

Pathway   Investigation
1: X → NAA   Risk factors for non-atopic asthma in non-atopic children
2: X → Y   Risk factors for atopy
3: Y → AA   Risk factors for atopic asthma among atopic children
4: X → AA   Risk factors for atopic asthma among non-atopic children

Figure 1: Pathways of possible asthma causation (Adapted from Barreto et al, 2010)

Figure 2: Estimated Odds ratios (ORs) of risk factors for atopic asthma compared to non-atopic asthma, excluding atopy risk (Pathways 3 v 1)

Figure 3: Estimated Odds ratios (ORs) of risk factors for asthma (in atopic individuals) compared to atopy (Pathways 3 v 2)

Figure 4: Estimated Odds ratios (ORs) of risk factors for atopic asthma compared to non-atopic asthma, excluding atopy risk (Pathways 3 v 1) stratified by country affluence

Figure 5: Estimated Odds ratios (ORs) of risk factors for asthma (in atopic individuals) compared to atopy (Pathways 3 v 2) stratified by country affluence

Table 1:   Exclusions

	WASP	ISAAC	SCAALA	ALSPAC	Total
Starting sample	1205	54 271	8266	8174[a]	71 916
Missing symptoms	0	5183	0	2090	7273
Not case/control	94	6469	2265	1926	10 754
No atopy data	22	18 789	231	1292	20 334
Missing sex/age	0	147	0	26	173
Not age 6-17	166	27	448	0	641
Final sample	923	23 656	5322	2840	32 741

[a]individuals who completed the age 7 questionnaire, without second twin; ISAAC=International Study of Asthma and Allergies in Childhood; WASP=World Asthma Phenotypes; SCAALA=Social Changes, Asthma and Allergy in Latin America; ALSPAC= Avon Longitudinal Study of Parents and Children;

Table 2:   Distribution of main outcome by study

Study	Non-Atopic Non-Asthmatic (X) (%)	Atopic Non-Asthmatic (Y) (%)	Atopic Asthmatic (AA) (%)	Non-Atopic Asthmatic (NAA) (%)	Total
WASP	171 (19%)	66 (7%)	422 (46%)	264 (29%)	923
ISAAC	16 239 (69%)	3922 (17%)	1584 (7%)	1911 (8%)	23 656
SCAALA	3860 (73%)	560 (11%)	203 (4%)	699 (13%)	5322
ALSPAC	1650 (58%)	322 (11%)	433 (15%)	435 (15%)	2840
TOTAL	21 920 (67%)	4870 (15%)	2642 (8%)	3309 (10%)	32 741

ISAAC=International Study of Asthma and Allergies in Childhood; WASP=World Asthma Phenotypes; SCAALA=Social Changes, Asthma and Allergy in Latin America; ALSPAC= Avon Longitudinal Study of Parents and Children

Table 3: **Prevalence of potential risk factors by centre / study**

Category	Covariable	n	ISAAC Multiple	WASP Brazil	WASP Ecuador	WASP New Zealand	WASP Uganda	WASP total	SCAALA Brazil	SCAALA Ecuador	SCAALA total	ALSPAC UK	Overall Total
Confounder	Male sex	32741	49%	38%	59%	49%	29%	45%	53%	51%	52%	49%	49%
House/ bedding	Feather bedding	14156	18%	4%	18%	37%	.	25%	.	.	.	.	18%
	Synthetic bedding	14155	43%	46%	4%	55%	.	35%	.	.	.	.	43%
	Blankets	15560	44%	14%	56%	47%	.	45%	22%	.	22%	.	43%
	Feather pillow	14780	17%	6%	55%	24%	.	32%	.	.	.	.	17%
	Synthetic pillow	14060	39%	71%	1%	72%	.	47%	.	.	.	.	40%
	Double glazing	20831	55%	.	.	.	.	.	.	.	.	81%	58%
	Double glazing in 1st year	20194	45%	.	.	.	.	.	.	.	.	59%	47%
	Urban	23452	92%	.	.	.	.	.	100%	36%	45%	88%	81%
	Urban in 1st year	16953	89%	.	.	.	.	.	.	.	.	93%	89%
Heating/ cooking	Heating	16979	68%	0%	0%	100%	.	48%	.	.	.	.	67%
	Wood heating	18386	14%	0%	0%	40%	0%	14%	.	.	.	.	14%
	Coal/coke heating	17603	4%	0%	0%	1%	0%	0%	.	.	.	.	4%
	Gas heating	16261	28%	.	.	34%	.	34%	.	.	.	.	28%
	Heating in 1st year	15523	68%	.	.	.	.	.	.	.	.	.	68%
	Wood heating in 1st year	16894	20%	.	.	.	.	.	.	.	.	.	20%
	Coal/coke heating in 1st year	16330	6%	.	.	.	.	.	.	.	.	.	6%
	Cooking electric	20728	45%	1%	15%	.	.	11%	0%	.	0%	.	43%
	Cooking gas	29549	54%	100%	95%	56%	.	77%	99%	96%	97%	61%	63%
	Cooking gas in 1st year	18776	19%	.	.	.	.	.	99%	.	99%	56%	27%
Animals	Pets	20342	43%	51%	63%	77%	30%	58%	37%	60%	57%	70%	51%
	Cats	30397	15%	19%	36%	48%	10%	31%	10%	37%	33%	30%	20%
	Dogs	30379	14%	32%	43%	41%	22%	36%	25%	44%	42%	20%	20%
	Birds	24130	14%	.	.	.	.	.	.	.	.	6%	13%
	Cat in 1st year	21091	9%	.	.	.	.	.	7%	.	7%	29%	12%
	Dog in 1st year	21091	8%	.	.	.	.	.	18%	.	18%	19%	10%
	Bird in 1st year	20399	7%	.	.	.	.	.	.	.	.	5%	7%
	Farm animals	22052	7%	.	77%	.	42%	60%	.	22%	22%	.	11%
Diet/ weight	Meat weekly	26025	88%	.	100%	92%	.	95%	61%	88%	85%	.	87%
	Seafood weekly	26507	57%	.	90%	54%	.	69%	10%	92%	81%	.	62%
	Green vegetables weekly	24540	83%	.	.	.	.	.	31%	73%	68%	78%	79%

Category	Variable	N											
	Fruit weekly	25019	90%	69%	96%	98%	.	93%	50%	93%	87%	96%	90%
	Fast food weekly	17878	25%	94%	11%	47%	.	42%	10%	6%	7%	.	20%
	Fruit juice weekly	18553	69%	.	.	.	.	.	55%	.	55%	69%	69%
	Overweight	14615	25%	21%	26%	23%	18%	22%	27%	8%	10%	14%	18%
	Obese	14615	7%	7%	6%	5%	3%	5%	13%	1%	3%	3%	4%
Family/ birth	Older siblings	29536	52%	.	.	.	.	.	62%	76%	75%	52%	56%
	Younger siblings	29470	49%	.	.	.	.	.	42%	77%	72%	52%	53%
	More than 1 sibling	29528	40%	.	.	.	.	.	47%	87%	82%	34%	47%
	Ever breastfed	29588	78%	.	.	.	.	.	96%	98%	97%	80%	82%
	Low birthweight	20176	18%	.	.	.	.	.	.	.	.	4%	16%
	Premature	16875	16%	.	.	.	.	.	8%	.	8%	9%	14%
	Parental allergic diseases	31175	36%	.	.	.	.	.	31%	41%	40%	74%	39%
	Maternal education	18189	79%	.	.	.	.	.	44%	32%	34%	100%	70%
	Paternal education	18389	80%	.	.	.	.	.	100%	38%	46%	100%	74%
Air-quality	Any smokers	29371	45%	.	.	.	.	.	21%	44%	41%	29%	43%
	Maternal smoking	30869	25%	.	.	.	.	.	11%	13%	13%	16%	22%
	Maternal smoking in 1st year	27364	19%	.	.	.	.	.	10%	11%	11%	17%	17%
	Maternal smoking during pregnancy	28545	12%	.	.	.	.	.	10%	10%	10%	16%	12%
	Damp	17670	14%	38%	38%	15%	30%	28%	.	.	.	90%	20%
	Mould	15021	9%	40%	22%	54%	35%	39%	.	.	.	52%	15%
	Damp or mould	15512	18%	50%	42%	55%	48%	49%	60%	.	60%	92%	28%
	Damp in 1st year	17763	15%	.	.	.	.	.	.	.	.	49%	20%
	Mould in 1st year	14599	10%	.	.	.	.	.	.	.	.	25%	13%
	Air conditioning	14910	29%	2%	.	3%	.	3%	1%	.	1%	.	27%

ISAAC=International Study of Asthma and Allergies in Childhood; WASP=World Asthma Phenotypes; SCAALA=Social Changes, Asthma and Allergy in Latin America; ALSPAC= Avon Longitudinal Study of Parents and Children