Understanding the impact of the COVID-19 pandemic and its control measures on women and children: A Zimbabwe case study

Tinotenda Taruvinga[1,2]; Rudo S. Chingono[1], Ioana D. Olaru[1,3]; Kenneth Masiye[4]; Claudius Madanhire[1,4]; Sharon Munhenzva[1,4]; Sibusisiwe Sibanda[1,4]; Lyton Mafuva[1,4], Natasha O'Sullivan[5.6]; Abdinasir Y. Osman[5,7,8]; Kevin Deane[5]; Tsitsi Brandson[1]; Manes Munyanyi[9]; Annamercy C. Makoni[10]; Solwayo Ngwenya[11,12]; Karren Webb[4], Theonevus T. Chinyanga[4]; Rashida A. Ferrand[1,3]; Justin Dixon[1,2], Katharina Kranzer[1,3,13], David McCoy[5,14]

1. The Health Research Unit Zimbabwe (THRU-ZIM), Biomedical Research & Training Institute (BRTI), Harare, Zimbabwe
2. Department of Global Health and Development, London School of Hygiene and Tropical Medicine (LSHTM), London, UK
3. Clinical Research Department, London School of Hygiene and Tropical Medicine, London, UK
4. Organisation for Public Health Interventions and Development (OPHID) Harare, Zimbabwe
5. Queen Mary University of London (QMUL), London, UK

6.   MRC Epidemiology Unit, University of Cambridge, Cambridge, UK

7.   Pathobiology and Population Science, Royal Veterinary College, London, UK

8.   Ministry of Health, Mogadishu, Somalia

9.   Ministry of Health and Childcare, Harare, Zimbabwe

10.   Harare City Council, Harare, Zimbabwe

11.   Mpilo Central Hospital, Bulawayo, Zimbabwe

12.   National University of Science and Technology (NUST), Bulawayo, Zimbabwe

13.   Division of Infectious Diseases and Tropical Medicine, LMU University Hospital, LMU

28   Munich,

14.   United Nations University, Shibuya, Tokyo, Japan

Corresponding author

Tinotenda Taruvinga

The Health Research Unit Zimbabwe (THRU ZIM)

Biomedical Research and Training Institute

10, Seagrave Road

Harare, Zimbabwe

E-Mail: Tinotenda.Taruvinga1@lshtm.ac.uk

Authors' contributions

TT, JD, KK, and DM conceptualized the study. IO, TB, TT, MM, and KM cleaned and curated

the data. TT, IO, RSC, AOY, and NO with support from KD, KK, JD, and DM analysed the data.

DM, KW, and RAF acquired the funds for the study. SS, LM, CM, and SM supported by RSC,

AYO, NO, AM, and TT performed the qualitative data collection. Study coordination and

supervision of field teams were done by RSC, TT, SN, ACM and TC. TT wrote the first draft of

the manuscript with feedback from JD, KK, and DM. All authors provided feedback on the

manuscript and read and approved the final version of the manuscript.

ABSTRACT

COVID-19 presented countries with unprecedented health policy challenges. For low-income

countries in particular, policymakers had to contend with both the direct threats posed by

COVID-19 as well as the social, educational, and economic harms associated with lockdown

and other infection prevention and control measures. We present a holistic and

contextualised case study of the direct and indirect impacts of COVID-19 on women and

children, with some assessment of their uneven distribution across socio-economic, age and

gender groups. We used different types of primary and secondary data from multiple sources

to produce a holistic descriptive analysis. Primary data included: qualitative data obtained

from 28 in-depth interviews of key informants, six focus group discussions; and 40 household

interviews. We also extracted data from government reports and announcements, the District

Health Information Software version 2 (DHIS2), newspaper articles and social media, as well

as from published research articles. Our findings show that the direct and indirect adverse

impacts of COVID-19 were compounded by many years of severe political economic

challenges, and consequent deterioration of the healthcare system. The indirect effects of the

pandemic had the most severe impacts on the poorest segment of society and widened age

and gender inequalities. The pandemic and its accompanying infection prevention and control

measures negatively affected health service delivery and uptake. The management of COVID-

19 presented enormous challenges to policymakers and public health specialists. These

included managing the greatest tension between direct and indirect harms; short-term and

long-term effects; and the unequal distribution of harms across different segments of society.

INTRODUCTION

The SARS-CoV-2 (COVID-19) pandemic presented all countries with major health challenges

[1,2]. These included ensuring adequate epidemiological surveillance of COVID-19,

implementing infection prevention and control (IPC) measures, providing care for both

COVID-19 and non-COVID-19 patients, and procuring including delivering COVID-19 vaccines.

Low-and-middle-income countries (LMICs) were also faced with having to implement

draconian lockdown measures without having the resources to mitigate their social,

educational, and economic harms [3,4], and the disruption caused to normal healthcare

provision [5,6].

Importantly, countries had to consider how the harms associated with the pandemic would

be unevenly distributed [1,7,8]. For example, wealthier households would be better able to

withstand the psychological and emotional stresses of lockdown compared to poorer

households [6,9]. Additionally, there were reasons to think that women and girls would

experience lockdown measures differently from men and boys [8]. Finally, the early

observation that infection fatality rates varied substantially across age groups raised

challenges to policymakers about balancing the different needs of children and adults [10,11].

We conducted research aimed at presenting a holistic and contextualised case study of the

direct and indirect impacts of COVID-19 on women and children in Zimbabwe, with some

assessment of their uneven distribution across socio-economic, age and gender groups. As is

typical of case studies seeking to understand complex phenomena, we used different types

of primary and secondary data from multiple sources [12] to produce an integrated

descriptive analysis of the COVID-19 epidemic, the policy responses to it, and their impacts

on maternal, child and women's health.

METHODS

To produce a chronological narrative account of policies related to COVID-19 from February

2020 to August 2021, we collected data from government policy documents and

announcements, newspaper articles, social media, and the Oxford COVID-19 response tracker

(OxCGRT), a project that collated data on COVID-19 control measures from countries across

the world [13].

To present an account of the COVID-19 pandemic, we obtained official data on the number

of COVID-19 tests done, confirmed cases and deaths from the daily situational reports and

the official Twitter (now X) handle of the Ministry of Health and Child Care (MoHCC). Given

the limited disease surveillance and low testing rates, we also drew on data from other

research to assess COVID-19 transmission in Zimbabwe.

To describe the effects and impacts of COVID-19 and its control measures on maternal and

child healthcare, we extracted routinely collected health service data on antenatal care (ANC),

growth monitoring of children and childhood vaccinations from the District Health

Information Software version 2 (DHIS2) for the period January 2016 to August 2021 in

Bulawayo and Harare metropolitan provinces. Harare the capital city and Bulawayo the

second biggest city in the country have estimated populations of 1,896,134 and 676,650,

respectively, and together accounted for about 30% of the country's recorded COVID-19 cases

[14]. Primary health services in the two cities are primarily administered by the city council

whilst tertiary and quaternary hospitals are administered by the central government. We

selected 20 public primary health clinics in Harare and 18 in Bulawayo to study trends in

healthcare provision and utilisation.

We selected two child and two maternal services: growth monitoring (GM), child vaccinations,

antenatal care (ANC) and HIV testing of pregnant women. Several data items are routinely

collected to monitor child vaccination services indicators, and we chose to analyse trends in

the number of children aged 9-12 months who have received their final measles, mumps,

rubella (MMR) vaccine which is captured in the DHIS2 as 'primary course complete' (PCC). For

maternal health services, we chose to analyse trends in the number of pregnant women

attending a 4th ANC visit (a proxy indicator of adequate ANC visits according to the WHO) as

well as the number of women who received an HIV test at their first ANC visit. However, data

for the latter was only available from 2019 onwards.

Changes over time in the four indicators of healthcare utilisation are shown graphically for

each clinic, stratified by city, with trend lines using loess smoothing and 95% confidence

intervals. Clinic-level changes in healthcare utilisation are shown as ratios computed by

dividing monthly counts by the clinic-specific monthly average for the period starting January

2016 up to the end of 2019. The ratios were colour-coded using shades of orange to mark

counts that were lower than the pre-COVID-19 average and shades of green for higher counts.

Some clinics had missing values for some months, especially in 2021, and these were coloured

in grey. These missing values include a combination of instances where there was no service

uptake (e.g. the clinic was open but there was no service utilisation, or the clinic was shut)

and when there was service uptake, but data were not recorded or uploaded onto the DHIS2.

We also collected primary qualitative data between April 2021 and December 2021 in Harare

and Bulawayo. We conducted a total of 28 in-depth interviews (IDIs) of key informants drawn

from community-based organisations (CBOs) (n=5), city health managers (n=5 Harare, n=3

Bulawayo), national programme managers and policymakers (n=5) and healthcare workers

(n=3 Harare, n=7 Bulawayo). We further conducted two focus group discussions (FGDs) with

healthcare workers from selected primary health facilities in Harare (n=12) and Bulawayo

(n=11), and four FGDs of members of the public (two in Harare with 9 and 10 participants,

and two in Bulawayo with 11 and 13 participants). In addition, we conducted 40 household

interviews, evenly distributed across three economic strata (n=20 Harare and n=20

Bulawayo). Topic guides were used for all the qualitative interviews. Written informed

consent was obtained from all the participants in the study.

Finally, we drew data from selected published literature to describe the historical and socio-

economic context of the pandemic and add further information about the impact of COVID-

19 on maternal and child health in Zimbabwe. We identified this literature from the database

of PubMed and Google Scholar for papers using search terms 'maternal and child health',

'COVID-19' and 'Africa' or 'Zimbabwe' up to the period January 2023.

FINDINGS

1. The context of Zimbabwe's COVID-19 epidemic

Zimbabwe is a low-income landlocked country of about 15 million people, with 42% of the population under the age of 15 [14]. In 2019, its Human Development Index was only 0.571, ranking 150[th] out of 189 countries [15]. Approximately 71% of the population lives below the international poverty line of $1.90 per day [16] leading to poor health indicators. Over 80% of income earners operate in the informal sector [17] which contributes to over 60% of gross domestic product (GDP) [18,19]. One in four of the urban population (about 1.25 million people) live in settlements with poor water and sanitation infrastructure as defined by UNICEF [20]. However, Zimbabwe has a highly literate population with 94% of men and women aged 15-49 years able to read, and 91% of all children of primary school age attending school [19]. Moreover, Zimbabwe's Gender Development Index (GDI) was 0.952 in 2018, suggesting relatively good gender equity, with secondary school attendance among girls being higher than boys [21].

The country had experienced severe political and economic challenges for the three decades before the COVID-19 pandemic. An unsustainable external debt burden and a structural adjustment programme resulted in a severe contraction of public expenditure in the 1990s which weakened most public services [22]. The turn of the millennium saw the country enter a period of protracted political turmoil and the imposition of targeted sanctions by several European countries [22,23] further undermined the economy. Just before the pandemic, Zimbabwe's GDP had declined by 11.3% in 2019 compared to 2018 and continued to contract in 2020 [24].

Although Zimbabwe's health system was once considered among the best in Africa, [18,25] it

deteriorated due to reductions in public spending, the out-migration of health professionals,

mismanagement, and industrial action [26,27]. By 2021 the doctor and nurse persons ratios

had deteriorated to 1:12,000 and 0.39:1,000 respectively, far below the World Health

Organization's (WHO) recommended developing countries' minimum threshold density of

2.28 doctors, nurses and midwives per 1,000 persons [28]. Today, there are high levels of out-

of-pocket payments and donor dependency [29]. The maternal mortality ratio was 458 per

100,000 live births [30] and the child mortality rate was 55 per 1000 live births [30] in 2020

[31,32].

On top of this chronic deterioration of the health system, there were a series of acute crises

in the run-up to the pandemic (see Figure 1). This included doctors and nurses engaging in

intermittent industrial action [33], typhoid and cholera outbreaks in 2018, a major natural

disaster (cyclone Idai) [34] in 2019, an acute worsening of inflation from April 2020 causing

shortages of fuel and water. The precarious state of the health system at the time of the

pandemic was noted in our qualitative data.

"Most of our stream mates have gone to the UK where there are better jobs. Others

are in the process of applying; we are all trying to leave for greener pastures (laughs).

Those of us who stay, we are mostly working out of duty...people are disgruntled, and

this can be sometimes seen through our attitudes and, umm, the strikes." (Nurse, HCW

FGD, Bulawayo)*.*

"Unfortunately, in Bulawayo when COVID-19 was declared and it hit us, we were facing

water challenges and other related things." (Doctor, KII, Bulawayo).

Figure 1: Timeline of key events concerning the period between January 2018 to December 2021

2. **Evolution of the COVID-19 epidemic and control measures**

(i) January to May 2020

Following WHO's call to prepare for the containment of COVID-19 in January 2020, the government of Zimbabwe quickly started developing a National Preparedness and Response (NPR) Plan and established an Interagency Coordination on Health and Epidemic Preparedness and Response Task Force [35]. Screening for SARS-CoV-2 infections at major entry points was quickly implemented for all incoming individuals. A series of inter-ministerial meetings on COVID-19 also took place in February and March with the involvement of the private sector, civil society, academia, professional associations, technical and donor partners and the media.

The NPR Plan was informed by a Preparedness and Response Plan for Pandemic Influenza A H1N1 formulated in 2009, recent simulation exercises, a comprehensive review of the 2018 cholera outbreak and a rapid assessment of the state of preparedness at major points of entry

into the country and major health facilities in February. The Plan was aligned with WHO's

COVID-19 Strategic Preparedness and Response Plan and finalised in March.

On March 17th, a state of national emergency was declared prior to any detectable surge in

SARS-CoV-2 infections. In fact, the first case of COVID-19 was only reported on 20th March in

an individual who had recently returned from the United Kingdom. Two days later, two

additional cases, imported from the United States, were documented. The first recorded

fatality occurred on 23rd March [36], the same day the Civil Protection (Declaration of State

of Disaster: Rural and Urban Areas of Zimbabwe) (COVID-19) Notice was issued [37].

A Public Health Act (COVID-19 Containment and Treatment) was also enacted on the 24th of

March, mandating compulsory testing of individuals suspected of infection and the

quarantine of those who tested positive [38]. Subsequently, a Statutory Instrument (SI) was

introduced imposing a 21-day lockdown from the 28th of March during which time public

gatherings of more than two people were prohibited. While 'essential services' were allowed

to operate, retail outlets were only allowed to open during specific hours, with mandatory

screening and hand sanitising at entrances. Although initially, schools and universities were

slated to remain open until the end of the term, a later announcement mandated the closure

of all educational institutions from 24th March 2020. The sale of alcohol was banned, and the

stockpiling of medical supplies and food was explicitly prohibited with strict penalties for any

violations.

During this phase, the government also designated certain health facilities as quarantine and

isolation centres, adopted and adapted WHO's COVID-19 case management guidelines [39],

organised training for health workers and upgraded several government and private

hospitals. Surveillance teams were created to facilitate case investigation and contact tracing.

Over 4,000 health sector posts were unfrozen, and an additional 200 new medical posts were

created and funded through a reallocation of funds [40] from other departments and from

the proceeds of the 2% Intermediated Money Transfer Tax which is normally used for social

protection and capital development projects. However, despite efforts to strengthen the

health system's capacity to respond to COVID-19, the shortage of PPE and inadequate pay led

many healthcare workers to feel vulnerable and undervalued [41] and precipitated a fresh

round of industrial action beginning on 25th March with a nurses' strike [42].

"An association representing the City of Harare nurses went around to check if there

was adequate PPE…After they observed that there was nothing that's when they called

for the nursing staff or for the healthcare worker to stop going to work, tools down

guys there is no PPE until proper adequate PPE is in place. So, you find that most nurses

of the city of Harare went home and stayed…The situation was also aggravated by the

fact that we were being poorly remunerated, we were not getting our salaries on

time." (Nurse, HCW FGD, Harare).

Lockdown measures were eventually extended for a further four weeks to 17th May, although

from 3rd May, the formal commercial and industrial sectors were allowed to operate between

08:00 and 15:00, provided IPC measures were in place. Throughout this period, numerous SIs

were gazetted to amend or extend the initial lockdown measures. For example, on April 3rd,

facemasks were made compulsory in public and transport services were required to check the

temperatures of boarding passengers and implement disinfection protocols on all vehicles.

While   initially,   residents   were   allowed   to   return   home   from   abroad   provided   they

quarantined at designated facilities, on May 4th all international travel was banned. Land

border crossings at Beitbridge (bordering South Africa) and Plumtree (bordering Botswana)

were also closed, and internal travel by road was restricted, with exemption letters required

at road checks.

Throughout this period, reported case numbers remained low. When lockdown was lifted on

May 17th, Zimbabwe had only recorded 46 cases and six deaths. However, because public

sector testing capacity was limited and the cost of private testing was unaffordable for most

people, the true number of cases during this period is unknown. Further, there were no

formal or informal reports of any surge in hospital admissions and mortality during the period.

(ii) June 2020 – November 2020

After the first lockdown was lifted, the number of recorded cases remained low for about

eight weeks. Despite this, the government imposed a two-week lockdown from June 20 to

July 4. This followed extensive civil society protests following the arrest of the health minister

on allegations of graft. It is widely believed that the lockdown was used to suppress political

opposition rather than the epidemic [43,44], with many protestors arrested for contravening

the new COVID-19 regulations.

However, when case numbers started rising, the government introduced a third lockdown on

July 24th. Night-time curfews (from 6 pm to 6 am) and travel restrictions were implemented.

However, essential services continued to operate as usual and non-essential businesses and

low-risk sports were allowed to operate from 9 am to 3 pm. Restaurants, hotels and other

tourism services were allowed to operate at 50% capacity; and schools were kept open until

July 30[th] to allow the completion of national examinations. Although the rise in case numbers

was partly due to increased testing capacity [45] there was some evidence of a real increase

in the incidence of infection with rising COVID-19 mortality and hospitalisations [46] including

some high-profile fatalities.

The third lockdown continued until September but with a progressive easing of restrictions

as case numbers started falling. Government quarantine centres were phased out in August

and replaced with self-isolation in private premises, and a phased opening of schools

commenced on the 8[th] of September and was completed in November. The sale of alcohol for

consumption off-premises was allowed on 30 September, and international airports started

operating again in October. While initially, only Zimbabwean residents were allowed to return

(provided they tested negative and self-quarantined at home for 14 days), entry was later

allowed for foreign travellers. In December land ports of entry were opened.

(iii) December 2020 – April 2021

In late November 2020, a new wave of cases emerged, driven by the Beta variant [47]. This

coincided with the reopening of schools, where some large outbreaks were reported [48]. As

numbers rose, and with several high-profile deaths including that of five cabinet ministers

[49], the government introduced a fourth national lockdown for 30-days on January 3, which

was extended to February 16 and then again to February 28. The government closed schools,

prohibited gatherings, reintroduced curfew hours and stopped intercity travel. Mask-wearing,

hand sanitisation/washing, and temperature checks remained mandatory in public. As the

country started recording a reduced number of cases, lockdown measures were eased in

March, inter-city travel resumed, and schools re-opened in a phased manner. April saw a

relaxation of most remaining IPC measures although international travel to and from

countries such as India that had reported the Delta variant were not permitted while

travellers from other countries were required to present a negative COVID-19 test result and

self-quarantine at home or in designated quarantine centres for 10 days [50].

During this period the country also began a COVID-19 vaccination program with a target of

vaccinating 10 million people by December 2021 [51]. The first phase prioritised frontline

healthcare workers, the elderly, and those with chronic medical conditions [52]. The second

phase which began in March 2021 extended coverage to uniformed forces, all civil servants

and those offering essential services in the private sector. The last phase was introduced in

July and extended coverage initially to everyone over 18 years, then to anyone over 16 years

and finally to anyone over 12 years [53].

(iv) May 2021 – September 2021: Third wave and fourth lockdown

In the middle of May, the new (Delta) variant was detected in the Kwekwe district, Midlands

province. A new SI was introduced to allow for a localised lockdown of the Kwekwe district

and then in two (Hurungwe and Kariba) other districts [54]. Eventually on June 27th, due to

rising cases, a fifth full lockdown was implemented across the whole country. However,

despite the increase in cases, the government came under pressure from parents and opened

schools in the first week of August 2021. This was followed on August 10 by the reopening of

some social activities with, for example, vaccinated congregants being allowed to attend

church services.

Meanwhile, by the end of September, 3,051,371 first vaccine doses and 2,211,880 second

vaccine doses had been administered nationally translating to national coverage of 35.7% and

25.8% respectively [55]. At about the same time the Global Fund pledged USD 75-150 million

[56] to be used to mitigate the impact of the COVID-19 pandemic on HIV, tuberculosis, and

malaria. From September onwards, the country gradually moved into a post-pandemic phase

even though there was no official declaration of the pandemic being over. However, the

government emphasised more on 'living with the new normal'.

3. Policy implementation

According to our key informants, a notable feature of the response to COVID-19 was that

policies were made and implemented in a centralised and top-down manner with little

bottom-up and contextualised input.

"There was centralisation of decision making and policy formulation …. everything was

moved to the centre …and ours was just to implement. Then we had a situation where

there was a take-over … of local authority institutions by central government. It had

its downside and upside. But the formulation of that policy, unfortunately, we didn't

input in." (Doctor, KII, Bulawayo).

"Let's not take that one size fits all or… maybe some African countries are doing this,

you just follow suit and do this …. We should design our measures to suit our people,

to suit our needs and to suit our country and our resources." (Nurse in charge, IDI,

Harare).

"I still insist that we must not have a lockdown that stretches from Zambezi to

Limpopo, that is being the same. Let's modify it and say we are now in Gweru. How do

we make this lockdown work? When we get to Chirundu, there it must not look like the

one in Chiredzi. There must be that distinction. I feel that with the localised input and

the modifications we make, the lockdowns will be more effective. Such that it will allow

us to even open the economy in one part of the country whilst the other part is under

lockdown." (Doctor, KII, Bulawayo).

Informants also noted that public health communication (mainly conducted through the

Zimbabwean broadcasting cooperation's television and radio channels) was not fully effective

and that misinformation and disinformation on various social media platforms contributed to

poor compliance with COVID-19 control measures, despite rapid response teams being

established to respond swiftly to rumours and false information, albeit limited capacities to

deal with all the rumours due to shortage of human and material resources. Others indicated

that many people felt overloaded with conflicting information [41].

"What was our challenge was people didn't have information. So, you would just see

like a uniformed person come to you to hit you and you don't have information,

especially at the beginning. People were just being hit for COVID, but if you would have

educated people … "this is COVID, people are affected this way, we need to prevent

using these methods" … I think we are an educated nation that can actually follow

through and take instructions" (CBO representative, KII, Bulawayo).

"We communicated, we talked, we tried to send messages, but I don't think it was

effective because people had their own expectations, and they had their own other

sources of information. And they did not trust the figures that were coming out of the

Ministry of Health and situational reports eeh however, localised or frequent, people

still did not feel that they, they were in as much danger as it was being portrayed."

(Doctor, KII, Bulawayo).

Another theme that emerged from our primary data was the impracticality and impossibility

of implementing social distancing measures, especially in high-density and overcrowded

informal settlements with inadequate water and sanitation infrastructure [57]. Other

measures such as the use of hand sanitisers were unaffordable for many. According to one

study in Harare and Mashonaland East, between 18% and 53% of healthcare workers

reported a lack of soap, water, and masks across the period from June 2020 to the middle of

2021 [58]. Other studies similarly described how communities could not comply with social

distancing measures because of the need to secure food, water, and income [41].

"We expect the government not to implement water cuts and to improve water supply

during such a time because we are having a difficult time, social distance is not possible

[in the queues], we are at risk of contracting COVID-19" (Participant, household IDI,

Harare).

"It was impossible to comply with the social distancing measure, at a time when there

were food shortages and one had to stand in long queues. I still remember there was

a time I went to a queue to get mealie-meal at Malbereign. The soldiers had to come

and space out the people." (Participant, household IDI, Harare).

Another cause for poor compliance with lockdown measures was the inability to monitor and

enforce travel restrictions. Despite the police and army being deployed, resources were

insufficient to enforce quarantine and ensure adherence to travel restrictions. For example,

despite major points of entry being manned by armed security, many people were able to

enter the country illegally without being quarantined [59].

The relative ineffectiveness of lockdown measures may also be deduced from seroprevalence

studies which found that by the end of March 2021, a high proportion of the population had

been infected. In one study of randomly selected households from three high-density

communities in Harare, seroprevalence was found to have risen from 19% in November-

December 2020 to 53% in Feb-April 2021 [59], suggesting high levels of community

transmission during the second wave. A survey of healthcare workers as early as June 2020

reported that 39.1% had a history of COVID-19 symptoms and a SARS-CoV-2 seroprevalence

of 8.9% [60].

A systematic review of seroprevalence studies in Africa between January 2020 and December

2021 also estimated that seroprevalence in Southern Africa was 56.1% (95% CI 44.6% - 66.9%)

by the third quarter of 2021, with lower rates in rural areas and high heterogeneity between

countries [61]. Of note is a study from Malawi that found seroprevalence increasing from 11%

in December 2020 to 65% in April 2021 [61]. A similar increase in seroprevalence was

observed in a study from Harare conducted between November 2020 and April 2021 [59].

The reported data also indicated a less severe disease profile in Africa with more

asymptomatic cases compared to other parts of the world [62].

Another feature of the policy response to COVID-19 was the government's efforts to mitigate

the social and economic harms of lockdown. Crucially, in May 2020 a COVID-19 Economic

Recovery and Stimulus Package worth about 9% of GDP was announced and included income

support for all vulnerable groups (US$5 monthly for everyone in a vulnerable household) and

fiscal support to key sectors of the economy such as manufacturing, agriculture, mining, and

tourism [63]. The government also put in temporary measures to defer rent and mortgage

payments during lockdown. However, the income support payments were not fully

implemented or implemented for long enough. Dzawanda citing Buckle and Mpofu reported

that the government's promise of a once-off allowance for vulnerable people was only

partially delivered and that four months into the lockdown, only 202,000 of the 1 million

eligible households had received the allowance [18]. Similarly, a promised compensation

scheme for civil servants who lost their lives while on duty was only partially delivered,

although all civil servants were given a US$75 COVID-19 monthly allowance for over 36

months in addition to their salaries.

"I think maybe what probably might have lacked…are the necessary support systems

to make sure that we take care of the most vulnerable in society, for example, if we

say there is no public transport that is going to be operating, we should then be able

to provide transport for those that want to seek health services." (NGO representative,

KII, Harare).

"For an economy that is largely informal, if we then say we are closing everything else

all the markets non-informal business and everything else, we should then put in place

a system where we can then support those in the informal sector so that at the end of

the day, they don't uhm starve." (CBO representative, KII, Bulawayo).

4. Impact of COVID-19 control measures

This section looks at the pandemic's impact on maternal and child health (MCH) services

provision and utilisation; interpersonal and household relations; social and economic well-

being; and child education and well-being.

(i) Maternal and child healthcare

Our analysis of routinely collected data from July 2016 to July 2021 in Harare and Bulawayo

found evidence of a reduction in healthcare utilisation. Figure 2 presents data on the number

of fourth antenatal care visits by pregnant women and the number of HIV tests conducted

among pregnant women at their first antenatal visit.

The data show a reduction in the number of ANC fourth visits in both provinces, with a steeper

decline in Harare compared to Bulawayo. However, there was no observable reduction in the

number of HIV tests on women attending antenatal care. Key informants had explained that

the HIV programme which is largely donor-funded andrelatively well resourced was better

able to maintain their pre-COVID-19 level of service. In one study conducted in Harare, the

number of pregnant women accessing HIV services was shown to have stayed the same

except for a slight decline during the first lockdown [5].

Figure 3 presents data on the number of GM visits by children under-5 years and the number

of child PCCs. The data show that the number of GM visits decreased in both Harare and

461 Bulawayo after the first COVID-19 case, and that in between lockdowns, GM visits rebounded

462 to some extent in Bulawayo but not in Harare. The number of PCCs decreased substantially

463 in Harare from March 2020 onwards but not in Bulawayo.

Figure 2: Source: Zimbabwe District Health Information Software Version 2 maternal

indicators

Figure 3: Source: Zimbabwe District Health Information Software Version 2 child indicators

data

Other studies also report that healthcare utilisation declined in 2020 and 2021 compared to

earlier years [64,65]. One study from Harare found that the number of people registered for

TB treatment had decreased by 34% and the number of individuals tested for HIV had

dropped by 63% in 2020 and 2021 compared to the period between 2016 and 2019 [66]. TB

treatment completion was found to have decreased from 81% to 70%, while treatment of

people living with HIV with antiretroviral therapy dropped from 96% to 92% [66,67]. In a study

at Mpilo Hospital in Bulawayo, although the mean number of monthly deliveries reduced

from 747 in the first quarter of 2020 to 681 in the second quarter, the overall reduction was

not substantial [68].

Our qualitative data pointed to several reasons for the reduction in healthcare utilisation. In

some places, clinics were closed because staff went off sick because of COVID-19 [69], or

because of industrial action. Some informants explained that healthcare utilisation also

dropped because health facilities were perceived as "hotspots" for COVID-19 transmission,

especially those that had been repurposed for COVID-19 services.

"People are not coming as they used to because staff tested positive and the

community heard about it, and so maybe they are fearing that maybe when they come

here, they might eh contract COVID-19" (Nurse, IDI, Harare).

"I have a young sister who was pregnant at the time… she had to go to Murehwa

because we thought it was safer than here in Harare, …" (Participant, Household IDI,

Harare).

Police roadblocks and checkpoints were highlighted as barriers to healthcare due to fear and

unwillingness to disclose one's health status. Healthcare workers reported finding it difficult

to access public transport because they were perceived to be a risk to others, especially whilst

wearing a uniform.

"You find someone could not access the hospital easily especially someone from the

rural areas … there was no transport for them to come. Even those who are in town

they couldn't access the hospital there was no transport… some mothers could not take

their babies for review even to the baby clinics, though it was said that the baby card

was also a pass for them." (Nurse, IDI, Harare).

"Then people started to have difficulties in attending health centres, funerals, no

hospital visits, this is the period from July to August. Whereby we are saying even HIV

status disclosure on a roadblock. People were now forced to disclose… So, this was a

forced disclosure…" (NGO representative, KII, Bulawayo).

"And other people ended up not even attempting to go …. we have other people who

are on ARVs whom we know that have defaulted because they don't want to be

harassed because you gave them a lift and you get there, they are asked to produce a

card by the police, at times they would want to hide their status and not produce that

card so end up not going to the hospital." (Nurse, HCW FGD, Harare).

Health services were especially disrupted by operating times being reduced from 07:30-17:30

to 08:00-14:00 and by the capping of the number of patients seen per day. In some instances,

operating hours were reduced even more because healthcare workers struggled to get to

work or had to leave early to comply with the curfew.

"Yes, at the clinic, they would serve a certain number of people per day, so that people

don't get crowded there. So, if they reach the number of people that they wanted per

day, some would be returned home and be advised to come the next day ..."

(Participant, Household IDI, Harare).

"Our staff couldn't get transport to work, no one was willing to give them a lift as

people considered them to be a high-risk group…We also had 6 staff members who

tested positive and the news spread in the local communities and people were avoiding

this facility for fear of getting the virus…" (Nurse in charge, IDI, Harare).

The pandemic diverted resources and attention away from MCH services. Informants

reported that in some clinics, GM was completely suspended to allow for prioritisation of

other health services. Stockouts particularly affected immunisation and family planning

services [70].

"Then in those findings, it was noticed that there was growth monitoring reduction.

Then we also go to maternity health services whereby during the COVID era, we are

saying only ANC bookings were done. No subsequent visits were done, limited

monitoring of BP checks. If you remember we were no longer doing these BP checks,

no physical examinations were done. Then om HIV and AIDS testing, the numbers

started to reduce, viral load collection we stopped doing it, CD4 count was stopped,

the updating of green books was also stopped." (Nurse, IDI, Bulawayo).

Access to healthcare was also restricted by some facilities implementing a policy of patients,

including pregnant women, showing a negative SARS-CoV-2 test before being admitted. This

led to patients being turned away from hospitals and ambulances refusing to transport

patients:

"Pregnant mothers were requested to have results for COVID-19 first before admission

to a facility and the test price was very high some of the mothers could not afford it…

and even the ambulance services would request you to produce results first before they

carry you. If you didn't have the results, they would test you at a fee. I remember it was

USD 60 for the COVID-19 test before you get into an ambulance." (Nurse, HCW FGD,

Harare).

Some healthcare workers believed that an increase in the number of home deliveries without

a skilled birth attendant resulted in an increased number of largely unreported maternal and

neonatal deaths.

"I don't have statistics, but as I was on night duty, you would see that some people did

not come for their regular reviews, some would come already complicated which if we

had seen them on time, we could have prevented those complications." (Nurse in

charge, IDI, Bulawayo).

"I'm not sure in terms of statistics of women who could have been unfortunate to die

during childbirth because of lack of access to health care, we don't really have access

to such information, but we know that happened." (CBO representative, KII,

Bulawayo).

"I know plenty who needed maternity services but could not get them…in fact, I know

someone who even died. They wanted to give birth…but they were supposed to get

COVID-19 tests before accessing health services…that process was costly and trying to

find money took long and she died before she could even be given medical attention."

(Participant, household IDI, Harare).

Other studies reported similar findings of women and children struggling to access MCH

services during the pandemic [10,68]. In Nyanga, Manicaland province, Nyashanu *et al* (2021)

describe transport problems, roadblocks, shortage of medication and personal protective

equipment, and lack of routine healthcare services affected routine health access such as

antiretroviral therapy [41]. Chimhunya *et al* (2021) viewed strikes by healthcare workers at

Sally Mugabe Hospital in Harare as an indirect consequence of COVID-19 which caused a

decline in neonatal admissions [65].

(ii) Inter-personal and household relations

Another theme explored in our qualitative research was the impact of lockdown on

interpersonal and domestic relationships against a backdrop of increasing stress and mental

illness due to prolonged periods of home confinement and increasing levels of poverty and

economic insecurity. Our findings indicated that gender-based violence (GBV) had increased.

"But with men, I would not say it's domestic violence, I would have said I think men

suffered emotional trauma because obviously when your woman is sitting at home,

you are not providing as much as you used to, they would not be violent, but they

would go through more of emotional trauma." (Participant, community FGD, Harare).

"It could have been there before COVID-19 but with the lockdown restrictions the GBV

got worse. So, those things are now like permanent scars in the community." (CBO

representative, KII, Bulawayo).

"We also saw an upsurge of gender-based violence cases, some of them were reported

through the normal channels, through the police and also health facilities but we also

see an upsurge of some of these cases that were reported in the media" (NGO

representative, KII, Harare).

"Yes. You would notice that where we would expect that maybe the children, the girls

are safe within the home, they were not even safe…We got reports of cases of rape by

family members… Mobility restriction to survivors to access services then resulted in

some cases of rape being reported late because people didn't know." (CBO

representative, KII, Bulawayo).

These findings are supported by other studies. A cross-sectional online survey conducted in

May 2020 among 507 Zimbabwean adults found signs and symptoms of generalised anxiety

disorder present in almost half (40%) of the participants [71]. A qualitative study among rural

women in Nyanga concluded that COVID-19 exacerbated domestic violence through the

effects of financial stress, hunger, frustrations and spending more time together and that

child marriages increased [72]. Others observed that as families spent more time in close

contact while experiencing economic and job losses, women were exposed to more violence

[73]. According to the Zimbabwe Gender Commission, domestic violence more than doubled

[74] and some security personnel abused their authority by requesting favours from women

in exchange for permission to sell their wares amidst lockdown restrictions [72].

Unfortunately, we were not able to obtain official data on reports of GBV to the police. To

date, there is no official account of a rise in domestic violence in Zimbabwe. But even if such

data showed a rise in the incidence of GBV, this would likely be an under-representation of

the true picture because lockdown measures had created barriers to survivors of GBV making

reports to the police or accessing services.

(iii) Social and economic well-being

Our primary research confirmed severe negative impacts on household income due to

reduced economic activity and rising unemployment, a situation that was accentuated by the

rising price of medicines, food, and other commodities [75]. One consequence was a

reduction in both the number of meals and the quantity of food, disproportionately affecting

those from the lowest socioeconomic strata, as well as women and children.

"On the issue of livelihoods, many people lost their jobs due to the closure of

companies, most companies closed, even those that have since opened they are only

taking a limited number. Most people were retrenched and came to just sit at home;

this affected food security". (Participant, Community FGD, Bulawayo).

"Then there was near starvation, we were not prepared for such a lockdown, it just

emerged from nowhere, we were no longer able to travel, we were not able to buy

enough food, some of us had no money, some of us were vendors we expected the day-

to-day income", (FGD participant, Bulawayo).

"Livelihoods were affected in a way such that in families there was a shortage of food

due to lack of money as some of the breadwinners became jobless…. Basic

commodities were in short supply especially mealie meal and cooking oil." (Nurse,

HCW FGD, Harare).

"Before the lockdown period, we would eat porridge in the morning, tea later, then

*have sadza [*Zimbabwe staple food made from maize meal flour*] for lunch. But because*

of lockdown we would eat once per day, or sometimes not find anything to eat at all."

(Participant, household IDI, Bulawayo).

According to the World Bank, the pandemic added 1.3 million to the global number of people

defined as extremely poor [76]. A cross-sectional online survey of 507 adults conducted at the

beginning of the COVID-19 pandemic in May 2020 found that lockdown had increased food

prices (95%) and decreased availability of nutritious foods (64%) [71]. The prevalence of

wasting increased from 3.6% in 2019 to 4.5% in 2020 [20]. Other qualitative studies confirmed

such negative impacts and noted a disproportionate negative impact on women whose

caregiving roles had increased during lockdown [18,72].

(iv) Education and child wellbeing

School closure meant that many children received little to no formal school education during

the COVID-19 lockdowns. Feelings of stress and anxiety about the lack of learning, especially

for those in exam years, were expressed by many participants. Many informants also

described how school and college/university closures widened existing educational

inequalities between private versus public institutions and urban and rural settings because

of the differential abilities to access teaching provided through online platforms and

television or radio broadcasts [72].

"The government introduced in some schools online learning which does not apply to

a rural child. And even if it's urban, electricity, or issues of data, that's a challenge as

well. So, there are selected few students that can access school online depending on

the background of the child…Then, there are radio programs that were scheduled and

shared to all the schools...That was a good initiative, but again the issue of how many

have radios, how many have electricity" (CBO representative, KII, Bulawayo).

"The closing of schools resulted in a big divide between the poor and the rich, so the

students who could afford extra lessons learnt and those who could not afford did not

and the results were devastating" (NGO representative, KII, Harare).

The long period of school closures, especially during the first lockdown, was also said to have

increased smoking, use of alcohol and illicit drugs and sexual behaviour, as well as

participation in so-called 'gang behaviour'.

"Now we have a surge of drug abuse, school dropouts and the like. To be honest crimes

have increased." (CBO representative, KII, Bulawayo).

"Since children were not going to school, they would spend time sitting at bridges,

smoking marijuana, yet if they were in school, they could have been busy with their

homework, but at that time they had no homework. The girls also were all over and

ended up being impregnated." (Participant, household IDI, Harare).

"We work with children who are disadvantaged as a charity organization, but we

noticed that during the lockdown, there were lots of unwanted pregnancies among

teenagers and also alcohol abuse among the boys" (Participant, household IDI,

Harare).

We were unable to obtain routinely collected raw data on school attendance before and

during the COVID-19 period. However, a 2022 Ministry of Primary and Secondary Education

report provided some details on school enrolments, dropouts, transition, completion and

examination pass rates over five years (2017-2021).[73] Gross Enrolment Ratio (GER) is a

measure of participation in the education system and defined as the number of students

enrolled at the beginning of the school year in specific levels of education, regardless of age,

expressed as a percentage of the official eligible school-age population of a given level of

education. Transition conveys information on the degree of access or transition from one

cycle or level of education to a higher one, namely from Grade 7 to Form 1 and from Form 4

to Form 5. Completion rates indicate the proportion of persons completing a given level of

education.

Primary education has two levels: infant (early child development (ECD) A/B and Grade 1/2)

and junior (Grade 3-7) [77]. Transition into secondary school is after sitting for the Grade 7

national examination. The secondary education has also two levels, which are lower

secondary (Form 1-4) and upper secondary (Form 5-6). Transition to upper secondary is only

after attaining good grades at competitive national examinations. National ordinary level ('O'

level) exams are completed after Form 4 and national advance ('A' level) exams are a

requirement for university admission. While the total number of pupils enrolled in ECD and

primary school have steadily increased from 2017-2021, the number of new entrants into

Grade 1 decreased by more than 30,000 between 2020 and 2021. The number enrolled in

higher secondary school also decreased by 14,941 (-14.7%) between 2020 and 2021. GER

decreased across primary, lower secondary and higher secondary in 2020 and 2021 compared

to previous years (Figure 4). Most other indicators also worsened in 2020 and 2021 compared

to years 2017-2019 (which had seen a steady improvement year after year) including

increased school dropouts, lower transition rates from Grade 7 to Form 1 and Form 4 to Form

6, lower completion rates across primary and secondary school and lower pass rates across

all national exams. Worsening indicators seem to have affected boys and girls similarly.

Figure 4: School indicators between 2017-2021

A: Number of children enrolled, new entrants and drops out across education levels

693   **B: Gross enrolment ratio and transition rates across education levels**

695   **C: Completion and pass rates across education levels**

DISCUSSION

The COVID-19 pandemic occurred in Zimbabwe at a time when the country had experienced decades of economic decline and a chronic deterioration of its health system. Around seven out of every ten people were living below the international poverty line and health status indicators were among the poorest in Sub-Saharan Africa [16]. Furthermore, in the years immediately preceding the COVID-19 pandemic, the health system experienced intermittent industrial action by nurses and doctors, a series of natural disasters, and outbreaks of cholera, typhoid and measles [78]. Although the government increased its budgets for healthcare and unfroze healthcare worker posts to help cope with the pandemic [40], the fragility of the health system also made it likely that lockdown measures would only further reduce both the supply and demand for healthcare [79].

Given this background, Zimbabwe faced considerable challenges when COVID-19 emerged. As with all countries, policymakers had to contend with a lack of knowledge about the natural history, virulence and transmissibility of the virus [80]. The high rates of hospitalisation and mortality observed in China, Italy, and other countries as well as WHO's declaration of a Public Health Emergency of International Concern on January 30th, 2020, led many countries to implement stringent lockdown measures before they experienced any upsurge in COVID-19 cases. Underpinning these decisions was the belief amongst public health specialists and policymakers that any harms produced through the imposition of lockdown measures would be less than the harms produced by the virus itself.

Early estimates of the infection fatality rate were such that the enforcement of lockdown, including the cessation of important social and economic activities, was felt to be justified.

However, from the middle of 2020, differences in the incidence of cases and deaths were

being observed across the world [62]. The relatively low incidence of cases and deaths in parts

of Africa was especially notable [62] leading to several hypotheses as to what might be the

cause. Among them was the low prevalence of risk factors such as obesity, low infection rate,

effective mitigation measures, youthful age structure, favourable warm weather, and

preexisting immunity from previous coronavirus infection. In Zimbabwe, by the end of 2020,

only 13,867 cases and 363 deaths had been officially recorded and by the end of 2021, the

number of deaths attributed to COVID-19 was only 5,004 [81]. While there would have been

many unrecorded deaths due to COVID-19, there was no period when hospitals were overrun

by COVID-19 patients or when morgues struggled to cope with a sudden rise in deaths as

experienced by many other countries. Furthermore, seroprevalence studies indicate that

substantial transmission had occurred by the end of 2020 [59], indicating that the relatively

low mortality rates were possibly not due to effective prevention of transmission.

The importance of reliable, timely and context-specific data to manage infectious diseases

outbreaks and epidemics cannot be understated. In Zimbabwe, the gaps in the health

information system meant that a timely, accurate and detailed account of the epidemic and

its direct impacts were unavailable. Throughout the pandemic, policymakers and public

health specialists had to make educated guesses as to how the epidemic was evolving and

what impact it was having. The not-so-perfect health information systems in the country also

meant that policymakers and public health specialists could not adequately assess the

harmful and sometimes devastating impact of the lockdown measures. Even now, estimates

of the size and the causes of the increase in all-cause mortality since the pandemic are

unclear.

A key question emerging from this research is whether Zimbabwe adopted the right set of

measures to effectively manage both the direct and indirect threats to health that were posed

by COVID-19, including the harms associated with lockdown. Here one must be wary of the

benefit of hindsight and the lack of reliable and timely data available to policymakers and

public health experts at the time. However, given the pre-existing and high levels of poverty

and food insecurity before the pandemic [17], the economic stagnation, reduced household

income and rising prices for basic commodities caused by the lockdowns were inevitably going

to be impactful. The inability of many households to generate an adequate income during

lockdown had drastic and detrimental effects on access to food and basic amenities and was

compounded by mounting loneliness, distress, and mental illness, fuelling a degradation of

social relations within households and an increase in interpersonal violence and antisocial

behaviour. Although the government created a support fund for poorer households, the

budget was inadequate, and many eligible households did not receive the promised support.

Studies from other LMICs have shown similar findings [6,8,82] and many countries are

currently still experiencing the long-term consequences of the economic contraction and

increased levels of indebtedness that occurred in 2020 and 2021 [83].

Similarly, a negative impact on an already fragile, under-resourced and under-utilised health

system was inevitable as were the significant and long-term negative effects on students and

their families due to the prolonged closure of education facilities. The disproportionate

impact on children is of particular importance. While COVID-19 was considerably much less

harmful to children, the converse is true for the effects of the lockdown measures, including

the cessation or reduction of essential child healthcare services; the heightened vulnerability

of infants and young children to reductions in food intake; and the cessation of school. Thus,

many children were deprived of essential care and services as a result of measures that mostly

benefited adults.

A further important point is that for large segments of the population, lockdown measures

were not feasible. Many informants explained how safe and adequate social distancing was

impractical in high-density areas and informal settlements. If true, for many households,

prolonged quarantine and social isolation were arguably unjustified, especially considering

the harmful effects of lockdown measures described above. Moreover, lockdown measures

of the kind experienced with COVID-19 involve extraordinary curtailments on civil liberties

that should only be implemented in extreme and valid circumstances. The coercive powers

claimed by governments across the world in the name of public safety may not only be abused

during a pandemic but also used to establish permanent systems and mechanisms of

surveillance and social control. Indeed, lockdown measures were used in Zimbabwe to curtail

political protest, and there were also accounts of security sector actors abusing their power

to commit corruption and sexual harassment to the population.

Firm conclusions about the full impact of COVID-19 and its associated communicable disease

control measures are not possible due to the lack of reliable and complete data. Crucially, any

future strengthening of the health information system must include a focus not just on the

future ability to conduct infectious diseases surveillance and control, but also on the ability

to avoid or mitigate negative social, economic, and educational impacts. Nonetheless, plenty

of data and evidence exist to suggest that the indirect impacts of COVID-19 were at least as

harmful if not greater than its direct impacts especially so for women and children.

Conflicts of interest

SN was one of the health experts working on the ground who called for the lockdowns as he

had first-hand information on the healthcare preparedness to face such a new novel

coronavirus.

Funding statement

This work was supported through the Queen Mary University of London (QMUL) with funding

from the United Kingdom Research Institute (UKRI) under reference number GCRF_NF391

and implemented through the Organisation of Public Health Interventions and Development

(OPHID) Zimbabwe. The views expressed do not necessarily reflect the policies of the

respective organisations. The funders had no role in study design, data collection and analysis,

the decision to publish, or manuscript preparation.

Tinotenda Taruvinga is supported by the Fogarty International Centre of the National

Institutes of Health (NIM; Bethesda, Maryland, MD, USA) under Award Number *D43*

TW009539. The content therein is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institutes of Health.

Acknowledgements

The study would like to acknowledge the support we got from the Zimbabwe research

authorities, government departments and local authorities. In addition, the study also

acknowledges Professor John Metcalfe, Thokozile Masha, Jane Mususa, and the OPHID and

BRTI team led by Dr Shungu Munyati, Thomas Godfrey-Faussett who helped with valuable

reference materials on education and most importantly the participants who spared their

time to share this valuable experience. AYO, OD, is a member of and acknowledges support

from, the Pan-African Network for Rapid Research, Response, Relief and Preparedness for

Infectious Disease Epidemics, funded by the European and Developing Countries Clinical Trials

Partnership, under the EU Horizon 2020 Framework Programme for Research and Innovation.

References

1. Zweig SA, Zapf AJ, Xu H, Li Q, Agarwal S, Labrique AB, et al. Impact of Public Health and
Social Measures on the COVID-19 Pandemic in the United States and Other Countries:
Descriptive Analysis. JMIR Public Health Surveill. 2021;7: e27917. doi:10.2196/27917

2. Haider N, Osman AY, Gadzekpo A, Akipede GO, Asogun D, Ansumana R, et al.
Lockdown measures in response to COVID-19 in nine sub-Saharan African countries.
BMJ Glob Health. 2020;5: e003319. doi:10.1136/bmjgh-2020-003319

3. Rosenberg J, Strauss I, Isaacs G. COVID-19 impact on SADC labour markets: Evidence
from high-frequency data and other sources. Afr Dev Rev Rev Afr Dev. 2021;33: S177–
S193. doi:10.1111/1467-8268.12528

4. ECLAC. The effects of the coronavirus disease (COVID-19) pandemic on international
trade and logistics. 2020; 7.

5. Chappell E, Chimwaza A, Manika N, Wedderburn CJ, Nenguke ZM, Gannon H, et al.
Impact of the COVID-19 pandemic on the provision and uptake of services for the
prevention of mother-to-child transmission of HIV in Zimbabwe. PLOS Glob Public
Health. 2023;3: e0002296. doi:10.1371/journal.pgph.0002296

6. Kotlar B, Gerson EM, Petrillo S, Langer A, Tiemeier H. The impact of the COVID-19
pandemic on maternal and perinatal health: a scoping review. Reprod Health. 2021;18:
10. doi:10.1186/s12978-021-01070-6

7. Chan JF-W, Yuan S, Kok K-H, To KK-W, Chu H, Yang J, et al. A familial cluster of
pneumonia associated with the 2019 novel coronavirus indicating person-to-person
transmission: a study of a family cluster. Lancet Lond Engl. 2020;395: 514–523.
doi:10.1016/S0140-6736(20)30154-9

8. Wenham C. The oversecuritization of global health: changing the terms of debate. Int
Aff. 2019;95: 1093–1110. doi:10.1093/ia/iiz170

9. Katsidzira L, Gwaunza L, Hakim JG. The Severe Acute Respiratory Syndrome
Coronavirus 2 (SARS-CoV-2) Epidemic in Zimbabwe: Quo Vadis? Clin Infect Dis Off Publ
Infect Dis Soc Am. 2020;71: 2180–2183. doi:10.1093/cid/ciaa552

10. Plotkin MK, Williams KM, Mbinda A, Oficiano VN, Nyauchi B, Walugembe P, et al.
Keeping essential reproductive, maternal and child health services available during
COVID-19 in Kenya, Mozambique, Uganda and Zimbabwe: analysis of early-pandemic
policy guidelines. BMC Public Health. 2022;22: 577. doi:10.1186/s12889-022-12851-4

11. Manyera A, Zinyama T, Sachikonye L. The Impact of Covid-19 Pandemic on Results-
Based Finance Programmes in Developing Countries: A Literature Review. J Public Adm
Dev Altern JPADA. 2021;6: 65–81. doi:10.55190/USGD9952

12.   Baxter P, Jack S. Qualitative Case Study Methodology: Study Design and
855   Implementation for Novice Researchers. Qual Rep. 2010;13. doi:10.46743/2160-
856   3715/2008.1573

857   13.   Oxford COVID-19 Government Response Tracker (OxCGRT) | Country Planning Cycle
858   Database. [cited 28 Apr 2024]. Available:
859   https://extranet.who.int/countryplanningcycles/reportsportal/oxford-covid-19-
860   government-response-tracker-oxcgrt

861   14.   Census2022_Preliminary_Report.pdf. Available: https://www.zimstat.co.zw/wp-
862   content/uploads/2022/07/Census2022_Preliminary_Report.pdf

863   15.   Nations U. Human Development Report 2020. Hum Dev Rep. United Nations; 2020
864   Dec. Available: https://hdr.undp.org/content/human-development-report-2020

865   16.   Chirisa I, Mavhima B, Nyevera T, Chigudu A, Makochekanwa A, Matai J, et al. The
866   impact and implications of COVID-19: Reflections on the Zimbabwean society. Soc Sci
867   Humanit Open. 2021;4: 100183. doi:10.1016/j.ssaho.2021.100183

868   17.   Scoones I. Surviving COVID-19: Fragility, Resilience and Inequality in Zimbabwe. In:
869   African Arguments [Internet]. 27 Mar 2020 [cited 19 Oct 2023]. Available:
870   https://africanarguments.org/2020/03/surviving-covid-19-fragility-resilience-and-
871   inequality-in-zimbabwe/

872   18.   Dzawanda B, Matsa M, Nicolau M. Poverty on the rise: The impact of the COVID-19
873   lockdown on the informal sector of Gweru, Zimbabwe. Int Soc Sci J. 2021;71: 81–96.
874   doi:10.1111/issj.12285

875   19.   Mpofu J, Chimhenga S. Unemployment and Implications to Social and Political Conflict:
876   Perspectives from Zimbabwe. IOSR J Res Method Educ IOSR-JRME. 2016;Vol 6.
877   Available: https://iosrjournals.org/iosr-jrme/papers/Vol-6%20Issue-3/Version-
878   1/B0603010813.pdf

879   20.   Humanitarian Action for Children 2021 Overview | UNICEF. 3 Dec 2020 [cited 19 Oct
880   2023]. Available: https://www.unicef.org/reports/humanitarian-action-children-2021-
881   overview

882   21.   Zimbabwe Demographic and Health Survey 2015: Final Report – TB DIAH. [cited 19 Oct
883   2023]. Available: https://www.tbdiah.org/resources/publications/zimbabwe-
884   demographic-and-health-survey-2015-final-report/

885   22.   Mazwi F, Chemura A, Mudimu GT, Chambati W. Political Economy of Command
886   Agriculture in Zimbabwe: A State-led Contract Farming Model. Agrar South J Polit Econ.
887   2019;8: 232–257. doi:10.1177/2277976019856742

23. Zvobgo, C.J.M. A History of Zimbabwe 1890-2000 and Postscript, Zimbabwe, 2000-
2008. Cambridge Scholars Publishers; 2009. Available:
https://www.cambridgescholars.com/resources/pdfs/978-1-4438-1360-0-sample.pdf

24. Macrotrends. Zimbabwe GDP Growth Rate 1961-2024. [cited 17 Jan 2024]. Available:
https://www.macrotrends.net/countries/ZWE/zimbabwe/gdp-growth-rate

25. Dixon J, Manyau S, Kandiye F, Kranzer K, Chandler CIR. Antibiotics, rational drug use
and the architecture of global health in Zimbabwe. Soc Sci Med 1982. 2021;272:
113594. doi:10.1016/j.socscimed.2020.113594

26. Castella T de. Health workers struggle to provide care in Zimbabwe. The Lancet.
2003;362: 46–47. doi:10.1016/S0140-6736(03)13848-2

27. Wakabi W. Zimbabwe's sick public health-care system fuels disease. Lancet Infect Dis.
2009;9: 147. doi:10.1016/S1473-3099(09)70033-7

28. World Health Organisation (WHO)-The State of Health Workforce in the African Region
2021. Available:
https://iris.who.int/bitstream/handle/10665/348855/9789290234555-
eng.pdf?sequence=1

29. Mangundu M, Roets L, van Rensberg EJ. Accessibility of healthcare in rural Zimbabwe:
The perspective of nurses and healthcare users. Afr J Prim Health Care Fam Med.
2020;12: 2245. doi:10.4102/phcfm.v12i1.2245

30. Zimbabwe (ZWE) - Demographics, Health & Infant Mortality. In: UNICEF DATA
[Internet]. [cited 19 Oct 2023]. Available: https://data.unicef.org/country/zwe/

31. SDG Target 3.1 Maternal mortality. [cited 16 Nov 2023]. Available:
https://www.who.int/data/gho/data/themes/topics/sdg-target-3-1-maternal-mortality

32. Under-five mortality rate - Sustainable Development Goals - United Nations Economic
Commission for Europe. [cited 16 Nov 2023]. Available:
https://w3.unece.org/SDG/en/Indicator?id=91

33. Mutizwa-Mangiza D. Doctors and the State: The Struggle for Professional Control in
Zimbabwe. Routledge; 2018.

34. Ian S. Surviving COVID-19: Fragility, Resilience and Inequality in Zimbabwe. African
Arguments 2020: 2020.

35. projects W-F web. UPDATED: President appoints Covid-19 taskforce. In: The Chronicle
[Internet]. 25 Mar 2020 [cited 9 Feb 2024]. Available:
https://www.chronicle.co.zw/president-launches-covid-19-ministerial-task-force/

36. Reporter S. Zororo Makamba dies of covid-19. In: The Zimbabwe Mail [Internet]. 23
Mar 2020 [cited 9 Feb 2024]. Available:
https://www.thezimbabwemail.com/zimbabwe/zororo-makamba-becomes-zimbabwe-
first-covid-19-death/

37. SI 2020-076 Civil Protection (Declaration of State of Disaster Rural and Urban Areas of
Zimbabwe) (COVID-19) Notice, 2020.pdf. Available:
https://www.veritaszim.net/sites/veritas_d/files/SI%202020-
076%20Civil%20Protection%20%28Declaration%20of%20State%20of%20Disaster%20R
ural%20and%20Urban%20Areas%20of%20Zimbabwe%29%20%28COVID-
19%29%20Notice%2C%202020.pdf

38. SI 2020-077 Public Health (COVID-19 Prevention, Containment and Treatment)
Regulations, 2020.pdf. Available:
https://www.veritaszim.net/sites/veritas_d/files/SI%202020-
077%20Public%20Health%20(COVID-
19%20Prevention,%20Containment%20and%20Treatment)%20Regulations,%202020.p
df

39. World Health Organization. Operational considerations for case management of
COVID-19 in health facility and community. Interim guidance. Pediatr Med Rodz.
2020;16: 27–32. doi:10.15557/PiMR.2020.0004

40. projects W-F web. Govt unfreezes 4000 posts – Avails $500 million, redirects 2020
budget allocations to fight Covid-19. In: The Chronicle [Internet]. 31 Mar 2020 [cited 19
Oct 2023]. Available: https://www.chronicle.co.zw/govt-unfreezes-4000-posts-avails-
500-million-redirects-2020-budget-allocations-to-fight-covid-19/

41. Mackworth-Young CR, Chingono R, Mavodza C, McHugh G, Tembo M, Chikwari CD, et
al. Community perspectives on the COVID-19 response, Zimbabwe. Bull World Health
Organ. 2021;99: 85–91. doi:10.2471/BLT.20.260224

42. Mavhunga C. Zimbabwe Doctors, Nurses Down Tools Over COVID-19 PPEs. In: VOA
[Internet]. 26 Mar 2020 [cited 19 Oct 2023]. Available:
https://www.voanews.com/a/africa_zimbabwe-doctors-nurses-down-tools-over-covid-
19-ppes/6186450.html

43. Kamuti T. Covidisation of oppression: COVID-19 and human rights violations in
Zimbabwe. Soc Sci Humanit Open. 2022;6: 100374. doi:10.1016/j.ssaho.2022.100374

44. Maulani N, Nyadera IN, Wandekha B. The generals and the war against COVID-19: The
case of Zimbabwe. J Glob Health. 10: 020388. doi:10.7189/jogh.10.020388

45. Murewanhema G. COVID-19 control pitfalls and challenges and drivers of SARS-CoV-2
transmission in Zimbabwe. Pan Afr Med J. 2021;38: 28.
doi:10.11604/pamj.2021.38.28.25758

46.   Tlotleng N, Cohen C, Made F, Kootbodien T, Masha M, Naicker N, et al. COVID-19
959   hospital admissions and mortality among healthcare workers in South Africa, 2020–
960   2021. IJID Reg. 2022;5: 54–61. doi:10.1016/j.ijregi.2022.08.014

961   47.   Chitungo I, Dzinamarira T, Tungwarara N, Chimene M, Mukwenha S, Kunonga E, et al.
962   COVID-19 Response in Zimbabwe: The Need for a Paradigm Shift? COVID. 2022;2: 895–
963   906. doi:10.3390/covid2070065

964   48.   Murewanhema G, Mukwenha S, Dzinamarira T, Mukandavire Z, Cuadros D, Madziva R,
965   et al. Optimising COVID-19 Vaccination Policy to Mitigate SARS-CoV-2 Transmission
966   within Schools in Zimbabwe. Vaccines. 2021;9: 1481. doi:10.3390/vaccines9121481

967   49.   Mavhunga C. Zimbabweans Blame Government for COVID-19 Deaths of Officials. In:
968   VOA [Internet]. 23 Jan 2021 [cited 23 Oct 2023]. Available:
969   https://www.voanews.com/a/covid-19-pandemic_zimbabweans-blame-government-
970   covid-19-deaths-officials/6201126.html

971   50.   Mhuruyengwe N. FACTSHEET: Zimbabwe's latest COVID-19 quarantine rules. In:
972   ZimFact [Internet]. 21 May 2021 [cited 23 Oct 2023]. Available:
973   https://zimfact.org/factsheet-zimbabwes-latest-covid-19-quarantine-rules/

974   51.   Zimbabwe pledges to start Covid-19 vaccination campaign next week. In: News24
975   [Internet]. [cited 2 Feb 2024]. Available:
976   https://www.news24.com/news24/africa/news/zimbabwe-pledges-to-start-covid-19-
977   vaccination-campaign-next-week-20210213

978   52.   Writer S. Ministry of Health information on COVID-19 vaccination in Zimbabwe. In:
979   Techzim [Internet]. 23 Jul 2021 [cited 2 Feb 2024]. Available:
980   https://www.techzim.co.zw/2021/07/ministry-of-health-information-on-covid-19-
981   vaccination-in-zimbabwe/

982   53.   Covid-19 vaccination extended to 12-year-olds. In: The Herald [Internet]. [cited 3 Feb
983   2024]. Available: https://www.herald.co.zw/covid-19-vaccination-extended-to-12-
984   year-olds/

985   54.   projects W-F web. Govt places Kwekwe under strict lockdown. In: The Chronicle
986   [Internet]. 22 May 2021 [cited 23 Oct 2023]. Available:
987   https://www.chronicle.co.zw/govt-places-kwekwe-under-strict-lockdown/

988   55.   Zimbabwe receives nearly one million COVID-19 vaccine doses from COVAX. [cited 23
989   Oct 2023]. Available: https://www.unicef.org/zimbabwe/press-releases/zimbabwe-
990   receives-nearly-one-million-covid-19-vaccine-doses-covax

991   56.   projects W-F web. Global Fund avails funds for Covid-19. In: The Chronicle [Internet].
992   11 Jun 2021 [cited 23 Oct 2023]. Available: https://www.chronicle.co.zw/global-fund-
993   avails-funds-for-covid-19/

57. ad365-zimbabwean_vulnerabilities_on_health_care_water-afrobarometer-
5june20.pdf. Available: https://www.afrobarometer.org/wp-
content/uploads/migrated/files/publications/Dispatches/ad365-
zimbabwean_vulnerabilities_on_health_care_water-afrobarometer-5june20.pdf

58. Kavenga F, Rickman HM, Chingono R, Taruvinga T, Marembo T, Manasa J, et al.
Comprehensive occupational health services for healthcare workers in Zimbabwe
during the SARS-CoV-2 pandemic. PLOS ONE. 2021;16: e0260261.
doi:10.1371/journal.pone.0260261

59. Fryatt A, Simms V, Bandason T, Redzo N, Olaru ID, Ndhlovu CE, et al. Community SARS-
CoV-2 seroprevalence before and after the second wave of SARS-CoV-2 infection in
Harare, Zimbabwe. eClinicalMedicine. 2021;41: 101172.
doi:10.1016/j.eclinm.2021.101172

60. Rusakaniko S, Sibanda EN, Mduluza T, Tagwireyi P, Dhlamini Z, Ndhlovu CE, et al. SARS-
CoV-2 Serological testing in frontline health workers in Zimbabwe. PLoS Negl Trop Dis.
2021;15: e0009254. doi:10.1371/journal.pntd.0009254

61. Lewis HC, Ware H, Whelan M, Subissi L, Li Z, Ma X, et al. SARS-CoV-2 infection in Africa:
a systematic review and meta-analysis of standardised seroprevalence studies, from
January 2020 to December 2021. BMJ Glob Health. 2022;7: e008793.
doi:10.1136/bmjgh-2022-008793

62. Sayed A, Challa KT, Arja S. Epidemiological Differences of COVID-19 Over the World.
Cureus. 12: e10316. doi:10.7759/cureus.10316

63. Details on the COVID-19 Economic Recovery and Stimulus Package | veritaszim. [cited
28 Apr 2024]. Available: https://www.veritaszim.net/node/4112

64. Seven babies stillborn in one night at Zimbabwe hospital as nurses strike. Reuters. 29
Jul 2020. Available: https://www.reuters.com/article/health-zimbabwe-
idUKL5N2F06H6. Accessed 23 Oct 2023.

65. Chimhuya S, Neal SR, Chimhini G, Gannon H, Cortina Borja M, Crehan C, et al. Indirect
impacts of the COVID-19 pandemic at two tertiary neonatal units in Zimbabwe and
Malawi: an interrupted time series analysis. BMJ Open. 2022;12: e048955.
doi:10.1136/bmjopen-2021-048955

66. Thekkur P, Takarinda KC, Timire C, Sandy C, Apollo T, Kumar AMV, et al. Operational
Research to Assess the Real-Time Impact of COVID-19 on TB and HIV Services: The
Experience and Response from Health Facilities in Harare, Zimbabwe. Trop Med Infect
Dis. 2021;6: 94. doi:10.3390/tropicalmed6020094

67. Timire C, Sandy C, Ferrand RA, Mubau R, Shiri P, Mbiriyawanda O, et al. Coverage and
effectiveness of conditional cash transfer for people with drug resistant tuberculosis in

Zimbabwe: A mixed methods study. PLOS Glob Public Health. 2022;2: e0001027.
doi:10.1371/journal.pgph.0001027

68.   Shakespeare C, Dube H, Moyo S, Ngwenya S. Resilience and vulnerability of maternity
1033   services in Zimbabwe: a comparative analysis of the effect of Covid-19 and lockdown
1034   control measures on maternal and perinatal outcomes, a single-centre cross-sectional
1035   study at Mpilo Central Hospital. BMC Pregnancy Childbirth. 2021;21: 416.
1036   doi:10.1186/s12884-021-03884-5

1037   69.   Suburban. Marlborough district office, clinic reopen. In: Suburban [Internet]. 12 Feb
1038   2021 [cited 19 Oct 2023]. Available: https://www.suburban.co.zw/marlborough-
1039   district-office-clinic-reopen/

1040   70.   Mavodza CV, Bernays S, Mackworth-Young CRS, Nyamwanza R, Nzombe P, Dauya E, et
1041   al. Interrupted Access to and Use of Family Planning Among Youth in a
1042   Community-Based Service in Zimbabwe During the First Year of the COVID-19
1043   Pandemic. Stud Fam Plann. 2022; 10.1111/sifp.12203. doi:10.1111/sifp.12203

1044   71.   Matsungo TM, Chopera P. Effect of the COVID-19-induced lockdown on nutrition,
1045   health and lifestyle patterns among adults in Zimbabwe. BMJ Nutr Prev Health. 2020;3:
1046   205–212. doi:10.1136/bmjnph-2020-000124

1047   72.   Nyahunda L, Chibvura S, Tirivangasi HM. Social Work Practice: Accounting for Double
1048   Injustices Experienced by Women Under the Confluence of Covid-19 Pandemic and
1049   Climate Change Impacts in Nyanga, Zimbabwe. J Hum Rights Soc Work. 2021;6: 213–
1050   224. doi:10.1007/s41134-021-00170-4

1051   73.   Providing services to survivors of gender-based violence during COVID-19 | United
1052   Nations in Zimbabwe. [cited 23 Oct 2023]. Available:
1053   https://zimbabwe.un.org/en/50660-providing-services-survivors-gender-based-
1054   violence-during-covid-19, https://zimbabwe.un.org/en/50660-providing-services-
1055   survivors-gender-based-violence-during-covid-19

1056   74.   Zimbabwe: Covid-19 Lockdown - Gender-Based Violence Cases Up By 70 Percent. New
1057   Zimbabwe. 28 Jul 2020. Available: https://allafrica.com/stories/202007280268.html.
1058   Accessed 23 Oct 2023.

1059   75.   The Zimbabwe Independent - The Leading Business Weekly. In: The Zimbabwe
1060   Indpendent [Internet]. [cited 19 Oct 2023]. Available:
1061   https://www.newsday.co.zw/theindependent/

1062   76.   Overview. In: World Bank [Internet]. [cited 19 Oct 2023]. Available:
1063   https://www.worldbank.org/en/country/zimbabwe/overview

77.   EMIS 2021 Annual Education Statistics Report.pdf. Available:
https://mopse.co.zw/sites/default/files/public/downloads/EMIS%202021%20Annual%
20Education%20Statistics%20Report.pdf

78.   Poncin M, Marembo J, Chitando P, Sreenivasan N, Makwara I, Machekanyanga Z, et al.
Implementation of an outbreak response vaccination campaign with typhoid conjugate
vaccine – Harare, Zimbabwe, 2019. Vaccine X. 2022;12: 100201.
doi:10.1016/j.jvacx.2022.100201

79.   Caribbean EC for LA and the. The effects of the coronavirus disease (COVID-19)
pandemic on international trade and logistics. Economic Commission for Latin America
and the Caribbean; 2020. Available: https://www.cepal.org/en/publications/45878-
effects-coronavirus-disease-covid-19-pandemic-international-trade-and-logistics

80.   Allington D, McAndrew S, Moxham-Hall V, Duffy B. Coronavirus conspiracy suspicions,
general vaccine attitudes, trust and coronavirus information source as predictors of
vaccine hesitancy among UK residents during the COVID-19 pandemic. Psychol Med.
2023;53: 236–247. doi:10.1017/S0033291721001434

81.   Zimbabwe Humanitarian Situation Report, 31 December 2021.pdf. Available:
https://www.unicef.org/media/114866/file/Zimbabwe%20Humanitarian%20Situation
%20Report,%2031%20December%202021.pdf

82.   (PDF) Gendered Socio-economic Implications of the COVID-19 Pandemic in Rural
Zimbabwe. [cited 23 Oct 2023]. Available:
https://www.researchgate.net/publication/344711278_Gendered_Socio-
economic_Implications_of_the_COVID-19_Pandemic_in_Rural_Zimbabwe

83.   Josephson A, Kilic T, Michler JD. Socioeconomic impacts of COVID-19 in low-income
countries. Nat Hum Behav. 2021;5: 557–565. doi:10.1038/s41562-021-01096-7

1088

1089

1090

1091

1092

1093

Fig 1: Timeline of key events

*BYO- Bulawayo; HAR- Harare

Fig 2: Maternal indicators

Fig 3: Child indicators

*BYO- Bulawayo; HAR- Harare

Fig 4A: Enrolled, new entrants and drop outs

Fig 4B: Gross enrolment ration and transition rates

Fig 4C: Completion and pass rates across education levels