Understanding Sociodemographic and Dietary Determinants of Cardiometabolic Risk: Cross-Sectional Evidence from the U.S. Healthy Eating Index to Inform Diet Quality Categories

Elise Sheinberg, MPH, RDN[1*]; Laura A. Schmidt, PhD, MSW, MPH[2]; Jerold R. Mande, MPH[1,3,4]; Euridice Martinez-Steele, PhD[5,6]; Deirdre K. Tobias, ScD[1,7]; Cindy W. Leung; ScD, MPH[1]

[1] Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA
[2] Philip R. Lee Institute for Health Policy Studies, School of Medicine, University of California, San Francisco, San Francisco, CA
[3] Nourish Science, Bethesda, MA
[4] Tisch College of Civic Life, Tufts University, Medford, MA
[5] Department of Nutrition, School of Public Health, University of Såo Paulo, São Paulo, Brazil
[6] Center for Epidemiological Studies in Health and Nutrition, University of São Paulo, São Paulo, Brazil
[7] Division of Preventive Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA

Corresponding Author:
Elise Sheinberg
Department of Nutrition
Harvard T.H. Chan School of Public Health
665 Huntington Avenue
Building 2
Boston, MA 02115
Phone: 402-719-4801
esheinberg@g.harvard.edu

ABSTRACT

Background: Rising rates of diet-related chronic diseases and disparities in disease prevalence

by sociodemographic factors highlight the need to improve the diet quality of Americans.

Understanding how the Healthy Eating Index-2020 (HEI-2020) can be used as a measure of diet

quality to create benchmarks for national nutrition monitoring can assist in surveillance and

improve the design of nutrition programs and policies.

Objective: To examine the utility of the HEI-2020 to create benchmarks for national diet quality

monitoring.

Methods: This serial cross-sectional study used data from the 2009-2018 National Health and

Nutrition Examination Surveys (NHANES). Nationally representative data for 22,168 US adults

(≥20 years) who completed two 24-hour recalls were analyzed. We derived HEI-2020 scores (0-

100) from participants' two 24-hour dietary recalls. Diet quality categories were established: high

diet quality (>70-100), marginal (>60-70), low (>50-60), and very low (0-50).

Results: Only 13% of US adults had high diet quality while nearly two-thirds had low or very

low diet quality**.** Diet quality was higher for older adults, female, "Other" race or ethnicity, born

outside of the US, have higher education attainment, higher income, and food security.

Compared to adults with high diet quality, adults with very low diet quality had lower intakes of

unprocessed or minimally processed foods, fruits, vegetables, whole grains, and seafood and the

highest intakes of ultra-processed foods, refined grains, and red and processed meats (all *P-*

trends <0.01). Adults with very low diet quality were more likely to have elevated adiposity,

lower HDL cholesterol, and higher triglycerides, fasting glucose, and hemoglobin A1c (all *P-*

trends ≤0.01).

Conclusions: The HEI-2020 is a robust measure of diet quality that can be directly linked to

biological measures associated with chronic disease risk. Using evidence-based HEI categories

could allow policymakers, public health practitioners, and nutrition professionals to set

benchmarks and nationwide targets for achieving improved diet quality.

Keywords: Healthy Eating Index, food security, diet quality, chronic disease prevention

INTRODUCTION

Poor diet quality is a risk factor for chronic diseases, including obesity, diabetes, cardiovascular disease, stroke, and some cancers.[1–3] Sociodemographic factors, including lower educational attainment, lower household income, minority racial and ethnic identity, and experiencing food insecurity, are associated with lower diet quality and higher chronic disease prevalence.[4–12] Food insecurity, defined as "the limited or uncertain availability of nutritionally adequate and safe foods, or limited or uncertain ability to acquire acceptable foods in socially acceptable ways,"[13] is associated with both poor diet quality and increased risk of chronic diseases.[14–16] Concern about the relationship between food insecurity, hunger, and malnutrition led to the national monitoring of food insecurity since 1995.[17] Validated food insecurity measurement tools can distinguish between individuals who have high food security, marginal food security, low food security, and very low food security.[13] National trends and differences by sub-groups have informed public health policy and program development.

Nationally representative dietary intake is available through the National Health and Nutrition Examination Survey (NHANES), and can be used to assess alignment to the Dietary Guidelines for Americans (DGA), which provide diet patterns that prevent disease and promote health and well-being, with the Healthy Eating Index (HEI). According to the 2020-2025 DGA, a high-quality diet includes adequate vegetables, fruits, whole grains, low- and non-fat dairy, and low-fat protein, including beans and nuts, and limits consumption of added sugar, sodium, and saturated fat.[18] The average HEI-2020 for adults was 56 out of 100 in 2023.[19] Despite most Americans' poor adherence to the DGA and rising rates of diet-related chronic diseases, diet quality categories that are associated with disease risk have not been established and used for national monitoring, as has been done for food insecurity.

There has been growing interest in the relationship between food security and diet

quality, as well as a recognition of the need to address diet quality to reduce chronic diseases,

constrain healthcare spending, and improve quality of life for all Americans.[20–22] This has

included bipartisan calls for policy action to monitor diet quality and consider the nutritional

impact of federal nutrition programs, like the Supplemental Nutrition Assistance Program

(SNAP).[23–25] Population-based analyses of HEI have typically presented score means, and

evidence to support the utility of specific thresholds (e.g. high vs. low diet quality) of risk for

nutrition-related disease and mortality is warranted to understand if categories using the HEI can

be used to monitor diet quality trends, similar to how food insecurity is monitored.[26–28]

Using nationally representative data, we created four diet quality categories of HEI-2020

scores: high (≥70), marginal (>60-70, low (>50-60), and very low (≤50) that were predictive of

sociodemographic disadvantage and biomarkers associated with cardiometabolic disease risk and

aligned with existing research.[3,29,30] Establishing objective, evidence-based diet quality

categories using the HEI can generate timely and critical data for policymakers, public health

practitioners, and nutrition professionals to tackle the systemic drivers of poor diet quality and

can promote long-term health for all Americans.

METHODS

Study Design and Population

We conducted our analyses using data from the 2009-2018 NHANES. NHANES is an

ongoing multistage, nationally representative survey conducted by the National Center for Health

Statistics (NCHS). In this study, the analytic sample included 22,168 adults ≥20 years with two

valid 24-hour recalls (**Supplemental Figure 1**). Data collection was approved by NCHS ethics

review board and all participants provided written consent.

Dietary Assessment and the HEI-2020 scoring

Dietary intake was assessed via two 24-hour dietary recalls administered by trained

interviewers using the USDA Automated Multi-Pass Method.[31] One 24-hour recall is collected

in-person, and a second recall is completed by telephone 3-10 days later. Methods and validation

studies for conducting 24-hour recalls have been previously published.[32] Data from the two 24-

hour recalls were converted into standard food groups and food pattern components using the

Food Patterns Equivalents Database (FPED) to calculate individual HEI-2020 scores.

Participants' HEI-2020 scores were calculated using the simple HEI scoring algorithm method

and averaged across the two days of intake (i.e., observed HEI). We also used the National

Cancer Institute (NCI) SAS Macros for the Bivariate Method (version 2.1) in order to account

for the within-person variability of dietary intake and to estimate predicted HEI scores (i.e., usual

HEI).[33]

HEI cut-points

The determination of the cut-points used to define categories of increasing diet quality

was informed on multiple criteria. First, we comprehensively reviewed prior findings from

prospective observational studies that examined HEI scores in relation to major chronic disease

outcomes (**Supplemental Table 1**).[2] Most studies derived population-specific quintiles of HEI

scores as the exposure, [34] and individuals in the highest HEI quintile consistently had lower risks

of cardiovascular disease, cancer, type 2 diabetes, and all-cause mortality compared to

individuals in the lowest quintile.[2,35] Second, we examined the distributions of observed and

usual HEI scores in NHANES to understand the extent to which US adults aligned with the cut

points in the studies above (**Supplemental Figure 2**). Finally, we considered NCI's suggested

HEI cut-points and corresponding grade.[36] Collectively, this evidence informed the following

category definitions: high diet quality (>70-100), marginal diet quality (>60-70), low diet quality

(>50-60), and very low diet quality (0-50). High diet quality indicates a favorable adherence to

the DGA and where the greatest risk reduction of morbidity and mortality from diet-related

chronic diseases has been observed.[2,29,37,38] In contrast, very low diet quality indicates poor

adherence to the DGA and is associated with the highest risks of morbidity and mortality from

diet-related chronic diseases. Marginal and low diet quality are intermediate risk levels.

Other Dietary Components

All dietary variables were derived from the average of two days of dietary recalls and

standardized to 1,000 kcal. Other dietary factors in the current analysis included total energy

intake (kcal/day). whole fruits (cup equivalents/day), vegetables excluding white potatoes (cup

equivalents), refined grains (ounce equivalents/day), whole grains (ounce equivalents/day), red

and processed meats (ounce equivalents/day), and seafood (ounce equivalents/day). Nutrients of

interest included saturated fat (g/day), dietary fiber (g/day), sodium (mg/day), potassium

(mg/day), calcium (mg/day), and added sugar (tsp equivalents/day). We also applied the NOVA

classification to derive intakes of unprocessed and minimally processed foods and ultra-

processed foods (% energy/day).[39,40]

Adiposity and cardiometabolic markers

Height, weight, and waist circumference (WC) were measured by trained personnel in the

Mobile Examination Center.[41] Body mass index was calculated as weight (kg) over height

squared (m^2). Obesity was defined as a BMI ≥ 30 kg/m^2.[42] High waist circumference was defined

as >88 centimeters for females and >102 centimeters for males.[43]

Cardiometabolic biomarkers in this analysis include blood levels of high-density

lipoprotein cholesterol (HDL-c; mg/dL), low-density lipoprotein cholesterol (LDL-c; mg/dL),

triglycerides (mg/dL), and glucose (mg/dL). Low HDL cholesterol was defined as <40 mg/dL

and elevated LDL cholesterol was defined as ≥100 mg/dL.[44] Elevated triglycerides were defined

as ≥150 mg/dL.[45] Elevated fasting glucose was defined as >100 mg/dL.[46]

Sociodemographic Characteristics

Characteristics that have been associated with diet quality in previous research were

included as covariates.[47–49] Self-reported sociodemographic variables were ascertained at patient

visits, including age, sex, race and ethnicity, place of birth, educational attainment, employment

status, marital status, family income to poverty ratio. Household food security status was derived

using the 18-item U.S. Food Security Survey Module.[50]

Statistical Analysis

Complex two-day dietary sampling weights were adjusted to reflect the 10-year analytic

period and applied to all analyses to make nationally representative inferences. We first

calculated the proportions US adults for each of our new HEI-based diet-quality categories

cumulatively and separately for each NHANES 2-year cycle. We evaluated whether diet quality

categories differed over time and by sociodemographic characteristics using the Satterwaite-

adjusted Chi-Square test. We generated radar plots to visualize the contribution of HEI-

components across diet quality categories, as a percentage of the maximum possible score for

that component. For other individual dietary factors and cardiometabolic biomarkers, we did

complete case analysis and estimated their least squared means across diet quality categories

using multivariable linear regression models adjusted for all sociodemographic characteristics to

improve precision. We estimate the predicted probabilities of the clinically relevant

cardiometabolic biomarkers in each diet quality category with logistic regression models,

adjusting for all sociodemographic characteristics. In sensitivity analyses, statistical tests were

repeated using diet quality categories derived from usual HEI scores. Statistical analyses were

performed with SAS 9.4 (SAS Institute Inc., Cary, NC).

RESULTS

Between 2009 and 2018, 12.7% of US adults had high diet quality (HEI ≥70), 19.9% had

marginal diet quality (HEI 60-<70), 26.9% had low diet quality (HEI 50-<60), and 40.5% had

very low diet quality (HEI ≤50) (**Figure 1**). When usual dietary intakes were used, only 0.8% of

US adults had high diet quality, 12.9% had marginal diet quality, 41.3% had low diet quality, and

45.0% of adults had very low diet quality. There were no significant differences over the 10-year

NHANES cycles in the study period *(***Supplemental Figure 3***, P>0.05)*

There were statistically significant differences in diet quality categories by all

sociodemographic characteristics except employment status (**Table 1**, *P-linear trend <0.01*).

Adults ≥51 years were more likely to have high diet quality (16.1%) than adults ≤30 years

(8.0%). Females were more likely to have high diet quality than males (14.9% compared with

10.4%, respectively). By race and ethnicity, 8.0% of non-Hispanic Black adults had high diet

quality compared to 11.0% of Hispanic adults, 13.1% of non-Hispanic white adults, and 19.4%

of adults in another racial or ethnic group (i.e., "Other"). Adults born outside of the US were

more likely to have high diet quality (19.2%) than adults born in the US (11.5%). Adults with

some college education or more were more likely to have high diet quality (15.5%) than those

with a high school education or less (8.0%). Married or partnered adults (13.7%) were more

likely to have high diet quality than single adults (11.1%). Adults with household incomes

>200% FPL were more likely to have high diet quality (14.6%) than adults with household

incomes ≤200% FPL (9.0%). Adults with high or marginal food security were more likely to

have high diet quality (13.9%) than adults with low or very low food security (6.3%).

A radar plot showing the proportion of the possible points for each HEI component for

each diet quality category is shown in **Figure 2**. Adults with high diet quality scored at least half

of the possible points for all HEI components. Adults with very low diet quality scored less than

half of the possible points for nine components. The greatest differences in HEI-2020 component

scores between adults with high versus very low diet quality were for whole grains (very low diet

quality scored 78.4% lower than high diet quality), whole fruit (very low diet quality scored

72.5% lower than high diet quality), total fruit (very low diet quality scored 70.9% lower than

high diet quality), greens and beans (very low diet quality scored 69.1% lower than high diet

quality), seafood and plant protein (very low diet quality scored 58.9% lower than high diet

quality) and fatty acids (very low diet quality scored 55.4% lower than high diet quality).

Multivariate-adjusted mean intakes of foods and nutrients of interest across diet quality

categories are shown in **Table 2**. Mean total energy intake was lowest among adults with high

diet quality (1976 kcal/day) and increased across diet quality categories (marginal: 2031

kcal/day, low: 2102 kcal/day; and very low: 2106 kcal/day) (*P-linear trend<0.01*). Adults with

high diet quality had the highest intakes of unprocessed or minimally processed foods (41.4%

energy) and adults with very low diet quality had the lowest intakes (25.8% energy) – a

difference of +15.6% (*P-linear trend<0.01*). Similarly, adults with high diet quality had the

lowest intakes of ultra-processed foods (42.5% energy) while adults with very low diet quality

had the highest intakes (61.7% energy)- a difference of -19.2% (*P-linear trend<0.01*).

Adults with high diet quality consumed significantly more fruits, vegetables, whole

grains, and seafood, and less refined grains and red and processed meats when compared to

adults with marginal, low, and very low diet quality (all *P-linear trends<0.01)* (**Table 2**). Adults

with high diet quality also consumed lower amounts of saturated fat, sodium, and added sugars

and higher amounts of dietary fiber, potassium, and calcium than adults with marginal, low, and

very low diet quality (all *P-linear trends<0.01*). Nearly all comparisons between the high diet

quality group with marginal, low, and very low diet quality were statistically significant, except

for seafood consumption between high and marginal diet quality groups. Sensitivity analyses

examining multivariate-adjusted means of usual intakes showed identical patterns

(**Supplemental Table 2**).

There were differences in BMI and obesity prevalence across diet quality categories

(**Table 3**). After adjusting for sociodemographic factors, the mean BMI for adults with high diet

quality was lower (27.4 kg/m^2) compared to adults with marginal (28.2 kg/m^2), low (29.2

kg/m^2), and very low diet quality (29.8 kg/m^2) (*P-linear trend<0.01*). Relatedly, 27.3% of adults

with high diet quality had obesity compared to 45.7% of adults with very low diet quality – a

difference of -18.4% (*P-trend<0.01*). Adults with high diet quality had a mean waist

circumference of 94.0 cm (49.3% with elevated waist circumference), compared to a mean waist

circumference of 100.4 cm for adults with very low diet quality (61.5% elevated waist

circumference) (*P-linear trends<0.01*).

Cardiometabolic biomarkers predictive of diet-related disease also differed across diet

quality categories (**Table 3**). Mean HDL cholesterol was higher among adults with high diet

quality (56.6 mg/dL) and decreased across diet quality categories (marginal: 54.1 mg/dL; low:

52.9 mg/dL; very low: 50.7 mg/dL) (*P-trend<0.01*). Clinically low HDL cholesterol was

observed in 13.5% of adults with high diet quality compared to 24.1% of adults with very low

diet quality – a difference of 10.6% (*P-linear trend<0.01*). Similarly, adults with high diet

quality had lower mean LDL cholesterol levels (107.8 mg/dL) compared to adults with marginal

(111.9 mg/dL), low (111.6 mg/dL), and very low diet quality (112.6 mg/dL) (*P-linear*

trend=0.01). There was no difference in the proportion of adults with elevated LDL cholesterol

by diet quality category (*P-linear trend=0.14*). Adults with high diet quality had a mean

triglyceride level of 103.6 mg/dL and 18.6% had clinically elevated triglycerides; adults with

very low diet quality had a mean triglyceride level of 123.7 mg/dL, and 25.7% had clinically

elevated triglycerides (*P-linear trends<0.01*). Adults with high diet quality had lower fasting

glucose (101.7 mg/dL) compared to adults with very low diet quality (mean fasting glucose

107.1 mg/dL) (*P-linear trends<0.01*). Approximately 43% of adults with high diet quality had

clinically elevated fasting glucose compared to 44.6% of adults with marginal, 48.6% with low,

and 47.7% with very low diet quality (*P-linear trend=<0.01*). In sensitivity analyses,

multivariate-adjusted differences in adiposity and cardiometabolic biomarkers by HEI categories

using usual intake revealed consistent patterns (**Supplemental Table 3**)

DISCUSSION

In this nationally representative ten-year study, only 13% of adults had high diet quality,

and most (67%) had low or very low diet quality. These proportions were more magnified when

considering usual HEI scores. This indicates that few Americans consume a diet aligned with the

DGA at levels that protect against diet-related chronic disease morbidity and mortality, with

notable disparities across sociodemographic factors. There were also no observable

improvements in the proportion of adults achieving high diet quality over time.[24]

We noted disparities in diet quality categories by key sociodemographic characteristics.

Non-Hispanic Black race/ethnicity, having a high school education or less, low household

income, and food insecurity were all significantly associated with higher prevalence of very low

diet quality. Although these are cross-sectional data, prior evidence indicates structural factors

contribute to disparities in diet quality including racism, limited educational opportunities,

inequitable food access, a lack of proper training, resources, or enough time to prepare healthy

meals.[10,51–53]

In this study, more healthful diet quality categories were associated with a healthy weight

status, lipid profile, and glycemic control in a dose-response manner. This is consistent with

meta-analyses of prospective observational studies, which found that individuals who score at

the top quintile of diet quality for their population have lower risks of all-cause mortality,

cardiovascular disease, cancer, and type 2 diabetes compared to individuals who score in the

bottom quintile.[2,38] This indicates that the proposed HEI cut-points reflect the underlying

cardiometabolic risk profile of US adults.

Significant differences in all foods, food groups, and nutrients of interest were observed

across diet quality categories. Shifting to a higher diet quality category can be achieved through

incremental dietary improvements. For example, the difference between high and marginal diet

quality categories could be translated to a daily 6% kcal reduction from ultra-processed foods,

0.5 cups more fruits, 0.3 cups more vegetables, 0.8 oz equivalents more whole grains, and 0.3 oz

equivalents more seafood. Similarly, the difference between marginal and low diet quality

categories was, on average, a daily 5.5% kcal reduction from ultra-processed foods, 0.4 cups

more fruits, 0.1 cups more vegetables 0.4 oz equivalents more whole grains and 0.7 oz

equivalents more seafood. Finally, the difference between low and very low diet quality was on

average an 8.7% reduction of energy/day from ultra-processed foods, 0.4 cups/day more fruit,

0.1 cups/day more vegetables, 0.5 oz equivalents/day more whole grains, and 0.3-ounce

equivalents/day more seafood.

There have been calls to supplement food security surveillance measures with diet quality

measures.[22,23,25,54] This study shows that defining cut-points of HEI-2020 scores adequately and

comprehensively captures underlying differences in key dietary, biological, and social risk

factors implicated in diet-related diseases and could be used to efficiently assess the diet quality

of the population. Nutrition interventions that work with individuals who have risk factors for

very low diet quality often monitor food insecurity status of participants, and using evidence-

based diet quality targets may enhance efforts to monitor and improve diet quality. This can

reduce disparities in diet-related chronic diseases by raising participants into higher diet quality

categories, in combination with other program evaluations and benchmarks.

Strengths of this study include our use of data from a large nationally representative

sample of US adults spanning recent years. Second, we identified the cut points for HEI

categories through multiple sources of health outcome evidence. Third, we leveraged the rich

data sources available from NHANES, validated dietary recalls, physical measurements, and

available biomarker data, to examine concurrent implications of the diet quality categories at the

individual level. However, this study is limited by its cross-sectional nature; prospective follow-

up is needed to capture clinical outcomes. The 24-hour recall is a validated approach but is

subject to measurement error and recall bias. Cut-points could result in misclassification of

individuals. Furthermore, we did not account for the presence of chronic diseases or use of

medications, as the goal was not to show causal effects of diet quality categories and biomarker

outcomes, but to characterize population-level differences across diet quality categories.

Conclusion:

Given that the HEI is already used to monitor the nutritional status of the US, using HEI cut-points to establish diet quality categories would be an efficient, evidence-based, and practical approach allow policy makers, public health practitioners and nutrition professionals to set benchmarks and targets for nutrition policies and interventions. Disaggregating continuous HEI scores into meaningful categories can better identify disparities in diet-related health outcomes and help tailor existing nutrition programs and policies to improve healthful food access, increase diet quality, and promote health for all Americans.

Author Contributions: CWL LAS, JRM and DKT designed research; EMS provided essential materials; CWL analyzed data; ES and CWL wrote paper; ES had primary responsibility for the final content; all authors read and approved the final manuscript; all authors have read and approved the final manuscript.

REFERENCES

1. Micha R, Peñalvo JL, Cudhea F, Imamura F, Rehm CD, Mozaffarian D. Association Between Dietary Factors and Mortality From Heart Disease, Stroke, and Type 2 Diabetes in the United States. *JAMA*. 2017;317(9):912-924. doi:10.1001/jama.2017.0947

2. Morze J, Danielewicz A, Hoffmann G, Schwingshackl L. Diet Quality as Assessed by the Healthy Eating Index, Alternate Healthy Eating Index, Dietary Approaches to Stop Hypertension Score, and Health Outcomes: A Second Update of a Systematic Review and Meta-Analysis of Cohort Studies. *J Acad Nutr Diet*. 2020;120(12):1998-2031.e15. doi:10.1016/j.jand.2020.08.076

3. Xu Z, Steffen LM, Selvin E, Rebholz CM. Diet quality, change in diet quality and risk of incident CVD and diabetes. *Public Health Nutr*. 2020;23(2):329-338. doi:10.1017/S136898001900212X

4. National Diabetes Statistics Report | Diabetes | CDC. November 20, 2023. Accessed November 24, 2023. https://www.cdc.gov/diabetes/data/statistics-report/index.html

5. Lopez-Neyman SM, Davis K, Zohoori N, Broughton KS, Moore CE, Miketinas D. Racial disparities and prevalence of cardiovascular disease risk factors, cardiometabolic risk factors, and cardiovascular health metrics among US adults: NHANES 2011–2018. *Sci Rep*. 2022;12(1):19475. doi:10.1038/s41598-022-21878-x

6. Schultz WM, Kelli HM, Lisko JC, et al. Socioeconomic Status and Cardiovascular Outcomes. *Circulation*. 2018;137(20):2166-2178. doi:10.1161/CIRCULATIONAHA.117.029652

7. Singh S, Sridhar P. A narrative review of sociodemographic risk and disparities in screening, diagnosis, treatment, and outcomes of the most common extrathoracic malignancies in the United States. *J Thorac Dis*. 2021;13(6):3827-3843. doi:10.21037/jtd-21-87

8. Agardh E, Allebeck P, Hallqvist J, Moradi T, Sidorchuk A. Type 2 diabetes incidence and socio-economic position: a systematic review and meta-analysis. *Int J Epidemiol*. 2011;40(3):804-818. doi:10.1093/ije/dyr029

9. Gupta A, Wilson LE, Pinheiro LC, et al. Association of educational attainment with cancer mortality in a national cohort study of black and white adults: A mediation analysis. *SSM - Popul Health*. 2023;24:101546. doi:10.1016/j.ssmph.2023.101546

10. Leung CW, Tester JM. The Association between Food Insecurity and Diet Quality Varies by Race/Ethnicity: An Analysis of National Health and Nutrition Examination Survey 2011-2014 Results. *J Acad Nutr Diet*. 2019;119(10):1676-1686. doi:10.1016/j.jand.2018.10.011

11. Leung CW, Wolfson JA, Brandt EJ, Rimm EB. Disparities in Cardiovascular Health by Food Security Status and Supplemental Nutrition Assistance Program Participation Using Life's Essential 8 Metrics. *JAMA Netw Open*. 2023;6(6):e2321375. doi:10.1001/jamanetworkopen.2023.21375

12. Johnson AE, Herbert BM, Stokes N, Brooks MM, Needham BL, Magnani JW. Educational Attainment, Race, and Ethnicity as Predictors for Ideal Cardiovascular Health: From the National Health and Nutrition Examination Survey. *J Am Heart Assoc Cardiovasc Cerebrovasc Dis*. 2022;11(2):e023438. doi:10.1161/JAHA.121.023438

13. United States Department of Agriculture. Food Insecurity in the U.S. - Measurement. Accessed November 26, 2023. https://www.ers.usda.gov/topics/food-nutrition-assistance/food-security-in-the-u-s/measurement/

14. Liu J, Yi SS, Russo RG, et al. Trends and disparities in prevalence of cardiometabolic diseases by food security status in the United States. *Nutr J*. 2024;23:4. doi:10.1186/s12937-023-00910-4

15. Leung CW, Epel ES, Ritchie LD, Crawford PB, Laraia BA. Food insecurity is inversely associated with diet quality of lower-income adults. *J Acad Nutr Diet*. 2014;114(12):1943-1953.e2. doi:10.1016/j.jand.2014.06.353

16. Laraia BA. Food Insecurity and Chronic Disease. *Adv Nutr*. 2013;4(2):203-212. doi:10.3945/an.112.003277

17. National Research Council. *Food Insecurity and Hunger in the United States: An Assessment of the Measure*. The National Academies Press; 2006. https://doi.org/10.17226/11578.

18. U.S. Department of Agriculture, U.S. Department of Health and Human Services. *Dietary Guidelines for Americans, 2020-2025*. DietaryGuidelines.gov

19. *Scientific Report of the 2025 Dietary Guidelines Advisory Committee: Advisory Report to the Secretary of Health and Human Services and Secretary of Agriculture*. HHS and USDA; 2024. doi:10.52570/DGAC2025

20. Herman PM, Nguyen P, Sturm R. Diet quality improvement and 30-year population health and economic outcomes: a microsimulation study. *Public Health Nutr*. 25(5):1265-1273. doi:10.1017/S136898002100015X

21. Jardim TV, Mozaffarian D, Abrahams-Gessel S, et al. Cardiometabolic disease costs associated with suboptimal diet in the United States: A cost analysis based on a microsimulation model. Basu S, ed. *PLOS Med*. 2019;16(12):e1002981. doi:10.1371/journal.pmed.1002981

22. Thorndike AN, Gardner CD, Kendrick KB, et al. Strengthening US Food Policies and Programs to Promote Equity in Nutrition Security: A Policy Statement From the American Heart Association. *Circulation*. 2022;145(24):e1077-e1093. doi:10.1161/CIR.0000000000001072

23. Booker C, Rubio M. *SNAP Nutrition Security Act*.; 2023. https://www.booker.senate.gov/imo/media/doc/snap_nutrition_security_act_of_2023.pdf

24. ODPHP. White House Conference on Hunger, Nutrition and Health. Accessed May 20, 2024. https://health.gov/our-work/nutrition-physical-activity/white-house-conference-hunger-nutrition-and-health

25. Thompson G. *Farm, Food, and National Security Act of 2024*.; 2024. https://www.congress.gov/bill/118th-congress/house-bill/8467

26. Liu J, Micha R, Li Y, Mozaffarian D. Trends in Food Sources and Diet Quality Among US Children and Adults, 2003-2018. *JAMA Netw Open*. 2021;4(4):e215262. doi:10.1001/jamanetworkopen.2021.5262

27. Tao MH, Liu JL, Nguyen USDT. Trends in Diet Quality by Race/Ethnicity among Adults in the United States for 2011–2018. *Nutrients*. 2022;14(19):4178. doi:10.3390/nu14194178

28. Shan Z, Rehm CD, Rogers G, et al. Trends in Dietary Carbohydrate, Protein, and Fat Intake and Diet Quality Among US Adults, 1999-2016. *JAMA*. 2019;322(12):1178-1187. doi:10.1001/jama.2019.13771

29. Hu EA, Steffen LM, Coresh J, Appel LJ, Rebholz CM. Adherence to the Healthy Eating Index–2015 and Other Dietary Patterns May Reduce Risk of Cardiovascular Disease, Cardiovascular Mortality, and All-Cause Mortality. *J Nutr*. 2020;150(2):312-321. doi:10.1093/jn/nxz218

30. Zhang Y, Lu C, Li X, et al. Healthy Eating Index-2015 and Predicted 10-Year Cardiovascular Disease Risk, as Well as Heart Age. *Front Nutr*. 2022;9:888966. doi:10.3389/fnut.2022.888966

31. Blanton CA, Moshfegh AJ, Baer DJ, Kretsch MJ. The USDA Automated Multiple-Pass Method accurately estimates group total energy and nutrient intake. *J Nutr*. 2006;136(10):2594-2599. doi:10.1093/jn/136.10.2594

32. Ahluwalia N, Dwyer J, Terry A, Moshfegh A, Johnson C. Update on NHANES Dietary Data: Focus on Collection, Release, Analytical Considerations, and Uses to Inform Public Policy12. *Adv Nutr*. 2016;7(1):121-134. doi:10.3945/an.115.009258

33. Usual Dietary Intakes: Details of the Method | EGRP/DCCPS/NCI/NIH. Accessed December 19, 2023. https://epi.grants.cancer.gov/diet/usualintakes/details.html

34. Wingrove K, Lawrence MA, McNaughton SA. A Systematic Review of the Methods Used to Assess and Report Dietary Patterns. *Front Nutr*. 2022;9:892351. doi:10.3389/fnut.2022.892351

35. Schwingshackl L, Bogensberger B, Hoffmann G. Diet Quality as Assessed by the Healthy Eating Index, Alternate Healthy Eating Index, Dietary Approaches to Stop Hypertension Score, and Health Outcomes: An Updated Systematic Review and Meta-Analysis of Cohort Studies. *J Acad Nutr Diet*. 2018;118(1):74-100.e11. doi:10.1016/j.jand.2017.08.024

36. Krebs-Smith SM, Pannucci TE, Subar AF, et al. Update of the Healthy Eating Index: HEI-2015. *J Acad Nutr Diet*. 2018;118(9):1591-1602. doi:10.1016/j.jand.2018.05.021

37. Liese AD, Krebs-Smith SM, Subar AF, et al. The Dietary Patterns Methods Project: Synthesis of Findings across Cohorts and Relevance to Dietary Guidance. *J Nutr*. 2015;145(3):393-402. doi:10.3945/jn.114.205336

38. Panizza CE, Shvetsov YB, Harmon BE, et al. Testing the Predictive Validity of the Healthy Eating Index-2015 in the Multiethnic Cohort: Is the Score Associated with a Reduced Risk of All-Cause and Cause-Specific Mortality? *Nutrients*. 2018;10(4):452. doi:10.3390/nu10040452

39. Monteiro CA, Cannon G, Levy RB, et al. Ultra-processed foods: what they are and how to identify them. *Public Health Nutr*. 2019;22(5):936-941. doi:10.1017/S1368980018003762

40. Steele EM, O'Connor LE, Juul F, et al. Identifying and Estimating Ultraprocessed Food Intake in the US NHANES According to the Nova Classification System of Food Processing. *J Nutr*. 2023;153(1):225-241. doi:10.1016/j.tjnut.2022.09.001

41. National Health and Examination Survey. *Anthropometry Procedures Manual*.; 2017.

42. CDC. Defining Adult Overweight and Obesity. Centers for Disease Control and Prevention. June 3, 2022. Accessed January 27, 2024. https://www.cdc.gov/obesity/basics/adult-defining.html

43. National Insitutes of Health, National Heart, Lung, and Blood Institute. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: The evidence report. *Obes Res*. 1998;6(S2):S51-S210. doi:10.1002/j.1550-8528.1998.tb00690.x

44. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the Management of Blood Cholesterol: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. doi:10.1161/CIR.0000000000000625

45. Miller M, Stone NJ, Ballantyne C, et al. Triglycerides and Cardiovascular Disease. *Circulation*. 2011;123(20):2292-2333. doi:10.1161/CIR.0b013e3182160726

46. American Diabetes Association. *Standards of Care in Diabetes—2023* Abridged for Primary Care Providers. *Clin Diabetes*. 2023;41(1):4-31. doi:10.2337/cd23-as01

47. McCullough ML, Chantaprasopsuk S, Islami F, et al. Association of Socioeconomic and Geographic Factors With Diet Quality in US Adults. *JAMA Netw Open*. 2022;5(6):e2216406. doi:10.1001/jamanetworkopen.2022.16406

48. Liu J, Mozaffarian D. Trends in Diet Quality Among U.S. Adults From 1999 to 2020 by Race, Ethnicity, and Socioeconomic Disadvantage. *Ann Intern Med*. 2024;177(7):841-850. doi:10.7326/M24-0190

49. Baraldi LG, Martinez Steele E, Canella DS, Monteiro CA. Consumption of ultra-processed foods and associated sociodemographic factors in the USA between 2007 and 2012: evidence from a nationally representative cross-sectional study. *BMJ Open*. 2018;8(3):e020574. doi:10.1136/bmjopen-2017-020574

50. Bickel G, Nord M, Price C, Hamilton W, Cook J. *Guide to Measuring Household Food Security, Revised 2000*. U.S. Department of Agriculture, Food and Nutrition Service; 2000.

51. Gregory CA, Coleman-Jensen A. *Food Insecurity, Chronic Disease, and Health Among Working-Age Adults*. U.S. Department of Agriculture, Economic Research Service; 2017.

52. Lee EK, Donley G, Ciesielski TH, et al. Health outcomes in redlined versus non-redlined neighborhoods: A systematic review and meta-analysis. *Soc Sci Med*. 2022;294:114696. doi:10.1016/j.socscimed.2021.114696

53. Madlala SS, Hill J, Kunneke E, Lopes T, Faber M. Adult food choices in association with the local retail food environment and food access in resource-poor communities: a scoping review. *BMC Public Health*. 2023;23(1):1083. doi:10.1186/s12889-023-15996-y

54. Seligman HK, Levi R, Adebiyi VO, Coleman-Jensen A, Guthrie JF, Frongillo EA. Assessing and Monitoring Nutrition Security to Promote Healthy Dietary Intake and Outcomes in the United States. *Annu Rev Nutr*. 2023;43(1):409-429. doi:10.1146/annurev-nutr-062222-023359

Figure 1: Proportion of diet quality categories using observed and usual HEI-2020 scores

Figure 2: HEI- 2020 component scores by diet quality categories, 2009-2018 NHANES

Table 1: Sociodemographic characteristics of US adults (≥20 years) stratified by diet quality categories using observed intake, NHANES 2009-2018[c]

	Overall N (weighted %)	Diet quality category			
		Very low (HEI 0-50) N (weighted %)	Low (HEI >50-60) N (weighted %)	Marginal (HEI >60-70) N (weighted %)	High (HEI >70-100) N (weighted %)
Age					
≤30 years	3917 (20.3)	2040 (51.6)	994 (25.3)	574 (15.1)	309 (8.0)
31-50 years	7396 (35.3)	3229 (42.8)	1949 (27.2)	1392 (18.7)	826 (11.2)
≥51 years	10855 (44.4)	3640 (33.5)	2945 (27.3)	2458 (23.1)	1812 (16.1)
Sex					
Male	10548 (48.1)	4616 (44.6)	2830 (27.1)	1943 (17.9)	1159 (10.4)
Female	11620 (51.9)	4293 (36.7)	3058 (26.7)	2481 (21.8)	1788 (14.9)
Race and ethnicity					
Hispanic	5331 (14.6)	1959 (40.9)	1488 (27.3)	1183 (20.8)	701 (11.0)
Non-Hispanic Black	4849 (11.3)	2176 (47.9)	1331 (27.5)	847 (16.5)	495 (8.0)
Non-Hispanic White	9219 (65.7)	3939 (40.0)	2376 (26.9)	1732 (20.0)	1172 (13.1)
Other[a]	2769 (8.4)	835 (33.0)	693 (24.9)	662 (22.7)	579 (19.4)
Birthplace					
United States	16599 (83.8)	7458 (42.7)	4407 (27.0)	2955 (18.8)	1779 (11.5)
Outside of the United States	5569 (16.2)	1451 (28.8)	1481 (26.4)	1469 (25.5)	1168 (19.2)
Educational attainment					
High school education or less	9723 (36.4)	4472 (49.4)	2615 (26.2)	1723 (16.4)	913 (8.0)
Any college or college graduate	12445 (63.6)	4437 (35.4)	3273 (27.3)	2701 (21.9)	2034 (15.5)
Employment status					

Employed	12172 (61.2)	5007 (41.3)	3225 (26.6)	2382 (19.9)	1558 (12.2)
Not employed	9996 (38.8)	3902 (39.2)	2663 (27.3)	2042 (20.0)	1389 (13.6)
Marital status					
Married or living with partner	13214 (63.1)	5000 (38.3)	3528 (26.9)	2786 (21.1)	1900 (13.7)
Single	8954 (36.9)	3909 (44.1)	2360 (26.8)	1638 (17.9)	1047 (11.1)
Household income					
≤200% federal poverty level	9690 (33.0)	4504 (48.7)	2533 (25.7)	1690 (16.6)	963 (9.0)
>200% federal poverty level	12478 (67.0)	4405 (36.4)	3355 (27.5)	2734 (21.6)	1984 (14.6)
Household food security		4405 (36.4)	3355 (27.5)	2734 (21.6)	1984 (14.6)
High/marginal food security	17534 (84.2)	6571 (37.8)	4745 (27.4)	3642 (20.9)	2576 (13.9)
Low/very low food security	4634 (15.8)	2338 (54.9)	1143 (24.1)	782 (14.6)	371 (6.3)

[a] Other included American Indiana or Alaskan Native, Asian, Native Hawaiian or Pacific Islander, and Other

[b] Household food security assessed using the U.S. Household Food Security Survey Module

Table 2: Multivariate-adjusted observed daily intakes of total energy, NOVA food groups, foods, and nutrients of interest stratified by diet quality categories (using observed HEI-2020 scores), NHANES 2009-2018

	Overall N	Diet quality category				P-linear trend
		Very low[a] Mean	Low[a] Mean	Marginal[a] Mean	High[a] Mean	
Total energy (kcal/day)	2085	2106[b]	2102[b]	2031[b]	1976	*<0.01*
Nova 1-2, Unprocessed/ minimally processed foods (% energy/day)	30.1	25.8[b]	32.4[b]	36.3[b]	41.4	*<0.01*
NOVA 4, Ultra-processed foods (% energy/day)	55.2	61.7[b]	53.0[b]	48.5[b]	42.5	*<0.01*
Foods (cup or oz eq per 1000 kcal)						
Fruits, excluding juice (cup eq/day)	0.7	0.3[b]	0.7[b]	1.1[b]	1.6	*<0.01*
Vegetables, excluding white potatoes (cup eq/day)	0.6	0.5[b]	0.6[b]	0.7[b]	1.0	*<0.01*
Refined grains (oz eq/day)	5.5	6.8[b]	5.7[b]	4.7[b]	3.7	*<0.01*
Whole grains (oz eq/day)	0.9	0.4[b]	0.9[b]	1.3[b]	2.1	*<0.01*
Red and processed meat (oz eq/day)	2.6	2.9[b]	2.5[b]	2.2[b]	1.6	*<0.01*
Seafood (oz eq/day)	0.6	0.4[b]	0.7[b]	1.4	1.7	*<0.01*
Nutrients per 1,000 calories						
Saturated fat (% energy/day)	26.3	29.1[b]	25.8[b]	22.1[b]	18.5	*<0.01*
Dietary fiber (g/day)	17.3	13.4[b]	17.4[b]	20.1[b]	25.5	*<0.01*
Sodium (mg/day)	3481	3663[b]	3528[b]	3307[b]	3007	*<0.01*
Potassium (mg/day)	2670	2249[b]	2676[b]	2907[b]	3221	*<0.01*
Calcium (mg/day)	967	908[b]	957[b]	982[b]	1049	*<0.01*
Added sugar (% energy/day)	16.4	20.3[b]	15.6[b]	13.2[b]	10.8	*<0.01*

[a] Values adjusted for age, sex, race and ethnicity, birthplace, educational attainment, employment status, marital status, household income, and household food security

[b] Significant difference compared to the high diet quality category

Table 3: Multivariate-adjusted adiposity and cardiometabolic markers stratified by diet quality categories (observed HEI-2020 scores), NHANES 2009-2018

| | Overall | Diet quality category | | | | P-linear trend |
		Very Low[a]	Low[a]	Marginal[a]	High[a]	
BMI (kg/m2)[c]	29.3	29.8 (0.1)[b]	29.2 (0.2)[b]	28.2 (0.2)[b]	27.4 (0.2)	<0.01
% with obesity[d]	*38.8*	*45.7 (8.4)[b]*	*41.5 (7.9)[b]*	*33.8 (7.2)[b]*	*27.3(6.1)*	*<0.01*
Waist circumference (cm)[c]	99.7	100.4 (0.3)[b]	98.6 (0.3)[b]	96.5 (0.4)[b]	94.0 (0.5)	<0.01
% with high WC[d]	*57.8*	*61.5 (16.4)[b]*	*60.2 (16.4)[b]*	*55.9 (16.6)*	*49.3 (16.1)*	*<0.01*
HDL cholesterol (mg/dL)[c]	53.6	50.7 (0.3)[b]	52.9 (0.4)[b]	54.1 (0.4)[b]	56.6 (0.5)	<0.01
% <40 mg/dL[d]	*18.6*	*24.1 (12.0)[b]*	*19.1 (10.4)*	*15.9(9.1)*	*13.5 (8.0)*	*<0.01*
LDL cholesterol (mg/dL)[c]	113.4	112.6 (0.7)[b]	111.6 (1.0)[b]	111.9 (1.0)[b]	107.8 (1.4)	*0.01*
% ≥100 mg/dL[d]	*62.8*	*63.1 (9.7)*	*63.1 (9.1)*	*65.3 (8.5)*	*61.6 (8.5)*	*0.14*
Triglycerides (mg/dL)[c]	120.4	123.7 (2.4)[b]	120.0 (2.0)[b]	118.4 (2.6)[b]	103.6 (2.6)	<0.01
% ≥150 mg/dL[d]	*23.2*	*25.7 (7.6)[b]*	*25.8 (7.3)[b]*	*23.9 (6.6)*	*18.6 (5.2)*	*<0.01*
Fasting glucose (mg/dL)[c]	106.8	107.1 (0.5)[b]	105.4 (0.6)[b]	106.1 (0.9)[b]	101.7 (0.8)	<0.01
% >100 mg/dL[d]	*42.9*	*47.7 (19.6)[b]*	*48.6 (19.1)[b]*	*44.6 (18.0)*	*43.0 (17.3)*	*<0.01*

[a] Values adjusted for age, sex, race and ethnicity, birthplace, educational attainment, employment status, marital status, household income, and household food security

[b] Significant difference compared to the high diet quality category

[c] Mean (Standard Error)

[d] Mean (Standard Deviation)