





1 **Association of Endovascular Thrombectomy Volume and Outcomes in Acute Ischemic**  
2 **Stroke: A National Inpatient Sample Study**

3  
4 Lane Fry, M.D.<sup>1</sup>; Aaron Brake, M.D.<sup>2</sup>; Cody Heskett, BS<sup>3</sup>; Frank A. De Stefano, D.O.<sup>4</sup>; Ari  
5 Williams, M.D.<sup>4</sup>; Nashaat Majo, BS<sup>3</sup>; Catherine Lei, BS<sup>3</sup>; Abdul-Rahman Alkiswani, BS<sup>3</sup>; Kevin  
6 Le, BS<sup>3</sup>; Adam G. Rouse, Ph.D, M.D.<sup>4</sup>; Jeremy Peterson, M.D.<sup>4</sup>; Koji Ebersole M.D.<sup>4</sup>

7  
8 <sup>1</sup>Department of Radiology, University of Kansas, Kansas City, Kansas

9 <sup>2</sup>Department of Neurology, University of Iowa Hospital and Clinics, Iowa City, Iowa

10 <sup>3</sup>University of Kansas School of Medicine, Kansas City, Kansas

11 <sup>4</sup>Department of Neurological Surgery, University of Kansas, Kansas City, Kansas

12

13 **Corresponding Author:**

14

15 Lane Fry

16 University of Kansas Medical Center

17 Department of Radiology

18 3901 Rainbow Blvd, MS 4032

19 Kansas City, Kansas, 66160

20 Phone: 816-294-4328

21 Email: rfry@kumc.edu

22

23 **Running Title:** *EVT Volume in National Inpatient Sample*

24

25 **Keywords:** Endovascular Thrombectomy, Acute Ischemic Stroke, National Inpatient Sample,  
26 Volume-Outcome Relationship

27

28

29 **Abstract:** 300

30 **Word count:** 5187

31 **Figures/Tables:** 2 Figures, 4 Tables

32 **References:** 23

33

34 **Disclosures:** None

35 **Funding:** None

36

37 **Abstract**

38 **Background:** Previous studies have reported a positive relationship between higher hospital  
39 endovascular thrombectomy (EVT) volume and shorter procedures, higher revascularization  
40 rates, and improved functional outcomes. We investigated the association between hospital EVT  
41 volume and clinical outcomes using the National Inpatient Sample (NIS) database from 2016-  
42 2020.

43  
44 **Methods:** A cross-sectional analysis of the NIS examining the relationship of hospital EVT  
45 volume and outcomes was performed. All relevant clinical and demographic information was  
46 collected. The outcomes were favorable functional outcome (home without assistance), inpatient  
47 mortality, and intracerebral hemorrhage (ICH). Centers were classified as high-volume if they  
48 were in the top quintile of annual EVT volume. We performed univariate, multivariate, nearest  
49 neighbor matched analysis, and an exploratory annual case volume cutoff analysis.

50  
51 **Results:** There were 114,640 patients who underwent EVT included in the sample. Of these,  
52 24,415 (21.3%) were in the high-volume group. High-volume centers had higher rates of  
53 favorable functional outcome in univariate (OR 1.20,  $p < 0.001$ ), multivariate (aOR 1.19,  $p =$   
54  $0.003$ ), and matched analysis (OR 1.14,  $p = 0.028$ ). Prior to matching, lower rates of inpatient  
55 mortality (OR 0.83,  $p < 0.001$ ). However, in univariate and matched analysis there were no  
56 differences between high and low-volume centers. There were no differences in ICH across all  
57 analyses. Functional benefit was first noted at  $\geq 50$  EVTs, but centers performing  $\geq 175$  EVTs  
58 had substantially higher functional benefit (aOR 1.42,  $p = 0.002$ ).

59  
60 **Conclusions:** Our analysis demonstrates increased hospital case volume is associated with a  
61 modest improvement in favorable functional outcomes in patients undergoing EVT for AIS.  
62 Attempts to identify procedural cut off values reveal likely improved functional outcomes  
63 beginning at 50 EVT per year, while this benefit seems to increase with increasing case volumes.  
64 These higher levels of case volumes do not lead to higher rates of inpatient mortality or ICH.

65  
66

67 **Introduction:**

68 Endovascular thrombectomy (EVT) has emerged as the standard of care for patients presenting  
69 with acute ischemic stroke due to large vessel occlusion (LVO), demonstrating significant  
70 benefits in terms of revascularization, neurological recovery, and functional outcomes.<sup>1,2</sup> The  
71 successful implementation of EVT relies on well-coordinated stroke systems of care,  
72 incorporating streamlined processes for patient selection, timely intervention, and optimal post-  
73 procedural management. Previous studies have reported a positive relationship between higher  
74 hospital EVT volume and shorter procedure times, higher revascularization rates, and improved  
75 functional outcomes.<sup>3-8</sup> As EVT becomes increasingly integrated into routine clinical practice,  
76 understanding the impact of hospital case volume on patient outcomes is essential for optimizing  
77 stroke care.

78  
79 Two prior administrative studies have examined the volume-outcome relationship for EVT in  
80 acute ischemic stroke.<sup>3,4</sup> Each of these studies demonstrated benefit for patients at high-volume  
81 thrombectomy centers in terms of favorable functional outcome and inpatient mortality.  
82 However, the patient population in both studies were limited to only patients on Medicare and  
83 only encompassed data from 1-2 years.

84  
85 In this study, we sought to investigate the association between hospital EVT volume and clinical  
86 outcomes using the National Inpatient Sample (NIS) database from 2016-2020. The NIS is a  
87 comprehensive and nationally representative dataset of inpatient hospital stays in the United  
88 States, including all payers. Our objective was to provide robust, population-level evidence to  
89 inform ongoing discussions regarding the optimization of EVT delivery and stroke care  
90 networks. We hypothesized that higher volume centers would have higher rates of favorable  
91 functional outcome when compared to lower volume centers.

92

93 **Methods:**

94 *Study Design*

95 This is a cross-sectional retrospective analysis of United States hospital inpatient admissions  
96 examining the relationship of hospital EVT volume and outcomes. This study was conducted in  
97 accordance with STROBE guidelines.

98

99 *Data Source*

100 The NIS is the largest national inpatient database in the United States, containing 7 million  
101 unweighted hospital admissions annually. It is developed and maintained by the Healthcare Cost  
102 and Utilization Project (HCUP) and consists of de-identified inpatient hospitalization data  
103 derived from billing and discharge information from the participating hospitals. Utilization of  
104 provided discharge weights reported by participating institutions allows for estimates of  
105 nationally representative statistics.

106

107 *Patient Selection*

108 Weighted discharge NIS data from 2016-2020 was surveyed for patients admitted with primary  
109 *International Classification of Disease, Tenth Version*, (ICD-10) admission diagnosis of AIS  
110 (cerebral infarction - ICD-10-CM I63.xx). Endovascular thrombectomy (EVT) (ICD-10-PR  
111 03CGxx, 03CHxx, 03CJxx, 03CKxx, 03CLxx, 03CPxx, 03CQxx, 037Gxx, 03Qxx) and  
112 intravenous tPA (IV-tPA) (ICD-10-CM Z92.82, ICD-10-PR 3E03316, 3E03317, 3E04316,

113 3E04317) procedural codes were then interrogated to determine treatment exposure. Patients  
114 were included if they were admitted for AIS and received EVT. An overview of patient selection  
115 is seen in **Supplemental Figure 1**.

116

### 117 *Demographic and Clinical Characteristics*

118 Baseline demographic data extracted from the NIS database was collected including age, sex,  
119 race, primary payer status, hospital location, and teaching status. Comorbidity burden was  
120 assessed by utilizing the Charlson comorbidity index.<sup>9,10</sup> We identified markers of stroke severity  
121 based on previous NIS studies analyzing EVT outcomes.<sup>11,12</sup> Markers of stroke severity included  
122 ICD-10 diagnoses related to level of consciousness (stupor and coma), infarct size (cerebral  
123 edema and herniation), neurological deficits (neglect, dysphagia, and aphasia), and reliance upon  
124 respiratory or nutritional support (mechanical ventilation, nasogastric tube, and parenteral  
125 nutrition). Additionally, the proportion of total AIS admissions undergoing EVT were collected.

126

### 127 *Outcome Measures*

128 The primary clinical end point of this study was favorable functional outcome, defined as a  
129 routine discharge to home without assistance. This discharge disposition has been demonstrated  
130 to have a strong correlation with mRS  $\leq 2$  at 90 days.<sup>12-14</sup> Secondary outcomes were the rates of  
131 ICH and inpatient mortality between the treatment groups. Additionally, we investigated all  
132 discharge destinations (Short-Term Rehab Hospital, Skilled Nursing/Intermediate Care Facility,  
133 Home Health Care) available in the NIS for each group. The length of stay (LOS) and the total  
134 charges from the admission were also analyzed.

135

### 136 *Determination of EVT Volume Status and Hospital Characteristics*

137 Utilizing the NIS HOSP\_NIS variable we counted the number of EVT procedures performed at  
138 each unique hospital for each year included in the analysis. High-volume centers were defined as  
139 those being in the top 20<sup>th</sup> percentile of EVT volume. This volume determination has been used  
140 prior in other national database analysis.<sup>15,16</sup>

141

142 To further explore potential volume cut-off values where outcomes were significantly improved,  
143 we used similar methodology as Stein et al.<sup>4</sup> We compared favorable functional outcomes and  
144 inpatient mortality measures between increasing intervals of EVT procedures performed at a  
145 hospital. We started at a volume of 25 EVTs and increased by 25 EVTs until we reached a  
146 threshold of 300 EVTs performed. Once we discovered statistical differences between the high  
147 and low-volume groups, we utilized increments of 5 EVTs to determine a more precise cutoff  
148 point.

149

### 150 *Propensity Score Matching*

151 Patients were matched 1:1 between high and low-volume centers utilizing nearest neighbor  
152 matching using logit distance. Patients were matched based on demographic (age, sex, minority  
153 status, and location/teaching status of the hospital), comorbidity burden (Charlson Comorbidity  
154 Index, Hypertension, Intracranial Atherosclerotic Disease (ICAD), Atrial Fibrillation, and  
155 Diabetes), stroke severity (as described above), and IV-tPA status.

156

### 157 *Statistical Analysis*

158 Statistical analyses were performed accounting for the sampling design of the NIS, with  
159 appropriate strata, weights, and clusters according to HCUP guidelines. Statistical analyses were  
160 conducted with RStudio (Posit Software, Boston, MA, USA). Propensity matching was  
161 performed with the “MatchIt” package.<sup>17</sup> All complex survey samples, including regression  
162 analysis and statistical tests, were performed with the “Survey” package.<sup>18</sup> Survey weighted t-  
163 tests and chi-squared analysis were performed to compare means and association of categorical  
164 variables, respectively. Univariate and multivariate models were used to compare the high-  
165 volume group to the low-volume group with all outcome measures and discharge dispositions.  
166 Variables with a significance level  $< 0.1$  were included in multivariate analysis. This was done  
167 with an unmatched cohort and then again with the matched cohort model. We assessed the trends  
168 of the number of EVT<sub>s</sub> performed nationwide and the number of centers offering EVT during  
169 the study period utilizing a Pearson’s correlation coefficient. We assumed the number of centers  
170 was distributed as a Poisson random variable, and thus performed a square root transformation to  
171 approximate a normal distribution before calculating the correlation. An alpha threshold of 0.05  
172 was selected to define significance, with all p-values being 2-sided.

173

## 174 **Results:**

175 Centers grouped in the top quintile of EVT volume performed  $\geq 115$  EVT<sub>s</sub> in a given year.  
176 Centers performing  $< 115$  EVT<sub>s</sub> in a given year were analyzed as the low-volume cohort. After  
177 aggregating the number of centers in individual years, there were 158 high-volume centers and  
178 2,580 low-volume centers during the study period. The total number of centers performing EVT  
179 increased over time ( $r = 0.960$ ,  $p = 0.009$ ). The numbers of high-volume ( $r = 0.961$ ,  $p = 0.009$ )  
180 and low-volume centers ( $r = 0.952$ ,  $p = 0.013$ ) both increased during the study period. The  
181 number of EVT<sub>s</sub> performed increased at both high ( $r = 0.963$ ,  $p = 0.008$ ) and low-volume centers  
182 ( $r = 0.988$ ,  $p = 0.002$ ). The trend of centers and number of EVT<sub>s</sub> performed is seen in **Figure 1**.

183

184 Before propensity score matching, there were 114,640 patients who underwent EVT included in  
185 the sample. Of these, 24,415 (21.3%) were in the high-volume group. An overview of patient  
186 groups is seen in **Supplemental Figure 1**. The mean ( $\pm$  SD) number of EVT<sub>s</sub> performed at high-  
187 volume centers was 166 ( $\pm 49$ ), while there was an average of 55 ( $\pm 28$ ) EVT<sub>s</sub> performed at low-  
188 volume centers. There were higher rates of IV-tPA at low-volume centers (36.73% vs. 32.77%,  $p$   
189  $< 0.001$ ). High-volume centers had lower rates of atrial fibrillation (41.80% vs. 44.95%,  $p$   
190  $< 0.001$ ) and younger patients (68.66 vs. 69.34,  $p = 0.022$ ). These findings are summarized in  
191 **Table 1**.

192

193 After nearest neighbor matching, there were 24,415 patients in each group. There were no  
194 statistical differences in age, sex, comorbid conditions, utilization of tPA, or markers of stroke  
195 severity in the matched groups. The nearest neighbor matching adequately matched  
196 comorbidities, markers of stroke severity, and utilization of IV-tPA. High-volume centers  
197 performed EVT in a higher proportion of stroke patients than low-volume centers (19% vs 12%,  
198  $p < 0.001$ ). The results after matching are summarized in **Table 2**.

199

200 Prior to matching, the high-volume centers had higher rates of favorable functional outcome, as  
201 defined by discharge to home without assistance (22.18% vs. 19.19%; OR 1.20, 95% CI [1.08-  
202 1.33];  $p < 0.001$ ) and lower rates of inpatient mortality (10.49% vs. 12.44%; OR 0.83, 95% CI  
203 [0.74-0.92];  $p < 0.001$ ), but there was no difference in rates of ICH (18.53% vs. 17.77%; OR

204 1.05, 95% CI [0.95-1.17];  $p = 0.340$ ). After matching, univariate analysis revealed high-volume  
205 centers outperformed low-volume centers in terms of favorable functional outcome (22.18% vs.  
206 19.97%; OR 1.14, 95% CI [1.01-1.29];  $p = 0.028$ ). There was no difference in inpatient mortality  
207 (10.49% vs. 10.73%; OR 0.97, 95% CI [0.85 - 1.12];  $p = 0.709$ ) or rates of ICH (18.53% vs.  
208 17.55%; OR 1.07, 95% CI [0.95-1.21];  $p = 0.287$ ) after matching. Multivariate analysis of  
209 unmatched data demonstrated a persistent benefit for favorable functional outcome for the high-  
210 volume group (aOR 1.19 95% CI [1.06 - 1.34],  $p = 0.003$ ) with no differences in inpatient  
211 mortality and ICH. Examining discharge dispositions, there were no differences in all other  
212 discharge destinations (Rehab, Skilled Nursing Facility, or Home with assistance) in univariate  
213 or multivariate models. There were no differences in LOS before and after matching. However,  
214 the mean total charges for patients at high-volume centers was significantly less in both the  
215 unmatched (\$180,336.91 vs. \$208,360,  $p < 0.001$ ) and matched (\$180,336.91 vs. \$198,974.41,  $p$   
216 = 0.013) cohorts. All outcome models are summarized in **Table 3**.

217  
218 In our exploratory volume cut-off analysis, a difference in the combined metric of favorable  
219 functional outcome and inpatient mortality manifested between 25 and 50 EVT's. A mortality  
220 benefit alone manifested at 45 EVT's while higher odds of favorable functional outcome alone  
221 manifested at 50 EVT's performed annually. The volume cut-off analysis is seen in **Figure 2**.

222  
223 There was a notable increase in the odds of favorable functional outcome in patients treated at  
224 centers performing  $\geq 175$  EVT's per year, identifying a group we labeled "super high-volume  
225 centers." This finding prompted an additional multivariate sub-analysis comparing three groups,  
226 low volume vs. high-volume vs. super high-volume centers. Compared to low-volume centers,  
227 super-high-volume centers had higher odds of favorable functional outcomes (aOR 1.42, 95% CI  
228 [1.13 - 1.78],  $p = 0.002$ ) while high-volume centers did not differ from the low-volume centers  
229 (aOR 1.09, 95% CI [0.97 - 1.22],  $p = 0.158$ ). Neither the super-high (aOR 0.91, 95% CI [0.75 -  
230 1.11],  $p = 0.341$ ) nor the high-volume (aOR 0.92, 95% CI [0.79 - 1.07],  $p = 0.291$ ) centers  
231 outperformed low-volume centers in terms of mortality. There was also no difference in rates of  
232 ICH between super-high (aOR 1.14, 95% CI [0.93-1.38],  $p = 0.209$ ) nor high-volume (aOR 0.99,  
233 95% CI [0.88-1.11],  $p=0.801$ ) centers compared to low-volume centers. These findings are  
234 summarized in **Table 4**.

## 235 236 **Discussion:**

237 An association between case volume and improved patient outcomes is well documented in the  
238 literature for percutaneous coronary intervention for acute coronary syndromes.<sup>19</sup> A similar  
239 finding has been demonstrated among centers performing mechanical thrombectomy for acute  
240 stroke.<sup>3,4</sup> For example, Stein et al., using Medicare/Medicaid data, estimated the annual hospital  
241 volume for outcome benefit and mortality benefit were 26.2 and 44.8, respectively.<sup>4</sup> In another  
242 analysis of CMS data, Waqas et al. demonstrated that hospital EVT volume was associated with  
243 lower mortality and higher discharge to home rates, identifying the two central quartiles of  
244 mortality and discharge to home rates to be 24 and 23 respectively.<sup>3</sup> Accordingly,  
245 neuroendovascular certification and training standards reflect the intuitive understanding that a  
246 relationship between case volume and outcomes likely exists. For example, as of 2021, The Joint  
247 Commission recommends 15 thrombectomies per year or 30 over 2 years to meet criteria for  
248 Thrombectomy-capable Stroke Center (TSC) or a Comprehensive Stroke Center (CSC)  
249 certification. The Society of Neurological Surgeons (SNS) and its Committee for Advanced



250 Subspecialty Training (CAST) has defined a minimum number of cases performed for  
251 competency in neuroendovascular surgery including at least 30 acute ischemic stroke  
252 interventions.<sup>20</sup>

253  
254 Consistent with prior studies and certification standards, our analysis of the NIS database also  
255 finds that higher hospital thrombectomy case volume is positively associated with higher rates of  
256 favorable functional outcome. Unlike prior studies, our analysis includes all patients who  
257 received EVT from 2016-2020, as reported in the database, regardless of payor source, yielding a  
258 larger sample size than previous studies and one that more closely reflects real-world  
259 populations. In our analysis, favorable functional outcome was defined as routine discharge to  
260 home without assistance, which strongly correlates to an mRS  $\leq 2$ .<sup>12-14</sup> Thus the present study  
261 analyzes patients achieving the very best outcomes after intervention. The positive association  
262 between case volume and favorable functional outcome was persistent across all analyses. The  
263 benefit first manifests at  $\geq 50$  EVTs and demonstrates increasing benefit with higher annual case  
264 volume. In a sub-analysis however, this effect appears driven by the outcomes achieved by  
265 super-high-volume centers, those performing  $\geq 175$  EVTs in a year. Centers performing  $\geq 175$   
266 EVTs annually outperformed lower volume centers in achieving very favorable outcomes by 6-  
267 7%. Further, the super-high volume high-volume centers referred a higher proportion of patients  
268 to mechanical thrombectomy, exposing more candidates to the benefit. The functional benefit is  
269 persistent after successful matching of patient groups and utilizing multivariate logistic  
270 regression. While a broader definition of “benefit” would undoubtedly yield a lower case volume  
271 requirement, perhaps even substantially so, our analysis suggests that the likelihood of achieving  
272 a very favorable outcome after mechanical thrombectomy is best at the very highest-volume  
273 centers.

274  
275 Consistent with prior studies, our analysis finds that mechanical thrombectomy is safe. In  
276 unmatched, univariate analysis, higher volume centers had lower rates of inpatient mortality, but  
277 this difference was not observed after matching nor in adjusted analyses. In matched univariate  
278 analysis, super-high-volume centers demonstrated a mortality benefit compared to low-volume  
279 centers, but the difference was not observed in matched multivariate analysis. Meanwhile, there  
280 was no identifiable association between rates of ICH and thrombectomy volume status in any  
281 analysis. In other words, our analysis demonstrates that low volume centers achieved the same  
282 safety outcomes as even the super-high-volume centers. Viewed another way, our study finds  
283 that the super-high-volume centers matched the safety outcomes of lower-volume centers,  
284 despite a higher proportion of stroke patients being referred from thrombectomy, and more  
285 patients treated outside the tPA window.

286  
287 Lastly, our analysis identified an apparent cost savings for patients treated at higher volume  
288 centers. Existing data reporting EVT procedure volume and incurred cost per patient does not  
289 exist to this point. Previous publications have only examined patient outcomes as a function of  
290 hospital volume.<sup>3-5</sup> Our matched analysis demonstrates a nearly \$18,000 savings per patient, on  
291 average, when treated at a high-volume center. Similar findings have been seen in other  
292 procedural disciplines with high-volume centers more likely to have lower costs and improved  
293 outcomes.<sup>21,22</sup>

294  
295 **Limitations:**

296 This study is primarily limited by the inherent nature of sampling from a large national database  
297 and retrospective study design. The NIS collects a sample of discharged patient data from each  
298 hospital partaking in the HCUP consortium. As a sample driven database, the NIS is not able to  
299 give exact numbers of cases per institution. This means that our value estimation for cutoff data  
300 is based on hospital sample data and is likely under-estimating the true number of cases  
301 performed by each hospital. This has significant implications in terms of numeric derivatives  
302 (such as cutoff values for outcomes). However, as a sample database we are still able to reliably  
303 group hospitals into quintiles based on relative case volume. The NIS is an administrative  
304 database, reliant on coding via the ICD-10 system. Patients that may not represent our intended  
305 sample population may inadvertently be included due to coding errors. While it is less likely that  
306 coding mistakes may occur for inpatient mortality and disposition status, our sample is not  
307 immune from potential errors. Additionally, the HOSP\_NIS variable does not carry over from  
308 year-to-year, so we are unable to assess if low-volume centers are increasing their volumes over  
309 time. Further, differing stroke triage systems exist and may alter the outcomes for stroke patients,  
310 but our study is not equipped to examine the effects of these systems as it relates to patient  
311 outcomes.<sup>23</sup>

312  
313 **Conclusion:**

314 Our analysis demonstrates increased hospital case volume is associated with a modest  
315 improvement in favorable functional outcomes in patients undergoing EVT for AIS. Attempts to  
316 identify procedural cut off values reveal likely improved functional outcomes beginning at 50  
317 EVT per year, while this benefit seems to increase with increasing case volumes. These higher  
318 levels of case volumes do not lead to higher rates of inpatient mortality or ICH. Higher volume  
319 centers were also associated with decreased costs per patient. These findings further validate the  
320 relationship between procedural volume and outcomes and should assist in construction of  
321 policies of optimal stroke-care management and credentialing of potential comprehensive stroke  
322 centers.

323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336

337 **References**

- 338 1. Badhiwala JH, Nassiri F, Alhazzani W, et al. Endovascular Thrombectomy for Acute  
339 Ischemic Stroke: A Meta-analysis. *JAMA*. 2015;314(17):1832. doi:10.1001/jama.2015.13767
- 340 2. Zhongxing Y, Zhiqiang L, Jiangjie W, et al. Efficacy and Safety of Endovascular Treatment  
341 for Acute Large-Vessel Ischemic Stroke Beyond 6 h After Symptom Onset: A Meta-  
342 Analysis. *Front Neurol*. 2021;12:654816. doi:10.3389/fneur.2021.654816
- 343 3. Waqas M, Tutino VM, Cappuzzo JM, et al. Stroke thrombectomy volume, rather than stroke  
344 center accreditation status of hospitals, is associated with mortality and discharge  
345 disposition. *J NeuroIntervent Surg*. 2023;15(3):209-213. doi:10.1136/neurintsurg-2021-  
346 018079
- 347 4. Stein LK, Mocco J, Fifi J, Jette N, Tuhim S, Dhamoon MS. Correlations Between Physician  
348 and Hospital Stroke Thrombectomy Volumes and Outcomes: A Nationwide Analysis.  
349 *Stroke*. 2021;52(9):2858-2865. doi:10.1161/STROKEAHA.120.033312
- 350 5. Shim DH, Kim Y, Roh J, et al. Hospital Volume Threshold Associated with Higher Survival  
351 after Endovascular Recanalization Therapy for Acute Ischemic Stroke. *J Stroke*.  
352 2020;22(1):141-149. doi:10.5853/jos.2019.00955
- 353 6. Kim BM, Baek JH, Heo JH, Kim DJ, Nam HS, Kim YD. Effect of Cumulative Case Volume  
354 on Procedural and Clinical Outcomes in Endovascular Thrombectomy. *Stroke*.  
355 2019;50(5):1178-1183. doi:10.1161/STROKEAHA.119.024986
- 356 7. Gupta R, Horev A, Nguyen T, et al. Higher volume endovascular stroke centers have faster  
357 times to treatment, higher reperfusion rates and higher rates of good clinical outcomes. *J*  
358 *NeuroIntervent Surg*. 2013;5(4):294-297. doi:10.1136/neurintsurg-2011-010245
- 359 8. Rinaldo L, Brinjikji W, Rabinstein AA. Transfer to High-Volume Centers Associated With  
360 Reduced Mortality After Endovascular Treatment of Acute Stroke. *Stroke*. 2017;48(5):1316-  
361 1321. doi:10.1161/STROKEAHA.116.016360
- 362 9. Prommik P, Tootsi K, Saluse T, Strauss E, Kolk H, Märtson A. Simple Excel and ICD-10  
363 based dataset calculator for the Charlson and Elixhauser comorbidity indices. *BMC Med Res*  
364 *Methodol*. 2022;22(1):4. doi:10.1186/s12874-021-01492-7
- 365 10. Sundararajan V, Henderson T, Perry C, Muggivan A, Quan H, Ghali WA. New ICD-10  
366 version of the Charlson comorbidity index predicted in-hospital mortality. *Journal of*  
367 *Clinical Epidemiology*. 2004;57(12):1288-1294. doi:10.1016/j.jclinepi.2004.03.012
- 368 11. Patel PD, Salwi S, Liles C, et al. Creation and Validation of a Stroke Scale to Increase Utility  
369 of National Inpatient Sample Administrative Data for Clinical Stroke Research. *J Stroke*  
370 *Cerebrovasc Dis*. 2021;30(4):105658. doi:10.1016/j.jstrokecerebrovasdis.2021.105658
- 371 12. Dicipinigaitis AJ, Dick-Godfrey R, Gellerson O, et al. Real-World Outcomes of  
372 Endovascular Thrombectomy for Basilar Artery Occlusion: Results of the BAronIS Study.  
373 *Ann Neurol*. Published online March 10, 2023. doi:10.1002/ana.26640
- 374 13. ElHabr AK, Katz JM, Wang J, et al. Predicting 90-day modified Rankin Scale score with  
375 discharge information in acute ischaemic stroke patients following treatment. *BMJ Neurol*  
376 *Open*. 2021;3(1):e000177. doi:10.1136/bmjno-2021-000177
- 377 14. Qureshi AI, Chaudhry SA, Sapkota BL, Rodriguez GJ, Suri MFK. Discharge destination as a  
378 surrogate for Modified Rankin Scale defined outcomes at 3- and 12-months poststroke  
379 among stroke survivors. *Arch Phys Med Rehabil*. 2012;93(8):1408-1413.e1.  
380 doi:10.1016/j.apmr.2012.02.032
- 381 15. Koester SW, Cole TS, Kimata AR, et al. Assessing the volume-outcome relationship of  
382 carotid artery stenting in nationwide administrative data: a challenge of patient population

- 383 bias. *J Neurointerv Surg*. 2023;15(e2):e305-e311. doi:10.1136/jnis-2022-019695
- 384 16. Leifer D, Fonarow GC, Hellkamp A, et al. Association Between Hospital Volumes and  
385 Clinical Outcomes for Patients With Nontraumatic Subarachnoid Hemorrhage. *JAHA*.  
386 2021;10(15):e018373. doi:10.1161/JAHA.120.018373
- 387 17. Ho D, Imai K, King G, Stuart EA. MatchIt: Nonparametric Preprocessing for Parametric  
388 Causal Inference. *J Stat Soft*. 2011;42(8):1-28. doi:10.18637/jss.v042.i08
- 389 18. Lumley T. Analysis of Complex Survey Samples. *J Stat Soft*. 2004;9(8):1-19.  
390 doi:10.18637/jss.v009.i08
- 391 19. Saito Y, Inohara T, Kohsaka S, et al. Volume-Outcome Relations of Percutaneous Coronary  
392 Intervention in Patients Presenting With Acute Myocardial Infarction (from the J-PCI  
393 Registry). *Am J Cardiol*. 2023;192:182-189. doi:10.1016/j.amjcard.2023.01.027
- 394 20. Day AL, Siddiqui AH, Meyers PM, et al. Training Standards in Neuroendovascular Surgery:  
395 Program Accreditation and Practitioner Certification. *Stroke*. 2017;48(8):2318-2325.  
396 doi:10.1161/STROKEAHA.117.016560
- 397 21. Li LT, Chuck C, Bokshan SL, et al. High-Volume and Privately Owned Ambulatory  
398 Surgical Centers Reduce Costs in Achilles Tendon Repair. *Orthop J Sports Med*.  
399 2020;8(4):2325967120912398. doi:10.1177/2325967120912398
- 400 22. Campos S, Diniz P, Ferreira FC, Voiosu T, Arvanitakis M, Deviere J. Assessing the impact  
401 of center volume on the cost-effectiveness of centralizing ERCP. *Gastrointest Endosc*.  
402 Published online December 5, 2023:S0016-5107(23)03135-8. doi:10.1016/j.gie.2023.11.058
- 403 23. Raychev R, Sun JL, Schwamm L, et al. Performance of Thrombectomy-Capable,  
404 Comprehensive, and Primary Stroke Centers in Reperfusion Therapies for Acute Ischemic  
405 Stroke: Report from the Get With The Guidelines Stroke Registry: Stroke Outcomes Per  
406 Hospital Certification Status. *medRxiv*. Published online July 6, 2023:2023.07.05.23292270.  
407 doi:10.1101/2023.07.05.23292270
- 408

409 **Table 1.** Demographic, clinical, and outcome measures of unmatched cohort.  
410

	Overall n = 114,640	Low Volume Center Patients n = 90,225	High Volume Center Patients n = 24,415	p-value
<b>Demographics</b>				
Age (± SD, Range)	69.19 ± 14.44 (18.00-90.00)	69.34 ± 14.34 (18.00-90.00)	68.66 ± 14.79 (18.00-90.00)	0.022
Sex (% Female)	58,135 (50.71%)	45,770 (50.73%)	12,365 (50.65%)	0.911
<b>Location and Teaching Status</b>				
Rural	320 (0.28%)	320 (0.35%)	0 (0.00%)	0.097
Urban, Non-Teaching	8,930 (7.79%)	7,600 (8.42%)	1,330 (5.45%)	
Urban, Teaching	105,390 (91.93%)	82,305 (91.22%)	23,085 (94.55%)	
<b>Minority Status</b>				
White	78,930 (68.85%)	61,445 (68.10%)	17,485 (71.62%)	0.035
Non-White	35,710 (31.15%)	28,780 (31.90%)	6,930 (28.38%)	
<b>Payer Status</b>				
Medicaid	11,310 (9.88%)	8,940 (9.92%)	2,370 (9.73%)	<0.001
Medicare	71,145 (62.14%)	56,050 (62.19%)	15,095 (61.94%)	
No Charge	310 (0.27%)	270 (0.30%)	40 (0.16%)	
Other	2,410 (2.10%)	1,930 (2.14%)	480 (1.97%)	
Private Insurance	24,940 (21.78%)	19,810 (21.98%)	5,130 (21.05%)	
Self-Pay	4,380 (3.83%)	3,125 (3.47%)	1,255 (5.15%)	
<b>Comorbidities</b>				
Diabetes	33,465 (29.19%)	26,315 (29.17%)	7,150 (29.29%)	0.884
ICAD	19,835 (17.30%)	15,445 (17.12%)	4,390 (17.98%)	0.258
Hypertension	60,375 (52.66%)	47,650 (52.81%)	12,725 (52.12%)	0.469
Atrial Fibrillation	50,765 (44.28%)	40,560 (44.95%)	10,205 (41.80%)	<0.001
Charlson Comorbidity Score	4.26 ± 1.95 (1.00-16.00)	4.26 ± 1.96 (1.00-16.00)	4.26 ± 1.92 (1.00-14.00)	0.942
<b>Stroke Presentation</b>				
Intravenous tPA	41,140 (35.89%)	33,140 (36.73%)	8,000 (32.77%)	<0.001
Average Proportion of Strokes taken to EVT	13% ± 6% (1% - 50%)	11% ± 5% (1% - 50%)	19% ± 7% (6% - 49%)	<0.001
<b>Markers of Stroke Severity</b>				
Cerebral Edema	24,280 (21.18%)	19,095 (21.16%)	5,185 (21.24%)	0.947
Herniation	9,235 (8.06%)	7,000 (7.76%)	2,235 (9.15%)	0.026
Dysphagia	37,870 (33.03%)	28,525 (31.62%)	9,345 (38.28%)	<0.001
Aphasia	57,520 (50.17%)	44,970 (49.84%)	12,550 (51.40%)	0.135
Neglect	12,685 (11.07%)	9,475 (10.50%)	3,210 (13.15%)	0.003
Coma	1,915 (1.67%)	1,640 (1.82%)	275 (1.13%)	0.005
Stupor	270 (0.24%)	225 (0.25%)	45 (0.18%)	0.395
Nasogastric Feeding Tube	550 (0.48%)	470 (0.52%)	80 (0.33%)	0.153
Parenteral Nutrition	175 (0.15%)	150 (0.17%)	25 (0.10%)	0.299
Mechanical Ventilation	28,925 (25.23%)	23,650 (26.21%)	5,275 (21.61%)	<0.001
<b>Length of Stay (± SD, Range)</b>	8.53 ± 10.15 (0.00-289.00)	8.54 ± 10.29 (0.00-289.00)	8.53 ± 9.59 (0.00-155.00)	0.974
<b>Total Charges (± SD)</b>	202,400.99 ± 172,509.20	208,360.04 ± 179,263.16	180,336.91 ± 142,649.79	<0.001

**Outcomes**

Favorable Functional Outcome	22,725 (19.82%)	17,310 (19.19%)	5,415 (22.18%)	<0.001
Inpatient Mortality	13,780 (12.02%)	11,220 (12.44%)	2,560 (10.49%)	<0.001
Intracerebral Hemorrhage	20,560 (17.93%)	16,035 (17.77%)	4,525 (18.53%)	0.34

411  
412 **Abbreviations:** EVT, Endovascular Thrombectomy; ICAD, Intracranial Atherosclerotic Disease;  
413 SD, Standard Deviation.

414 **Table 2.** Demographic, clinical, and outcome measures after nearest neighbor matching.

	Overall n = 48,830	Low Volume Center Patients n = 24,415	High Volume Center Patients n = 24,415	p- value
<b>Demographics</b>				
Age (± SD, Range)	68.84 ± 14.68 (18.00-90.00)	69.02 ± 14.57 (18.00-90.00)	68.66 ± 14.79 (18.00-90.00)	0.295
Sex (% Female)	25,085 (51.37%)	12,720 (52.10%)	12,365 (50.65%)	0.129
<b>Location and Teaching Status</b>				
Rural	0 (0.00%)	0 (0.00%)	0 (0.00%)	> 0.999
Urban, Non-Teaching	2,635 (5.40%)	1,305 (5.35%)	1,330 (5.45%)	
Urban, Teaching	46,195 (94.60%)	23,110 (94.65%)	23,085 (94.55%)	
<b>Minority Status</b>				
White	35,245 (72.18%)	17,760 (72.74%)	17,485 (71.62%)	0.505
Non-White	13,585 (27.82%)	6,655 (27.26%)	6,930 (28.38%)	
<b>Comorbidities</b>				
Diabetes	14,165 (29.01%)	7,015 (28.73%)	7,150 (29.29%)	0.58
ICAD	8,495 (17.40%)	4,105 (16.81%)	4,390 (17.98%)	0.191
Hypertension	25,380 (51.98%)	12,655 (51.83%)	12,725 (52.12%)	0.8
Atrial Fibrillation	20,760 (42.51%)	10,555 (43.23%)	10,205 (41.80%)	0.182
Charlson Comorbidity Score (± SD, Range)	4.26 ± 1.95 (1.00-15.00)	4.25 ± 1.98 (1.00-15.00)	4.26 ± 1.92 (1.00-14.00)	0.864
<b>Stroke Presentation</b>				
Intravenous tPA	16,190 (33.16%)	8,190 (33.54%)	8,000 (32.77%)	0.52
Average Proportion of Strokes taken to EVT	15% ± 7% (1% - 49%)	12% ± 5% (1% - 33%)	19% ± 7% (6% - 49%)	<0.001
<b>Markers of Stroke Severity</b>				
Cerebral Edema	10,075 (20.63%)	4,890 (20.03%)	5,185 (21.24%)	0.312
Herniation	4,410 (9.03%)	2,175 (8.91%)	2,235 (9.15%)	0.741
Dysphagia	18,420 (37.72%)	9,075 (37.17%)	9,345 (38.28%)	0.467
Aphasia	25,085 (51.37%)	12,535 (51.34%)	12,550 (51.40%)	0.959
Neglect	6,560 (13.43%)	3,350 (13.72%)	3,210 (13.15%)	0.593
Coma	540 (1.11%)	265 (1.09%)	275 (1.13%)	0.858
Stupor	75 (0.15%)	30 (0.12%)	45 (0.18%)	0.43
Nasogastric Feeding Tube	145 (0.30%)	65 (0.27%)	80 (0.33%)	0.617
Parenteral Nutrition	50 (0.10%)	25 (0.10%)	25 (0.10%)	>0.999
Mechanical Ventilation	10,220 (20.93%)	4,945 (20.25%)	5,275 (21.61%)	0.265
<b>Length of Stay (± SD, Range)</b>	8.36 ± 9.11 (0.00-155.00)	8.20 ± 8.61 (0.00-147.00)	8.53 ± 9.59 (0.00-155.00)	0.153
<b>Total Charges (± SD)</b>	189,654.70 ± 153,334.78	198,974.41 ± 162,806.47	180,336.91 ± 142,649.79	0.013
<b>Outcomes</b>				
Favorable Functional Outcome	10,290 (21.07%)	4,875 (19.97%)	5,415 (22.18%)	0.028
Inpatient Mortality	5,180 (10.61%)	2,620 (10.73%)	2,560 (10.49%)	0.709
Intracerebral Hemorrhage	8,810 (18.04%)	4,285 (17.55%)	4,525 (18.53%)	0.287

415 **Abbreviations:** EVT, Endovascular Thrombectomy; ICAD, Intracranial Atherosclerotic Disease;  
 416 SD, Standard Deviation.

417

418 **Table 3. Univariate, Multivariate, and Matched Logistic Regression Models of Outcome**  
 419 **Measures and Discharge Dispositions comparing High Volume Centers to Low Volume Centers.**

			Odds Ratios (95% Confidence Interval)					
	Low Volume Center Patients	High Volume Center Patients	Unmatched Univariate	p-value	Matched Univariate	p-value	Unmatched Multivariate*	p-value
Favorable Functional Outcome (High vs. Low Volume)	17,310 (19.19%)	5,415 (22.18%)	1.20 (1.06 - 1.33)	< 0.001	1.14 (1.01 - 1.29)	0.028	1.19 (1.06 - 1.34)	0.003
Inpatient Mortality (High vs. Low Volume)	11,220 (12.44%)	2,560 (10.49%)	0.83 (0.74 - 0.92)	< 0.001	0.97 (0.85 - 1.12)	0.709	0.92 (0.81 - 1.04)	0.184
Intracerebral Hemorrhage (High vs. Low Volume)	16,035 (17.77%)	4,525 (18.53%)	1.05 (0.95 - 1.17)	0.34	1.07 (0.95 - 1.21)	0.287	1.04 (0.93 - 1.15)	0.574
Short-term Rehab Hospital	3,020 (3.35%)	660 (2.70%)	0.80 (0.46 - 1.40)	0.439	0.93 (0.52 - 1.65)	0.798	0.87 (0.50 - 1.52)	0.629
Skilled Nursing Facility	49,630 (55.01%)	13,180 (53.98%)	0.96 (0.88 - 1.04)	0.328	0.92 (0.83 - 1.01)	0.08	0.92 (0.84 - 1.00)	0.053
Home Health	9,045 (10.02%)	2,600 (10.65%)	1.07 (0.95 - 1.21)	0.268	1.04 (0.90 - 1.20)	0.59	1.08 (0.96 - 1.22)	0.205

\*Adjusted for Age, Atrial Fibrillation, Intravenous tPA, Location and Teaching Status, Minority Status, Herniation, Dysphagia, Neglect, Coma, Mechanical Ventilation, and Proportion of Strokes treated with EVT at a specific Hospital

420

421



422 **Table 4.** Univariate and Multivariate Logistic Regression Models comparing Super-High Volume  
 423 vs. High Volume vs. Low Volume Centers.

424

	Low Volume Center Patients n = 90,225	High Volume Center Patients n = 16,635	Super-High Volume Center Patients n = 7,780	Odds Ratios (95% Confidence Interval)							
				Univariate High Volume	p- val	Univariate Super- High Volume	p- va lu e	Multivariat e* High- Volume	p- va lu e	Multivaria te* Super- High Volume	p- va lu e
Good Functional Outcome***	17,310 (19.19%)	3,370 (20.26%)	2,045 (26.29%)	1.07 (0.97 - 1.18)	0.1 73	1.50 (1.23 - 1.84)	0.00 1	1.08 (1.97 - 1.22)	0.15 8	1.42 (1.13 - 1.78)	0.00 2
Inpatient Mortality	11,220 (12.44%)	1,805 (10.85%)	755 (9.70%)	0.86 (0.76 - 0.97)	0.0 16	0.76 (0.62 - 0.93)	0.00 7	0.92 (0.79 - 1.07)	0.29 1	0.91 (0.75 - 1.11)	0.34 1
Intracerebral Hemorrhage	16,035 (17.77%)	3,010 (18.09%)	1,515 (19.47%)	1.02 (0.91 - 1.15)	0.7 08	1.12 (0.92 - 1.36)	0.26 4	0.99 (0.88 - 1.11)	0.80 1	1.14 (0.93 - 1.38)	0.20 9

\*Adjusted for Age, Atrial Fibrillation, Intravenous tPA, Location and Teaching Status, Minority Status, Herniation, Dysphagia, Neglect, Coma, Mechanical Ventilation, and Proportion of Strokes treated with EVT at a specific Hospital

\*\* All Odds Ratios in comparison to Low-Volume Centers

\*\*\* As defined by routine discharge to home without assistance

425

426

427 **Figure Legends**

428

429 **Figure 1.** Line plots demonstrating the increase in total number of endovascular thrombectomies  
430 (EVT) performed and total centers performing EVT in the United States from 2016 to 2020.

431

432 **Figure 2.** Line plots demonstrating between dichotomized volume thresholds and outcomes in  
433 multivariate models for hospital case volume. Models adjusted age, atrial fibrillation, intravenous  
434 tPA, location and teaching status, minority status, herniation, dysphagia, neglect, coma,  
435 mechanical ventilation, and proportion of strokes treated with endovascular thrombectomy at a  
436 specific hospital.

437 \* Denotes a statistically difference in adjusted odds ratios

438

439

440

441