Health coaching 2.0: redefining a core lifestyle medicine intervention through a structured,

BCT-based model

Short title: Health coaching 2.0: a BCT-based model

Authors and affiliations

Ari Langinkoski, Graduate Student of Health Sciences[1,2]*

Johanna Hämäläinen, MSc[1]

Krista Rapo, MSc[1]

Mikko Rinta, MSc, RD[3]

Pekka Aroviita, MD, PhD, DiplBLM[3]

Mika Venojärvi, PhD, Title of Docent[1]

[1] School of Medicine, Institute of Biomedicine: Sports and Exercise Medicine, University of

Eastern Finland, Kuopio, Finland

[2] Nordic Health Academy, Helsinki, Finland

[3] Terveystalo Oyj, Helsinki, Finland

* Corresponding author: Ari Langinkoski, University of Eastern Finland, Email:

arilan@student.uef.fi

Word count (main text only): [2,254]

Keywords: health coaching; behavior change; lifestyle medicine; digital health; preventive care

Abstract

Background:

Preventing non-communicable diseases (NCDs) is a global health priority, as they remain the

leading cause of mortality globally, responsible for over 40 million deaths each year. Health

coaching has demonstrated potential to improve lifestyle behaviors and health outcomes, yet its

practical application is limited by inconsistent definitions and training standards.

Objective:

This brief report evaluates the feasibility and preliminary effectiveness of a systematically

defined health coaching model—PrioMed®—that translates the coaching process into evidence-

based Behavior Change Techniques (BCTs), providing a structured and teachable framework for

healthcare delivery.

Methods:

A single-arm pilot study (n = 25) was conducted in private healthcare in Finland. Two registered

nurses without prior coaching experience were trained for three half-day sessions and used a 25-

page manual to deliver PrioMed® coaching across seven phone-based sessions over six months.

Participants completed baseline and follow-up surveys, including the PROMIS-10 Global Health

instrument. Paired-sample t tests and Cohen's d were used to assess changes in physical and

mental health T-scores.

Results:

Participants (mean age 61.9 years, 76% female) showed significant improvements in self-

reported health. Physical health increased by +5.72 T-points ($p < .001$, $d = 1.31$) and mental

health by +4.43 T-points ($p < .001$, $d = 0.83$), both indicating large effect sizes.

Conclusions:

Redefining health coaching through an explicit BCT framework appears feasible, teachable, and

capable of generating meaningful health improvements within limited coaching time. The

PrioMed® model offers a replicable approach for digital and nurse-led lifestyle interventions that

warrants further evaluation in controlled trials and more diverse real-world populations.

Introduction

Non-communicable diseases (NCDs) remain the leading cause of global mortality, accounting

for approximately 43 million deaths annually and 75% of non-pandemic deaths worldwide (1).

The majority of these conditions—such as cardiovascular diseases, type 2 diabetes, chronic

respiratory diseases, and certain cancers—are strongly associated with modifiable lifestyle

factors including diet, physical inactivity, alcohol use, smoking, and stress-related behaviors (2).

Addressing these determinants through effective, scalable, and sustainable interventions remains

a cornerstone of preventive health and lifestyle medicine.

Health coaching has emerged as one of the most promising methods for facilitating sustained

lifestyle change in both preventive and clinical contexts. Systematic reviews and meta-analyses

(3–5) indicate that health coaching can enhance self-efficacy, improve health behaviors, and

contribute to better clinical outcomes, often with favorable cost-effectiveness profiles (6).

However, despite its growing popularity, *health coaching* remains ambiguously defined across

the literature and professional practice. The concept has been variably operationalized—from

motivational interviewing and counseling approaches to comprehensive lifestyle medicine

interventions—resulting in considerable heterogeneity in both methodology and reported

outcomes (4, 5, 7).

This lack of a shared operational definition poses two key challenges. First, it impedes the

reproducibility and scientific validation of health coaching interventions. Second, it hinders the

ability to train health professionals consistently and efficiently. Without a structured and

evidence-based framework, coaching programs often rely on individual experience rather than

standardized methods, limiting their scalability within healthcare systems.

To address these challenges, we developed and tested a new coaching model—PrioMed® Health

Coaching—that systematically defines the coaching process using the Behavior Change

Technique (BCT) taxonomy v1 (8). The model is grounded in the six pillars of lifestyle medicine

(9) and integrates only those BCTs with established evidence of effectiveness (10). By

translating the traditionally abstract concept of "coaching" into a set of clearly defined,

teachable, and measurable behavior change components, the PrioMed® model aims to make the

coaching process both scientifically replicable and practically scalable.

In contrast to previous coaching programs that require extensive training, PrioMed® has been

specifically engineered for rapid integration into clinical practice. In this pilot, two nurses with

no prior coaching experience received concise theoretical and practical training and implemented

the intervention with their clients in a private healthcare environment. The training was delivered

by the Nordic Health Academy, in collaboration with the University of Eastern Finland and

Terveystalo Oyj, a large private healthcare provider. Supplementary data from a related study

(11) support the acceptability and feasibility of the PrioMed® training framework, demonstrating

that both coaches and clients viewed the integration of the health coaching into routine clinical

practice as practical and well-received.

From a digital health perspective, the PrioMed® model is particularly relevant because it

represents a structured, data-compatible coaching process that can be readily integrated with

digital monitoring tools and remote care pathways. Each phase of the coaching intervention—

goal setting, needs analysis, feedback, and review—corresponds to distinct, codifiable BCT

elements. This opens the door to future digital augmentation through wearable data, decision-

support algorithms, and artificial intelligence–assisted behavior tracking, enabling personalized

and continuous lifestyle support at scale.

The present study reports the findings from the initial feasibility and effectiveness evaluation of

the PrioMed® pilot program. Conducted in a real-world private healthcare setting, this pilot

explored whether health coaching, redefined through a structured BCT-based framework, could

produce measurable improvements in self-reported physical and mental health within a limited

intervention time. Specifically, the study aimed to (A) assess the feasibility of implementing the

PrioMed® model in practice, and (B) evaluate preliminary health outcomes following a six-

month intervention delivered by newly trained nurses.

By addressing long-standing issues in the conceptual clarity and standardization of health

coaching, the PrioMed® model seeks to provide a universal, evidence-based structure that

bridges the gap between lifestyle medicine theory and scalable digital health implementation.

Methods

Study design and setting

This single-arm, six-month pilot study evaluated the feasibility and preliminary effectiveness of

the PrioMed® Health Coaching model in a real-world private healthcare environment. The study

was conducted at Terveystalo Oyj, one of Finland's largest private healthcare providers, in

collaboration with the Nordic Health Academy and the University of Eastern Finland.

The study adhered to the Declaration of Helsinki and the ICMJE recommendations for research

involving human participants. Ethical approval for the study was obtained from The Regional

Medical Research Ethics Committee of Eastern Finland Collaborative Area and all participants

provided written informed consent prior to participation.

Participants and recruitment

Participants were recruited from Terveystalo's existing client base, including those who had

previously participated in an annual health check-up or a digital wellness program. Inclusion

criteria were age ≥18 years, adequate Finnish language skills, and willingness to participate in

remote coaching. Exclusion criteria included acute medical conditions or cognitive impairments

preventing informed participation.

A total of 25 participants completed both baseline and six-month follow-up assessments. The

demographic characteristics are presented in Table 1.

Intervention

The PrioMed® model operationalizes health coaching through evidence-based Behavior Change

Techniques (BCTs) drawn from the taxonomy by Michie et al. (8) and informed by the six pillars

of Lifestyle Medicine (9). It provides a structured and teachable framework emphasizing self-

assessment, needs analysis, goal setting, graded progression, and feedback.

Two registered nurses with no prior coaching experience were trained by the first author in three

four-hour sessions using a 25-page instructor manual. The training combined theoretical

instruction and practical exercises focusing on motivational communication, goal-oriented

dialogue, and structured session management.

Each participant received up to seven phone-based sessions across six months, combining an

initial assessment and goal-setting phase with five to six follow-up sessions focused on feedback

and reinforcement. Total contact time averaged less than four hours per participant.

A detailed description of the PrioMed® coaching process, including session structure, BCT

mapping, and sample tools (PrioMed® Lifestyle Medicine Vital Signs [PMLMVS] and

PrioMed® Needs Analysis [PMNA]), is provided in Supplementary Material 1.

Outcome measures

The primary outcomes were self-reported physical and mental health, measured using the

PROMIS-10 Global Health questionnaire, which produces standardized T-scores (mean = 50, SD

= 10; higher scores indicate better health). Surveys were completed electronically at baseline and

at six-month follow-up.

Secondary outcomes included demographic data and descriptive summaries of lifestyle self-

assessments (PMLMVS) and applied BCT frequencies recorded by the coaches.

Data analysis

PROMIS-10 raw scores were converted to T-scores using the *HealthMeasures Scoring Service*.

Data normality was verified with the Shapiro–Wilk test, and pre-post differences were analyzed

using paired-sample t-tests. Effect sizes were expressed as Cohen's d, using the standard

deviation of the mean difference as the denominator. The threshold for statistical significance

was $p < 0.05$ (two-tailed). Analyses were conducted in IBM SPSS Statistics v29.

Complementary qualitative findings related to training and user experience are presented in a

companion study (11).

Results

Participant characteristics

Twenty-five participants completed both baseline and six-month follow-up surveys.

Demographic characteristics are summarized in Table 1. Participants were predominantly older,

well-educated adults, mostly women, and nearly half were retired or not currently working.

Primary outcomes

Statistically significant improvements were observed in both physical and mental health domains

of the PROMIS-10 Global Health measure after six months of PrioMed® health coaching.

•   Physical Health T-score: Baseline mean = 45.51 ± 5.59; follow-up mean = 51.23 ± 5.04;

mean change = + 5.72 points (95 % CI 3.92–7.51; $p < .001$).

•   Mental Health T-score: Baseline mean = 50.20 ± 6.91; follow-up mean = 54.63 ± 5.47;

mean change = + 4.43 points (95 % CI 2.24–6.62; $p < .001$).

Effect sizes were large for both scales (Cohen's $d = 1.31$ for physical health; $d = 0.83$ for mental

health), suggesting clinically meaningful changes in perceived wellbeing.

Secondary observations

Analysis of coaching session documentation indicated high adherence to the intended PrioMed®

protocol. The most frequently used Behavior Change Techniques (BCTs) were:

• *Goal setting (behavior) – 100 % of participants*

• *Goal setting (outcome) – 88 %*

• *Self-monitoring of behavior – 68 %*

• *Social reward – 64 %*

• *Review behavior goals – 60 %*

• *Comparative imagining of future outcomes – 60 %*

• *Action planning – 56 %.*

Other BCTs were applied in fewer than half of the coaching relationships, reflecting the limited

duration and goal-oriented nature of the intervention. Average total contact time remained below

four hours per participant, confirming the intended efficiency of the model.

Feasibility and implementation findings

From an implementation standpoint, the PrioMed® model proved feasible for delivery by nurses

with no prior coaching experience. Both coaches successfully completed the training, adhered to

the structured protocol, and maintained consistent client engagement throughout the study. No

technical or logistical difficulties were reported in the telephone-based format.

Complementary qualitative findings, reported in a companion study (11), reinforce these

findings: Coachees characterized the sessions as supportive, personalized, and motivating, while

coaches reported marked improvements in their confidence and competence over the six-month

period.

Discussion

This pilot study evaluated the feasibility and preliminary effectiveness of a newly developed

health coaching model grounded in BCTs, within a real-world private healthcare setting. Despite

the brief duration and limited contact time, the intervention was associated with notable

improvements in participants' self-reported physical and mental health outcomes. These findings

indicate that a structured, behaviorally defined coaching framework can feasibly be implemented

and may contribute to enhanced subjective wellbeing, even when delivered with a minimal

resource investment.

Interpretation of main findings

The observed effect sizes in the present study are substantially larger than those typically

reported in lifestyle intervention meta-analyses (12, 13). From clinical perspective, changes in

this magnitude are typically regarded as meaningful, reflecting noticeable enhancement in

functional capacity and reduction symptom burden. The observed gains are particularly notable

considering the brevity of the intervention (< 4 h per participant) and the fact that coaches were

newly trained nurses with no prior experience in behavioral counseling. Together, these results

suggest that redefining health coaching through clearly operationalized BCTs can enhance both

its efficiency and trainability.

Documentation from the coaching sessions demonstrated high protocol adherence, with frequent

use of core BCTs such as goal setting, self-monitoring, and action planning. This supports the

fidelity of the intervention and reinforces its theoretical grounding.

Comparison with previous literature

Previous systematic reviews have identified health coaching as a potentially effective yet highly

heterogeneous intervention domain (3, 4, 5). Variability in definitions, training, and delivery

formats has hindered replication and large-scale implementation. By explicitly structuring the

PrioMed® model around empirically supported BCTs (8, 10), this study addresses a long-

standing gap in the literature: the need for a reproducible, evidence-linked framework that

preserves the person-centered ethos of coaching while enabling systematic evaluation.

The present results are consistent with emerging evidence showing that targeted, time-limited

coaching interventions can yield measurable health benefits, particularly when they include goal

setting, self-monitoring (12–14), and graded tasks (13). The structured yet autonomy-supportive

design of PrioMed® may explain the strong adherence observed among participants. Additional

qualitative insights from the companion study (11) further supports the feasibility of

intervention. Both coaches and coachees reported high levels of satisfaction and perceived the

sessions as relevant and beneficial, even delivered via remote modalities (11).

Practical implications and digital health relevance

From an implementation perspective, these findings support the integration of structured health

coaching models into both primary care and digital health ecosystems. The PrioMed®

framework—by defining the coaching process in modular, codifiable BCT units—creates a

foundation for digitally augmented coaching, in which wearable-derived data and algorithmic

decision support can inform personalized behavioral feedback loops. Such standardization also

enables quality assurance and competence-based training for health professionals, addressing a

key barrier to scaling lifestyle interventions in healthcare systems.

In practical terms, the model demonstrates that brief, remote coaching interactions can be

sufficient to produce meaningful improvements in perceived health status. This efficiency is

particularly relevant for preventive care models that seek to reach large populations at moderate

cost. The structured PrioMed® model further allows delegation of coaching to non-specialist

healthcare staff, such as nurses or health advisors, thereby expanding access while maintaining

methodological fidelity.

Within the broader digital health landscape, the PrioMed® approach aligns with current priorities

in data-driven, personalized preventive healthcare—an area increasingly supported by wearable

sensors, continuous monitoring, and AI-enabled behavior analytics. The BCT-based logic of the

model provides an interpretable bridge between behavioral science and digital system design,

facilitating future research on how human and automated coaching elements can complement

each other.

Strengths, limitations, and future directions

The main strengths of this pilot include its pragmatic design, use of a validated outcome measure

(PROMIS-10), and detailed documentation of coaching content using BCT taxonomy.

Conducting the study within an active private healthcare setting enhances ecological validity and

demonstrates feasibility under real-world conditions.

However, some limitations must be acknowledged. The study lacked a control group and

randomization, limiting causal inference. The relatively small, self-enrolled sample—

predominantly older, well-educated women—restricts generalizability. Furthermore, the

coaches' dual roles as healthcare employees may have introduced unmeasured contextual effects.

Outcomes were self-reported, and objective behavioral or physiological measures (e.g., activity

tracking, biometrics) were not included.

Future studies should therefore employ randomized controlled designs, larger and more diverse

samples, and multimodal outcome measures integrating both subjective and digital health data.

Longer-term follow-up would also clarify the sustainability of observed changes. Additional

research could evaluate the PrioMed® framework's integration with wearable technologies and

adaptive algorithms, as well as its cost-effectiveness in large-scale preventive health programs.

Conclusion

Overall, the pilot demonstrated that a standardized, BCT-based coaching model can be

effectively implemented in private healthcare, achieving meaningful improvements in health-

related quality of life with minimal time investment. The findings support continued

development of standardized and scalable coaching frameworks that can effectively integrate

principles of lifestyle medicine with emerging innovations in digital health.

Declarations

Funding

This research received no specific grant from any funding agency in the public, commercial, or

not-for-profit sectors.

Conflicts of Interest

The authors declare the following potential conflicts of interest with respect to the research,

authorship, and/or publication of this article:

•   *PrioMed®* is a trademark registered in Finland by Nordic Health Academy.

•   Ari Langinkoski is employed by Nordic Health Academy as Head of Education and is a

shareholder of the company.

•   Ari Langinkoski, Mikko Rinta, Pekka Aroviita, and Mika Venojärvi serve as members of the

Scientific Advisory Board of Nordic Health Academy.

•   Mikko Rinta is employed by Terveystalo as Lead Nutritionist and is a minority shareholder in

Nordic Health Academy.

•   Pekka Aroviita works as an independent private lifestyle medicine practitioner and is a minority

shareholder in Nordic Health Academy.

•   Ari Langinkoski is also Chair of the EuropeActiveTechnical Expert Group (TEG), which is

developing European standards to support the health and lifestyle coaching role.

All other authors declare no additional conflicts of interest.

Ethics approval and consent to participate

The study was approved by The Regional Medical Research Ethics Committee of Eastern

Finland Collaborative Area (approval number: [1121/13.02.00/2022]). All participants provided

written informed consent prior to enrollment, and the study was conducted in accordance with

the principles of the Declaration of Helsinki.

Consent for publication

Not applicable.

Author contributions

Acknowledgements

The authors thank the participating nurses and study participants for their valuable contributions.

The authors also acknowledge the use of OpenAI's ChatGPT (GPT-5, 2025) to assist in language

editing, reference formatting, and structural alignment with the *Digital Health* author guidelines.

The authors remain fully responsible for the content and accuracy of this manuscript.

Availability of data and materials

Deidentified participant data are available from the corresponding author upon reasonable

request. Supplementary materials describing the intervention are available as Supplementary

Material 1.

References

1. World Health Organization. Noncommunicable diseases. Geneva: WHO; 2025. Available

from: https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases

[accessed 2025 Oct 6].

2. Global Burden of Disease Collaborators. Global, regional, and national age-sex-specific

mortality for 282 causes of death, 1980–2019: a systematic analysis. Lancet

2020;396:1160–1204. doi:10.1016/S0140-6736(20)30925-9

3. Kivelä K, Elo S, Kyngäs H, et al. The effects of health coaching on adult patients with

chronic diseases: a systematic review. Patient Educ Couns 2014;97:147–157.

doi:10.1016/j.pec.2014.07.026.

4. Sherifali D, Viscardi V, Bai JW, Ali RMU. Evaluating the effect of a diabetes health

coach in individuals with type 2 diabetes. Can J Diabetes 2016;40(1):84–94.

doi:10.1016/j.jcjd.2015.10.006

5. Pirbaglou M, Katz J, Motamed M, Pludwinski S, Walker K, Ritvo P. Personal health

coaching as a type 2 diabetes mellitus self-management strategy: a systematic review and

meta-analysis of randomized controlled trials. Am J Health Promot 2018;32:1613–1626.

doi:10.1177/0890117118758234

6. Hale R, Giese J. Cost-effectiveness of health coaching: an integrative review. Prof Case

Manag 2017;22(5):228–238. doi:10.1097/NCM.0000000000000223

7. Mendes Sieczkowska S, Padilha de Lima A, Swinton PA, Dolan E, Roschel H, Gualano

B. Health coaching strategies for weight loss: a systematic review and meta-analysis. Adv

Nutr 2021;12:1449–1460. doi:10.1093/advances/nmaa159

8. Michie S, Richardson M, Johnston M, et al. The behavior change technique taxonomy
(v1) of 93 hierarchically clustered techniques: building an international consensus for the
reporting of behavior change interventions. Ann Behav Med 2013;46:81–95.
doi:10.1007/s12160-013-9486-6

9. American College of Lifestyle Medicine. About lifestyle medicine. Available from:
https://lifestylemedicine.org/about-lifestyle-medicine/ [accessed 2025 Jan 20].

10. Guerreiro M, Strawbridge J, Cavaco AM, Félix IB, Marques MM, Cadogan C.
Development of a European competency framework for health and other professionals to
support behaviour change in persons self-managing chronic disease. BMC Med Educ
2021;21:287. doi:10.1186/s12909-021-02720-w

11. Hämäläinen J, Rapo K, Langinkoski A, Rinta M, Aroviita P, Haaranen A, Kettinen J,
Venojärvi M. Health coaching as a tool for lifestyle changes: insights from the PrioMed®
training and pilot intervention. medRxiv. 2025. doi:10.1101/2025.11.15.25340170. Epub
ahead of print.

12. Ashton LM, Sharkey T, Whatnall MC, Haslam RL, Bezzina A, Aguiar EJ, Collins CE,
Hutchesson MJ. Which behaviour change techniques within interventions to prevent
weight gain and/or initiate weight loss improve adiposity outcomes in young adults? A
systematic review and meta-analysis of randomized controlled trials. *Obes Rev.*
2020;21(6):e13009. doi:10.1111/obr.13009

13. Carraça E, Encantado J, Battista F, Beaulieu K, Blundell J, Busetto L, van Baak M,
Dicker D, Ermolao A, Farpour-Lambert N, Pramono A, Woodward E, Bellicha A, Oppert
J-M. Effective behavior change techniques to promote physical activity in adults with

352 overweight or obesity: A systematic review and meta-analysis. *Obesity Reviews*

353 2021;22(Suppl 4):e13258. doi:10.1111/obr.13258

354 14. Samdal GB, Eide GE, Barth T, Williams G, Meland E. Effective behaviour change

355 techniques for physical activity and healthy eating in overweight and obese adults:

356 systematic review and meta-regression analyses. Int J Behav Nutr Phys Act.

357 2017;14(1):42. doi:10.1186/s12966-017-0494-y

359 **Table 1. Participants demographics**

Characteristic	Value
Participants, n	25
Age, mean ± SD (range)	61.9 ± 10.8 (33–75)
Female, n (%)	19 (76%)
Married or partnered, n (%)	18 (72%)
Retired or not currently working, n (%)	12 (48%)
Post-secondary or higher education, n (%)	23 (92%)

361