Spatial data analysis and geodemographics for the prevalence of diabetes among adult population in Bangladesh

Chowdhury Mashrur Mahdee[1*¶], Anika Tabashsum[1], Most. Jannatul Ferdous Asha[1], Md. Ashraful Islam Khan[1], Md. Aminur Rahman[1¶]

[1] Department of Population Science and Human Resource Development, University of Rajshahi, Rajshahi, Bangladesh

* Corresponding author

E-mail: mashrur.mahdee10@gmail.com

¶ These authors contributed equally to this work.

Abstract

Bangladesh, like most of the developing countries, is experiencing quite a substantial rise in the prevalence of noncommunicable diseases, like diabetes. For both men and women, the prevalence of diabetes has been increasing in recent decades. The objective of this study is to determine if there are statistically significant geospatial inconsistencies in the prevalence of diabetes. Cross-sectional and spatial analysis was concluded using data on 12100 adults aged 18 years and above from the Bangladesh Demographic and Health Survey (BDHS) 2017-18. The prevalence of diabetes is analyzed and visualized. Spatial autocorrelation was identified using Moran's I. Hotspot analysis was also done using Moran's I. Men had a higher prevalence of diabetes (10.52%) than women do (9.50%); yet, in the sample group, more women than men have diabetes. Among diabetic men and women, 37.11% (95% CI, 32.04-42.48) and 39.71% (95% CI, 35.47-44.12) respectively, are aware of their condition. Among them 33.29% men and 36.44% women are being treated for it. People living in Dhaka division (men: 15.62%, women: 13.31%) are more affected by diabetes than people living in other divisions. Increasing age, living in the highest wealth quintile, and obesity and being overweight are positively associated with the increasing prevalence of diabetes. There also exists positive spatial autocorrelation (Moran's I = 0.215, p=0.001). According to the Local Indicators of Spatial Association (LISA) cluster map, the hotspots are Dhaka and the central districts (10 districts), and the cold spots are the central of northside districts (7 districts). The findings draw attention to the advantages of spatial analysis in the healthcare system. And this research could help policymakers and healthcare organizations to plan and implement policies with the objective of attenuating the prevalence as well as the risk of diabetes in Bangladesh.

Introduction

A significant result of the 21st-century demographic and epidemiological shifts has been the "health transition," defined by an unprecedented increase in the prevalence of chronic illnesses among middle-aged and older adults, such as diabetes, cancer, hypertension, and so on[1,2]. Diabetes is a chronic illness in which there is a deviation from normal glucose tolerance, typically because the body requires more insulin than the pancreas is able to produce or manage. This may occur as a result of inadequate naturally occurring insulin production or because body cells become resistant to the effects of insulin[3,4]. Previously known as "adult-onset diabetes," it was once thought to be an illness that only affected older people. But it is currently the most prevalent metabolic disease in the world and a major cause

of death for young and middle-aged adults[5,6]. Despite the fact that there are many biological, behavioral, and

environmental risk factors, we still don't fully understand the reasons, both direct and indirect[7]. It is believed that

local surroundings can either facilitate or restrict individual choices for both nutritional intake and physical activity

patterns. Local environments are regarded to function as key factors of individualistic behaviors[8,9]. Such

surroundings affect people's vulnerability to unfavorable health consequences as well as their capacity to effectively

control it.

Globally, the estimated number of adults with diabetes in 2021 is 536.6 million, and by 2045, that number is

estimated to increase to 783.2 million[7]. By 2045, it is predicted that low and middle-income countries (LMIC),

where population growth is anticipated to be higher, would account for 94% of the growth in the number of individuals

with diabetes worldwide[7]. It is estimated that 22.3 million adults in Bangladesh will suffer from diabetes by 2045,

an increase from 13.1 million in 2021[7].

Geodemographics is the study of socioeconomic and behavioral information about individuals in relation to

location (i.e., geography) and neighborhood settlements[10]. Public health research uses geodemographic data and

related lifestyle segmentation methods somewhat rarely[11]. Geodemography's strength is in its ability to produce

fresh perspectives on spatial effects that may be researched and discussed in more depth[12]. So, it is an exploratory

method. The application of geodemographics as a tool for health informatics is still intriguing, despite the reasonable

criticisms of its flaws[13,14]. It is also an effective tool for evaluating health disparities at the local level[14]. The

social context and the environment in which risky behavior occurs influence many of the changeable risk factors for

diabetes, like sedentary behavior and food habits. Additionally, many of the risk variables that cannot be changed,

such as age, income, and education, have an impact on the regions where diabetes is most prevalent[15,16].

The purpose of this study is to assess the prevalence of diabetes, along with awareness, treatment, and control of

diabetes; as well as the factors connected with diabetes among Bangladeshi people aged 18 and above using the

Bangladesh Demographic and Health Survey (BDHS) 2017–18 data.

Methods

Data source

This cross-sectional study examined data from the BDHS for 2017–18[17]. Major health indicators for Bangladesh, such as fasting blood glucose (FBG) biomarker measurements, were reported by BDHS 2017–18. The survey was taken between October 2017 and March 2018. The stratified sample of households used for the survey was created in two stages. A sampling frame was created using enumeration areas (EAs) taken from the 2011 Bangladesh Population and Housing Census. An average of about 120 houses make up the survey's EA, or primary sampling unit (PSU). Initially, 675 EAs were selected, among which 425 and 250 were from rural and urban regions, respectively. Data were collected from 672 PSUs, and 3 PSUs were excluded due to flooding. A systematic sample of, on average, 30 households was selected from each EA in the second stage in order to produce precise estimates of the demographic and health characteristics for each of the eight divisions, for both urban as well as rural areas, and for the country as a whole. Out of the 20160 households chosen using this method, interviews were completed in 19457 households[17]. Out of these, 4864 households were chosen for the biomarker collection. In the selected 4864 households, a total of 14,704 respondents (8013 women, 6691 men), aged 18 years and older, were available for blood glucose measurement. A blood glucose test was administered to 12,100 respondents who were 18 years of age and older (6919 women and 5181 men)[17] (Fig 1).

Fig 1: Flowchart of study sample.

Outcome variables

The outcome variables included awareness, treatment, and control of diabetes. At the time of the survey, individuals were deemed to have diabetes if their fasting plasma glucose (FPG) equivalent level was 7 mmol/L or higher, or if they disclosed that they were currently taking a prescribed medicine to treat their high blood sugar[17]. Prior to the test, respondents were instructed to fast for at least eight hours and only consume plain water. Using the HemoCue Glucose 201 DM system, which has plasma conversion, a capillary blood sample from willing and qualified respondents was tested[17]. The measures of fasting whole blood glucose recorded throughout the survey were

85   automatically translated to corresponding fasting plasma glucose values. Respondents are considered to be aware of

86   their condition if they have previously known their glucose level and/or have been told they have diabetes by a

87   physician or nurse. Those who were receiving medication at the time of the survey are regarded as being under

88   treatment for diabetes. Participants who had their FPG reading at or below 7.0 mmol/L at the time of the survey were

89   considered to be in control of their diabetes[17].

Explanatory variables

Following explanatory variables were incorporated into the study: age (18-34, 35-39, 40-44, 45-49, 50-54, 55-59,

60-64 and 65 and above), sex (men, women), place of residence (urban, rural), division of residence (Barisal,

Chittagong, Dhaka, Khulna, Mymensingh, Rajshahi, Rangpur and Sylhet), level of education (no education, primary,

secondary, college or higher), wealth quintile (poorest, poorer, middle, richer, richest), nutritional status or BMI

(underweight, normal, overweight, obese), hypertension (yes, no), marital status (never married, married, widowed,

divorced), current employment status (yes, no)[17]. Principal component analysis was used to determine the wealth

quintile based on the participants' durable and nondurable household items. BMI was classified using the WHO-

recommended Asian cut-off[18]. Indicators of hypertension were a systolic blood pressure of ≥140 mmHg and/or a

diastolic blood pressure of ≥90 mmHg and/or currently using medication to treat hypertension [17].

Statistical analysis

101   For the entire study population as well as population subgroups, the crude diabetes prevalence was estimated.

102   Prevalence was estimated taking into consideration the weighted sample and complex survey design. Crude prevalence

103   estimates of diabetes awareness, treatment, and control were derived. The data were presented with a 95% confidence

104   interval (CI), and statistical significance was established by using a p-value of <0.05. The statistical software program

105   Stata (version 13·1; Stata Corp LP, College Station, Texas) was used for all analyses.

Spatial analysis

This research employs a number of techniques to deconstruct the spatial distribution of diabetes prevalence in

Bangladesh. To measure the spatial proximity between each potential pair of locations, a spatial weight matrix was

generated. Various methods, including the Queen's, Rook's, and Bishop's methods, can be used to calculate the matrix

depending on how the neighbors are defined[19]. This study uses Queen's approach to construct a 1^{st} order spatial

weight matrix. Additionally, a spatial weight matrix was generated using the inverse distance wighting method. When

working with standard square grids, this approach is more advantageous. The prevalence and clustering of diabetes in

Bangladesh are examined using Moran's I, a global indicator of spatial autocorrelation[20]. The values of Moran's I

range from 1 to -1. When similar values cluster together, the value "1" denotes perfect positive spatial autocorrelation.

The close spatial distribution of regions with identical attribute values would be indicated by a positive spatial

autocorrelation. And "-1" denotes perfect negative spatial autocorrelation. Negative spatial autocorrelation would

suggest differences in locations that are closely related to one another. The 999 iterations of the Monte Carlo

simulation were chosen as the threshold for testing significance. It was determined how similar or different the data

points were from their neighbors using the local Moran's I, which also measures spatial autocorrelation[21].

Neighborhood values that are near a specific spatial location are correlated, and this is measured by Local Indicators

of Spatial Association (LISA). It represents how randomly and spatially clustered the data are. Furthermore, using

five fundamental classifications, the local Moran's I is utilized to determine the spatial distribution of diabetes clusters

and outliers in Bangladesh:

1. High-High (hot spots): A district is classified as High-High if it has a high prevalence of diabetes and is

surrounded by other districts that also have high prevalence rates.

2. High-Low: A district is classified as High-Low if it has a high prevalence of diabetes and is surrounded by

other districts that have low prevalence rates.

3. Low-High: A district is classified as Low-High if it has a low prevalence of diabetes and is surrounded by

other districts that have high prevalence rates.

4. Low-Low (cold spots): A district is classified as Low-low if it has a low prevalence of diabetes and is

surrounded by other districts that also have low prevalence rates.

5. Not Significant: Districts with statistically insignificant spatial patterns.

For clusters found by the local Moran's I, the significance level can also be altered. For instance, analysts can be more

selective by choosing $p<0.01$ or $p<0.001$ as the significant level instead of $p<0.05$. If regions (districts) be discovered

to be statistically significant within the boundaries of these more strict p-values, there's a possibility that the outliers

and/or clusters will be of higher quality[22]. R (version 4.2.1) was used to conduct all of the analyses.

Results

Table-1 provides information about the houses and localities of the study population's participants, as well as their sociodemographic and health characteristics. Out of the 12100 individuals that were analyzed, 1202 of them had diabetes. The participants' average (SE) fasting plasma glucose values were 56.74 (0.15) mmol/dL, their average (SE) age was 39.85 (0.15) years, and their average (SE) BMI was 22.39 (0.04) kg/m2. 26.5% of participants lived in urban regions; 57.2% of participants were female; 26.6% had no formal education; and 39.6% were overweight or obese.

Table 1: Distribution of the study sample according to presence of diabetes.

Variables	Total (n=12100)	Diabetes		p-value
		Yes (n=1202)	No (n=10898)	
Mean (SE) of fasting plasma glucose, mmol/dL	56.74 (0.15)			
Age (in years)				
Mean (SE)	39.85 (0.15)			
18 to 34	5415	284	5130	<0.001
35 to 39	1384	141	1243	
40 to 44	1044	130	914	
45 to 49	1008	133	875	
50 to 54	677	116	561	
55 to 59	685	104	581	
60 to 64	679	112	568	
65 or more	1208	182	1026	
Sex				
Female	6919	657	6261	0.0715
Male	5181	545	4636	
Nutritional status (BMI)				
Mean (SE)	22.39 (0.04)			
Underweight	2068	128	1940	<0.001
Normal	5098	402	4696	
Overweight	1917	204	1713	
Obese	2879	448	2431	
Hypertension				
No	8755	648	8107	<0.001
Yes	3346	555	2791	
Currently working status				
No	4738	561	4177	<0.001
Yes	7362	641	6721	
Education level				
No formal education	3218	314	2903	0.8991
Primary	3614	363	3250	
Secondary	3569	364	3205	
College or above	1701	161	1539	
Wealth quintile				
Poorest	2338	134	2204	<0.001
Poorer	2391	142	2248	

Middle	2496	198	2297	
Richer	2410	276	2134	
Richest	2466	452	2014	
Place of residence				
Urban	3211	426	2785	<0.001
Rural	8889	777	8113	
Division of residence				
Dhaka	2792	400	2392	<0.001
Barisal	668	64	603	
Chattogram	2084	233	1850	
Khulna	1512	125	1387	
Mymensingh	989	78	911	
Rajshahi	1751	141	1610	
Rangpur	1516	85	1431	
Sylhet	789	77	712	

Table 2 and 3 give a summary of the prevalence of diabetes and its awareness, treatment, and control by sex of respondents. Diabetes is more prevalent in men (10.52, 95% CI: 9.59-11.54) compared to women (9.50, 95% CI: 8.66-10.42). Older individuals exhibited a higher crude prevalence rate than younger individuals among both women and men. The lowest prevalence of diabetes is found in adults 18 to 34 years old, regardless of sex. For both men (12.95, 95% CI: 11.29-14.80) and women (13.52, 95% CI: 11.69-5.58), living in an urban area is associated with a greater prevalence of diabetes than living in a rural one. The prevalence of diabetes was the maximum among those with obesity (men: 18.00, 95% CI: 15.42-20.89; women: 14.48, 95% CI: 12.78-16.35); hypertension (men: 16.97, 95% CI: 14.87-19.30; women: 16.31, 95% CI: 14.43-18.38); and in the highest quintile of wealth (men: 21.08, 95% CI: 18.57-23.84; women: 16.28, 95% CI: 14.15-18.65). Diabetes awareness increased for both sexes with each wealth quintile. High awareness and treatment rates are also present in hypertensive men and women. Overweight and obese individuals of both sexes had a greater prevalence of diabetes awareness and treatment. Diabetes awareness (61.22, 95% CI: 45.78-47.70) and treatment (54.62, 95% CI: 39.21-69.19) were highest among women aged 50-54 years. Men aged 55-59 years had the highest prevalence of diabetes awareness (63.17, 95% CI: 43.72-79.12) and treatment (62.12, 95% CI: 42.82-78.20). Women 65 years and older and men aged 45-49 years have the highest diabetes control rates: 44.38 (95% CI: 27.96-62.13) and 43.49 (95% CI: 20.13-70.16) respectively. Furthermore, diabetes treatment and control is higher for both sexes in rural regions than in urban ones.

Table-2: Crude prevalence of diabetes and its awareness, treatment and control status of both sex for different age groups.

Characteristics	Prevalence of diabetes (95% Confidence Interval)	
	Women	**Men**

	Diabetes (%)	Aware (%)	Treated (%)	Controlled (%)	Diabetes (%)	Aware (%)	Treated (%)	Controlled (%)
Total	9.50(8.66-10.42)	39.71(35.47-44.12)	36.44(32.37-40.71)	29.70(24.11-35.97)	10.52(9.59-11.54)	37.11(32.04-42.48)	33.29(28.54-38.40)	32.40(25.69-40.08)
Age								
18-34	5.26 (4.43-6.24)	19.28 (13.99-25.99)	16.35 (11.47-22.78)	25.06 (12.33-44.30)	5.23 (4.21-6.48)	6.23 (2.87-13.00)	5.90 (2.62-12.74)	9.84 (1.25-48.53)
35-39	11.50 (9.00-14.58)	33.23 (22.96-45.38)	32.11 (21.97-44.28)	21.59 (9.37-42.33)	8.74 (6.66-11.39)	27.54 (16.81-41.69)	25.45 (15.08-39.62)	20.22 (6.92-46.37)
40-44	11.89 (9.28-15.10)	44.11 (32.06-56.89)	38.30 (27.05-50.96)	26.73 (13.64-45.73)	13.13 (9.86-17.26)	21.20 (11.65-35.43)	19.49 (10.30-33.78)	29.77 (8.44-66.10)
45-49	14.74 (11.74-18.36)	52.78 (40.01-65.20)	50.75 (38.49-62.91)	34.55 (21.18-50.91)	11.22 (8.29-15.02)	45.96 (30.50-62.24)	36.73 (22.83-53.26)	43.49 (20.13-70.16)
50-54	16.67 (12.60-21.72)	61.22 (45.78-74.70)	54.62 (39.21-69.19)	21.15 (9.41-40.93)	17.66 (13.54-22.70)	33.80 (21.91-48.16)	31.40 (19.96-45.66)	37.96 (18.16-62.78)
55-59	16.44 (12.85-20.80)	50.03 (37.42-62.64)	44.36 (31.88-57.61)	27.99 (14.25-47.62)	13.62 (9.52-19.11)	63.17 (43.72-79.12)	62.12 (42.84-78.20)	24.75 (10.93-46.86)
60-64	15.03 (11.30-19.72)	52.14 (36.54-67.32)	49.11 (33.80-64.59)	31.21 (14.99-53.86)	17.71 (13.71-22.59)	58.99 (44.63-71.96)	54.15 (40.02-67.64)	33.34 (18.32-52.74)
65 and above	13.55 (10.67-17.06)	46.32 (33.91-59.20)	44.88 (32.62-57.78)	44.38 (27.96-62.13)	16.31 (13.45-19.64)	55.27 (44.06-65.96)	47.15 (36.37-58.20)	36.70 (23.66-52.03)

Table-3: Crude prevalence of diabetes and its awareness, treatment and control status of both sex by different factors.

Characteristics	Prevalence of diabetes (95% Confidence Interval)							
	Women				Men			
	Diabetes (%)	Aware (%)	Treated (%)	Controlled (%)	Diabetes (%)	Aware (%)	Treated (%)	Controlled (%)
Residence								
Urban	13.52 (11.69-15.58)	39.90 (33.94-46.19)	36.29 (30.69-42.29)	26.80 (19.03-36.31)	12.95 (11.29-14.80)	33.14 (26.13-40.98)	30.86 (24.34-38.25)	28.59 (17.64-42.81)
Rural	9.50 (8.66-10.42)	39.60 (33.99-45.50)	36.53 (31.10-42.32)	31.37 (24.00-39.98)	9.61 (8.49-10.84)	39.14 (32.44-46.27)	33.29 (28.54-38.40)	34.25 (25.94-43.65)
Division								
Barisal	9.49 (7.14-12.52)	34.42 (23.91-46.73)	32.39 (22.49-44.17)	44.08 (27.32-62.31)	9.85 (7.23-13.28)	43.96 (28.27-60.95)	40.47 (25.62-57.31)	40.65 (19.49-65.95)
Chittagong	10.63 (8.60-13.06)	49.03 (38.62-59.52)	43.34 (33.52-53.70)	28.03 (16.85-42.81)	12.08 (9.50-15.23)	46.84 (34.47-59.62)	43.91 (32.30-56.23)	32.57 (18.39-50.86)
Dhaka	13.31 (10.82-16.27)	27.56 (20.28-36.26)	25.34 (18.34-33.90)	21.11 (10.35-38.28)	15.62 (13.18-18.42)	26.95 (17.87-38.48)	22.79 (14.45-34.03)	30.59 (16.29-49.95)
Khulna	7.82 (6.27-9.72)	51.54 (41.45-61.51)	49.03 (39.70-58.42)	33.84 (20.41-50.50)	8.85 (6.92-11.26)	33.26 (22.90-45.52)	31.13 (21.06-43.37)	23.92 (10.05-46.93)
Mymensingh	8.54 (6.65-10.90)	34.34 (22.29-48.81)	32.40 (20.70-46.82)	36.93 (16.83-62.87)	6.95 (5.12-9.37)	35.28 (20.87-52.97)	32.02 (18.44-49.54)	65.08 (37.37-85.34)

Rajshahi	7.52 (5.88-9.57)	48.88 (36.49-61.40)	42.50 (30.78-55.13)	30.12 (16.15-49.10)	8.74 (6.49-11.68)	46.19 (31.02-62.10)	37.90 (24.79-53.06)	22.27 (9.87-42.85)
Rangpur	5.01 (3.67-6.79)	49.51 (34.70-64.41)	46.86 (32.60-61.65)	18.25 (6.71-40.91)	6.30 (4.62-8.54)	53.25 (36.36-69.44)	52.19 (35.32-68.58)	39.15 (22.18-59.22)
Sylhet	10.22 (7.71-13.42)	36.65 (24.95-50.17)	36.65 (24.95-50.17)	50.36 (35.86-64.80)	9.08 (6.58-12.39)	30.07 (20.14-42.30)	27.18 (17.92-38.94)	25.62 (9.26-53.77)
Education								
No education	10.28 (8.85-11.91)	44.41 (36.52-52.60)	41.38 (33.68-49.54)	35.09 (24.92-46.83)	8.92 (7.14-11.09)	28.59 (19.26-40.19)	26.84 (17.80-38.33)	43.18 (24.85-63.58)
Primary	10.22 (8.63-12.07)	34.34 (27.48-41.93)	31.12 (24.73-38.33)	28.32 (18.84-40.21)	9.84 (8.30-11.63)	27.62 (20.43-36.20)	22.62 (16.12-30.77)	42.01 (25.44-60.60)
Secondary	8.96 (7.58-10.56)	42.10 (34.85-49.72)	38.40 (31.32-46.01)	19.51 (11.55-31.04)	12.03 (10.17-14.17)	43.69 (34.21-53.65)	39.35 (30.32-49.18)	21.45 (12.84-33.60)
College or higher	7.12 (5.51-9.16)	34.13 (23.24-47.01)	31.26 (20.70-44.19)	51.61 (29.81-72.81)	11.49 (9.46-13.90)	49.25 (39.27-59.29)	45.87 (36.08-55.99)	34.45 (22.37-48.94)
Wealth quintile								
Poorest	6.34 (4.98-8.05)	34.87 (24.54-46.85)	34.41 (24.11-46.43)	37.73 (22.81-55.41)	4.90 (3.51-6.80)	21.27 (10.47-38.43)	21.27 (10.47-38.43)	83.31 (43.92-96.95)
Poorer	5.68 (4.46-7.21)	29.49 (19.54-41.85)	27.52 (18.00-39.65)	35.87 (18.54-57.90)	6.33 (4.81-8.29)	16.69 (8.76-29.48)	13.56 (6.55-25.97)	26.04 (6.89-62.63)
Middle	7.74 (6.30-9.49)	35.74 (25.36-47.66)	30.32 (20.79-41.91)	31.38 (16.96-50.58)	8.21 (6.55-10.26)	37.62 (27.08-49.48)	32.68 (22.77-44.43)	47.21 (28.64-66.59)
Richer	11.29 (9.54-13.32)	35.21 (27.05-44.66)	32.85 (24.80-42.06)	28.54 (17.57-42.81)	11.62 (9.63-13.96)	38.93 (29.08-49.78)	35.16 (26.18-45.35)	23.81 (12.45-40.73)
Richest	16.28 (14.15-18.65)	49.84 (43.01-56.67)	45.48 (39.04-52.07)	26.16 (18.49-35.63)	21.08 (18.57-23.84)	45.27 (37.14-53.66)	40.85 (32.93-49.27)	26.84 (17.89-38.19)
Nutritional status (BMI)								
Underweight	6.15 (4.59-8.18)	16.03 (8.52-28.13)	16.03 (8.52-28.13)	41.10 (16.54-71.07)	6.27 (4.72-8.29)	22.17 (12.39-36.47)	22.17 (12.39-36.47)	69.71 (36.02-90.39)
Normal	7.18 (6.15-8.37)	43.76 (36.33-51.47)	40.97 (33.64-48.73)	34.84 (24.38-47.00)	8.65 (7.43-10.06)	30.15 (23.52-37.73)	26.02 (19.95-33.16)	30.69 (19.28-45.08)
Overweight	8.97 (7.29-11.01)	41.30 (31.16-52.22)	37.60 (27.73-48.62)	27.48 (17.35-33.35)	13.09 (10.74-15.85)	43.01 (32.99-53.64)	38.76 (29.14-49.35)	29.40 (16.21-47.27)
Obese	14.48 (12.78-16.35)	42.12 (35.86-48.65)	37.81 (31.78-44.25)	24.48 (17.35-35.54)	18.00 (15.42-20.89)	47.29 (38.41-56.34)	43.56 (34.78-52.76)	28.38 (17.95-41.78)
Currently working status								
No	11.56 (10.35-12.88)	39.02 (33.91-44.38)	36.36 (31.43-41.60)	28.06 (21.98-35.07)	13.36 (10.87-16.31)	42.74 (32.64-53.48)	37.97 (28.74-48.15)	35.16 (21.80-51.35)
Yes	6.73 (5.85-7.74)	41.32 (34.22-48.80)	36.63 (29.80-44.03)	33.45 (22.93-35.07)	10.03 (9.05-11.10)	35.80 (30.36-41.65)	32.20 (27.00-37.89)	31.74 (24.14-40.45)
Hypertension								
No	6.78 (5.98-7.68)	26.32 (21.45-31.84)	22.94 (18.29-28.35)	29.53 (19.87-41.47)	8.20 (7.20-9.34)	26.29 (20.71-32.75)	23.80 (18.52-30.03)	38.73 (27.53-51.25)
Yes	16.31 (14.43-18.38)	53.65 (47.80-59.40)	50.49 (44.64-56.33)	29.77 (23.42-37.02)	16.97 (14.87-19.30)	51.66 (43.77-59.46)	46.05 (38.30-53.89)	28.13 (19.59-38.62)

Fig 2 and 3 depict the awareness of diabetes and the status of its treatment for both men and women, respectively.

Separate pie charts are employed to illustrate the awareness and treatment status of diabetes for each of the eight

divisions. Four categories are displayed in the pie charts: "Aware, not treated," "Aware, treated, and controlled,"

"Aware, treated, and not controlled," and "Unaware". Each division's sample population is shown by a pie chart whose

size corresponds to its overall size. Fig 2 and 3 illustrates that, among all the divisions, Dhaka and Chittagong have

the biggest sample sizes for both men and women.

Fig 2: Pie chart of diabetes awareness, treatment status by division for men in Bangladesh.

Fig 3: Pie chart of diabetes awareness, treatment status by division for women in Bangladesh.

Diabetes hotspot maps were produced using the Local Moran's I along with univariate local indicator of spatial

association (LISA) clustering methods to investigate the spatial dependence and clustering of diabetes in Bangladesh.

The calculated Moran's I value (0.215, p = 0.001) suggests that there is positive spatial autocorrelation, which suggests

that values that are similar, whether high or low, are clustered. The univariate LISA cluster map (Fig 4) and LISA

significance map (Fig 5) showed that Dhaka, Gazipur, Comilla, Chandpur, Narayanganj, and other locations were

home to all of the hotspots. Panchagarh, Rangpur, Dinajpur, Feni, Lalmonirhat, and some other districts were where

the cold spots were identified. The statistical significance level at which each region can be considered to have

meaningfully contributed to the findings of the global spatial autocorrelation is shown by the LISA significance map.

Fig 4: LISA cluster map of diabetes hotspots and cold spots by division.

Fig 5: LISA significance map of diabetes hotspots and cold spots by division.

Discussion

The study specifically aims to investigate the spatial distribution of diabetes prevalence across 64 districts in

Bangladesh using data from a nationally representative sample survey (BDHS 17-18). In this study, the relationship

between age, sex, nutritional status, residency, level of education, employment status, socio-economic status, and

hypertension is also examined in relation to diabetes prevalence. According to the findings, those who are older, more

educated, in better socioeconomic standing, have higher BMIs, and residing in urban areas have a much greater chance

of developing diabetes than those who do not meet those criteria. The research's conclusions are in line with the results

of numerous other studies carried out in comparable circumstances[23,24].

Our study shows that 9.50% of women and 10.52% of men have diabetes. In other Asian nations, comparable

figures have been noted: China (women: 8.8%, men: 10.6%)[25], Nepal (women: 10.0%, men: 15.3%)[26], Korea

(women: 7.5%, men: 8.1%)[27], Pakistan (women: 15.8%, men: 14.8%)[28]. Additionally, the findings demonstrate

that in terms of men and women, residents of urban regions had significantly higher diabetes prevalence rates than

those in rural areas[29]. Like Bangladesh, other countries on the Indian subcontinent, such as India, Bhutan, Nepal,

and Sri Lanka, have a higher prevalence of diabetes in urban areas[30].

Diabetes, and BMI, and older age are also related. Men and women who are older in age have a higher prevalence

of diabetes than those who are younger. This has also been noticed on other studies done in Bangladesh[31,32],

China[25,33], India[34]. This reflects the characteristics of diabetes, a chronic illness that typically manifests in later

life[7]. Furthermore, overweight and obese individuals are more susceptible to diabetes than others. The results of

additional research that has been undertaken also support these findings[35,36]. Higher socioeconomic class, as

indicated by the wealth quintile and education level, was associated with a higher prevalence of diabetes. Many other

research conducted in developing nations identified that socioeconomic status raised the likelihood of obesity and

diabetes[37–39]. Moreover, people tend to live more sedentary lifestyles and consume more calorie-dense diets, which

may increase their chance of developing diabetes along with other chronic illnesses[7,40]. Our findings show that the

prevalence of diabetes is higher in unemployed people than the employed individuals. In our study, individuals with

hypertension had a higher possibility of developing diabetes than those without. This has been observed in other

studies as well[25,39,41,42].

39.71% of women and 37.11 % of men are aware of their condition, according to the findings. Among them,

36.44% females and 33.29% males are taking medication for diabetes. And of those taking medication, only 29.70%

of women and 32.40% of men have their diabetes under control. 37.11% of men and 39.71% of women in Bangladesh

are aware that they have diabetes, which is lower than awareness in Nepal (men: 68%, women: 62%)[26], China (men:

38%, women: 41%)[43]. These findings suggest that diabetes management in Bangladesh is worse than in other

countries of South and East Asia[44]. Moreover, our study indicates that women have a greater likelihood to be aware

of, take medication for, and control their diabetes than men. This is also found in other studies done in

Bangladesh[45,46], Nepal[26] and China[47].

This study revealed that awareness and treatment increases with age, which is consistent with findings from earlier

research carried out in LMICs in Asia, including Bangladesh, Nepal, and China[26,45,47]. Our findings also point to

relatively low rates of diabetes awareness, treatment, and control among young individuals, both for men and women.

According to our study, people in the two quintiles of highest wealth are more likely to be aware of being diabetic

and receive medical treatment for it than people in other quintiles. Findings from additional studies conducted in

Bangladesh give support to this[46].

Additionally, a more detailed geographic description of diabetes hotspots and diabetes cold spots in Bangladesh

were defined. Hotspots and cold spots were mapped according to the prevalence of diabetes within the sample

population using Moran's I[20]. Moran's I is a popular geographical statistic for identifying global spatial trends[48].

It evaluates how closely related one object is to those around it. An approach to checking for autocorrelation in data

is the Moran's I test. Spatial autocorrelation denotes the presence of systematic spatial variation in a mapped

variable[49]. Geographically close values of a variable on a map have a tendency to be similar when there is positive

spatial autocorrelation: high values tend to exist in proximity to other high values; medium values are generally located

closer to medium values; and low values are generally found nearby other low values[50]. As a result, a clustering of

similar values in terms of diabetes prevalence can be seen across districts in Bangladesh.

LISA provides a measure of how much the distribution of values surrounding a specific location differs from spatial

randomization [51]. The LISA significance map displays the areas with significant local Moran statistics, and the

LISA cluster map classifies these locations according to the type of association[19]. The presence of spatial clusters

(high values surrounded by similar high values and low values surrounded by similar low values) and spatial outliers

(high values surrounded by low values along with low values surrounded by high values) are illustrated on the LISA

cluster map[19]. Cold spots denote regions with low level of diabetes prevalence that are also surrounded by regions

with low level of diabetes prevalence, while hot spots denote regions with high level of diabetes prevalence that are

surrounded by other regions with high level of diabetes prevalence. The statistical level at which any region can be

considered to be significantly contributing to the findings of the global spatial autocorrelation is illustrated on the

LISA significance map[21]. We can identify which areas are contributing most significantly to the overall result and

in which direction by utilizing the LISA cluster map and the LISA significance map.

This analysis has several limitations that should be stated. In addition to the factors that our analysis identified to

be linked to diabetes, there are a number of other factors. Diabetes is linked to dietary patterns, smoking behaviors,

alcohol use, physical activity levels, insulin resistance, and more[52–56]. Information about these factors is absent

from the dataset. Because this study is cross-sectional, it can only give a partial picture of spatial accessibility over a

specific period of time and may also be vulnerable to problems with residential self-selection. Despite the fact that

there are many different indicators of global and local spatial association, only the global and local Moran's I statistic

was utilized in our investigation[57]. This is another limitation of our study.

The study's advantages include the use of a sizable, nationally representative dataset, which suggests the

conclusions are reliable. The application of precise techniques for determining clinical parameters like blood pressure,

weight, height, and fasting blood glucose is another advantage of this study. Diabetes was categorized according to

WHO standards. With the use of geographical analysis, we were able to research and examine the regional differences

in diabetes in Bangladesh.

254   Conclusion

This study raises concerns about the need to better comprehend the major and predictable effects that geography

has on population health. According to this study, initiatives in Bangladesh to increase public awareness of and control

of diabetes need to be reinforced and optimized, with additional funding. This might necessitate alterations to the

health care system, including a focus on non-communicable disease prevention and a reassessment of medical care

payments to cut down on out-of-pocket costs. Our data further suggests that efforts to prevent diabetes should center

on lowering obesity and controlling hypertension. There is a requirement for improved preventative strategies in

addition to better management of obesity and hypertension. Diabetes will continue to escalate in Bangladesh if

appropriate prevention measures are not taken.

Furthermore, the prevalence of diabetes with regard to geographic clusters can be examined using spatial analysis.

Additionally, hot spot analysis will enable public health professionals to implement suitable strategies and initiatives

in light of specific geographical regions. As regions encompasses more than just a place's physical features; it also

refs to the cultures, institutions, customs, and way of life that its inhabitants are regularly exposed to[58]. The LISA

cluster and significance maps can also be useful in detecting diabetes risk factors in various geographic locations,

which might further boost the preventive and control measures of diabetes with reference to various associated factors.

Acknowledgement

The authors thank MEASURE DHS for allowing permission for using the BDHS 17-18 dataset.

Reference

1.   Caldwell J, Caldwell P. What have we learnt about the cultural, social and behavioural determinants of health? From Selected Readings to the first Health Transition Workshop. Health Transit Rev. 1991;1(1):3–19.

2.   Omran AR. eweb:279147. 2005 [cited 2022 Dec 5]. The Epidemiologic Transition: A Theory of the Epidemiology of Population Change. Available from: https://repository.library.georgetown.edu/handle/10822/985755

3.   Kahn SE, Hull RL, Utzschneider KM. Mechanisms linking obesity to insulin resistance and type 2 diabetes. Nature. 2006 Dec;444(7121):840–6.

4.   Scobie IN, Samaras K. Fast Facts: Diabetes Mellitus. Karger Medical and Scientific Publishers; 2014. 138 p.

5.   Chatterjee S, Khunti K, Davies MJ. Type 2 diabetes. The Lancet. 2017 Jun 3;389(10085):2239–51.

6.   WHO. Diabetes [Internet]. [cited 2022 Dec 5]. Available from: https://www.who.int/news-room/fact-sheets/detail/diabetes

7.   IDF. IDF Diabetes Atlas 2021 | IDF Diabetes Atlas [Internet]. 2021 [cited 2022 Nov 28]. Available from: https://diabetesatlas.org/atlas/tenth-edition/

8.   Morland K, Diez Roux AV, Wing S. Supermarkets, Other Food Stores, and Obesity: The Atherosclerosis Risk in Communities Study. Am J Prev Med. 2006 Apr 1;30(4):333–9.

288   9.   Swinburn BA, Sacks G, Hall KD, McPherson K, Finegood DT, Moodie ML, et al. The global obesity
289   pandemic: shaped by global drivers and local environments. The Lancet. 2011 Aug 27;378(9793):804–14.

290   10.   Mesev V. Remotely-Sensed Cities. CRC Press; 2003. 416 p.

291   11.   Grubesic TH, Miller JA, Murray AT. Geospatial and geodemographic insights for diabetes in the United
292   States. Appl Geogr. 2014 Dec 1;55:117–26.

293   12.   Harris R, Sleight P, Webber R. Geodemographics, GIS and Neighbourhood Targeting. John Wiley and Sons;
294   2005. 368 p.

295   13.   Twigg L, Moon G, Jones K. Predicting small-area health-related behaviour: a comparison of smoking and
296   drinking indicators. Soc Sci Med. 2000 Apr 1;50(7):1109–20.

297   14.   Abbas J, Ojo A, Orange S. Geodemographics – a tool for health intelligence? Public Health. 2009 Jan
298   1;123(1):e35–9.

299   15.   Barquera S, Tovar-Guzmán V, Campos-Nonato I, González-Villalpando C, Rivera-Dommarco J. Geography
300   of diabetes mellitus mortality in Mexico: an epidemiologic transition analysis. Arch Med Res. 2003 Sep
301   1;34(5):407–14.

302   16.   Green C, Hoppa RD, Young TK, Blanchard JF. Geographic analysis of diabetes prevalence in an urban area.
303   Soc Sci Med. 2003 Aug 1;57(3):551–60.

304   17.   Niport NI of PR and T, Welfare M of H and F, ICF. Bangladesh Demographic and Health Survey 2017-18.
305   2020 Oct 1 [cited 2022 Nov 6]; Available from: https://dhsprogram.com/publications/publication-FR344-
306   DHS-Final-Reports.cfm

307   18.   WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for
308   policy and intervention strategies. Lancet Lond Engl. 2004 Jan 10;363(9403):157–63.

309   19.   Anselin L, Syabri I, Kho Y. GeoDa: An Introduction to Spatial Data Analysis. In: Fischer MM, Getis A,
310   editors. Handbook of Applied Spatial Analysis: Software Tools, Methods and Applications [Internet]. Berlin,
311   Heidelberg: Springer; 2010 [cited 2022 Nov 28]. p. 73–89. Available from: https://doi.org/10.1007/978-3-
312   642-03647-7_5

20. Moran PAP. Notes on Continuous Stochastic Phenomena. Biometrika. 1950;37(1/2):17–23.

21. Anselin L. Local Indicators of Spatial Association—LISA. Geogr Anal. 1995;27(2):93–115.

22. Anselin L. Exploring Spatial Data with GeoDaTM : A Workbook. 2004.

23. Saquib N, Saquib J, Ahmed T, Khanam MA, Cullen MR. Cardiovascular diseases and Type 2 Diabetes in

Bangladesh: A systematic review and meta-analysis of studies between 1995 and 2010. BMC Public Health.

2012 Jun 13;12(1):434.

24. Akter S, Rahman MM, Abe SK, Sultana P. Prevalence of diabetes and prediabetes and their risk factors

among Bangladeshi adults: a nationwide survey. Bull World Health Organ. 2014 Mar 1;92(3):204-213A.

25. Yang W, Lu J, Weng J, Jia W, Ji L, Xiao J, et al. Prevalence of Diabetes among Men and Women in China.

N Engl J Med. 2010 Mar 25;362(12):1090–101.

26. Gyawali B, Hansen MRH, Povlsen MB, Neupane D, Andersen PK, McLachlan CS, et al. Awareness,

prevalence, treatment, and control of type 2 diabetes in a semi-urban area of Nepal: Findings from a cross-

sectional study conducted as a part of COBIN-D trial. PLOS ONE. 2018 Nov 2;13(11):e0206491.

27. Kim SM, Lee JS, Lee J, Na JK, Han JH, Yoon DK, et al. Prevalence of Diabetes and Impaired Fasting

Glucose in Korea: Korean National Health and Nutrition Survey 2001. Diabetes Care. 2006 Feb 1;29(2):226–

31.

28. Akhtar S, Nasir JA, Abbas T, Sarwar A. Diabetes in Pakistan: A systematic review and meta-analysis. Pak J

Med Sci. 2019;35(4):1173–8.

29. Kibria GMA, Swasey K, Gupta RD, Choudhury A, Nayeem J, Sharmeen A, et al. Differences in prevalence

and determinants of hypertension according to rural–urban place of residence among adults in Bangladesh. J

Biosoc Sci. 2019 Jul;51(4):578–90.

30. Wild S, Roglic G, Green A, Sicree R, King H. Global Prevalence of Diabetes: Estimates for the year 2000

and projections for 2030. Diabetes Care. 2004 May 1;27(5):1047–53.

31.   Sayeed MA, Ali L, Hussain MZ, Rumi MA, Banu A, Azad Khan AK. Effect of Socioeconomic Risk Factors

on the Difference in Prevalence of Diabetes Between Rural and Urban Populations in Bangladesh. Diabetes

Care. 1997 Apr 1;20(4):551–5.

32.   Abdul Baker Chowdhury M, Uddin MJ, Khan H, Haque MR. Type 2 diabetes and its correlates among adults

in Bangladesh: a population based stud. Dep Biostat Fac Publ [Internet]. 2015 Oct 19; Available from:

https://digitalcommons.fiu.edu/biostatistics_fac/2

33.   Pan XR, Yang W, Li GW. Prevalence of Diabetes and Its Risk Factors in China, 1994 | Diabetes Care |

American Diabetes Association [Internet]. [cited 2022 Nov 30]. Available from:

https://diabetesjournals.org/care/article/20/11/1664/21163/Prevalence-of-Diabetes-and-Its-Risk-Factors-in

34.   Ramachandran A, Snehalatha C, Kapur A, Vijay V, Mohan V, Das AK, et al. High prevalence of diabetes

and impaired glucose tolerance in India: National Urban Diabetes Survey. Diabetologia. 2001 Sep

1;44(9):1094–101.

35.   Abarca-Gómez L, Abdeen ZA, Hamid ZA, Abu-Rmeileh NM, Acosta-Cazares B, Acuin C, et al. Worldwide

trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of

2416 population-based measurement studies in 128·9 million children, adolescents, and adults. The Lancet.

2017 Dec 16;390(10113):2627–42.

36.   Chowdhury MAB, Adnan MM, Hassan MZ. Trends, prevalence and risk factors of overweight and obesity

among women of reproductive age in Bangladesh: a pooled analysis of five national cross-sectional surveys.

BMJ Open. 2018 Jul 1;8(7):e018468.

37.   Jebb SA, Moore MS. Contribution of a sedentary lifestyle and inactivity to the etiology of overweight and

obesity: current evidence and research issues. Med Sci Sports Exerc. 1999 Nov 1;31(11 Suppl):S534-41.

38.   Misra A, Khurana L. Obesity and the Metabolic Syndrome in Developing Countries. J Clin Endocrinol

Metab. 2008 Nov 1;93(11_supplement_1):s9–30.

39.   Ahasan HN, Islam MZ, Alam MB, Miah MT, Nur Z, Mohammed FR, et al. Prevalence and Risk Factors of

Type 2 Diabetes Mellitus Among Secretariat Employees of Bangladesh. J Med. 2011;12(2):125–30.

40.   Ahmed SM, Hadi A, Razzaque A, Ashraf A, Juvekar S, Ng N, et al. Clustering of chronic non-communicable

362   disease risk factors among selected Asian populations: levels and determinants. Glob Health Action. 2009

363   Nov 1;2(s4):1986.

364   41.   Rahim MA, Hussain A, Azad Khan AK, Sayeed MA, Keramat Ali SM, Vaaler S. Rising prevalence of type

365   2 diabetes in rural Bangladesh: A population based study. Diabetes Res Clin Pract. 2007 Aug 1;77(2):300–

366   5.

367   42.   Bhowmik B, Afsana F, Diep LM, Munir SB, Wright E, Mahmood S, et al. Increasing Prevalence of Type 2

368   Diabetes in a Rural Bangladeshi Population: A Population Based Study for 10 Years. Diabetes Metab J. 2013

369   Feb 18;37(1):46–53.

370   43.   Li M zhi, Su L, Liang B yun, Tan J jing, Chen Q, Long J xiong, et al. Trends in Prevalence, Awareness,

371   Treatment, and Control of Diabetes Mellitus in Mainland China from 1979 to 2012. Int J Endocrinol. 2013

372   Oct 28;2013:e753150.

373   44.   Islam A, Biswas T. Health System in Bangladesh: Challenges and Opportunities. Am J Health. 2014;

374   45.   Rahman MS, Akter S, Abe SK, Islam MR, Mondal MNI, Rahman JAMS, et al. Awareness, Treatment, and

375   Control of Diabetes in Bangladesh: A Nationwide Population-Based Study. PLOS ONE. 2015 Feb

376   18;10(2):e0118365.

377   46.   Khan N, Oldroyd JC, Hossain MB, Islam RM. Awareness, Treatment, and Control of Diabetes in Bangladesh:

378   Evidence from the Bangladesh Demographic and Health Survey 2017/18. Seixas A, editor. Int J Clin Pract.

379   2022 Apr 22;2022:1–9.

380   47.   Wang Q, Zhang X, Fang L, Guan Q, Guan L, Li Q. Prevalence, awareness, treatment and control of diabetes

381   mellitus among middle-aged and elderly people in a rural Chinese population: A cross-sectional study. PLOS

382   ONE. 2018 Jun 1;13(6):e0198343.

383   48.   Jackson MC, Huang L, Xie Q, Tiwari RC. A modified version of Moran's I. Int J Health Geogr. 2010 Jun

384   29;9(1):33.

385   49.   Haining RP. Spatial Autocorrelation. In: Smelser NJ, Baltes PB, editors. International Encyclopedia of the

386   Social & Behavioral Sciences [Internet]. Oxford: Pergamon; 2001 [cited 2022 Dec 4]. p. 14763–8. Available

387   from: https://www.sciencedirect.com/science/article/pii/B0080430767025110

388   50.   Li L, Jiang Z, Duan N, Dong W, Hu K, Sun W. Chapter 8 - An Approach to Optimize Police Patrol Activities

389   Based on the Spatial Pattern of Crime Hotspots11This chapter is adapted from Li, L., Jiang, Z., Duan, N.,

390   Dong, W., & Sun, W., Police patrol service optimization based on the spatial pattern of hotspots, in 2011

391   IEEE International Conference on Service Operations and Logistics, and Informatics. In: Xiong G, Liu Z,

392   Liu XW, Zhu F, Shen D, editors. Service Science, Management, and Engineering: [Internet]. Boston:

393   Academic   Press;   2012   [cited   2022   Dec   4].   p.   141–63.   Available   from:

394   https://www.sciencedirect.com/science/article/pii/B9780123970374000089

395   51.   Anselin L, Cohen J, Cook D, Gorr W, Tita G. Spatial Analyses of Crime. Crim JUSTICE. 2000;51.

396   52.   Wiki J, Kingham S, Campbell M. A geospatial analysis of Type 2 Diabetes Mellitus and the food environment

397   in urban New Zealand. Soc Sci Med. 2021 Nov 1;288:113231.

398   53.   Will JC, Galuska DA, Ford ES, Mokdad A, Calle EE. Cigarette smoking and diabetes mellitus: evidence of

399   a positive association from a large prospective cohort study. Int J Epidemiol. 2001 Jun 1;30(3):540–6.

400   54.   Howard AA, Arnsten JH, Gourevitch MN. Effect of Alcohol Consumption on Diabetes Mellitus. Ann Intern

401   Med. 2004 Feb 3;140(3):211–9.

402   55.   Plotnikoff RC, Lippke S, Courneya K, Birkett N, Sigal R. Physical activity and diabetes: An application of

403   the theory of planned behaviour to explain physical activity for Type 1 and Type 2 diabetes in an adult

404   population sample. Psychol Health. 2010 Jan 1;25(1):7–23.

405   56.   Seidell JC. Obesity, insulin resistance and diabetes — a worldwide epidemic. Br J Nutr. 2000 Jun;83(S1):S5–

406   8.

407   57.   Bivand RS, Wong DWS. Comparing implementations of global and local indicators of spatial association.

408   TEST. 2018 Sep 1;27(3):716–48.

409   58.   Cresswell T. Place: An Introduction. John Wiley & Sons; 2014. 242 p.

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5