Title: Targeting Molecular Collagen Defects from the Initiation of Knee Osteoarthritis

Authors: Huang Kui[1], Rongmao Qiu[1], Yijie Fang[1,2], Dantian Zhu[1,2], Xiaojing Li[1], Zhongyang Lv[3,4], Taoyu Jia[2], Yuxiang Fei[3], Daoning Zhou[1], Wenjun Wu[1], Yongjie Huang[1], Suwen Zhao[1], Yongqiao Zhu[5], Shaolin Li[2], Dongquan Shi[3,*], and Yang Li[1,*]

Affiliations:

[1]Guangdong Provincial Engineering Research Center of Molecular Imaging, Guangdong-Hong Kong-Macao University Joint Laboratory of Interventional Medicine, the Fifth Affiliated Hospital, Sun Yat-sen University; Zhuhai 519000, China.

[2]Department of Radiology, the Fifth Affiliated Hospital, Sun Yat-sen University; Zhuhai 519000, China.

[3]Division of Sports Medicine and Adult Reconstructive Surgery, Department of Orthopedic Surgery, Nanjing Drum Tower Hospital, Affiliated Hospital of Medical School, Nanjing University; Nanjing 210008, China.

[4]Department of Orthopedics, Nanjing Jinling Hospital, Affiliated Hospital of Medical School, Nanjing University; Nanjing 210002, China.

[5]Cardiac Surgery and Structural Heart Disease Unit of Cardiovascular Center, the Fifth Affiliated Hospital, Sun Yat-sen University, Zhuhai 519000, China.

*Corresponding author. Email: shidongquan@nju.edu.cn (Dongquan Shi); liyang266@mail.sysu.edu.cn (Yang Li).

Abstract

Knee osteoarthritis (OA) is the most prevalent degenerative joint disease. When morphological changes become apparent on radiographs, no approved treatment can reverse the disease process. Early diagnosis is an unmet need demanding new molecular and imaging biomarkers to define OA from the earliest stages. In this context, we focus on collagen, the most basic building block of all joint tissues, and interrogate how OA development affects collagen's molecular folding, a previously underexplored area. Here, through whole-joint mapping with a peptide that recognizes unfolded collagen molecules, we report the discovery of collagen denaturation in cartilage before proteolysis and major histopathological degeneration in animal models and patients. Mechanistically, we reveal that such molecular collagen defects can be driven by mechanical overloading without collagenase degradation and are intimately associated with glycosaminoglycan loss. We showcase the advantages of using collagen denaturation as an early-stage OA hallmark for in vivo therapeutic evaluation and molecular magnetic resonance imaging (MRI) of subtle joint defects that are challenging to detect with conventional morphology-based MRI. These results highlight biomolecular integrity as a crucial dimension for characterizing joint degeneration and a molecular foundation for diagnosing early-stage OA and beyond.

Main Text:

INTRODUCTION

Osteoarthritis (OA) is the most prevalent joint disease affecting over 500 million individuals

worldwide, including over 365 million with knee OA.[1 2] OA is a leading cause of disability in

elders and its incidence is increasing globally with aging populations and rising rates of

obesity and injury.[1-3] There is no approved disease-modifying treatment for OA.[4] Standard

radiographic changes are indirect and insensitive,[5] such as joint-space narrowing, which may

not appear for years after the disease onset.[6] Routine OA management often focuses on

progressing and late-stage patients, when joint replacement is often the ultimate

recommendation.[7] Nonetheless, OA is a slowly progressing disease; a 'window of

opportunity' for early diagnosis and treatment does exist but remains to be precisely defined.[7]

[8] Therefore, new molecular and imaging biomarkers that enable accurate characterization and

non-invasive examination of early-stage OA are critically needed to support a paradigm shift

from reaction to late disease toward prevention.[8]

The destruction of articular cartilage is the defining characteristic of OA.[9] The rich

collagen network primarily maintains the structural integrity and mechanical functioning of

cartilage and other joint tissues, including the menisci and ligaments.[10] As OA advances,

mechanical insults and increased matrix-protease activities induced by inflammatory

cytokines and abnormal cell metabolism disrupt the collagen matrix progressively, leading to

structural failure of the tissues and painful disorders.[1 9 11] The collagen molecule is the basic

building block of the hierarchical collagen architecture in cartilage.[10] We hypothesized that

the structural disruption to the collagen matrix occurs at and progresses from the molecular

level in OA from the initiation.

The molecular defects in collagen structure, namely the denaturation and unwinding of its

unique triple-helix folding, cannot be directly detected by regular imaging, histology, and

electron microscopies.[12] Alternatively, current research for collagen changes in OA focuses

on matrix proteases,[13] OA-specific collagen subtypes,[14] and cryptic immuno-epitopes in the

matrix or body fluids resulting from collagen degradation.[15-17] However, the vital relationship

between the structural integrity of collagen and OA development remains to be established at

the molecular level. Here, we examine and delineate the spatiotemporal changes in the

unfolding of collagen molecules in cartilage and surrounding knee joint tissues during early

OA progression using a CHP that we developed to specifically bind denatured collagen

chains through triple-helical hybridization.[12] With investigation into its underlying driving

forces, we propose that the denatured collagen molecule is a crucial hallmark of early OA

damage in the joint, and showcase its advantages in therapeutic evaluation and molecular

magnetic resonance imaging for early OA.

METHODS

This study aims to address the unmet clinical need for defining and noninvasively detecting

OA from the pre-osteoarthritic stages. These goals were addressed by (i) interrogating the

molecular integrity of the collagen and demonstrating collagen denaturation as the structural

hallmark of early OA through histological and in vivo collagen hybridization and ex vivo 3D

whole-mount joint imaging in rodents, (ii) investigating the mechanical and biological

driving forces of molecular collagen defects in OA cartilage damage, (iii) evaluating the

therapeutic efficacy of agents for early OA in vivo in rat models, (iv) developing an MR

contrast agent and demonstrating its capacity and specificity in identifying early OA changes

in rats, and (v) initiating a pilot study to translate these findings using OA patients' life-size

cartilage specimens. All procedures for animal experiments were reviewed and approved by

the Experimental Animal Ethics Committee of the Fifth Affiliated Hospital of Sun Yat-sen

University (approval no. 00151). All mice received humane care under the criteria set out in

the *Guide for the Care and Use of Laboratory Animals* published by the National Institutes of

Health. In all animal experiments, mice of similar age and size were randomly assigned to

each group. The investigators were blinded for histological and MRI scoring. Because

collagen-hybridizing MRI is a new imaging technology, it is difficult to estimate sample size

with adequate power. All in vivo and in vitro experiments were replicated at least three

independent times. Images acquired from tissue staining (e.g., figure 1A,B), in vivo or ex

vivo fluorescence imaging (e.g., figure 1E, figure 3B, and online supplemental figure S5),

and light-sheet fluorescence microscopy (e.g., figure 2, online supplemental figure S7 and

S8) are representative of similar results from at least three mice from each group. The details

of the study design, sample size, experimental replicates, and statistics are provided in the

corresponding figures and figure legends. Human OA tissue samples were from Nanjing

Drum Tower Hospital. The Ethics Committee of Nanjing Drum Tower Hospital of Nanjing

University approved these studies (approval no. K228-1). Informed consent was obtained

from the patients. The online supplementary materials provide details about peptide synthesis

and labeling, near-infrared fluorescence imaging, histology and microscopy, MR imaging,

and statistical analysis.

RESULTS

Collagen denaturation in mouse knee cartilage during early OA.

We utilized a widely adopted OA model prepared by surgical destabilization of the medial meniscus (DMM) in the right knee of a C57BL/6 mouse paired with a sham operation in the left knee.[18] [19] To examine collagen denaturation in cartilage histologically during OA progression, we harvested knee joints from these mice weekly post-operation and stained them with fluorescently labeled Cy3-CHP in vitro, following a customized decalcification procedure that leaves the folding of collagen molecules intact for CHP staining (figure 1A, online supplemental figure S1 and S2). The cartilage of the DMM joints exhibited only subtle loss of glycosaminoglycan (GAG) without signs of erosion within the first 3 weeks post-operation; yet, there was already notable CHP binding at the cartilage surface, often in locations with mild GAG loss (figure 1B, arrowheads, and online supplemental figure S3) even as early as 1-week post-DMM (figure 1C). The size and fluorescence intensity of CHP-stained cartilage increased steadily during this stage (OARSI grade 0-3, figure 1B,D), with the tibial side showing stronger CHP signals than the femoral side (figure 1B). The CHP binding level and area increased drastically and peaked at OARSI grade 3.5-4 (usually 4 weeks post-DMM, figure 1B,D), when the total CHP fluorescence signal from the cartilage section approximated at 10 times that of the sections of OARSI grade 1 (figure 1D). Beyond 4 weeks post-DMM, substantial cartilage erosion was seen histologically (OARSI grade $\geq$ 4, figure 1B) with immunostaining showing great loss of collagen II down to deep cartilage layers (online supplemental figure S4). Meanwhile, the area and fluorescence of the CHP-stained cartilage were also diminishing accordingly (figure 1B,D, and online supplemental figure S3).

To verify our findings in vivo, we performed near-infrared fluorescence imaging of the mice weekly post-operation (p.o.) following tail vein injection of a Cy5-labeled CHP. Consistently, the DMM-injured knee exhibited stronger fluorescence from in vivo uptake of Cy5-CHP (but not the sequence-scrambled control peptide Cy5-[S]CHP) than the sham-operated knee of the same mouse each week p.o. (online supplemental figure S5A,B). The DMM-injured knee joints collected 3 h post-CHP-injection showed steadily increasing ex vivo fluorescence until 4 weeks p.o. (figure 1E,F, and online supplemental figure S5C,D) when the CHP uptake peaked in the diseased joints (figure 1F, online supplemental figure S5D). This trend strongly correlated with the above CHP histology findings (figure 1B,D).

139 Furthermore, similar results were obtained in rats post-DMM operation with intra-articular
140 injection of Cy5-CHP (online supplemental figure S6). Overall, these results (figure 1)
141 demonstrate that collagen denaturation may occur much earlier than histomorphologic
142 changes in the DMM-induced OA joints and can be targeted in vivo by CHP hybridization.
143

Landscape of molecular collagen defects

Using light-sheet microscopy, we fluorescently imaged the CHP uptake in the DMM mouse
knee joints collected 3 h post-intravenous-injection of Cy5-CHP (figure 2A,B, online
supplemental figure S7 and S8), and performed 3D reconstruction (figure 2C-F, online
supplemental figure S9) and spatial measurements of the molecular lesions with higher-than-
normal CHP fluorescence (figure 2G). Such molecular mapping reveals distinct, meaningful
spatiotemporal features of collagen defects in major anatomical components during early OA
progression. (1) Higher-than-sham CHP fluorescence could be found in the medial cartilage
of the DMM joints as early as 1 week p.o. (figure 2B, yellow arrowheads); the fluorescence
and volume of the CHP-labeled cartilage lesion (figure 2B,G) peaked at 4 weeks p.o. (figure
2C,G). From 1 to 4~5 weeks p.o., the molecular cartilage defect was greater on the medial
side than the lateral (especially in the tibial plateau, figure 2B,C, online supplemental figure
S9A). (2) Collagen denaturation was modest in the anterior and posterior cruciate ligaments
1-3 weeks p.o. (figure 2D, online supplemental figure S9C) but became extensive in the
medial collateral ligaments after 4 weeks p.o. (figure 2D) increasing the volume of ligament
collagen defects (figure 2G). (3) The region rich in denatured collagen appeared to rotate
counterclockwise in the superior view of the post-DMM-surgery meniscus (figure 2E, region
$i \rightarrow vi$, online supplemental figure S9D) with the destabilized medial anterior horn next to the
transected ligament first to suffer from collagen denaturation (region i).
163

Driving forces of collagen denaturation in early post-DMM cartilage

Next, we examined the driving forces of collagen denaturation in OA knee cartilage. Matrix
metalloproteinases (MMPs), especially MMP13, are considered the main collagen catabolic
factor in OA pathogenesis.[20] [21] We thereby first tested the correlation between MMP
activities and collagen denaturation by in vivo imaging the DMM mice administered with
both Cy5-CHP and MMPSense750, a fluorescence probe for broad-spectrum MMP activities
(figure 3A).[22] Unexpectedly, the fluorescence of Cy5-CHP and MMPSense750 showed
distinct localizations in the same OA knee joints (figure 3B and online supplemental figure

S10A). Moreover, the normalized intensity of MMPSense750 from the DMM-injured knees

(i.e., fold change in fluorescence of each mouse's DMM-injured knee over the sham-operated

knee) did not increase until 6 weeks post-DMM surgery (online supplemental figure S10B),

displaying a temporal profile of in vivo MMP activity distinct from that of Cy5-CHP uptake

(online supplemental figure S10B,C). Immunohistochemical staining showed that MMP13

remained non-detectable in the cartilage until 5 weeks p.o. when it became highly

concentrated in the superficial zone; however, the Cy3-CHP fluorescence in the same

cryosections was localized to the middle and superficial zones during 3-4 weeks p.o.,

overlapping minimally with the MMP13-positive coloration (figure 3C and online

supplemental figure S11A). Also, the number of MMP13-positive cells correlated poorly

with the level of Cy3-CHP binding in the same cartilage tissues (figure 3D, online

supplemental figure S11B). Overall, the inconsistency between MMP expression and CHP

binding across multiple spatiotemporal profiles, suggests that the early cartilage collagen

denaturation in the DMM-injured mice may not be adequately explained by MMP

degradation.

The above results prompted us to investigate whether and how mechanical insults can

cause molecular collagen denaturation in cartilage. We stained osteochondral plug specimens

excised from normal porcine patella-femoral joints with Cy3-CHP, which had been

compressed in vitro (figure 3E) with increasing levels (online supplemental figure S12A) or

cycles (figure 3F). The images and fluorescence quantification indicated that loading caused

collagen denaturation that allowed triple-helix specific CHP hybridization (online

supplemental figure S12B) and that the denaturation increased with the level of the applied

stress and cycle (figure 3F, online supplemental figure S12A,C). Cross-sectional confocal

microscopy showed that collagen denaturation was concentrated in the superficial layer of

these loaded cartilage tissues to ~150 μm deep (online supplemental figure S12D,E).

We often noted that whether it is introduced in vivo or ex vivo, CHP binds to cartilage

more prominently in locations with even subtle loss of GAGs (e.g., figure 1B,C, online

supplemental figure S6D). Considering GAGs' vital contribution to swelling and

compressive resistance in cartilage,[10][23] we hypothesized that GAG loss intensifies

mechanical disruption to collagen molecules. To test this, we treated porcine cartilage

samples with chondroitinase ABC (ChABC) to selectively reduce their GAG content before

loading.[24] Post-loading CHP fluorescence stain showed more collagen denaturation with

increasing ChABC concentrations under the same loading (figure 3G); for samples treated

with 0.5 U/mL ChABC, loading of merely 100 N yielded a degree of collagen denaturation

that would require loading of 400 N on normal untreated samples (figure 3G and online supplemental figure S12F). Cross-sectional fluorescence images verified that the same loading caused increasing collagen denaturation within deeper zones in cartilage with declining GAG content (figure 3H).[25] Finally, successive CHP staining before and after loading revealed that a mild process (100 N, 5 cycles) that induced no collagen unfolding to normal, GAG-rich cartilage resulted in substantial collagen denaturation and CHP fluorescence increment in samples with GAG removal (figure 3I). These results demonstrated GAGs' protection to collagen molecules against mechanical denaturation in cartilage. All combined (figure 3), the data suggest that the molecular integrity of collagen in cartilage can be compromised by mechanical factors during early OA progression.

Therapeutic assessment for early OA

To showcase the potential of using collagen denaturation in therapeutic assessment for early OA, we live imaged the knee joints of DMM-operated rats weekly with intra-articularly dosed Cy5-CHP to monitor two treatments in vivo: oral glucosamine sulfate (GlcN) and intra-articular hyaluronic acid (HA) (figure 4A). Both treatments have shown preclinical efficacy against OA development.[26-29] According to the in vivo Cy5-CHP fluorescence, both treatments reduced collagen denaturation within the DMM joints at 4-7 weeks p.o., keeping the CHP signal at a minimal level approximating the sham-operated knees in individual rats (figure 4B,C, online supplemental figure S13). At 7 weeks p.o., the in vivo CHP uptake in the DMM-knee was significantly lower in the two treatment groups than in the untreated group (figure 4C). The therapies' attenuating effect on cartilage destruction was further corroborated by the endpoint histopathology (figure 4D and online supplemental figure S14) and in situ fluorescence examination of the in vivo Cy5-CHP uptake in the cartilage tissue (figure 4E).

In vivo magnetic resonance imaging

To enable in vivo detection of molecular collagen defects in the joints by MRI, we synthesized Gd_n-Cy5-CHP by conjugating Cy5-CHP with multiple Gd-ion chelates, creating a contrast agent with sensitivity severalfold that of Magnevist, a common clinical MR contrast agent (figure 5A, detailed in online supplemental figure S15 and methods). Compared to the pre-contrast scans, intra-articularly administered Gd_n-Cy5-CHP markedly enhanced the T1 signals in the tibia cartilage and ligaments of the DMM-injured (as opposed

to the sham-operated) rat knee joints (figure 5B,C). The high signal sites in the cartilage
shown in T1WI imaging matched closely with the postmortal fluorescence results of the
isolated joints, both indicating Gd_n-Cy5-CHP's in vivo binding to cartilage locations with
GAG loss (figure 5D,E). Furthermore, longitudinal Gd_n-Cy5-CHP enhanced T1WI scans of
the same individuals depicted changes in the level and location of collagen defects in
cartilage and ligaments over weeks during early OA progression (figure 5F); these subtle
damages were not readily visible in the T2WI scan routinely used for clinical examination of
soft tissue degeneration (figure 5D,F).

Ex vivo study of clinical cartilage tissues
We prepared osteochondral plug specimens from different regions of femoral cartilage
surgically removed from OA patients during knee replacement (figure 6A). We grouped these
samples into three [group i: mild disruption (OARSI grade $\leq$ 2), ii: moderate degeneration (2
< OARSI grade $\leq$ 4), and iii: severe damage (OARSI grade > 4)] based on the cartilage
morphology (Fig. 6A) and histopathology (figure 6B) and labeled them with Gd_n-Cy5-CHP
ex vivo. The T1WI magnetic resonance and fluorescence imaging showed that the cartilage
plugs in group ii exhibited the most pronounced CHP signals near the less-smooth cartilage
surface (figure 6C-E, online supplemental figure S16A,B). In situ fluorescence and histology
from the post-MRI cryosections revealed that Gd_n-Cy5-CHP accumulated strongly to regions
of significant GAG loss close to the cartilage surface in group ii while being less effective in
targeting the cartilage in group i in which collagen was not yet disrupted and in group iii,
where collagen had been substantially eroded (figure 6F and online supplemental figure S16).
Three-dimensional light-sheet fluorescence imaging of CHP-labeled, bulk knee cartilage
from OA patients delineated the heterogeneous spatial distribution of denatured collagen in
the tissue (figure 6G, online supplemental figure S17). The above results are highly
consistent with our findings in animal models (figures 1-5), suggesting that CHP can
effectively detect molecular collagen defects in the modestly damaged cartilage of OA
patients.

DISCUSSION
OA typically develops over decades, presenting a window of opportunity for intervention. A
major undertaking in current OA management is to transition from late disease alleviation
toward prevention.[3 7 8] To empower and accelerate this paradigm shift, the pre-osteoarthritic

stages of joint degeneration must be defined with new biomarker discovery and diagnosis

development.[8 30 31] Therefore, we interrogated the integrity of the most basic building block

of joint anatomy - the molecular folding of the collagen architecture - from OA onset and

established collagen denaturation as a disease hallmark preceding collagen proteolysis (figure

3, online supplemental figure S10 and S11) and loss (figure S4) that result in apparent

histopathologic and radiographic changes (OARSI grade 1-4; figure 1 and 6).[32] We further

mapped and measured the molecular lesions three-dimensionally in full joint anatomy along

the course of OA progression in mice (figure 2), delineating the molecular landscape of

collagen destruction in early OA pathogenesis and biomechanics.[33]

This study exemplifies a new approach to characterizing tissue degeneration from the

perspective of material science[34] and structural biology. The collagen matrix is otherwise

equally abundant in normal and pre-osteoarthritic joints and therefore can be overlooked in

search of molecular biomarkers, yet it may be the first to suffer from structural alteration in

degeneration and injury. Herein, the disease characteristic arises from the abnormality in

protein structure instead of quantity, offering a new category of biomarkers not easily

identified by the conventional 'candidate protein' approach.

Investigating the mechanisms driving early-stage OA progression is crucial for

developing effective treatments and risk management.[35] We identified that mechanical

overload, more than protease-degradation, is an underlying driving force of molecular

collagen denaturation in early cartilage degeneration (figure 3); we further showed that the

compressive-resistant GAG component plays a decisive protective role against the

mechanical collagen denaturation in cartilage (figure 3).[36-38] These findings provide

molecular structural evidence that closes the gap in early OA biomechanics and

pathophysiology and underlines weight, joint overloading, and GAG loss as modifiable risk

factors to reduce the disease burden.[35 39 40] Furthermore, GAG loss is one of the first signs of

early OA but is challenging to image and target.[13 41] Our study uncovered a strong

spatiotemporal and logical association between collagen denaturation and GAG loss (figure

1B,C, figure 3G-I, figure 6, online supplemental figure S6D and S16), providing an indirect

strategy for detecting and targeting GAG loss within the degenerating cartilage.

To date, no validated diagnostic criteria are available for early-stage knee OA.[8] The

detection, monitoring, and therapy development of early OA have been hampered by the

insensitivity of the conventional radiographic measures of joint health.[7] We demonstrated that

the CHP (even intravenously administered) can target the denatured collagen and deliver

diagnostic modalities to the avascular articular cartilage in vivo (figures 1-3),[42][43] allowing

straightforward monitoring of knee joint integrity and therapeutic assessments through live

imaging (figure 4). Such noninvasive detection of early joint defects will facilitate the

development of new disease-modifying therapies targeting the pre-osteoarthritic condition[44]

and reduce the reliance on post-mortal histology. MRI is currently the most common

modality for direct imaging of soft tissues in the clinic and research.[45][46] As such, we

developed contrast agent Gd_n-Cy5-CHP and showcased its sensitivity by highlighting the

subtle cartilage defects beyond the capacity of routine T2-based morphological assessment

(figures 5). We further demonstrated that the MR-fluorescence CHP agent can visualize

molecular collagen defects in human knee cartilage with only mild-to-moderate degeneration

(i.e., OARIS grades 1-4) and depict the defect heterogeneity in 3D over life-size patient

specimens (figures 6). OA is a heterogeneous disease; such precision in illustrating collagen

defects may contribute to the much-desired molecular stratification of early OA population.[47-

49] Overall, these translational findings set the stage for further clinical development of

collagen hybridization in the early diagnosis and treatment of OA among a range of

musculoskeletal degenerative diseases and injuries.

321
322

References and Notes

1. Hunter DJ, Bierma-Zeinstra S Osteoarthritis. *Lancet* 2019;393:1745-59.
2. World Health Organization Osteoarthritis. 2023;https://www.who.int/news-room/fact-sheets/detail/osteoarthritis.
3. Hunter DJ, March L, Chew M Osteoarthritis in 2020 and beyond: a Lancet Commission. *Lancet* 2020;396:1711-12.
4. Duong V, Oo WM, Ding C, et al. Evaluation and treatment of knee pain: a review. *JAMA* 2023;330:1568-80.
5. Jiang T, Lau S-H, Zhang J, et al. Radiomics signature of osteoarthritis: Current status and perspective. *J Orthop Translat* 2024;45:100-06.
6. Roemer FW, Guermazi A, Demehri S, et al. Imaging in osteoarthritis. *Osteoarthritis Cartilage* 2022;30:913-34.
7. Chu CR, Williams AA, Coyle CH, et al. Early diagnosis to enable early treatment of pre-osteoarthritis. *Arthrit Res Ther* 2012;14:212.
8. Mahmoudian A, Lohmander LS, Mobasheri A, et al. Early-stage symptomatic osteoarthritis of the knee — time for action. *Nat Rev Rheumatol* 2021;17:621-32.
9. Loeser RF, Collins JA, Diekman BO Ageing and the pathogenesis of osteoarthritis. *Nat Rev Rheumatol* 2016;12:412-20.
10. Bielajew BJ, Hu JC, Athanasiou KA Collagen: quantification, biomechanics and role of minor subtypes in cartilage. *Nat Rev Mater* 2020;5:730-47.
11. Sandell LJ, Aigner T Articular cartilage and changes in arthritis: Cell biology of osteoarthritis. *Arthrit Res Ther* 2001;3:107.
12. Li X, Zhang Q, Yu SM, et al. The chemistry and biology of collagen hybridization. *J Am Chem Soc* 2023;145:10901-16.
13. Kaufmann JO, Brangsch J, Kader A, et al. ADAMTS4-specific MR probe to assess aortic aneurysms in vivo using synthetic peptide libraries. *Nat Commun* 2022;13:2867.
14. Lin C-Y, Wang Y-L, Chen Y-J, et al. Collagen-binding peptides for the enhanced imaging, lubrication and regeneration of osteoarthritic articular cartilage. *Nat Biomed Eng* 2022;6:1105-17.
15. Poole AR Biochemical/immunochemical biomarkers of osteoarthritis: utility for prediction of incident or progressive osteoarthritis. *Rheumatic Disease Clinics* 2003;29:803-18.
16. Billinghurst RC, Dahlberg L, Ionescu M, et al. Enhanced cleavage of type II collagen by collagenases in osteoarthritic articular cartilage. *J Clin Invest* 1997;99:1534-45.
17. Rousseau J-C, Chapurlat R, Garnero P Soluble biological markers in osteoarthritis. *Ther Adv Musculoskelet Dis* 2021;13:1759720X211040300.
18. Fang H, Beier F Mouse models of osteoarthritis: modelling risk factors and assessing outcomes. *Nat Rev Rheumatol* 2014;10:413-21.
19. Glasson SS, Blanchet TJ, Morris EA The surgical destabilization of the medial meniscus (DMM) model of osteoarthritis in the 129/SvEv mouse. *Osteoarthritis Cartilage* 2007;15:1061-69.
20. Li H, Wang D, Yuan Y, et al. New insights on the MMP-13 regulatory network in the pathogenesis of early osteoarthritis. *Arthrit Res Ther* 2017;19:248.
21. Mehana E-SE, Khafaga AF, El-Blehi SS The role of matrix metalloproteinases in osteoarthritis pathogenesis: An updated review. *Life Sci* 2019;234:116786.
22. Bremer C, Tung C-H, Weissleder R In vivo molecular target assessment of matrix metalloproteinase inhibition. *Nat Med* 2001;7:743-48.
23. Han EH, Chen SS, Klisch SM, et al. Contribution of proteoglycan osmotic swelling pressure to the compressive properties of articular cartilage. *Biophys J* 2011;101:916-24.
24. Bansal PN, Joshi NS, Entezari V, et al. Contrast enhanced computed tomography can predict the glycosaminoglycan content and biomechanical properties of articular cartilage. *Osteoarthritis Cartilage* 2010;18:184-91.
25. Inagawa K, Oohashi T, Nishida K, et al. Optical imaging of mouse articular cartilage using the glycosaminoglycans binding property of fluorescent-labeled octaarginine. *Osteoarthritis Cartilage* 2009;17:1209-18.
26. Wen ZH, Tang CC, Chang YC, et al. Glucosamine sulfate reduces experimental osteoarthritis and nociception in rats: association with changes of mitogen-activated protein kinase in chondrocytes. *Osteoarthritis Cartilage* 2010;18:1192-202.
27. Xie R, Yao H, Mao AS, et al. Biomimetic cartilage-lubricating polymers regenerate cartilage in rats with early osteoarthritis. *Nat Biomed Eng* 2021;5:1189-201.
28. Singh A, Corvelli M, Unterman SA, et al. Enhanced lubrication on tissue and biomaterial surfaces through peptide-mediated binding of hyaluronic acid. *Nat Mater* 2014;13:988-95.

29. Salamanna F, Giavaresi G, Parrilli A, et al. Effects of intra-articular hyaluronic acid associated to
Chitlac (arty-duo®) in a rat knee osteoarthritis model. *J Orthop Res* 2019;37:867-76.
30. Kraus VB, Collins JE, Hargrove D, et al. Predictive validity of biochemical biomarkers in knee
osteoarthritis: data from the FNIH OA Biomarkers Consortium. *Ann Rheum Dis* 2017;76:186-95.
31. Hunter DJ, Deveza LA, Collins JE, et al. Multivariable modeling of biomarker data from the phase I
foundation for the national institutes of health osteoarthritis biomarkers consortium. *Arthritis Care Res*
2022;74:1142-53.
32. Pritzker KPH, Gay S, Jimenez SA, et al. Osteoarthritis cartilage histopathology: grading and staging.
Osteoarthritis Cartilage 2006;14:13-29.
33. Englund M. The role of biomechanics in the initiation and progression of OA of the knee. *Best Pract*
Res Clin Rheumatol 2010;24:39-46.
34. Wang X, Wu Q, Zhang R, et al. Stage-specific and location-specific cartilage calcification in
osteoarthritis development. *Ann Rheum Dis* 2023;82:393.
35. O'Neill TW, McCabe PS, McBeth J Update on the epidemiology, risk factors and disease outcomes of
osteoarthritis. *Best Pract Res Clin Rheumatol* 2018;32:312-26.
36. Nia HT, Bozchalooi IS, Li Y, et al. High-bandwidth AFM-based rheology reveals that cartilage is most
sensitive to high loading rates at early stages of impairment. *Biophys J* 2013;104:1529-37.
37. Han G, Boz U, Eriten M, et al. Glycosaminoglycan depletion increases energy dissipation in articular
cartilage under high-frequency loading. *J Mech Behav Biomed Mater* 2020;110:103876.
38. Elahi SA, Castro-Viñuelas R, Tanska P, et al. Contribution of collagen degradation and proteoglycan
depletion to cartilage degeneration in primary and secondary osteoarthritis: an in silico study.
Osteoarthritis Cartilage 2023;31:741-52.
39. Diamond LE, Grant T, Uhlrich SD Osteoarthritis year in review 2023: Biomechanics. *Osteoarthritis*
Cartilage 2024;32:138-47.
40. Guilak F Biomechanical factors in osteoarthritis. *Best Pract Res Clin Rheumatol* 2011;25:815-23.
41. Zhou X, Anne KH, Benjamin ES, et al. Detection of glycosaminoglycan loss in articular cartilage by
fluorescence lifetime imaging. *J Biomed Opt* 2018;23:126002.
42. Deng R, Zhao R, Zhang Z, et al. Chondrocyte membrane–coated nanoparticles promote drug retention
and halt cartilage damage in rat and canine osteoarthritis. *Sci Transl Med*16:eadh9751.
43. Zhou L, Xu J, Schwab A, et al. Engineered biochemical cues of regenerative biomaterials to enhance
endogenous stem/progenitor cells (ESPCs)-mediated articular cartilage repair. *Bioact Mater*
2023;26:490-512.
44. Cho Y, Jeong S, Kim H, et al. Disease-modifying therapeutic strategies in osteoarthritis: current status
and future directions. *Exp Mol Med* 2021;53:1689-96.
45. Roemer FW, Eckstein F, Hayashi D, et al. The role of imaging in osteoarthritis. *Best Pract Res Clin*
Rheumatol 2014;28:31-60.
46. Hayashi D, Roemer FW, Guermazi A Imaging of osteoarthritis—recent research developments and
future perspective. *Br J Radiol* 2018;91:20170349.
47. Lv Z, Yang YX, Li J, et al. Molecular classification of knee osteoarthritis. *Front Cell Dev Biol* 2021;9.
48. Yuan C, Pan Z, Zhao K, et al. Classification of four distinct osteoarthritis subtypes with a knee joint
tissue transcriptome atlas. *Bone Res* 2020;8:38.
49. Mobasheri A, Saarakkala S, Finnilä M, et al. Recent advances in understanding the phenotypes of
osteoarthritis. *F1000Res* 2019;8.
425
426

Acknowledgments We thank Profs. Jiang Xia, Chao Liu, and Hang Lin for the informative
discussion, Dr. Xianghe Xu for the guidance in producing the DMM model, and Yuqi Lin for
assistance during MR imaging. Online supplemental figure S2 was created with
BioRender.com.

Contributors Conceptualization: YL, DS. Methodology: HK, YL, RQ, YF, XL, ZL, TJ.
Investigation: HK, RQ, YF, DZ, XL, ZL, TJ, YXF (Yuxiang Fei), YH. Visualization: HK,
RQ, YF, DZ, DNZ (Daoning Zhou), WW, YZ, YL. Funding acquisition: YL, DS. Project
administration: DS, YL. Supervision: SL, DS, YL. Writing—original draft: HK, RQ, DZ, YF,
XL, YL.Writing—review & editing: YL, DS.

Funding This work was supported by the National Natural Science Foundation of China
(92059104, 82071977, 82325035, 82172481, 32271409), the 2018 High-level Health Team
of Zhuhai, the Six Talent Peaks Project of Jiangsu Province (WSW-079), the Innovation
Project of National Orthopedics and Sports Medicine Rehabilitation Clinical Medical
Research Center (2021-NCRC-CXJJ-ZH-16), and the Guangdong-Hong Kong-Macao
University Joint Laboratory of Interventional Medicine Foundation of Guangdong Province
(2023LSYS001).

Competing interests The authors declare no competing interests.

Ethics approval In the study, all animal experiments were approved by the experimental
animal use and ethics committee of the Fifth Affiliated Hospital of Sun Yat-sen University
(protocol number: 00151), and all experimental plans involving knee specimens from patients
were approved by the Ethics Committee of Nanjing Drum Tower Hospital (number: K228-1).

Data availability statement All data are available in the main text or the supplementary
materials. The online supplementary materials include supplementary materials and methods,
supplementary table S1, supplementary figures S1-S17, and supplementary references.

456
457

Figures

Figure 1   Molecular collagen denaturation in the knee articular cartilage during early osteoarthritis (OA) revealed through histologic and in vivo collagen hybridization. (A) Unlike standard hard tissue processing [decalcification (decal) before CHP staining] that results in extensive artificial collagen denaturation and Cy3-CHP staining (top) in bone (*) in cryosections of normal mouse knee joints, our customized processing (staining before decalcification) leaves the triple-helical folding of collagen molecules intact (bottom, detailed in online supplemental figure S1). (B) Representative images of cryosections from the knee joints of C57BL/6 mice with surgical destabilization of the medial meniscus (DMM, right knee) or sham operation (left knee), harvested weekly post-operation ($n = 3$ mice per week) for histology with Safranin O/fast green (top) and Cy3-CHP (bottom). (C) Up-close histology images of a mouse knee joint one-week post-DMM surgery (OARSI = 0.5) exhibiting overall intact cartilage morphology but with Cy3-CHP fluorescence localizing to micrometer-size surface defects with minor loss in Safranin-O coloration (arrowheads in B, C). (D) Total Cy3-CHP fluorescence intensities quantified from all knee joint sections (images in B and fig. S3), plotted against their corresponding OARSI grades. (E) Representative fluorescence images of the mice's knee joints collected (3 h post intravenous injection of 1 nmol of Cy5-CHP) weekly post-DMM-operation ($n = 3$ mice per week). (F)

478   Excess Cy5-CHP fluorescence signals from the DMM-injured knee compared to the

479   paired sham-operated knee in individual mice measured each week post-operation.

480   Additional images and data are shown in online supplemental figures S1-S6. Data are

481   mean ± s.e.m. and were statistically analyzed using one-way ANOVA followed by

482   Tukey's multiple-comparison tests (F). Scale bars: 300 µm (A-C).

Figure 2 Spatiotemporal landscape of molecular collagen defects in knee joint components during early OA progression. (A) Anatomy of the right knee joint in the anterior and superior views. Scissors indicate the location of DMM surgery. (B) Light-sheet fluorescence microscopy images of the sham-operated mouse knee joints compared to the joints collected 1 to 6 weeks post-DMM-operation. The specimens were harvested for clearing 3 h post-intravenous injection of Cy5-CHP, which was uptaken by the

joint tissues in vivo. Arrowheads indicate locations with notable collagen denaturation in cartilage (yellow) and ligaments (white). (C to F) Three-dimensional (3D) reconstruction of the mice's joints showing Cy5-CHP fluorescence signals from cartilage (C), ligament (D), and meniscus (E, *i*→*iv*: counterclockwise) overlaid in (F, G), Changes in the lesion volume with molecular collagen defects in the cartilage, ligaments, and meniscus post-DMM operation, measured by the volumes of CHP-positive tissues identified in the 3D fluorescence reconstruction (C to E). Additional images and results are in online supplemental figures S7-S9. Data are mean ± s.e.m. ($n = 3$ mice per time point) and were statistically analyzed using one-way ANOVA followed by Tukey's multiple-comparison tests (G). Scale bars: 500 μm (B), 300 μm (C-F).

Figure 3 Early collagen denaturation post-DMM-injury is driven by mechanical insults, not MMPs. (A) Schematic of the DMM-injured mice (n = 3 mice per time point) administered in vivo with both Cy5-CHP and MMPSense750 for knee joint imaging. (B) Representative up-close ex vivo fluorescence images of the mice's knee joints 4- and 6-weeks post-operation showing distinct localization of Cy5-CHP (3 h p.i.) and MMPSense750 (24 h p.i.) signals. (C) Paired images of cryosections from the DMM (or sham) operated knee joints harvested weekly post-operation and co-stained with an anti-MMP13 antibody (immunohistochemistry) and Cy3-CHP. (D) Cy3-CHP

fluorescence quantified for each cartilage section showed poor correlation with the number of MMP13-positive cells, as interpreted by the Pearson correlation coefficient. (E) Schematic of the setup for the mechanical loading tests on an articular osteochondral plug specimen (diameter: 1 cm) excised from the porcine tibia plateau. (F) Representative fluorescence images (left) and quantification (right) of normal cartilage samples stained with Cy3-CHP after 0-20 cycles of 400 N (i.e., 5.1 MPa) loading. Data: median with interquartile range, statistics: Kruskal-Wallis, Dunn's. (G) Representative fluorescence images (left) and quantification (right) of the normal or chondroitinase ABC (ChABC) treated cartilage samples after 100 or 400 N loading and staining with Cy3-CHP. (H) Representative fluorescence images of the cryosections of the Cy3-CHP-stained cartilage samples from (G). A neighboring section was stained by a highly cationic peptide Cy5-Ahx-RRRRRRRR (Cy5-R8) to verify the GAG loss due to ChABC treatments. (I) Experimental design and outcome, recorded as the quantified fluorescence increment from sequential Cy3-CHP staining of normal or ChABC-treated porcine cartilage samples before and after 5 cycles of 100 N loading. Additional images and results are in online supplemental figures S10-S12. Data are mean ± s.e.m. Statistical analysis was performed using one-way ANOVA followed by Tukey's multiple-comparison tests (G and I). Scale bars: 34 μm (C), 250 μm (H).

Figure 4 Evaluation of treatments for early OA by collagen hybridization. (A) DMM-operated SD rats (*n* = 3–4 mice per group) were treated with glucosamine sulfate (GlcN, daily, oral) or hyaluronic acid (HA, weekly, intra-articular injection) and monitored longitudinally through in vivo Cy5-CHP fluorescence imaging. (B) Representative in vivo fluorescence images obtained 4–7 weeks post-operation showing minimal Cy5-CHP uptake in each DMM-operated knee joint for the GlcN and HA groups compared to the untreated DMM group. The fluorescence display of each image has been individually adjusted such that a minimal intensity is shown for the sham-operated knee. (C) Serial measurements of the ratio of the in vivo Cy5-CHP fluorescence signals between the DMM- and sham-operated knee joints of the rats in each group. (D and E) Representative micrographs of endpoint histology at 7 weeks post-operation showing the knee cartilage integrity of the rats from each group (D, Safranin O stain) and the matching fluorescence (E) due to the in vivo cartilage uptake of Cy5-CHP (harvested 3 h post-CHP-injection). The tibial plateau cartilage of the sham knee was intact with low CHP binding to the surface; for the GlcN- or HA-treated DMM cartilage, minor GAG loss with modest surface irregularity and fibrillation was found in locations with moderate CHP fluorescence. In contrast, the cartilage of the untreated DMM joints showed severe defects extending to the subchondral bone with substantial CHP uptake in the remaining surrounding cartilage.

552   Additional images are in online supplemental figure S13 and S14. Data are mean ±

553   s.e.m. and analyzed using repeated measures two-way ANOVA followed by Tukey's

554   multiple-comparison tests (C). Scale bars: 250 μm (D and E).

Figure 5 In vivo magnetic resonance (MR) imaging of molecular collagen defects in DMM-injured rat knee joints. (A) Chemical structure of contrast agent Gd_n-Cy5-CHP and its measured longitudinal relaxivity compared to Gd-DTPA (Magnevist). (B) Representative in vivo T1WI images of the sham- or DMM-operated knee joints of the same rat 7 weeks post-surgery. The rats were scanned before and after intra-articular injection of 250 nmol of Gd_n-Cy5-CHP. (C) Contrast-to-noise ratio (CNR, detailed in online supplementary materials) quantified from regions of interest in the medial tibial plateau cartilage of the knee pairs before and after contrast enhancement in individual rats imaged in c ($n = 4$, statistically analyzed using paired t-test). (D) Representative in vivo T1WI images of the DMM-injured rat knee joints 7 weeks post-operation pre- and post-contrast enhancement with Gd_n-Cy5-CHP, compared to

the T2WI image of the same joint. (E) Images of the Safranin O stain (bottom) and in situ fluorescence (top) of the sections from the cartilage within the boxed region in (D), highlighting Gd_n-Cy5-CHP's selective binding to spots with GAG loss in vivo. (F) Repeated in vivo T1WI (with or without Gd_n-Cy5-CHP contrast enhancement) and T2WI scans of the same DMM-injured rat knee joint 3- and 7-week post-operation. Additional data in online supplemental figure S15. Scale bars: 250 µm (E).

Figure 6 Ex vivo identification of molecular collagen defects in degenerated cartilage from OA patients. (A) Photograph of representative femoral cartilage specimens surgically removed from OA patients during knee replacement. i-iii: Regions with mild-to-severe levels of cartilage degeneration and OARSI grades. Osteochondral plug specimens were excised from regions i-iii ($n = 5$ plugs per group, diameter: 6 mm). (B to D) Representative images of the histology (B, Safranin O stain), T1WI magnetic resonance (C), and fluorescence imaging (D) of the plug specimens labeled with Gd_n-Cy5-CHP ex vivo. (E) Quantification of the fluorescence intensity from Gd_n-Cy5-CHP in the cartilage samples of groups i-iii. (F) In situ Gd_n-Cy5-CHP fluorescence in cryosections adjacent to the ones in (B), cut from the plug specimens of groups i-iii. (F) Three-dimensional light-sheet fluorescence scans of a life-size femoral knee cartilage specimen from an OA patient labeled with Cy5-CHP. Cell nuclei were stained with YO-PRO-1. The boxed area is zoomed in on the right. Data are mean ± s.e.m. and statistical analysis was performed using one-way ANOVA followed by Tukey's multiple-comparison test (E). Additional images in online supplemental figure S16 and S17. Scale bars: 250 μm (B and F), 3 mm (G).