JAK inhibition decreases the autoimmune burden in Down syndrome

Author list: Angela L. Rachubinski[1,2*], Elizabeth Wallace[3], Emily Gurnee[3], Belinda A. Enriquez Estrada[1], Kayleigh R. Worek[1], Keith P. Smith[1], Paula Araya[1], Katherine A. Waugh[1,4], Ross E. Granrath[1], Eleanor Britton[1], Hannah R. Lyford[1], Micah G. Donovan[1,5], Neetha Paul Eduthan[1], Amanda A. Hill[1], Barry Martin[6], Kelly D. Sullivan[1,7], Lina Patel[1,8], Deborah J. Fidler[9], Matthew D. Galbraith[1,5], Cory A. Dunnick[3], David A. Norris[3], Joaquin M. Espinosa[1,5*].

Affiliations:

[1]Linda Crnic Institute for Down Syndrome, University of Colorado Anschutz Medical Campus, Aurora, CO, USA.

[2]Department of Pediatrics, Section of Developmental Pediatrics, University of Colorado Anschutz Medical Campus, Aurora, CO, USA.

[3]Department of Dermatology, University of Colorado Anschutz Medical Campus, Aurora, CO, USA.

[4]Current address: Department of Cell Biology and Physiology, University of Kansas Medical Center, Kansas City, KS, USA.

[5]Department of Pharmacology, University of Colorado Anschutz Medical Campus, Aurora, CO, USA

[6]Department of Internal Medicine, University of Colorado Anschutz Medical Campus, Aurora, CO, USA.

[7]Department of Pediatrics, Section of Developmental Biology, University of Colorado Anschutz Medical Campus, Aurora, CO, USA.

[8]Department of Psychiatry, Child and Adolescent Division, University of Colorado Anschutz Medical Campus, Aurora, CO, USA.

[9]Department of Human Development and Family Studies, Colorado State University, Fort Collins, CO, USA.

24   *Corresponding authors:

25   joaquin.espinosa@cuanschutz.edu / angela.rachubinski@cuanschutz.edu

Abstract.

Individuals with Down syndrome (DS), the genetic condition caused by trisomy 21 (T21), display clear signs of immune dysregulation, including high rates of autoimmune disorders and severe complications from infections. Although it is well established that T21 causes increased interferon responses and JAK/STAT signaling, elevated autoantibodies, global immune remodeling, and hypercytokinemia, the interplay between these processes, the clinical manifestations of DS, and potential therapeutic interventions remain ill defined. Here, we report a comprehensive analysis of immune dysregulation at the clinical, cellular, and molecular level in hundreds of individuals with DS. We demonstrate multi-organ autoimmunity of pediatric onset concurrent with unexpected autoantibody-phenotype associations. Importantly, constitutive immune remodeling and hypercytokinemia occur from an early age prior to autoimmune diagnoses or autoantibody production. We then report the interim analysis of a Phase II clinical trial investigating the safety and efficacy of the JAK inhibitor tofacitinib through multiple clinical and molecular endpoints. Analysis of the first 10 participants to complete the 16-week study shows a good safety profile and no serious adverse events. Treatment reduced skin pathology in alopecia areata, psoriasis, and atopic dermatitis, while decreasing interferon scores, cytokine scores, and levels of pathogenic autoantibodies without overt immune suppression. Additional research is needed to define the effects of JAK inhibition on the broader developmental and clinical hallmarks of DS. ClinicalTrials.gov identifier: NCT04246372.

Introduction.

Trisomy of human chromosome 21 (T21) occurs at a rate of ~1 in 700 live births, causing Down syndrome (DS)[1,2]. Individuals with DS display a distinct clinical profile including developmental delays, stunted growth, cognitive impairments, and increased risk of leukemia, autism spectrum disorders, seizure disorders, and Alzheimer's disease[2,3]. People with DS also display widespread immune dysregulation, which manifests through severe complications from respiratory viral infections and high prevalence of myriad immune conditions, including autoimmune thyroid disease (AITD)[4-6], celiac disease[7,8], and skin conditions such as atopic dermatitis, alopecia areata, hidradenitis suppurativa (HS), vitiligo, and psoriasis[9-11]. Furthermore, people with DS display signs of neuroinflammation from an early age[12-14]. Although it is now well accepted that immune dysregulation is a hallmark of DS, the underlying mechanisms and therapeutic implications are not yet fully defined.

We previously reported that T21 causes consistent activation of the interferon (IFN) transcriptional response in multiple immune and non-immune cell types with concurrent hypersensitivity to IFN stimulation and hyperactivation of downstream JAK/STAT signaling[15-17]. Plasma proteomics studies identified dozens of inflammatory cytokines with mechanistic links to IFN signaling that are elevated in people with DS[18]. A large metabolomics study revealed that T21 drives the production of neurotoxic tryptophan catabolites via the IFN-inducible kynurenine pathway[19]. Deep immune profiling revealed global immune remodeling with hypersensitivity to IFN across all major branches of the immune system[15], and dysregulation of T cell lineages toward a hyperactive, autoimmunity-prone state[16]. These results could be partly explained by the fact that four of the six IFN receptors (IFNRs) are encoded on chr21, including Type I, II and III IFNR subunits[20]. In a mouse model of DS, normalization of *IFNR* gene copy number rescues multiple phenotypes of DS, including lethal immune hypersensitivity, congenital heart defects (CHDs), cognitive impairments, and craniofacial anomalies[21]. JAK inhibition rescues lethal immune hypersensitivity in these mouse models[22] and attenuates the global dysregulation of gene expression caused by the trisomy across multiple murine tissues[23]. Furthermore, prenatal JAK inhibition in pregnant

mice prevents the appearance of CHDs[24]. Altogether, these results support the notion that T21 elicits an

interferonopathy in DS, and that pharmacological inhibition of IFN signaling could have multiple

therapeutic benefits in this population.

Although it is now well established that T21 disrupts immune homeostasis toward an autoimmunity-

prone state, the interplay between overexpression of chromosome 21 genes, hyperactive interferon

signaling, dysregulation of immune cell lineages, autoantibody production, hypercytokinemia, and the

various developmental and clinical features of DS remain to be elucidated. Previous studies established

similarities between the immune profiles of typical aging, autoimmunity in the general population, and

DS, proposing a role for accelerated immune aging in the pathophysiology of DS[25-27]. Other studies

indicate a role for elevated cytokine production, hyperactivated T cells, and ongoing B cell activation as

drivers of autoimmunity in DS[15,16,28]. However, given the relatively small sample sizes and observational

nature of these studies, it has not been possible to define the contribution of specific dysregulated events

to breach of tolerance leading to clinically evident autoimmunity in DS. Therefore, additional research is

needed to define driver versus bystander events that could illuminate therapeutic strategies to decrease the

burden of autoimmunity in DS.

Within this context, we report here a comprehensive analysis of the immune disorder of DS, including

detailed annotation of autoimmune and inflammatory conditions and quantification of autoantibodies in

hundreds of research participants, which reveals widespread autoimmune attack on all major organ

systems in DS from an early age, including unexpected autoantibody-phenotype associations. Then, using

deep immune mapping and quantitative proteomics, we demonstrate that T21 causes widespread immune

remodeling toward an autoimmunity-prone state accompanied by hypercytokinemia prior to clinically

evident autoimmunity or autoantibody production. Lastly, we report the interim analysis of a clinical trial

investigating the safety and efficacy of the JAK1/3 inhibitor tofacitinib (Xeljanz, Pfizer) in DS. These

results demonstrate that JAK inhibition improves multiple immunodermatological conditions in DS,

normalizes interferon scores, decreases levels of major pathogenic cytokines (e.g., TNF-α, IL-6), and

reduces levels of pathogenic autoantibodies [e.g., anti-thyroid peroxidase (anti-TPO)]. Altogether, these

results point to hyperactive JAK/STAT signaling as driver of autoimmunity in DS and justify the ongoing

trials of JAK inhibitors in DS for multiple clinical endpoints.

Results.

Widespread multi-organ autoimmunity and autoantibody production in Down syndrome.

Previous studies have documented increased rates of diverse autoimmune conditions in DS relative to the general population including autoimmune thyroid disease (AITD)[29], celiac disease[30], autoimmune skin conditions[9-11], and type I diabetes[31,32]. However, many of these studies were limited by relatively small sample sizes, independent analysis of individual autoimmune conditions, or a focus on specific age ranges. In order to complete a more comprehensive analysis of autoimmune conditions in DS across the lifespan, we analyzed the harmonized clinical profiles of 441 research participants with DS, aged 6 months to 57 years, enrolled in the Human Trisome Project cohort study (HTP, NCT02864108), which annotates clinical data through a combination of participant/caregiver surveys and expert abstraction of electronic health records (EHRs) (see **Materials and Methods**, **Supplementary file 1**). In this analysis, the most common autoimmune condition is AITD, affecting 53.1% of the total cohort (**Figure 1a**, **Figure 1 – figure supplement 1a**). Grouped together, autoimmune and inflammatory skin conditions represent the second most common category, affecting 43% of the cohort, including: atopic dermatitis / eczema (27.9%), hidradenitis suppurativa / folliculitis / boils (20.6%), alopecia areata (7.7%), psoriasis (6.1%), and vitiligo (1.9%) (**Figure 1a**, **Figure 1 - supplement 1b**). The rate of celiac disease (9.6%) is also highly elevated over that of the general population[33]. We observed 10 cases (2.2%) of juvenile Type I diabetes, which has been reported to be more common in DS[31,32]. Other autoimmune conditions common in the general population, such as systemic lupus erythematosus or multiple sclerosis, were not observed in the HTP cohort.

In the general population, the risk of autoimmune conditions increases with age and is higher in females, with autoimmune conditions tending to cluster, whereby occurrence of one autoimmune condition predisposes to a second condition[34,35]. Within the HTP cohort, analysis of age trajectories of immune-related conditions in DS revealed early onset, with >80% of AITD, autoimmune/inflammatory skin conditions, and celiac disease being diagnosed in the first two decades of life (**Figure 1 - supplement**

1c-e). The cumulative burden of autoimmunity and autoinflammation is similar in males versus females

with DS, albeit with slightly increased rates of AITD and hidradenitis suppurativa in females (**Figure 1 -**

supplement 1f). In terms of co-occurrence, when evaluating the adult population (18+ years old) for

AITD, autoimmune/inflammatory skin conditions and celiac disease, we found that 75% of participants

had a history of at least one condition, 38.4% had at least two, and 13.6% had three or more conditions

(**Figure 1b**).

Interestingly, analysis of medical records found an unexpectedly low number of individuals with

records of autoantibodies against the thyroid gland [4.3%, e.g., anti-thyroid peroxidase (TPO), anti-

thyroglobulin (TG)] within the HTP cohort (**Figure 1 - supplement 1a**). This could be explained by the

fact that thyroid disease is commonly diagnosed through measurements of thyroid-relevant hormones

(TSH, T3, T4) without concurrent testing of autoantibodies. To investigate further, we measured anti-TPO

levels as well as levels of anti-nuclear antibodies (ANA), a more general biomarker of autoimmunity

(**Supplementary file 2**). Remarkably, 82.4% of adults with DS show positivity for at least one of these

autoantibodies, with 41.2% being positive for both (**Figure 1c**). Indeed, 62% of individuals with history

of hypothyroidism were TPO+, whereby anti-TPO is just one of the possible autoantibodies associated

with AITD. Prompted by these results, we next completed a more comprehensive analysis of

autoantibodies in DS using protein array technology, with a focus on ~380 common autoepitopes from

270 proteins (see **Materials and Methods, Supplementary file 2**). These efforts identified 25

autoantibodies significantly over-represented in people with DS relative to age- and sex-matched controls

(**Figure 1d**), with 98.3% of individuals with DS being positive for at least one of these autoantibodies,

and 63.3% being positive for six or more (**Figure 1d-e**). In addition to autoantibodies against TPO, which

is expressed exclusively in the thyroid gland, we identified autoantibodies targeting proteins that are either

broadly expressed across multiple tissues (e.g., TOP1, UBA1, LAMP2) or preferentially expressed in

specific organs across the human body, including liver (e.g., CYP1A2), pancreas (e.g., SLC30A8), skin

(e.g., DSG3), bone marrow (e.g. SRP68), and brain tissue (e.g., AIMP1) (**Figure 1d, f**).

Analysis of autoantibody positivity relative to history of co-occurring conditions produced several interesting observations. Expectedly, individuals with hypothyroidism are more likely to be positive for anti-TPO antibodies (**Figure 1g-h**). However, unexpectedly, TPO+ status also associates with higher rates of use of pressure equalizing (PE) tubes employed to alleviate the symptoms of recurrent ear infections and otitis media with effusion (OME), which is common in DS[36] (**Figure 1g-h**). Possible interpretations for this result are provided in the Discussion. Positivity for additional autoantibodies was more common in those with other co-occurring neurological conditions, a broad classification encompassing various seizure disorders, movement disorders, and structural brain abnormalities (**Figure 1g-h, Figure 1 – figure supplement 1g**). Salient examples are antibodies against MUSK, a muscle-associated receptor tyrosine kinase involved in clustering of the acetylcholine receptors in the neuromuscular junction[37]; UBA1, a ubiquitin conjugating enzyme involved in antigen presentation[38]; and MYH6, a cardiac myosin heavy chain isoform (**Figure 1g-h**, **Figure 1 – figure supplement 1g**). Individuals with history of tricuspid valve regurgitation display higher rates of four different autoantibodies, most prominently against WARS1, a tryptophan tRNA synthetase mutated in various neurodevelopmental disorders[39], and SRP68, a protein commonly targeted by autoantibodies in necrotizing myopathies[40] (**Figure 1 – figure supplement 1g**). Individuals with a history of frequent pneumonia present a higher frequency of autoantibodies against DSG3 (desmoglein 3), a cell adhesion molecule targeted by autoantibodies in paraneoplastic pemphigus (PNP), an autoimmune disease of the skin and mucous membranes that can involve fatal lung complications[41] (**Figure 1g-h**).

Altogether, these results demonstrate widespread multi-organ autoimmunity across the lifespan in people with DS, with production of multiple autoantibodies that could potentially contribute to a number of co-occurring conditions more common in this population.

Figure 1. Multi-organ autoimmunity and widespread autoantibody production in Down syndrome.

a, Overview of autoimmune and inflammatory conditions prevalent in persons with Down syndrome (DS) enrolled in the Human Trisome Project (HTP) cohort study. Percentages indicate the fraction of participants (n=441, all ages) with history of the indicated conditions. **b,** Pie chart showing

autoimmune/inflammatory condition burden in adults (n=278, 18+ years old) with DS. **c**, Pie chart

showing rates of positivity for anti-TPO and/or anti-nuclear antibodies (ANA) in adults (n=212, 18+ years

old) with DS. **d**, Bubble plot displaying odds-ratios and significance for 25 autoantibodies with elevated

rates of positivity in individuals with DS (n=120) versus 60 euploid controls (D21). q values calculated

by Benjamini-Hochberg adjustment of p-values from Fisher's exact test. **e**, Pie chart showing fractions of

adults with DS (n=120, 18+ years old) testing positive for various numbers of the autoantibodies identified

in d. **f**, Representative examples of autoantibodies more frequent in individuals with T21 (n=120) versus

euploid controls (D21, n=60). MAD: median absolute deviation. Dashed lines indicate the positivity

threshold of 90^{th} percentile for D21. Data are presented as modified sina plots with boxes indicating

quartiles. **g**, Bubble plots showing the relationship between autoantibody positivity and history of various

clinical diagnoses in DS (n=120). Size of bubbles is proportional to -log-transformed p values from

Fisher's exact test. **h**, Sina plots displaying the levels of selected autoantibodies in individuals with DS

with or without the indicated co-occurring conditions. MAD: median absolute deviation. Dashed lines

indicate the positivity threshold of 90^{th} percentile for D21. Sample sizes are indicated under each plot. q

values calculated by Benjamini-Hochberg adjustment of p-values from Fisher's exact tests.

Figure 1 – figure supplement 1. Early onset multi-organ autoimmunity and autoantibody production in Down syndrome. a-b, Upset plots showing overlap between various reported diagnoses indicative of autoimmune thyroid disease (a) or autoimmune/inflammatory skin conditions (b) in research participants with Down syndrome (DS, all ages, n=441) enrolled in the Human Trisome Project (HTP). **c-**

e, Plots showing the percentages of cases by age at diagnosis for AITD (c), autoimmune/inflammatory

skin conditions (d), and celiac disease (e). Sample sizes indicated in each chart. **f**, Odds ratio plot for

Fisher's exact test of proportions (cases vs. controls in males vs. females) for history of co-occurring

conditions in individuals with DS (all ages, total n=441). Conditions with q < 0.1 (10% FDR) are

highlighted in red. The size of square points is inversely proportional to q value; error bars represent 95%

confidence intervals. **g**, Sina plots displaying the levels of select autoantibodies in individuals with DS,

with or without history of the indicated co-occurring conditions. MAD: median absolute deviation.

Horizontal dashed lines indicate 90th percentiles for the D21 group. Sample sizes are indicated under each

plot. q values calculated by Benjamini-Hochberg adjustment of p-values from Fisher's exact tests.

Trisomy 21 causes global immune remodeling regardless of evident clinical autoimmunity.

Several immune cell changes have been proposed to underlie the autoimmunity-prone state of DS[15,16,25,42], but specific immune cell-to-phenotype associations have not been established in previous studies using smaller sample sizes. Therefore, we next investigated immune cell changes associated with various clinical and molecular markers of autoimmunity in DS. Toward this end we analyzed mass cytometry data from 292 individuals with DS relative to 96 euploid controls and tested for potential differences in immune cell subpopulations, identified using FlowSOM[43], within the DS cohort based on number of autoimmune/inflammatory disease diagnoses, ANA positivity, TPO positivity, and positivity for additional autoantibodies. In agreement with previous analyses[15,16,25,42], we observed massive immune remodeling in all major myeloid and lymphoid subsets, including increases in basophils, along with depletion of eosinophils and total B cells (**Figure 2a-c, Figure 2 – figure supplement 1a-c**). When comparing various subgroups within the DS cohort based on autoimmunity status, we observed that these global immune changes are largely independent of the presence of clinical diagnoses or autoantibody positivity, with very few additional changes significantly associated with these measures of autoimmunity (**Figure 2 – figure supplement 1d**). For example, the significant depletion of B cells and enrichment of basophils in DS is not significantly different among the various subgroups (**Figure 2c**, **Figure 2 – figure supplement 1c**). Among CD45+ CD66^{lo} non-granulocytes, most changes are conserved among subgroups, with the sole of exception of non-classical monocytes, which are further elevated in the ANA+ group (**Figure 2 – figure supplement 1d-e**). Among T cells, the overall pattern of depletion of naïve subsets and enrichment of differentiated subsets characteristic of DS[15,16,28,42] is conserved across subgroups, as illustrated by consistent depletion of CD8+ naïve subsets along with increases in the CD8+ terminally differentiated effector memory (TEMRA) subset (**Figure 2f, Figure 2 – figure supplement 1d**). Notably, we observed depletion of γδ T cells (both total and CD8+) in those with multiple autoimmune diagnoses (**Figure 2 – figure supplement 1d, f**), a result that is in line with reports documenting depletion of these subsets from peripheral circulation toward sites of active autoimmunity[44].

We also observed slight elevation of CD4+ T central memory cells (TCM) (**Figure 2 – figure supplement**

1d, f). Among B cells, the overall shift toward more differentiated states such as plasmablasts, age-

associated B cells (ABCs), and IgM+ memory cells is also conserved among subgroups, with the sole

exception of ABCs, which tend to be further elevated in the TPO+ group (**Figure 2g-i**, **Figure 2 – figure**

supplement 1d, g).

Altogether, these results indicate that T21 causes global remodeling of the immune system toward an

autoimmunity-prone and pro-inflammatory state, prior to clinically evident autoimmunity, and dwarfing

any additional effects associated with confirmed diagnoses of autoimmune/inflammatory conditions or

common biomarkers of autoimmunity.

Figure 2. Trisomy 21 causes global immune remodeling regardless of clinically evident

autoimmunity. a, t-distributed Stochastic Neighbor Embedding (t-SNE) plot displaying major immune

populations identified by FlowSOM analysis of mass cytometry data for all live cells (left) and color

coded by significant impact of T21 (beta regression q<0.1) on their relative frequency (right). Red

indicates increased frequency and blue indicates decreased frequency among research participants with

T21 (n=292) versus euploid controls (D21, n=96). **b**, Volcano plot showing the results of beta regression

analysis of major immune cell populations among all live cells in research participants with T21 (n=292)

versus euploid controls (D21, n=96). The dashed horizontal line indicates a significance threshold of

10% FDR (q<0.1) after Benjamini-Hochberg correction for multiple testing. **c**, Frequencies of B cells

among all live cells in euploid controls (D21, n=96) versus individuals with T21 and history of 0 (n=69),

1 (n=102) or 2+ (n=121) autoimmune/inflammatory conditions. Data is displayed as modified sina plots

with boxes indicating quartiles. **d-f**, Description as in a-c, but for subsets of T cells. **g-i**, Description as

in a-c, but for subsets of B cells.

Figure 2 – figure supplement 1. Consistent remodeling of the peripheral immune system in Down syndrome. a, t-distributed Stochastic Neighbor Embedding (t-SNE) plot displaying major immune populations identified by FlowSOM analysis of mass cytometry data for CD45+ CD66[lo] non-granulocytes (left) and color coded by the impact of trisomy 21 (T21) on their relative frequency (right). Red indicates increased frequency and blue indicates decreased frequency among research participants with T21 (n=292) versus euploid controls (D21, n=96). **b**, Volcano plot showing the results of beta regression analysis of immune cell populations among CD45+ CD66[lo] non-granulocytes from research participants with T21 (n=292) versus euploid controls (D21, n=96). The dashed horizontal line indicates

a significance threshold of 10% FDR (q<0.1) after Benjamini-Hochberg correction for multiple testing.

c, Frequencies of basophils among all live cells in euploid controls (D21, n=96) versus individuals with

T21 and history of 0 (n=44), 1 (n=71) or 2+ (n=88) autoimmune/inflammatory conditions. Data is

displayed as modified sina plots with boxes indicating quartiles. **d**, Heatmap summarizing the results of

beta regression testing for differences in frequencies of indicated immune cell populations among all

live cells, CD45+ CD66^{lo} non-granulocytes, T cells, and B cells by T21 (n=292) versus D21 (n=96)

status, or by different subgroups within the T21 cohort: 2+ (n=88) versus 0 (n=44)

autoimmune/inflammatory conditions; TPO+ (n=144) versus TPO- (n=148); ANA+ (n=124) versus

ANA- (n=49); or positivity for 8-20 (n=49) versus 0-7 (n=54) autoantibodies elevated in DS. Asterisks

indicate significance after Benjamini-Hochberg correction for multiple testing (q<0.1, 10% FDR). **e-g**,

Representative examples of immune cell populations from d, showing effects of ANA positivity (e),

number of autoimmune conditions (f), and TPO status (g). Data are presented as modified sina plots

with boxes indicating quartiles, with q-values indicating beta regression significance after Benjamini-

Hochberg correction for multiple testing.

Trisomy 21 causes hypercytokinemia from an early age independent of autoimmunity status.

It is well established that individuals with DS display elevated levels of many inflammatory markers, including several interleukins, cytokines, and chemokines known to drive autoimmune conditions, such as IL-6 and TNF-α[18,28,42,45]. However, the interplay between hypercytokinemia, individual elevated cytokines, and development of autoimmune conditions in DS remains to be elucidated. Therefore, we analyzed data available from the HTP cohort for 54 inflammatory markers in plasma samples from 346 individuals with DS versus 131 euploid controls and cross-referenced these data with the presence of autoimmune conditions and autoantibodies. These efforts confirmed the notion of profound hypercytokinemia in DS[18,28,42,45], with significant elevation of multiple acute phase proteins (e.g. CRP, SAA, IL1RA), pro-inflammatory cytokines (TSLP, IL-17C, IL-22, IL-17D, IL-9, IL-6, TNF-α) and chemokines (IP-10, MIP-3a, MIP-1a, MCP-1, MCP-4, Eotaxin), as well as growth factors associated with inflammation and wound healing (FGF, PlGF, VEGF-A) (**Figure 3a**). However, when evaluating for differences within the DS cohort based on various metrics of autoimmunity, we did not observe important differences based on number of autoimmune/inflammatory conditions, ANA or TPO positivity status, or number of other autoantibodies (**Figure 3a**). For example, CRP, IL-6, and TNF-α are equally elevated across all these subgroups (**Figure 3b-d, Figure 3 - figure supplement 1a**).

Previous studies have reported signs of early immunosenescence and inflammaging in DS, including accelerated progression of immune lineages toward terminally differentiated states, early thymic atrophy, and elevated levels of pro-inflammatory markers associated with age in the typical population[25,27,46,47]. However, the extent to which the inflammatory profile of DS represents accelerated ageing versus other processes remains ill-defined. To address this, we first identified age-associated changes in immune markers within the euploid and DS cohorts separately (**Figure 3 - figure supplement 1b-c**). This exercise identified multiple immune markers that were up- or down-regulated with age, with an overall conserved pattern of age trajectories in both groups (**Figure 3 - figure supplement 1c**). For example, increased age is associated with increased CRP levels and decreased IL-17B levels in both cohorts (**Figure 3 - figure**

supplement 1d). We then compared the effects of age versus T21 status on cytokine levels in the DS

cohort, which identified many inflammatory factors elevated in DS across the lifespan that do not display

a significant increase with age, such as IL-9 and IL-17C, or that increase with age only in the DS cohort,

such as IP-10 (**Figure 3e-f, Figure 3 - figure supplement 1e**).

Altogether, these results indicate that T21 induces a constitutive hypercytokinemia from early

childhood, with only a fraction of these inflammatory changes being exacerbated with age.

Figure 3. Trisomy 21 causes constitutive hypercytokinemia independent of autoimmunity status

from an early age. a, Heatmap displaying log2-transformed fold-changes for plasma immune markers

with significant differences in trisomy 21 (T21, n=346) versus euploid (D21, n=131), and between

different subgroups within the T21 cohort: history of 2+ (n=139) versus 0 (n=87)

autoimmune/inflammatory conditions (AI conds.); TPO+ (n=133) versus TPO- (n=162); ANA+ (n=100)

versus ANA- (n=39); or positivity for 8-20 (n=57) versus 0-7 (n=62) autoantibodies (AutoAbs) elevated

in DS. Asterisks indicate linear regression significance after Benjamini-Hochberg correction for multiple

testing (q<0.1, 10% FDR). **b-d**, Comparison of CRP, IL-6 and TNF-α levels in euploid controls (D21,

n=131) versus subsets of individuals with T21 based on number of autoimmune/inflammatory

conditions (**b**), ANA positivity (**c**) or TPO positivity (**d**). Data are presented as modified sina plots with

boxes indicating quartiles. Samples sizes as in a. q-values indicate linear regression significance after

Benjamini-Hochberg correction for multiple testing. **e**, Scatter plot comparing the effect of T21

karyotype versus the effect of age in individuals with T21 (n=54 immune markers in 346 individuals

with T21), highlighting immune markers that are significantly different by T21 status, age, or both. ns:

not significantly different by T21 status or age. **f**, Scatter plots for example immune markers that are

significantly elevated in T21, but which are either not elevated with age in the euploid (D21) cohort (i.e.,

IP-10), or in either the T21 (n=346) or D21 (n=131) cohorts. Lines represent least-squares linear fits

with 95% confidence intervals in grey.

Figure 3 – figure supplement 1. Consistent hypercytokinemia from an early age in Down syndrome. a, Comparison of CRP, IL-6, and TNF-α levels in euploid controls (D21, n=131) versus subsets of individuals with T21 based on number of autoantibodies commonly elevated in Down syndrome: 0-7 autoantibodies (n=62) versus 8-20 autoantibodies (n=57). Data are presented as modified

sina plots with boxes indicating quartiles. q-values indicate linear regression significance after

Benjamini-Hochberg correction for multiple testing. **b**, Volcano plots presenting the results of linear

regression testing for association between age and the levels of 54 immune markers in the plasma of

euploid controls (left, D21, n=131) and individuals with trisomy 21 (right, T21, n=346) enrolled in the

Human Trisome Project (HTP) study. Horizontal dashed lines indicate a significance threshold of 10%

FDR (q<0.1) after Benjamini-Hochberg correction for multiple testing. **c**, Heatmap comparing the effect

of age on levels of immune markers in D21 and T21. Heatmap color scale represents log2-transformed

mean fold-change per year of age; asterisks indicate significance (q < 0.1) for linear regression testing.

d, Scatter plots showing the age trajectories of select immune markers in D21 versus T21. Sample sizes

as in c. Lines represent least squares linear fits with shaded areas indicating 95% confidence interval. **e**,

Diagram representing the overlap between immune markers elevated in T21 versus D21 and those

elevated with age in T21.

A clinical trial for JAK inhibition in Down syndrome.

Several lines of evidence support the notion that IFN hyperactivity and downstream JAK/STAT signaling are key drivers of immune dysregulation in DS[15-19,22,24,42,48]. In mouse models of DS, both normalization of *IFNR* gene copy number and pharmacologic JAK1 inhibition rescue their lethal immune hypersensitivity phenotypes[22,48]. Furthermore, we recently demonstrated that IFN transcriptional scores derived from peripheral immune cells correlate significantly with the degree of immune remodeling and hypercytokinemia in DS[42], and we and others have reported the safe use of JAK inhibitors for treatment of diverse immune conditions in DS, including alopecia areata[49], psoriatic arthritis[50] and hemophagocytic lymphohistocytosis[51] through small case series. Encouraged by these results, we launched a clinical trial to assess the safety and efficacy of the JAK inhibitor tofacitinib (Xeljanz, Pfizer) in DS, using moderate-to-severe autoimmune/inflammatory skin conditions as a qualifying criterion (NCT04246372). This trial is a single-site, open-label, Phase II clinical trial enrolling individuals with DS between the ages of 12 and 50 years old affected by alopecia areata, hidradenitis suppurativa, psoriasis, atopic dermatitis, or vitiligo (see qualifying disease scores in **Supplementary file 3**). After screening, qualifying participants are prescribed 5 mg of tofacitinib twice daily for 16 weeks, with an optional extension to 40 weeks (**Figure 4a**, see **Materials and Methods**). After enrollment and assessments at a baseline visit, participants attend five safety monitoring visits during the main 16-week trial period. The recruitment goal for this trial is 40 participants who complete 16 weeks of tofacitinib treatment, with an IRB-approved qualitative interim analysis triggered when the first 10 participants completed the main 16-week trial (**Figure 4b**). Among the first 13 participants enrolled, one participant withdrew shortly after enrollment, one was excluded from analyses due to medication non-compliance (i.e., >15% missed doses), and one participant had not yet completed the trial at the time of the interim analysis (**Figure 4b**). Demographic characteristics of the 10 participants included in the interim analysis are shared in **Supplementary file 3**. Baseline qualifying conditions of the 10 participants included in the interim analysis were alopecia areata (n=6), hidradenitis suppurativa (n=3), and psoriasis (n=1) (open circles in **Figure 4c**). Two participants presented with

concurrent atopic dermatitis, two with concurrent vitiligo, and two with concurrent hidradenitis suppurativa, albeit below the severity required to be the qualifying conditions (see closed circles in **Figure 4c**). In addition, seven participants had AITD/TPO+ and three had a celiac disease diagnosis (**Figure 4c**).

Tofacitinib is well tolerated in Down syndrome.

Analysis of adverse events (AEs) recorded for the 10 first participants over 16 weeks did not identify any AEs considered definitely related to tofacitinib treatment or classified as severe. Several AEs were annotated as 'possibly related' to treatment (**Figure 4d, Supplementary file 4**). Five episodes of upper respiratory infections (URIs) affecting five different participants were observed. Based on the safety data for tofacitinib in the general population[52], all episodes of URIs were annotated as possibly related to treatment. Participant AA2 developed occasional cough and rhinorrhea that resolved with over-the-counter medication. Two other participants reported transient rhinorrhea (AA3, AA5). Participant AA4 developed a nasal congestion, with chest pain and a productive cough. This participant tested negative for SARS-Co-V2, Flu A-B, and RSV. Tofacitinib was not paused during this episode, and symptoms resolved with over-the-counter medication. Participant HS2 experienced a sore throat with middle ear inflammation that resolved with over-the-counter treatment. This participant also presented with folliculitis, which resolved with antibiotic treatment. Participant HS1 experienced a short transient elevation (<3 days) in creatine phosphokinase (CPK) that resolved spontaneously, and rash acneiform. Participant Ps1 experienced a transient and asymptomatic decrease in white blood cell (WBC) counts that resolved by the end of the trial.

Overall, tofacitinib treatment was not discontinued for any of the 10 participants over the 16-week study period, and seven participants eventually obtained off-label prescriptions after completing the trial and are currently taking the medicine. Based on these interim results, recruitment resumed and is ongoing.

Figure 4. Clinical trial for JAK inhibition in Down syndrome. a, Schedule of activities for clinical trial

of JAK inhibition in Down syndrome (NCT04246372). **b,** Consort chart for first 13 participants enrolled

in the clinical trial. **c,** Upset plot displaying the qualifying and co-occurring autoimmune/inflammatory

conditions for the 10 participants included in the interim analysis. **d,** Upset plots summarizing the adverse

events annotated for the first 10 participants over a 16-week treatment period.

Tofacitinib improves diverse autoimmune/inflammatory skin conditions in Down syndrome.

In the clinical trial, skin pathology is monitored using global metrics of skin health, including the Investigator's Global Assessment (IGA) and the Dermatology Life Quality Index (DLQI), as well as disease-specific scores, such as the severity of alopecia tool (SALT), the psoriasis area and severity index (PASI), or the eczema area and severity index (EASI) (see **Materials and Methods, Supplementary file 5**. The interim analysis showed that seven of the ten participants had an improvement in the IGA score and eight of the ten reported some improvement on their life quality related to their skin condition as measured by the DLQI (**Figure 5a-b**). The most striking effects were observed for alopecia areata (**Figure 5c-d, Figure 5 – figure supplement 1a**). Five of six participants with alopecia areata showed scalp hair regrowth, with the exception being a male participant (AA1) with history of alopecia totalis for 20+ years who only showed facial hair and eyelash re-growth. One participant presented with psoriasis due to psoriatic arthritis and experienced an almost complete remission of psoriatic arthritis symptoms (Ps1, **Figure 5 – figure supplement 1b-c**). For the two participants that presented with atopic dermatitis, the clinical manifestations were markedly reduced during tofacitinib treatment (**Figure 5e-f**). A total of five participants were affected by HS, three of them as the qualifying condition (HS1-3). No clear trend was seen in the Modified Sartorius Scale (MSS) score used to monitor HS (**Figure 5 – figure supplement 1d-e**).

Altogether, these results indicate that JAK inhibition could provide therapeutic benefit for several autoimmune/inflammatory skin conditions more common in DS.

Figure 5. Tofacitinib improves diverse immune skin pathologies in Down syndrome. a-b, Investigator global assessment (IGA) scores (a) and Dermatological Life Quality Index (DLQI) scores (b) for the first 10 participants at baseline visit (B), mid-point (8 weeks) and endpoint (16 weeks) visits. MD: median difference. **c,** Severity of Alopecia Tool (SALT) scores for the first seven participants with alopecia areata in the trial. **d,** Images of participant AA6 at baseline versus week 16. **e,** Eczema Area and Severity Index (EASI) scores for two participants with mild atopic dermatitis. **f,** Images of participant AA2 showing improvement in atopic dermatitis upon tofacitinib treatment. p values not shown as per interim analysis plan.

Figure 5 – figure supplement 1. Tofacitinib improves diverse skin pathologies in Down syndrome.

a, Images of five participants with alopecia areata at baseline and after 16 weeks of tofacitinib treatment.

b-c, Psoriasis Area and Severity Index score (b) and images (c) for participant with psoriatic arthritis. **d,**

Modified Sartorius Scale (MSS) scores for five participants with hidradenitis suppurativa (HS). MD:

444   median difference. **e**, Images for participant affected by HS at baseline and 16-week endpoint visit. p

445   values not shown as per interim analysis plan.

Tofacitinib normalizes IFN scores and decreases pathogenic cytokines and autoantibodies.

It is well demonstrated that individuals with DS display elevated IFN signaling across multiple immune and non-immune cell types[15-17,19]. Using an IFN transcriptional score composed of 16 interferon-stimulated genes (ISGs)[53] measured via bulk RNA sequencing of peripheral blood RNA, individuals with DS in the HTP cohort study show a significant increase in these scores[48] (**Figure 6a**). Reduction of IFN scores is designated as a primary endpoint in the trial. At baseline, clinical trial participants show IFN scores within the typical range for DS, but values are decreased at 2, 8, and 16 weeks of tofacitinib treatment (**Figure 6a, Supplementary file 6**). Time course analysis revealed that most participants show a decrease in IFN scores as soon as two weeks of treatment which is sustained over time, with two clear exceptions (**Figure 6 – figure supplement 1a**). At the 8-week study midpoint, 9 of 10 participants had decreased IFN scores relative to baseline, except participant AA2 who reported a COVID-19 vaccination three days prior to the visit and was pausing tofacitinib at the time of the blood draw (**Figure 6 – figure supplement 1a**). At the 16-week time point, nine of ten participants had decreased IFN scores, with the exception being AA4, who developed an URI in the week prior to the blood draw (**Figure 6 – figure supplement 1a**). Therefore, although all participants displayed decreased IFN scores at one or more time points during the treatment, IFN scores could be sensitive to immune triggers. Analysis of individual ISGs composing the IFN score revealed that whereas many ISGs elevated in DS display reduced expression upon tofacitinib treatment (e.g., *RSAD2*, *IFI44L*), others do not (e.g., *BPGM*) (**Figure 6b, Figure 6 – figure supplement 1b**). To investigate this further, we defined the impact of tofacitinib on all 136 ISGs significantly elevated in DS that are not encoded on chr21[42] (**Figure 6c**). Collectively, ISGs as a group are significantly downregulated upon tofacitinib treatment, but the effect is not uniform across all ISGs (**Figure 6c**), indicating that JAK1/3 inhibition does not reduce all IFN signaling elevated in DS, which could be explained by the fact that the IFN pathways also employ JAK2 for signal transduction[54,55]. Global analysis of transcriptome changes revealed that tofacitinib treatment reverses the dysregulation of many gene signatures observed in DS, effectively attenuating many pro-inflammatory signatures beyond IFN

gamma and alpha responses, such as Inflammatory Response, TNF-α signaling via NFkB, IL-2 STAT5 signaling, and IL-6 JAK STAT3 signaling (**Figure 6 – figure supplement 1c**). Tofacitinib also reversed elevation of genes involved in Oxidative Phosphorylation and dampened downregulation of gene sets involved in Wnt/Beta Catenin and Hedgehog Signaling (**Figure 6 – figure supplement 1c-d**). Conversely, tofacitinib did not rescue elevation of genes involved in Heme Metabolism or Mitotic Spindle (**Figure 6 – figure supplement 1c-d**), suggesting that these transcriptome changes are not tied to the inflammatory profile of DS.

A secondary endpoint in the trial is decrease of peripheral inflammatory markers as defined by a composite cytokine score derived from measurements of TNF-α, IL-6, CRP, and IP-10, and which is significantly increased in participants with DS in the HTP study (**Figure 6d**). At baseline, clinical trial participants show cytokine scores within the range observed for DS, but these values decrease at 2, 8 and 16 weeks relative to baseline (**Figure 6d-e, Figure 6 – figure supplement 1e**). The decreases in TNF-α and IL-6 observed upon tofacitinib treatment indicates that elevation of these potent inflammatory cytokines requires sustained JAK/STAT signaling in DS (**Figure 6e**). As for the IFN scores assessment, time course analysis revealed that most participants show decreases in cytokine scores within two weeks of treatment that are sustained over time, again with the exception of AA2 at week 8 and AA4 at week 16. This reveals a correspondence between RNA-based transcriptional IFN scores and circulating levels of these cytokines in plasma, while also illustrating that both metrics may remain sensitive to immune triggers (**Figure 6 – figure supplement 1f-g**).

One tertiary endpoint of the trial investigates the impact of tofacitinib treatment on levels of autoantibodies and markers employed to diagnose AITD [e.g., anti-TPO, anti-TG, anti-thyroid stimulating hormone receptor (TSHR)] and celiac disease [e.g., anti-tissue transglutaminase (tTG), anti-deamidated gliadin peptide (DGP)]. Seven of the 10 participants presented at baseline with anti-TPO levels above the upper limit of normal (ULN, 60U/mL), and all seven experienced a decrease in these auto-antibodies at 8 weeks and 16 weeks relative to baseline (**Figure 6f**). In fact, for one participant (HS1) the levels decreased

below the ULN while on the trial. All seven of these participants had a history of thyroid disease (**Figure**

4c), which was being medically managed and/or clinically monitored with acceptable TSH and T4 values.

Additionally, three of these seven participants also had anti-TG levels above the ULN (4 IU/mL) and all

three showed a decrease from baseline levels while on tofacitinib at both 8 and 16 weeks, with one

participant (AA6) falling below the ULN upon treatment (**Figure 6f**). Three participants also had anti-

TSHr levels above the ULN, but no clear changes were observed upon treatment (**Supplementary file 5**).

None of the 10 participants displayed anti-tTG or anti-DGP levels detected above ULN at screening.

Altogether, these results indicate that tofacitinib treatment decreases IFN scores, levels of key

pathogenic cytokines, and key autoantibodies involved in AITD. Importantly, tofacitinib treatment lowers

IFN scores and cytokine levels to within the range observed in the general population, not below,

indicating that this immunomodulatory strategy can provide therapeutic benefit in DS without overt

immune suppression.

Figure 6. Tofacitinib reduces IFN scores, hypercytokinemia, and pathogenic autoantibodies in Down syndrome. a, Comparison of interferon (IFN) transcriptional scores derived from whole blood transcriptome data for research participants in the Human Trisome Project (HTP) cohort study by karyotype status (D21, grey; T21, green) and the clinical trial cohort at baseline (B), and weeks 2, 8 and

16 of tofacitinib treatment. Data are represented as modified sina plots with boxes indicating quartiles. Sample sizes are indicated below the x-axis. Horizontal bars indicate comparisons between groups with median differences (MD) with p-values from Mann-Whitney U-tests (HTP cohort) or q-values from paired Wilcox tests (clinical trial). q value for the 16-week endpoint is not shown as per interim analysis plan. **b,** Heatmap displaying median z-scores for the indicated groups (as in a) for the 16 interferon-stimulated genes (ISGs) used to calculate IFN scores. **c,** Analysis of fold changes for 136 ISGs not encoded on chr21 that are significantly elevated in Down syndrome (T21 versus D21) at 2, 8 and 16 weeks of tofacitinib treatment relative to baseline. Sample sizes as in a. q-values above each group indicate significance of Mann-Whitney U-tests against log2-transformed fold-change of 0 (no-chance), after Benjamini-Hochberg correction for multiple testing. **d,** Comparison of cytokine score distributions for the HTP cohort by karyotype status (D21, T21) versus the clinical trial cohort at baseline (B) and 2, 8 and 16 weeks of tofacitinib treatment. Data are represented as modified sina plots with boxes indicating quartiles. Sample sizes are indicated below the x-axis. Horizontal bars indicate comparisons between groups with median differences (MD) with p-values from Mann-Whitney U-tests (HTP cohort) and q-values from paired Wilcox tests (clinical trial). q value for the 16-week endpoint is not shown as per interim analysis plan. **e,** Comparison of plasma levels of cytokines in the HTP cohort by karyotype status (D21, T21) and the clinical trial cohort at baseline (B) versus 2, 8 and 16 weeks of tofacitinib treatment. Data are represented as modified sina plots with boxes indicating quartiles. Sample sizes are indicated below x-axis. Horizontal bars indicate comparisons between groups with median differences (MD) with p-values from Mann-Whitney U-tests (HTP cohort) and q values from paired Wilcox tests (clinical trial). q value for the 16-week endpoint is not shown as per interim analysis plan. **f,** Plots showing levels of autoantibodies against thyroid peroxidase (TPO) and thyroglobulin (TG) at baseline versus 8 and 16 weeks of tofacitinib treatment. Sample sizes are indicated in each plot.

Figure 6 – figure supplement 1. JAK inhibition reduces multiple markers of inflammation and

autoimmunity in Down syndrome. a, Plot showing trajectory of IFN scores derived from whole blood

transcriptome for 10 clinical trial participants at baseline (B), versus 2, 8 and 16 weeks of tofacitinib

treatment. **b**, Comparison of ISG expression in the whole blood transcriptome data from research

participants in the Human Trisome Project (HTP) cohort study by karyotype status (D21, grey; T21, green)

and the clinical trial cohort at baseline (B), and weeks 2, 8 and 16 of tofacitinib treatment. Data are

represented as modified sina plots with boxes indicating quartiles. Sample sizes are indicated below x-

axis. Horizontal bars indicate comparisons between groups with median differences (MD) with p-values

from Mann-Whitney U-tests (HTP cohort) and q-values from paired Wilcox tests (clinical trial). **c,**

Heatmap displaying the results of Gene Set Enrichment Analysis (GSEA) of global transcriptome changes

in the whole blood RNA of research participants in the HTP cohort (T21, n=304; D21, n=96) versus the

clinical trial cohort at 2 (n=10), 8 (n=9), and 16 weeks (n=10) of tofacitinib treatment relative to baseline

(n=10). Asterisks indicate significance after correction by Benjamini-Hochberg method for multiple

testing (q<0.1, 10% FDR). NES: normalized enrichment score. **d,** Analysis of fold changes for 109 genes

involved in oxidative phosphorylation and 120 genes involved in heme metabolism significantly elevated

in Down syndrome (T21 versus D21 in the HTP cohort) versus the clinical trial cohort at 2, 8 and 16

weeks of tofacitinib treatment relative to baseline. Sample numbers as in c. **e,** Comparison of CRP levels

in the HTP cohort by karyotype status (D21, grey; T21, green) versus the clinical trial cohort at baseline

(B) and 2, 8 and 16 weeks of tofacitinib treatment. Data are represented as modified sina plots with boxes

indicating quartiles. Sample sizes are indicated below x-axis. Horizontal bars indicate comparisons

between groups with median differences (MD) with p-values from Mann-Whitney U-tests (HTP cohort)

and q-values from paired Wilcox tests (clinical trial). **f,** Plot showing trajectory of cytokine scores for 10

clinical trial participants at baseline (B), versus 2, 8 and 16 weeks of tofacitinib treatment. **g,** Plots showing

Spearman correlations between fold changes in IFN scores versus cytokine scores at 8 and 16 weeks of

tofacitinib treatment versus baseline. Sample size is n=10.

Discussion.

An increasing body of evidence indicates that immune dysregulation contributes to the pathophysiology of DS and that immunomodulatory therapies could provide multidimensional benefits in this population. In mouse models, triplication of four *IFNR* genes contributes to multiple hallmarks of DS[48,56] and JAK inhibition attenuates global dysregulation of gene expression[42] while rescuing key phenotypes, such as lethal immune hypersensitivity[22] and CHDs[24]. The fact that gene signatures of IFN hyperactivity are present in human embryonic tissues with T21[57] and embryonic tissues from mouse models of DS[48,58] indicates that the harmful effects of IFN hyperactivity could start *in utero*, supporting the notion that DS could be understood, in part, as an inborn error of immunity with similarities to monogenic interferonopathies[59].

Results presented here demonstrate that T21 causes widespread multi-organ autoimmunity of pediatric onset, with production of autoantibodies targeting every major organ system. These results justify additional efforts to define the key pathogenic autoantibodies in DS beyond those commonly associated with AITD and celiac disease. Our analysis found significant associations between specific autoantibodies and some conditions more common in DS, but the diagnostic value of these observations will require validation efforts in much larger cohorts, which could lead to a personalized medicine approach for the management of autoimmunity in DS. For example, we found autoantibodies associated with various forms of auditory dysfunction (**Figure 1g**), suggesting the possibility of autoimmune hearing loss in DS[60]. Elevated levels of anti-TPO in individuals with history of use of ear tubes suggests an interplay between otitis media and endocrine dysfunction in DS[61]. For example, it is possible that recurrent ear infections cause a chronic immune stimulus that lead to eventual breach of tolerance in this autoimmunity-prone population, even perhaps through epitope mimicry[62]. Antibodies targeting MUSK, which we found to be elevated in DS and associated with co-occurring neurological phenotypes (**Figure 1g-h**), have been linked to development of myasthenia gravis, a chronic autoimmune neuromuscular disease that causes weakness in the skeletal muscles[63]. Whether MUSK antibodies associate with similar phenotypes in DS will require

further investigation. Elevation of SRP68 autoantibodies in DS (**Figure 1d,f**), which are common in

necrotizing myopathies with cardiovascular involvement[40], suggests a potential autoimmune basis for

musculoskeletal and cardiovascular complications in DS, which also warrants additional research.

We observed constitutive global immune remodeling and hypercytokinemia regardless of reported

diagnoses of autoimmune disease or measurable autoantibody production from an early age, indicative of

an autoimmunity-prone state throughout the lifespan. Therefore, measurements of specific immune cell

types or cytokines in the bloodstream are unlikely to provide diagnostic value for autoimmunity in DS.

However, antigen-specific immune assays, such as T cell or B cell activation assays, may reveal the

specific timing of loss of tolerance and transition to clinical phenotypes. Future studies should also include

analysis of tissue-resident immune cells, which may identify sites of local autoimmune attack in DS.

Among the many strategies that could be used to attenuate IFN hyperactivity, JAK inhibitors are the

most well-studied and have the most approved indications[64]. Of the more than ten globally-approved JAK

inhibitors[64], we chose to employ in our clinical trial the JAK1/3 inhibitor tofacitinib, which is used to treat

diverse autoimmune/inflammatory conditions and which was approved in 2020 for treatment of

polyarticular course juvenile idiopathic arthritis (pcJIA) in children 2 years and older[64,65]. Notably, all four

IFNRs encoded on chr21 utilize JAK1 for signal transduction in combination with either JAK2 or TYK2,

making JAK1 inhibitors the most logical choice to dampen the effects of *IFNR* gene triplication. As part

of the clinical trial protocol, the approved interim analysis was designed to qualitatively evaluate

feasibility and initial safety data on the first 10 participants completing a 16-week course of tofacitinib

treatment. This analysis established that there were no AEs that required a change or cessation of

tofacitinib dosing and that this medicine is well tolerated in individuals with DS. The clear benefits

observed for diverse autoimmune skin conditions align with an increasing body of evidence supporting

the use of JAK inhibition for immunodermatological conditions, including their recent approval for

alopecia areata and atopic dermatitis in the general population[66,67]. At this sample size, the effects of

tofacitinib on HS are inconclusive. Although some participants and caregivers reported benefits in terms

of fewer flares of lesser severity, the MSS metric did not show a clear trend, which may reveal the need

for more frequent or different types of monitoring for HS, a condition that cycles periodically in severity.

Our results indicate that tofacitinib does not fully suppress the immune response in people with DS,

but rather attenuates IFN scores and cytokine scores to levels observed in the general population, which

is an important consideration given the likely requirement for long-term use of the drug in this population.

Furthermore, the effects of the drug are clearly gene-specific, highlighting the presence of inflammatory

processes that may not be attenuated with this inhibitor, which could be beneficial in terms of preserving

immune activity. Importantly, during treatment, both IFN scores and cytokine scores remain sensitive to

immune stimuli, as evidenced by participants who had received a vaccine or experienced an URI before a

blood draw (**Figure 6a**, **Figure 6 – figure supplement 1f**). Overall, it is encouraging that key

inflammatory markers decreased in a relatively short timeframe, likely offering systemic benefits beyond

skin pathology. Importantly, the fact that levels of IL-6 and TNF-α are reduced upon tofacitinib treatment

supports the use of JAK inhibitors over TNF-blockers or anti-IL-6 agents in this population. Although

TNF-α-blockers are recommended to be used first in the treatment of rheumatoid arthritis in the general

population[68], the value of this recommendation in people with DS remains to be defined. The clear

decrease in anti-TPO and anti-TG levels indicates that autoreactive B cell function requires elevated

JAK/STAT signaling, but whether this effect is cell-autonomous versus a consequence of a reduced

systemic inflammatory milieu will require further investigation. Defining the effect of tofacitinib on other

autoantibodies elevated in DS will also require a larger sample size and may be revealed in the full dataset

after completion of this trial.

Lastly, this ongoing clinical trial includes measurements of various dimensions of neurological

function not reported here. Although the absence of a placebo control arm may impede a clear

interpretation of any effect of JAK inhibition on cognitive function, preliminary results have prompted

the design and launch of a second trial (NCT05662228) aimed at defining the relative safety and efficacy

of tofacitinib, intravenous immunoglobulin (IVIG), and the benzodiazepine lorazepam for Down

syndrome Regression Disorder (DSRD), a condition characterized by sudden loss of neurological

function[69].

Altogether, these findings justify both a deeper investigation of all the deleterious effects of

autoimmunity and hyperinflammation in DS and the expanded testing of immunomodulatory strategies

for diverse aspects of DS pathophysiology, even perhaps from an early age.

Materials and Methods.

Human Trisome Project (HTP) study.

All aspects of this study were conducted in accordance with the Declaration of Helsinki under protocols

approved by the Colorado Multiple Institutional Review Board. Results and analyses presented herein are

part of a nested study within the Crnic Institute's Human Trisome Project (HTP, NCT02864108, see also

www.trisome.org) cohort study. All study participants, or their guardian/legally authorized representative,

provided written informed consent. The HTP study has generated multiple multi-omics datasets on

hundreds of research participants, some of which have been analyzed in previous studies, including whole

blood transcriptome data[42], white blood cell transcriptome data[19], plasma proteomics[42], plasma

metabolomics[19,42], and immune mapping via flow cytometry[16] and mass cytometry[42,48]. This paper reports

new analyses of select previous datasets (transcriptome, mass cytometry, MSD immune markers) within

the larger multi-omics dataset of the HTP study, as well as analyses of new datasets (e.g., anti-TPO, ANA,

autoantibodies), as described in detail below.

Annotation of co-occurring conditions.

Within the HTP, a clinical history for each participant is curated from both medical records and

participant/family reports. Both surveys are set up as REDCap[70] instruments that collect information as a

review of systems (e.g., cardiovascular, immunity, endocrine). Expert data curators complete the medical

record review and evaluate answers provided by self-advocates and caregivers. In cases of discordant

answers across the two instruments, medical records take precedence. De-identified demographic and

clinical metadata obtained is then linked to de-identified biospecimens used to generate the various -omics

(e.g., RNA sequencing) and targeted assay datasets (e.g., anti-TPO assays). For annotation of AITD,

several possible entries were considered as shown in **Figure 1 – figure supplement 1a**, including history

of hypothyroidism, hyperthyroidism, Hashimoto's disease, Grave's disease, anti-TPO or -TG antibodies,

and subclinical hypothyroidism. For annotation of immune skin conditions, atopic dermatitis and eczema

were combined and counted in a single group, as were hidradenitis suppurativa (HS), folliculitis, and

'boils'.

Blood sample collection and processing.

The biological datasets analyzed herein were derived from peripheral blood samples collected using

PAXgene RNA Tubes (Qiagen) and BD Vacutainer K2 EDTA tubes (BD). Whole blood from PAXgene

collection tubes was processed for RNA sequencing as described below. Two 0.5 mL aliquots of whole

blood were withdrawn from each EDTA tube and processed for mass cytometry as described below. The

remaining EDTA blood samples were centrifuged at 700 x g for 15 min to separate plasma, buffy coat

containing white blood cells (WBC), and red blood cells (RBCs). Samples were then aliquoted, flash

frozen and stored at -80°C until subsequent processing and analysis. Centrifugation and storage of samples

took place within 2 hours of collection.

Measurements of autoantibodies.

Anti-TPO status was determined from plasma samples using an electrochemiluminescence-based assay

[71], and carried out by the Autoantibody/HLA Core Facility of the Barbara Davis Center for Childhood

Diabetes at the University of Colorado Anschutz Medical Campus. Sample values were calculated as

(sample signal – negative control signal) / (positive control signal – negative control signal), with the

threshold (upper limit of normal) for TPO positivity based on the 95th percentile of healthy control

samples.

Anti-nuclear antigen (ANA) status was determined from plasma samples using a qualitative ELISA kit

(MyBioSource, cat. no. 702970) according to manufacturer instructions, with a sample OD_{450nm} / negative

control OD_{450nm} ratio $\geq$ 2.1 evaluated as positive and a ratio < 2.1 evaluated as negative.

Autoantigen profiling of EDTA plasma samples (50 µL each; T21, n = 120; D21, n = 60) was performed

by the Affinity Proteomics unit at SciLifeLab (KTH Royal Institute of Technology, Stockholm, Sweden)

using peptide arrays. Antigens were selected to cover potential associations to autoimmune diseases and

consisted of 380 peptide fragments covering ~270 unique proteins (1-5 fragments per protein). Fragments

were ~20-163 amino acids long (median 82). All antigens were expressed in E. coli with a hexahistidyl

and albumin binding protein tag (His6ABP). Using, COOH-NH2 chemistry, the analyzed antigens, in

addition to controls, were immobilized on color-coded magnetic beads (MagPlex, Luminex). Controls

consisted of His6ABP, buffer, rabbit anti-human IgG (loading control, Jackson ImmunoResearch), and

Epstein-Barr nuclear antigen 1 (EBNA1, Abcam). Research samples and technical controls (commercial

plasma; Seralab) were diluted (1:250) in assay buffer, which consisted of 3% BSA, 5% milk, 0.05%

Tween-20, and 160 µg/ml His6ABP tag in PBS. Diluted samples and controls were incubated for 1 hour

at room temperature then subsequently incubated with the antigen bead array for 2 hour. The reactions

were then fixed for 10 minutes using 0.02% paraformaldehyde, then incubated for 30 minutes with goat

Fab specific for human IgG Fc-γ tagged with the fluorescent marker R-phycoerythrin (Invitrogen). Median

fluorescence intensity (MFI) and number of beads for each reaction was analyzed using a FlexMap 3D

instrument (Luminex Corp.). Quality control was performed using MFI and bead count to exclude antigens

and samples not passing technical criteria including minimal bead counts and antigen coupling efficiency.

To adjust for sample specific backgrounds, MFI values were transformed per reaction median absolute

deviations (MADs) using the following calculation:

$$MADs_{sample} = (MFI - median_{sample(MFI)}) / MAD_{sample(MFI)}$$

Subsequent data analysis and handling was performed using R. For each antigen, positivity was defined

as >90th percentile MAD value for D21 samples only. Overrepresentation of positivity for each antigen

in the T21 versus D21 group was determined using Fisher's exact test, excluding antigens detected in <18

samples (<10% of total experiment). Correction for multiple testing was performed using the Benjamini-

Hochberg approach and significance defined as q<0.1 (10% FDR). Similarly, within the T21 group,

Fisher's exact test was used to test for overrepresentation of antigen positivity in cases versus controls for

co-occurring conditions, with only those with at least 5 cases considered in the analysis.

Immune profiling via mass cytometry.

Generation of the mass cytometry dataset was described previously[42], but a full description is included

here for reference. Two 0.5 mL aliquots of EDTA whole blood samples underwent RBC lysis and white

blood cell fixation using TFP FixPerm Buffer (Transcription Factor Phospho Buffer Set, BD Biosciences).

WBCs were then washed in 1x in PBS (Rockland), resuspended in Cell Staining Buffer (Fluidigm) and

stored at −80°C. For antibody staining, samples were thawed at room temperature, washed in Cell Staining

Buffer, barcoded using a Cell-ID 20-Plex Pd Barcoding Kit (Fluidigm), and combined per batch. Each

batch was able to accommodate 19 samples with a common reference sample. Antibodies were either

purchased pre-conjugated to metal isotopes or conjugation was performed in-house using a Maxpar

Antibody Labeling Kit (Fluidigm). See **Supplementary file 7** for antibodies. Working dilutions for

antibody staining were titrated and validated using the common reference sample and comparison to

relative frequencies obtained by independent flow cytometry analysis. Surface marker staining was carried

out for 30 min at 4°C in Cell Staining Buffer with added Fc Receptor Binding Inhibitor

(eBioscience/ThermoFisher Scientific). Staining was followed by a wash in Cell Staining Buffer. Next,

cells were permeabilized in Buffer III (Transcription Factor Phospho Buffer Set, BD Pharmingen) for 20

min at 4°C followed by washing with perm/wash buffer (Transcription Factor Phospho Buffer Set, BD

Pharmingen). Intracellular transcription factor and phospho-epitope staining was carried out for 1 hour at

4°C in perm/wash buffer (Transcription Factor Phospho Buffer Set, BD Pharmingen), followed by a wash

with Cell Staining Buffer. Cell-ID Intercalator-Ir (Fluidigm) was used to label barcoded and stained cells.

Labeled cells were analyzed on a Helios instrument (Fluidigm). Mass cytometry data were exported as

v3.0 FCS files for pre-processing and analysis.

Analysis of mass cytometry data.

Pre-processing. Bead-based normalization via polystyrene beads embedded with lanthanides, both within

and between batches, followed by bead removal was carried out as previously described using the Matlab-

based Normalizer tool[72]. Batched FCS files were demultiplexed using the Matlab-based Single Cell

Debarcoder tool[73]. Reference-based normalization of individual samples across batches against the

common reference sample was then carried out using the R script *BatchAdjust()*. For the analyses

described in this manuscript, CellEngine (CellCarta) was used to gate and export per-sample FCS files at

four levels: Firstly, CD3+CD19+ doublets were excluded and remaining cells exported as 'Live' cells;

Live cells were then gated for hematopoietic lineage (CD45-positive) non-granulocytic (CD66-low) cells

and exported as CD45+CD66low. Lastly, CD45+CD66low cells were gated on CD3-positivity and CD19-

positivity and exported as T- and B-cells, respectively. Per-sample FCS files were then subsampled to a

maximum of 50,000 events per file for subsequent analysis.

Unsupervised clustering. For each of the four levels (live, non-granulocytes, T cells, and B cells), all 388

per-sample FCS files were imported into R as a flowSet object using the *read.flowSet()* function from the

flowCore R package[74]. Next a SingleCellExperiment object was constructed from the flowSet object using

the *prepData()* function from the CATALYST package[75]. Arcsinh transformation was applied to marker

expression data with cofactor values ranging from ~0.2 to ~15 to give optimal separation of positive and

negative populations for each marker, using the *estParamFlowVS()* function from the flowVS R package[76]

and based on visual inspection of marker histograms (see **Supplementary file 7)**. Quality control and

diagnostic plots were examined with the help of functions from CATALYST and the

tidySingleCellExperiment R package. Unsupervised clustering using the FlowSOM algorithm[43] was

carried out using the *cluster()* function from CATALYST, with grid size set to 10 x 10 to give 100 initial

clusters and a maxK value of 40 was explored for subsequent meta-clustering using the

ConsensusClusterPlus algorithm. Examination of delta area and minimal spanning tree plots indicated that

30-40 meta clusters gave a reasonable compromise between gains in cluster stability and number of

clusters for each level. Each clustering level was re-run with multiple random seed values to ensure

consistent results.

Visualization using t-distributed stochastic neighbor imbedding (tSNE). Dimensionality reduction to two

dimensions was carried out using the *runDR()* function from the CATALYST package, with 500 cells per

sample, and using several random seed values to ensure consistent results. Multiple values of the

perplexity parameter were tested, with a setting of 440, using the formula Perplexity=N^(1/2) as suggested

at https://towardsdatascience.com/how-to-tune-hyperparameters-of-tsne-7c0596a18868, providing a

visualization with good agreement with the clusters defined by FlowSOM.

Cell type classification. To aid in assignment of clusters to specific lineages and cell types, the MEM

package (marker enrichment modeling) was used to call positive and negative markers for each cell cluster

based on marker expression distributions across clusters. Manual review and comparison to marker

expression histograms, as well as minimal spanning tree plots and tSNE plots colored by marker

expression, allowed for high-confidence assignment of most clusters to specific cell types. Clusters that

were insufficiently distinguishable were merged into their nearest cluster based on the minimal spanning

tree. Relative frequencies for each cell type / cluster were calculated for each sample as a percentage of

total live cells and as a percentage of cells used for each level of clustering: total CD45+CD66low cells,

total T cells, or total B cells.

Beta regression analysis. To identify cell clusters for which relative frequencies are associated with either

trisomy 21 status or with various clinical subgroups (e.g. ANA+) among individuals with trisomy 21, beta

regression analysis was carried out using the betareg R package, with each model using cell type cluster

proportions (relative frequency) as the outcome/dependent variable and either T21 status or clinical

subgroups as independent/predictor variables, along with adjustment for age and sex, and a logit link

function. Extreme outliers were classified per-karyotype and per-cluster as measurements more than three

times the interquartile range below or above the first and third quartiles, respectively (below Q1 - 3*IQR

or above Q3 + 3*IQR) and excluded from beta regression analysis. Correction for multiple comparisons

was performed using the Benjamini-Hochberg (FDR) approach. Effect sizes (as fold-change in T21 vs.

euploid controls or among T21 subgroups) for each cell type cluster were obtained by exponentiation of

beta regression model coefficients. Fold-changes were visualized by overlaying on tSNE plots using

ggplot2. For visualization of individual clusters, data points were adjusted for age and sex, using the

adjust() function from the datawizard R package, and visualized as sina plots (separated by T21 status or

clinical subgroup).

Measurement of immune markers and calculation of cytokine scores.

Briefly, from each EDTA plasma sample, two replicates of 12-25 µL were analyzed using the Meso Scale

Discovery (MSD) multiplex immunoassay platform V-PLEX Human Biomarker 54-Plex Kit (HTP

cohort) or U-PLEX Human Biomarker Group 1 71-Plex and V-PLEX Human Vascular Injury Panel 2

Kits (clinical trial cohort) on a MESO QuickPlex SQ 120 instrument. Assays were carried out as per

manufacturer instructions. Concentration values were calculated against a standard curve with provided

calibrators. MSD data are reported as concentration values in picograms per milliliter of plasma.

Analysis of immune marker data. Plasma concentration values (pg/mL) for each of the cytokines and

related immune factors measured across multiple MSD assay plates was imported to R, combined, and

analytes with >10% of values outside of detection or fit curve range flagged. For each analyte, missing

values were replaced with either the minimum (if below fit curve range) or maximum (if above fit curve

range) calculated concentration per plate/batch and means of duplicate wells used for subsequent analysis.

For the HTP study analysis, extreme outliers were classified per-karyotype and per-analyte as

measurements more than three times the interquartile range below or above the first and third quartiles,

respectively, and excluded from further analysis. Differential abundance analysis for inflammatory

markers measured by MSD was performed using mixed effects linear regression as implemented in the

lmer() function from the lmerTest R package (v3.1-2) with log2-transformed concentration as the

outcome/dependent variable, T21 status or clinical subgroup (e.g., ANA+) as the predictor/independent

variable, age and sex as fixed covariates, and sample source as a random effect. Multiple hypothesis

correction was performed with the Benjamini-Hochberg method using a false discovery rate (FDR)

threshold of 10% (q < 0.1). Prior to visualization or correlation analysis, MSD data were adjusted for age,

sex, and sample source using the *removeBatchEffect()* function from the limma package (v3.44.3).

Calculation of cytokine scores. For comparison of clinical trial samples across time points, cytokine scores

were calculated as the sum of the Z-scores for TNF-α, IL-6, CRP and IP-10. For comparison of clinical

trial samples to the HTP cohort, Z-scores were first calculated from age-, sex, and batch-adjusted values

for each sample, based on the mean and standard deviation of the HTP euploid control samples.

Whole blood transcriptome analysis and calculation of IFN scores.

Strand-specific sequencing libraries were prepared from globin-depleted, polyA-enriched whole blood

RNA and sequenced on the Illumina NovaSeq platform (2x 150 bases). Data quality was assessed using

FASTQC (v0.11.5) and FastQ Screen (v0.11.0). Trimming and filtering of low-quality reads was

performed using bbduk from BBTools (v37.99) and fastq-mcf from ea-utils (v1.05). Alignment to the

human reference genome (GRCh38) was carried out using HISAT2 (v2.1.0) in paired, spliced-alignment

mode against a GRCh38 index and Gencode v33 basic annotation GTF, and alignments were sorted and

filtered for mapping quality (MAPQ > 10) using Samtools (v1.5). Gene-level count data were quantified

using HTSeq-count (v0.6.1) with the following options (--stranded=reverse –minaqual=10 –type=exon --

mode=intersection-nonempty) using a Gencode v33 GTF annotation file. Differential gene expression in

T21 versus D21 was evaluated using DESeq2 (version 1.28.1)[77], with q < 0.1 (10% FDR) as the threshold

for differential expression.

DS IFN scores. RNA-seq-based 'Down syndrome interferon scores' (DS IFN scores) were calculated as

follows: for comparison of clinical trial samples across time points, DS IFN scores were calculated as the

sum of Z-scores across 16 interferon-stimulated genes (ISGs) genes with significant mean fold-change of

at least 1.5 in the HTP T21 group vs. the euploid control group, excluding *IFNAR2*, *MX1*, and *MX2* which

are encoded on chromosome 21. For comparison of clinical trial samples to the HTP cohort, gene-wise Z-

scores were first calculated from age-, sex, and sequencing batch-adjusted FPKM values for each sample,

based on the mean and standard deviation of the HTP euploid control samples.

Gene set enrichment analysis (GSEA). GSEA[78] was carried out in R using the fgsea package (v1.14.0),

using Hallmark gene sets, $\log_2$-transformed fold-change values as the ranking metric.

Clinical trial design and oversight.

All aspects of this study were conducted in accordance with the Declaration of Helsinki. All study

activities were approved by the Colorado Multiple Institutional Review Board (COMIRB, protocol # 19-

1362, NCT04246372) with an independent Data and Safety Monitoring Board (DSMB) appointed by the

National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS). Written consent was

obtained from all participants, or their legally authorized representative if the participant was unable to

provide consent, in which case participant assent was obtained. The Clinical Trial Protocol is provided in

the **Supplementary file 8**. We report here interim results of a single-site, open-label phase 2 clinical trial

enrolling individuals with DS between the ages of 12 and 50 years old with moderate-to-severe alopecia

areata, hidradenitis suppurativa, psoriasis, atopic dermatitis, or vitiligo. Qualifying disease scores are

shown in **Supplementary file 3**. After screening, qualifying participants are prescribed 5 mg tofacitinib

twice daily for 16 weeks, with an optional extension arm to week 40. During the main 16-week trial,

participants attend five safety monitoring visits after enrollment at the Baseline visit.

Trial Population.

The recruitment goal for this trial is 40 participants completing 16 weeks of tofacitinib treatment, with a

qualitative interim analysis triggered when 10 participants completed the main 16-week trial. Of the 10

participants included in the interim analysis, 4 are female, 100% identify as White/Caucasian, 3 identify

as Hispanic or Latino, and mean age at enrollment was 23.1 years old (range 15-38.1 years old)

(**Supplementary file 3**). Baseline qualifying conditions of the 10 participants were alopecia areata: n=6

(46.1%), hidradenitis suppurativa: n=3 (30.8%), or psoriasis: n=1 (7.7%). Two participants also had atopic

dermatitis, and two others had vitiligo, albeit below the severity required to be the qualifying condition.

Outcome Measures.

Primary endpoints. The two primary outcome measures for this trial are safety and reduction in IFN

transcriptional scores derived from peripheral whole blood. Based on the safety profile for tofacitinib in

the general population[68], the safety primary endpoint was defined as no more than two serious adverse

events (SAEs) definitely attributable to tofacitinib over the course of 16 weeks for 40 participants. Adverse

events were classified based using Common Terminology Criteria for Adverse Events 5.0 (CTCAE 5.0).

IFN scores are commonly used to monitor disease severity and response to treatment in IFN-driven

pathologies[79,80] and their calculation form RNAseq data is described above.

Secondary endpoints. The secondary outcome measures for this trial include improvements in skin health

as defined by a global assessment, the Investigator Global Assessment (IGA), as well as the disease-

specific assessments. Overall skin pathology, accounting for all present skin conditions regardless of

severity, was assessed using a modified IGA which scores on a five-point scale for each skin condition

(six points for HS) with a range of 0-21. Another secondary endpoint assessing global skin health is a

change in the Dermatological Quality of Life Index (DLQI), used to assess participant-reported impact of

skin conditions on self-image, relationships, and daily activities. Possible total scores range from 0-30,

with higher scores indicating a more impaired quality of life. Condition-specific assessments used are

Severity of Alopecia Tool (SALT) for AA affecting at least 25% of the scalp (qualifying score is ≥ 25);

Hidradenitis Suppurativa-Physicians Global Assessment (HS-PGA) to define eligibility (qualifying score

≥ 3) and Modified Sartorius Scale (MSS) to monitor changes throughout the study for HS; Psoriasis Area

and Severity Index (PASI, qualifying score is ≥ 10) for psoriasis; Vitiligo Extent Tensity Index (VETI,

qualifying score is ≥ 2), for moderate-to-severe vitiligo; and Eczema Area and Severity Index (EASI,

qualifying EASI score ≥ 16) for moderate-to-severe atopic dermatitis. The last secondary endpoint is

reduction in a cytokine score coalescing information on four inflammatory markers elevated in DS: Tumor

Necrosis Factor Alpha (TNF-α), interleukin 6 (IL-6), C-reactive protein (CRP), and IFN-inducible protein

10 (IP10, CXCL10)[18]. Measurement of these proteins and calculation of the cytokine score is described

above.

Tertiary endpoints. This clinical trial includes multiple exploratory tertiary endpoints (see full protocol in

Supplementary file 8), including reduction in autoantibodies related to AITD (anti-TPO, anti-TG, and

anti-TSHr) and celiac disease (anti-tTG, anti-DGP) . In the clinical trial, these autoantibodies were

assessed using established clinical assays.

Statistical Analysis.

The Statistical Analysis Plan (SAP) approved by the appointed DSMB is included with the Clinical Trial

Protocol in the **Supplementary file 8**. This report includes analysis of the time points used to assess

endpoints (baseline and 16 weeks), as well research-only time points at 2 and 8 weeks of treatment. Given

the qualitative nature of this interim analysis, statistical analysis is not completed for changes observed

between baseline and the 16-week endpoint. Data may be displayed as $\log_2$ transformed for clarity in

viewing the graphs.

Data Availability Statement:

A   collection   with   datasets   used   in   this   study   is   available   in   Synapse

(https://doi.org/10.7303/syn53185135), with the individual datasets also accessible as detailed below.

Demographic and health history data for research participants in the HTP study are available on both the

Synapse data sharing platform (https://doi.org/10.7303/syn31488784) and through the INCLUDE Data

Hub (https://portal.includedcc.org/). Mass cytometry data for 380+ HTP research participants are

available both in Synapse (https://doi.org/10.7303/syn53185253) and FlowRepository, Study ID FR-

FCM-Z5GE,

https://flowrepository.org/id/RvFrQaBqhe8TGyko1OMdQKtR7HN8nulAnHW0PJkm1CEyyo8fnJg2rHr

WvNrhE8xu. Targeted plasma proteomics for inflammatory markers using Meso Scale Discovery (MSD)

assays   for   470+   HTP   research   participants   can   be   accessed   through   Synapse

(https://doi.org/10.7303/syn31475487) and the INCLUDE Data Hub. Whole blood transcriptome data for

400 HTP research participants can be accessed through Synapse (https://doi.org/10.7303/syn31488780),

the INCLUDE Data Hub, and NCBI Gene Expression Omnibus (GSE190125). Whole blood

transcriptome data for 10 clinical trial participants at baseline and after 2, 8, and 16 weeks of tofacitinib

treatment can be accessed through Synapse (https://doi.org/10.7303/syn53185250), and NCBI Gene

Expression Omnibus (GSE PENDING). Targeted plasma proteomics for inflammatory markers using

Meso Scale Discovery (MSD) assays for 10 clinical trial participants can be accessed through Synapse

(https://doi.org/10.7303/syn53185252).

Code Availability Statement: No custom code or algorithms were developed during the course of this

study. R analysis scripts will be made available upon request.

References.

Lejeune, J., Turpin, R. & Gautier, M. [Mongolism; a chromosomal disease (trisomy)]. *Bull Acad Natl Med* **143**, 256-265 (1959).

Antonarakis, S. E. *et al.* Down syndrome. *Nature reviews. Disease primers* **6**, 9 (2020). https://doi.org:10.1038/s41572-019-0143-7

Chicoine, B. *et al.* Prevalence of Common Disease Conditions in a Large Cohort of Individuals With Down Syndrome in the United States. *J Patient Cent Res Rev* **8**, 86-97 (2021). https://doi.org:10.17294/2330-0698.1824

Amr, N. H. Thyroid Disorders in Subjects with Down Syndrome: An Update. *Acta Biomed* **89**, 132-139 (2018). https://doi.org:10.23750/abm.v89i1.7120

Pierce, M. J., LaFranchi, S. H. & Pinter, J. D. Characterization of Thyroid Abnormalities in a Large Cohort of Children with Down Syndrome. *Hormone research in paediatrics* **87**, 170-178 (2017). https://doi.org:10.1159/000457952

Iughetti, L. *et al.* Ten-year longitudinal study of thyroid function in children with Down's syndrome. *Hormone research in paediatrics* **82**, 113-121 (2014). https://doi.org:10.1159/000362450

Book, L. *et al.* Prevalence and clinical characteristics of celiac disease in Downs syndrome in a US study. *American journal of medical genetics* **98**, 70-74 (2001).

Zachor, D. A., Mroczek-Musulman, E. & Brown, P. Prevalence of celiac disease in Down syndrome in the United States. *Journal of pediatric gastroenterology and nutrition* **31**, 275-279 (2000).

Sureshbabu, R. *et al.* Phenotypic and dermatological manifestations in Down Syndrome. *Dermatology online journal* **17**, 3 (2011).

10 Madan, V., Williams, J. & Lear, J. T. Dermatological manifestations of Down's syndrome.

Clinical and experimental dermatology **31**, 623-629 (2006). https://doi.org:10.1111/j.1365-

2230.2006.02164.x

11 Lam, M., Lai, C., Almuhanna, N. & Alhusayen, R. Hidradenitis suppurativa and Down

syndrome: A systematic review and meta-analysis. *Pediatric dermatology* **37**, 1044-1050 (2020).

https://doi.org:10.1111/pde.14326

12 Araya, P. *et al.* IGF1 deficiency integrates stunted growth and neurodegeneration in Down

syndrome. *Cell reports* **41**, 111883 (2022). https://doi.org:10.1016/j.celrep.2022.111883

13 Flores-Aguilar, L. *et al.* Evolution of neuroinflammation across the lifespan of individuals with

Down syndrome. *Brain* **143**, 3653-3671 (2020). https://doi.org:10.1093/brain/awaa326

14 Wilcock, D. M. & Griffin, W. S. T. Down's syndrome, neuroinflammation, and Alzheimer

neuropathogenesis. *Journal of neuroinflammation* **10**, 864 (2013). https://doi.org:10.1186/1742-

2094-10-84

15 Waugh, K. A. *et al.* Mass Cytometry Reveals Global Immune Remodeling with Multi-lineage

Hypersensitivity to Type I Interferon in Down Syndrome. *Cell reports* **29**, 1893-1908 e1894

(2019). https://doi.org:10.1016/j.celrep.2019.10.038

16 Araya, P. *et al.* Trisomy 21 dysregulates T cell lineages toward an autoimmunity-prone state

associated with interferon hyperactivity. *Proc Natl Acad Sci U S A* **116**, 24231-24241 (2019).

https://doi.org:10.1073/pnas.1908129116

17 Sullivan, K. D. *et al.* Trisomy 21 consistently activates the interferon response. *Elife* **5** (2016).

https://doi.org:10.7554/eLife.16220

18 Sullivan, K. D. *et al.* Trisomy 21 causes changes in the circulating proteome indicative of

chronic autoinflammation. *Scientific reports* **7**, 14818 (2017). https://doi.org:10.1038/s41598-

017-13858-3

19   Powers, R. K. *et al.* Trisomy 21 activates the kynurenine pathway via increased dosage of

interferon receptors. *Nat Commun* **10**, 4766 (2019). https://doi.org:10.1038/s41467-019-12739-9

20   Secombes, C. J. & Zou, J. Evolution of Interferons and Interferon Receptors. *Front Immunol* **8**,

209 (2017). https://doi.org:10.3389/fimmu.2017.00209

21   Waugh, K. A., Minter, R., Baxter, J., Chi, C., Tuttle, K.D., Eduthan, N.P., Galbraith, M.D.,

Kinning, K.T., Andrysik, Z., Araya, P., Dougherty, H., Dunn, L.N., Ludwig, M., Schade, K.A.,

Tracy, D., Smith, K.P., Granrath, R.E., Busquet, N., Khanal, S., Anderson, R.D., Cox, L.L.,

BEnriquez Estrada, B., Rachubinski, A.L., Lyford, H.R., Britton, E.C., Orlicky, D.J., Matsuda,

J.L., Song, K., Cox, T.C., Sullivan, K.D., and Espinosa, J.M. Interferon receptor gene dosage

determines diverse hallmarks of Down syndrome. *BioRxiv* (2022).

https://doi.org:https://doi.org/10.1101/2022.02.03.478982

22   Tuttle, K. D. *et al.* JAK1 Inhibition Blocks Lethal Immune Hypersensitivity in a Mouse Model

of Down Syndrome. *Cell reports* **33**, 108407 (2020).

https://doi.org:10.1016/j.celrep.2020.108407

23   Galbraith, M. D., Rachubinski, A.L., Smith, K.P., Araya, P., Waugh, K.A., Enriquez-Estrada, B.,

Worek, K., Granrath, R.E., Kinning, K.T., Eduthan, N.P., Ludwig, M.P. Hsieh, E.W.Y., Sullivan,

K.D., Espinosa, J.M. Multidimensional definition of the interferonopathy of Down syndrome and

its response to JAK inhibition. *Science Advances* **9** (2023).

24   Chi, C., Knight, W.E., Riching, A.S., Zhang, Z., Tatavosian, R., Zhuang, Y., Moldovan, R.,

Rachubinski, A.L., Gao, D., Xu H., Espinosa, J.M., Song, K. . Interferon hyperactivity impairs

cardiogenesis in Down syndrome via downregulation of canonical Wnt signaling. *iScience* **26,**

107012 (2023). https://doi.org:https://doi.org/10.1016/j.isci. 2023.107012

25   Lambert, K. *et al.* Deep immune phenotyping reveals similarities between aging, Down

syndrome, and autoimmunity. *Science translational medicine* **14**, eabi4888 (2022).

https://doi.org:10.1126/scitranslmed.abi4888

26   Khor, B. & Buckner, J. H. Down syndrome: insights into autoimmune mechanisms. *Nature*

reviews. Rheumatology, 1-2 (2023). https://doi.org:10.1038/s41584-023-00970-0

27   Gensous, N., Bacalini, M. G., Franceschi, C. & Garagnani, P. Down syndrome, accelerated aging

and immunosenescence. *Seminars in immunopathology* (2020). https://doi.org:10.1007/s00281-

020-00804-1

28   Malle, L. *et al.* Autoimmunity in Down's syndrome via cytokines, CD4 T cells and CD11c(+) B

cells. *Nature* **615**, 305-314 (2023). https://doi.org:10.1038/s41586-023-05736-y

29   Aversa, T. *et al.* Epidemiological and clinical aspects of autoimmune thyroid diseases in children

with Down's syndrome. *Ital J Pediatr* **44**, 39 (2018). https://doi.org:10.1186/s13052-018-0478-9

30   Liu, E. *et al.* Routine Screening for Celiac Disease in Children With Down Syndrome Improves

Case Finding. *Journal of pediatric gastroenterology and nutrition* **71**, 252-256 (2020).

https://doi.org:10.1097/mpg.0000000000002742

31   Johnson, M. B. *et al.* Trisomy 21 Is a Cause of Permanent Neonatal Diabetes That Is

Autoimmune but Not HLA Associated. *Diabetes* **68**, 1528-1535 (2019).

https://doi.org:10.2337/db19-0045

32   Aitken, R. J. *et al.* Early-onset, coexisting autoimmunity and decreased HLA-mediated

susceptibility are the characteristics of diabetes in Down syndrome. *Diabetes Care* **36**, 1181-

1185 (2013). https://doi.org:10.2337/dc12-1712

33   Singh, P. *et al.* Global Prevalence of Celiac Disease: Systematic Review and Meta-analysis. *Clin*

Gastroenterol Hepatol **16**, 823-836 e822 (2018). https://doi.org:10.1016/j.cgh.2017.06.037

34   Markle, J. G. *et al.* Sex differences in the gut microbiome drive hormone-dependent regulation of

autoimmunity. *Science* **339**, 1084-1088 (2013). https://doi.org:10.1126/science.1233521

35   Molano-Gonzalez, N. *et al.* Cluster analysis of autoimmune rheumatic diseases based on

autoantibodies. New insights for polyautoimmunity. *Journal of autoimmunity* **98**, 24-32 (2019).

https://doi.org:10.1016/j.jaut.2018.11.002

36 Elling, C. L. *et al.* Otitis Media in Children with Down Syndrome Is Associated with Shifts in

the Nasopharyngeal and Middle Ear Microbiotas. *Genet Test Mol Biomarkers* **27**, 221-228

(2023). https://doi.org:10.1089/gtmb.2023.0132

37 Ghazanfari, N., Morsch, M., Reddel, S. W., Liang, S. X. & Phillips, W. D. Muscle-specific

kinase (MuSK) autoantibodies suppress the MuSK pathway and ACh receptor retention at the

mouse neuromuscular junction. *J Physiol* **592**, 2881-2897 (2014).

https://doi.org:10.1113/jphysiol.2013.270207

38 Poulter, J. A. *et al.* Novel somatic mutations in UBA1 as a cause of VEXAS syndrome. *Blood*

137, 3676-3681 (2021). https://doi.org:10.1182/blood.2020010286

39 Lin, S. J. *et al.* Biallelic variants in WARS1 cause a highly variable neurodevelopmental

syndrome and implicate a critical exon for normal auditory function. *Hum Mutat* **43**, 1472-1489

(2022). https://doi.org:10.1002/humu.24435

40 Allenbach, Y., Benveniste, O., Stenzel, W. & Boyer, O. Immune-mediated necrotizing

myopathy: clinical features and pathogenesis. *Nature reviews. Rheumatology* **16**, 689-701

(2020). https://doi.org:10.1038/s41584-020-00515-9

41 Amagai, M., Nishikawa, T., Nousari, H. C., Anhalt, G. J. & Hashimoto, T. Antibodies against

desmoglein 3 (pemphigus vulgaris antigen) are present in sera from patients with paraneoplastic

pemphigus and cause acantholysis in vivo in neonatal mice. *The Journal of clinical investigation*

102, 775-782 (1998). https://doi.org:10.1172/JCI3647

42 Galbraith, M. D. *et al.* Multidimensional definition of the interferonopathy of Down syndrome

and its response to JAK inhibition. *Science Advances* **9**, eadg6218 (2023).

https://doi.org:doi:10.1126/sciadv.adg6218

43 Van Gassen, S. *et al.* FlowSOM: Using self-organizing maps for visualization and interpretation

of cytometry data. *Cytometry A* **87**, 636-645 (2015). https://doi.org:10.1002/cyto.a.22625

44   Amini, A., Pang, D., Hackstein, C. P. & Klenerman, P. MAIT Cells in Barrier Tissues: Lessons

047   from Immediate Neighbors. *Front Immunol* **11**, 584521 (2020).

048   https://doi.org:10.3389/fimmu.2020.584521

049   45   Zhang, Y. *et al.* Aberrations in circulating inflammatory cytokine levels in patients with Down

050   syndrome: a meta-analysis. *Oncotarget* **8**, 84489-84496 (2017).

051   https://doi.org:10.18632/oncotarget.21060

052   46   Kusters, M. A., Gemen, E. F., Verstegen, R. H., Wever, P. C. & E, D. E. V. Both normal

053   memory counts and decreased naive cells favor intrinsic defect over early senescence of Down

054   syndrome T lymphocytes. *Pediatr Res* **67**, 557-562 (2010).

055   https://doi.org:10.1203/PDR.0b013e3181d4eca3

056   47   Trotta, M. B. *et al.* Inflammatory and Immunological parameters in adults with Down syndrome.

057   *Immun Ageing* **8**, 4 (2011). https://doi.org:10.1186/1742-4933-8-4

058   48   Waugh, K. A. *et al.* Triplication of the interferon receptor locus contributes to hallmarks of

059   Down syndrome in a mouse model. *Nature genetics* (2023). https://doi.org:10.1038/s41588-023-

060   01399-7

061   49   Rachubinski, A. L. *et al.* Janus kinase inhibition in Down syndrome: 2 cases of therapeutic

062   benefit for alopecia areata. *JAAD Case Rep* **5**, 365-367 (2019).

063   https://doi.org:10.1016/j.jdcr.2019.02.007

064   50   Pham, A. T. *et al.* JAK inhibition for treatment of psoriatic arthritis in Down syndrome.

065   *Rheumatology* (2021). https://doi.org:10.1093/rheumatology/keab203

066   51   Guild, A., Fritch, J., Patel, S., Reinhardt, A. & Acquazzino, M. Hemophagocytic

067   lymphohistocytosis in trisomy 21: successful treatment with interferon inhibition. *Pediatr*

068   *Rheumatol Online J* **20**, 104 (2022). https://doi.org:10.1186/s12969-022-00764-w

52   Cohen, S. B. *et al.* Long-term safety of tofacitinib for the treatment of rheumatoid arthritis up to

8.5 years: integrated analysis of data from the global clinical trials. *Ann Rheum Dis* **76**, 1253-

1262 (2017). https://doi.org:10.1136/annrheumdis-2016-210457

53   Honda, K., Takaoka, A. & Taniguchi, T. Type I interferon [corrected] gene induction by the

interferon regulatory factor family of transcription factors. *Immunity* **25**, 349-360 (2006).

https://doi.org:10.1016/j.immuni.2006.08.009

54   Schwartz, D. M. *et al.* JAK inhibition as a therapeutic strategy for immune and inflammatory

diseases. *Nature reviews. Drug discovery* **17**, 78 (2017). https://doi.org:10.1038/nrd.2017.267

55   Schwartz, D. M., Bonelli, M., Gadina, M. & O'Shea, J. J. Type I/II cytokines, JAKs, and new

strategies for treating autoimmune diseases. *Nature reviews. Rheumatology* **12**, 25-36 (2016).

https://doi.org:10.1038/nrrheum.2015.167

56   Maroun, L. E., Heffernan, T. N. & Hallam, D. M. Partial IFN-alpha/beta and IFN-gamma

receptor knockout trisomy 16 mouse fetuses show improved growth and cultured neuron

viability. *Journal of interferon & cytokine research : the official journal of the International*

Society for Interferon and Cytokine Research **20**, 197-203 (2000).

https://doi.org:10.1089/107999000312612

57   Bhattacharya, S., Cherry, C., Deutsch, G., Birth Defects Research Laboratory (BDRL), Glass,

I.A., Mariani, T.J., Al Alam, D., Danopoulos, S. A Trisomy 21 Lung Cell Atlas. *bioRxiv* **doi:**

https://doi.org/10.1101/2023.03.30.534839 (2023). https://doi.org:doi:

https://doi.org/10.1101/2023.03.30.534839

58   Aziz, N. M. *et al.* Lifespan analysis of brain development, gene expression and behavioral

phenotypes in the Ts1Cje, Ts65Dn and Dp(16)1/Yey mouse models of Down syndrome. *Dis*

Model Mech **11** (2018). https://doi.org:10.1242/dmm.031013

59   Rodero, M. P. & Crow, Y. J. Type I interferon-mediated monogenic autoinflammation: The type

I interferonopathies, a conceptual overview. *The Journal of experimental medicine* **213**, 2527-

2538 (2016). https://doi.org:10.1084/jem.20161596

60   Breslin, N. K., Varadarajan, V. V., Sobel, E. S. & Haberman, R. S. Autoimmune inner ear

disease: A systematic review of management. *Laryngoscope Investig Otolaryngol* **5**, 1217-1226

(2020). https://doi.org:10.1002/lio2.508

61   Kocyigit, M. *et al.* Association Between Endocrine Diseases and Serous Otitis Media in

Children. *J Clin Res Pediatr Endocrinol* **9**, 48-51 (2017). https://doi.org:10.4274/jcrpe.3585

62   Ercolini, A. M. & Miller, S. D. The role of infections in autoimmune disease. *Clin Exp Immunol*

155, 1-15 (2009). https://doi.org:10.1111/j.1365-2249.2008.03834.x

63   Dresser, L., Wlodarski, R., Rezania, K. & Soliven, B. Myasthenia Gravis: Epidemiology,

Pathophysiology and Clinical Manifestations. *J Clin Med* **10** (2021).

https://doi.org:10.3390/jcm10112235

64   Shawky, A. M., Almalki, F. A., Abdalla, A. N., Abdelazeem, A. H. & Gouda, A. M. A

Comprehensive Overview of Globally Approved JAK Inhibitors. *Pharmaceutics* **14** (2022).

https://doi.org:10.3390/pharmaceutics14051001

65   Ruperto, N. *et al.* Tofacitinib in juvenile idiopathic arthritis: a double-blind, placebo-controlled,

withdrawal phase 3 randomised trial. *Lancet* **398**, 1984-1996 (2021).

https://doi.org:10.1016/S0140-6736(21)01255-1

66   King, B. A. & Craiglow, B. G. Janus kinase inhibitors for alopecia areata. *Journal of the*

American Academy of Dermatology **89**, S29-S32 (2023).

https://doi.org:10.1016/j.jaad.2023.05.049

67   Tampa, M., Mitran, C. I., Mitran, M. I. & Georgescu, S. R. A New Horizon for Atopic

Dermatitis Treatments: JAK Inhibitors. *J Pers Med* **13** (2023).

https://doi.org:10.3390/jpm13030384

68   Ytterberg, S. R. *et al.* Cardiovascular and Cancer Risk with Tofacitinib in Rheumatoid Arthritis.

The New England journal of medicine **386**, 316-326 (2022).

https://doi.org:10.1056/NEJMoa2109927

69   Santoro, J. D. *et al.* Assessment and Diagnosis of Down Syndrome Regression Disorder:

International Expert Consensus. *Front Neurol* **13**, 940175 (2022).

https://doi.org:10.3389/fneur.2022.940175

70   Harris, P. A. *et al.* Research electronic data capture (REDCap)--a metadata-driven methodology

and workflow process for providing translational research informatics support. *J Biomed Inform*

42, 377-381 (2009). https://doi.org:10.1016/j.jbi.2008.08.010

71   Gu, Y. *et al.* High-throughput multiplexed autoantibody detection to screen type 1 diabetes and

multiple autoimmune diseases simultaneously. *EBioMedicine* **47**, 365-372 (2019).

https://doi.org:10.1016/j.ebiom.2019.08.036

72   Finck, R. *et al.* Normalization of mass cytometry data with bead standards. *Cytometry A* **83**, 483-

494 (2013). https://doi.org:10.1002/cyto.a.22271

73   Zunder, E. R. *et al.* Palladium-based mass tag cell barcoding with a doublet-filtering scheme and

single-cell deconvolution algorithm. *Nature protocols* **10**, 316-333 (2015).

https://doi.org:10.1038/nprot.2015.020

74   Hahne, F. *et al.* flowCore: a Bioconductor package for high throughput flow cytometry. *BMC*

bioinformatics **10**, 106 (2009). https://doi.org:10.1186/1471-2105-10-106

75   Chevrier, S. *et al.* Compensation of Signal Spillover in Suspension and Imaging Mass

Cytometry. *Cell Syst* **6**, 612-620 e615 (2018). https://doi.org:10.1016/j.cels.2018.02.010

76   Azad, A., Rajwa, B. & Pothen, A. flowVS: channel-specific variance stabilization in flow

cytometry. *BMC Bioinformatics* **17**, 291 (2016). https://doi.org:10.1186/s12859-016-1083-9

77   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome biology* **15**, 550 (2014). https://doi.org:10.1186/s13059-

014-0550-8

78   Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for

interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550

(2005). https://doi.org:10.1073/pnas.0506580102

79   de Jesus, A. A. *et al.* Distinct interferon signatures and cytokine patterns define additional

systemic autoinflammatory diseases. *The Journal of clinical investigation* **130**, 1669-1682

(2020). https://doi.org:10.1172/JCI129301

80   Banchereau, R. *et al.* Personalized Immunomonitoring Uncovers Molecular Networks that

Stratify Lupus Patients. *Cell* **165**, 1548-1550 (2016). https://doi.org:10.1016/j.cell.2016.05.057

Article and author information.

Acknowledgements: This work was supported primarily by NIH grant R61AR077495. Additional funding was provided by NIH grants R01AI150305 (J.M.E.), T32CA190216 (K.A.W.), 2T32AR007411-31 (K.A.W.), UM1TR004399 (data generation and REDCap support), P30CA046934 (support of shared resources), the Linda Crnic Institute for Down Syndrome, the Global Down Syndrome Foundation, the Anna and John J. Sie Foundation, the Human Immunology and Immunotherapy Initiative, the University of Colorado School of Medicine, the Boettcher Foundation, and Fast Grants. We are grateful to all research participants and their families involved in the Human Trisome Project and the clinical trial. We thank Lyndy Bush for administrative support, Dr. Kim Jordan and her team at the Human Immune Monitoring Shared Resource for outstanding service in generation of the immune marker dataset, and Dr. Eric Clambey and his team at the Flow Cytometry Shared Resource for outstanding service in generation of the mass cytometry dataset. We are also grateful to the Colorado Translational and Sciences Institute and the Colorado Multiple Institutional Review Board for assistance in all clinical research projects involving the Crnic Institute. Special thanks to Michelle Sie Whitten, the team at the Global Down Syndrome Foundation, Dr. John Reilly, and Dr. Ron Sokol for logistical support at multiple stages of the project.

Author Contributions Statement: A.L.R.: conceptualization, methodology, formal analysis, project administration, data curation, visualization, investigation, resources, writing of manuscript; E.W.: methodology, data curation, investigation, writing of manuscript; E.G.: methodology, data curation, investigation, writing of manuscript; BEE: methodology, project administration, data curation, investigation, writing of manuscript; K.R.W.: methodology, data curation, investigation, writing of manuscript; K.P.S.: methodology, project administration, formal analysis, data curation, visualization, investigation, writing of manuscript; P.A.: conceptualization, methodology, formal analysis, data curation, visualization, investigation, writing of manuscript; K.A.W.: conceptualization, methodology,

investigation, writing of manuscript; R.E.G.: methodology, investigation, writing of manuscript; E.B.: methodology, investigation, writing of manuscript; H.R.L.: methodology, investigation, writing of manuscript; M.G.D.: methodology, investigation, formal analysis, data curation, visualization, writing of manuscript; N.P.E.: methodology, investigation, formal analysis, data curation, visualization, writing of manuscript; A.A.H.: conceptualization, methodology, project administration, investigation, writing of manuscript; B.M.: methodology, data curation, investigation, writing of manuscript; K.D.S.: methodology, investigation, visualization, writing of manuscript; L.P.: methodology, investigation, formal analysis, data curation, project administration, writing of manuscript; D.F.: methodology, investigation, formal analysis, data curation, project administration, writing of manuscript; M.D.G.: methodology, investigation, formal analysis, data curation, visualization, project administration, writing of manuscript; C.A.D.: conceptualization, methodology, data curation, investigation, project administration, writing of manuscript; D.A.N.: conceptualization, methodology, data curation, investigation, resources, project administration, writing of manuscript; J.M.E.: conceptualization, methodology, project administration, visualization, investigation, resources, writing of manuscript.

Competing Interests Statement: J.M.E. has provided consulting services for Eli Lilly Co., Gilead Sciences Inc., and Biohaven Pharmaceuticals and serves on the advisory board of Perha Pharmaceuticals. The remaining authors declare no competing interests.

Supplementary File Legends.

Supplementary file 1. (A) Cohort characteristics and (B) autoimmune conditions for Human Trisome Project participants analyzed in this study.

Supplementary file 2. Autoantibody measurements of Human Tisome Project participants: (A) anti-thyroid peroxidase (TPO) reactivity; (B) anti-nuclear antigen (ANA) reactivity; (C) SciLifeLabs autoantigen peptide array data.

Supplementary file 3. (A) Minimum qualifying scores for skin conditions. (B) Cohort characteristics for clinical trial participants.

Supplementary file 4. Adverse events for clinical trial participants.

Supplementary file 5. Skin pathology metrics for clinical trial participants: (A) Investigator's Global Assessment (IGA); (B) Dermatology Life Quality Index (DLQI); (C) Severity of Alopecia Tool (SALT); (D) Psoriasis Area and Severity Index (PASI); and (E) Eczema Area and Severity Index (EASI).

Supplementary file 6. (A) DS IFN scores; (B) Cytokine scores; (C) anti-thyroid peroxidase (TPO) titers; (D) anti-transglutaminase (TG) titers; and (E) anti-thyroid stimulating hormone receptor (TSHR) titers for clinical trial participants.

Supplementary file 7. Marker information for mass cytometry analysis.

Supplementary file 8. Clinical trial protocol.