MUTATE: A Human Genetic Atlas of Multi-organ AI Endophenotypes using GWAS Summary Statistics

Aleix Boquet-Pujadas[1], Jian Zeng[2], Ye Ella Tian[3], Zhijian Yang[4], Li Shen[5], the MULTI consortium[#], Andrew Zalesky[3], Christos Davatzikos[6], Junhao Wen[1*],

[1]Laboratory of AI and Biomedical Science (LABS), Columbia University, New York, NY, USA
[2]Institute for Molecular Bioscience, University of Queensland, Brisbane, QLD 4072, Australia
[3]Melbourne Neuropsychiatry Centre, Department of Psychiatry, Melbourne Medical School, The University of Melbourne, Melbourne, Victoria, Australia
[4]GE Healthcare, Bellevue, WA, USA
[5]Department of Biostatistics, Epidemiology and Informatics, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA
[6]Artificial Intelligence in Biomedical Imaging Laboratory (AIBIL), Center for AI and Data Science for Integrated Diagnostics (AI²D), Perelman School of Medicine, University of Pennsylvania, Philadelphia, USA
[7]Center for Innovation in Imaging Biomarkers and Integrated Diagnostics (CIMBID), Department of Radiology, Columbia University, New York, NY, USA
[8]New York Genome Center (NYGC), New York, NY, USA

[*]Corresponding authors:
Junhao Wen, junhao.wen89@gmail.com
622 W 168th St, New York, NY 10032
Consortium representative: Dr. Junhao Wen

Graphical abstract

Keywords

Multi-organ AI endophenotypes, genetic correlation, Mendelian randomization, polygenic risk score

Key points

- Two AI- and neuroimaging-derived subtypes of schizophrenia (MAE-SCZ1 and MAE-SCZ2) show lower polygenicity and weaker negative selection signatures than the disease endpoint/diagnosis of schizophrenia, supporting the endophenotype hypothesis.
- Brain AI endophenotypes are more polygenic than other organ systems.
- Most multi-organ AI endophenotypes exhibit negative selection signatures, whereas a small proportion of brain patterns of structural covariance networks exhibit positive selection signatures.
- The 2024 multi-organ AI endophenotypes are genetically and causally associated with within-organ and cross-organ disease endpoints/diagnoses.

Abstract

Artificial intelligence (AI) has been increasingly integrated into imaging genetics to provide intermediate phenotypes (i.e., endophenotypes) that bridge the genetics and clinical manifestations of human disease. However, the genetic architecture of these AI endophenotypes remains largely unexplored in the context of human multi-organ system diseases. Using publicly available GWAS summary statistics from the UK Biobank, FinnGen, and the Psychiatric Genomics Consortium, we comprehensively depicted the genetic architecture of 2024 multi-organ AI endophenotypes (MAEs). Two AI- and imaging-derived subtypes[1] showed lower polygenicity and weaker negative selection effects than schizophrenia disease diagnoses[2], supporting the endophenotype hypothesis[3]. Genetic correlation and Mendelian randomization analyses reveal both within-organ relationships and cross-organ interconnections. Bi-directional causal relationships were established between chronic human diseases and MAEs across multiple organ systems, including Alzheimer's disease for the brain, diabetes for the metabolic system, asthma for the pulmonary system, and hypertension for the cardiovascular system. Finally, we derived polygenic risk scores for the 2024 MAEs for individuals not used to calculate MAEs and returned these to the UK Biobank. Our findings underscore the promise of the MAEs as new instruments to ameliorate overall human health. All results are encapsulated into the MUTATE genetic atlas and are publicly available at https://labs-laboratory.com/mutate.

Introduction

Multi-organ research[1,4–11] represents a pivotal frontier in advancing our understanding of human aging and disease. In particular, integrating artificial intelligence (AI) into multi-organ imaging genetics[1,4,12,6] has emerged as a novel approach, offering potential promise in advancing precision medicine[13]. This integration introduces a new array of endophenotypes[14,15], serving as intermediate, often quantitative, phenotypes, potentially reshaping how we perceive and approach medical AI[16] in imaging and genetic research.

In recent years, three primary catalysts have significantly advanced the field of genetics. The first pivotal factor stems from the extensive collaborative efforts in consolidating large-scale multi-omics datasets, which has endowed researchers with unprecedented statistical power previously inaccessible. As an illustration, the UK Biobank (UKBB) study[17] stands out for its comprehensive collection of multi-organ imaging[18], genetics[19], and proteomics[20,21] data within the United Kingdom. Similarly, the FinnGen study[22], conducted in Finland, has amassed extensive clinical and genetic data. Secondly, efforts toward open science have propelled the field, especially emphasizing the significance of publicly available resources, such as genome-wide association study (GWAS) summary statistics and widespread scientific dissemination. Notably, the FinnGen study and Psychiatric Genomics Consortium (PGC[23]) have publicly made all the GWAS summary statistics accessible[22]. Public GWAS platforms such as the GWAS Catalog[24], OpenGWAS[25], and GWAS ATLAS[26] have consolidated and harmonized vast GWAS datasets, rendering them suitable for subsequent genetic analyses. Likewise, such good practice was also employed in the newly burgeoning field of brain imaging genetics[27], including the BIG40 (https://open.win.ox.ac.uk/ukbiobank/big40/), the BIG-KP (https://bigkp.org/), BRIDGEPORT (https://labs-laboratory.com/bridgeport), and MEDICINE (https://labs-laboratory.com/medicine) knowledge portals. Finally, advanced computational genomics statistical methods using solely GWAS summary statistics, along with sufficient linkage disequilibrium information, have been developed, presenting an unparalleled chance to comprehend the genetic architecture of highly polygenic disease traits. For example, LDSC[28] has been extensively utilized to estimate single-nucleotide polymorphism (SNP)-based heritability and genetic correlations. Mendelian randomization[29] is a statistical method to dissect associations further, probing potential causal relationships among these complex human disease traits, although these methods often rely on several sensitive model assumptions[30].

Despite these advancements, the intricate genetic foundation shaping these AI endophenotypes in the context of pleiotropic human disease endpoints (DE) within multi-organ systems remains largely uncharted. We previously applied AI to imaging genetic data and derived 2024 multi-organ AI endophenotypes (MAE). These encompassed 2003 multi-scale brain patterns of structural covariance (PSC) networks generated through a deep learning-analogy non-negative matrix factorization method[12] (visualization for C32_1 encompassing deep subcortical structures: https://labs-laboratory.com/bridgeport/MuSIC/C32_1), 9 dimensional neuroimaging endophenotypes (DNE) quantifying neuroanatomical heterogeneity (also known as disease subtype) within 4 common brain diseases[1], and 12 biological age gap (BAG) assessing the individual deviation in typical aging (i.e., acceleration or deceleration from the chronological age) across 9 human organ systems[4,6] (**Supplementary eTable 1a**). To contribute to open science[31], we made all the GWAS summary statistics derived from UKBB data publicly available at the MEDICINE knowledge portal: https://labs-laboratory.com/medicine. In addition, FinnGen analyzed genetic data for 2269 binary and 3 quantitative DEs from 377,277 individuals and 20,175,454 variants. They made these massive GWAS summary statistics publicly available

to the community at https://finngen.gitbook.io/documentation/ (**Supplementary eTable 1b**).
Finally, PGC consolidated GWAS results focused on neurological disorders worldwide and
made the GWAS summary statistics accessible to the research community (https://pgc.unc.edu/,
Supplementary eTable 1c).
This study harnesses the extensive GWAS summary resources made publicly available
by us on behalf of UKBB, FinnGen, and PGC (**Method 1**), along with the utilization of several
advanced computational genomics statistical methods (refer to **Code Availability**), to thoroughly
depict the genetic architecture of the 2024 MAEs (**Method 2**) and 525 DEs (>5000 cases) in the
context of multi-organ investigations. Importantly, our previous research explored the genetic
foundation of the 2024 MAEs but did not systematically encompass the FinnGen or PGC data.
Specifically, we included 521 DEs released by the FinnGen study, accessible at
https://finngen.gitbook.io/documentation/v/r9/, and 4 brain DEs (Alzheimer's disease (AD),
Attention-deficit/hyperactivity disorder (ADHD), bipolar disorder (BIP), and schizophrenia
(SCZ)) from PGC (https://pgc.unc.edu/). This study expanded on this by systematically
benchmarking the genetic analyses and comprehensively comparing various statistical
methodologies[28,30,32–38] (**Method 3**). Specifically, we aimed to compute the SNP-based
heritability (h^2_{SNP}), polygenicity (π), the relationship between SNP effect size and minor allele
frequency (S: signature of natural selection, genetic correlation (r_g), causality, and polygenic risk
score (PRS) between the 2024 MAEs and 525 DEs. These findings were encapsulated within the
MUTATE (**MU**l**T**i-organ **A**I endopheno**T**yp**E**) genetic atlas, which is publicly available at
https://labs-laboratory.com/mutate.

Results

The genetic architecture of the 2024 MAEs and 525 DEs

We computed three parameters to fully depict the genetic architecture of the 2024 MAEs (**Method 3a**). For the SNP-based heritability (h^2_{SNP}), SBayesS[39] obtained the highest h^2_{SNP} for the 2016 brain MAEs (mean $\hat{h}^2_{SNP}$=0.13 [0.01, 0.38]), followed by the pulmonary BAG (0.16±0.004), the eye BAG (0.14±0.009), the cardiovascular BAG (0.12±0.003), the renal BAG (0.10±0.003), and the musculoskeletal BAG (0.10±0.003) (**Fig. 1a** and **Supplementary eFile 1**). It is worth noting that SNP-based heritability varies across methods and depends on the input data, i.e., summary data or individual-level genotype data used in the method[40]. We aimed to benchmark the summary data-based methods by comparing the results from SBayesS with those of LDSC[28] and SumHer[33]. Overall, while the estimates from the three methods were highly correlated (r=0.97 between LDSC and SumHer; r=0.99 between SBayesS and SumHer; r=0.99 between SBayesS and LDSC; **Supplementary eFigure 1**), SumHer (0.23±0.14) generally yielded larger h^2_{SNP} estimates than both LDSC (0.16±0.10) and SBayesS (0.13±0.08) (**Supplementary eFile 1**). We present the h^2_{SNP} estimate of the 525 DEs and 2024 MAEs in **Supplementary eFigure 2**. **Supplementary eFile 2** presents the results of the 525 DEs. For the 525 DEs, we converted the h^2_{SNP} estimates from the observed scales to the liability scales, following the recommendations of Ojavee et al[41]. It's important to clarify that we did not intend to compare the h^2_{SNP} estimates of the two data sources due to differences in genotype coverage, sample sizes, allele frequencies, and other factors.

We then computed the natural selection signature (S) for the 2024 MAEs. The metabolic BAG showed a strong negative selection (S=-0.82±0.10), followed by the pulmonary BAG (S=-0.79±0.05), the hepatic BAG (S=-0.74±0.09), the renal BAG (S=-0.68±0.08), and the immune BAG (S=-0.66±0.11). For the brain MAEs (S=-0.33 [-1, 0.43]), the brain BAG and (S=-0.70±0.12) the subtype (ASD1) for autism spectrum disorder[42] (S=-0.90±0.11) showed strong negative selection effects (**Fig. 1b** and **Supplementary eFile 3**).

Finally, we calculated the polygenicity (π) for the 2024 MAEs. We found that brain MAEs (0.040 [0.003, 0.072]) showed higher polygenicity than other organ systems (t-statistic=5.75; P-value=1.03x10^{-8}), followed by the pulmonary BAG (0.018±0.001), the musculoskeletal BAG (0.013±0.001), and the cardiovascular BAG (0.011±0.001) (**Fig. 1C** and **Supplementary eFile 4**). The PSC (C128_115: https://labs-laboratory.com/bridgeport/MuSIC/C128_115) showed the highest polygenicity estimate (0.072±0.002).

Potential evidence for the endophenotype hypothesis

Previous studies[43,44] have found supporting evidence for the endophenotype hypothesis[14,15] using traditional brain map-based signatures, showing that more genetic variants are associated with disease endpoints than imaging-derived signatures (i.e., endophenotypes). Of note, considering genetic differences between FinnGen and UKBB samples, SBayesS with the UKBB as LD reference may give biased estimates of S and π (LD from FinnGen not fully available; **Method 3a**). Therefore, we used the GWAS summary data for PGC schizophrenia (SCZ[2]) and two subtypes of SCZ (SCZ1 and SCZ2[1]) from our UKBB analysis to demonstrate this. The advantage of using PGC data is that the GWAS summary statistics are better powered (large sample sizes), and the data were from European ancestry groups across different countries. A data harmonization procedure is outlined in **Supplementary eMethod 1** to ensure a fair

comparison of these estimates, which led to the utilization of a common set of SNPs and linkage disequilibrium information for computing the *S* and parameters. Our results showed that MAE-SCZ1 (=0.048±0.002; *S*=-0.61±0.09) and MAE-SCZ2 (=0.047±0.002; *S*=-0.54±0.12) had lower polygenicity signals and weaker negative selection effects than PGC-SCZ (=0.055±0.003; *S*=-0.82±0.04) (**Fig. 1d**). **Supplementary eFigure 3** shows the Manhattan plot of the harmonized summary data for MAE-SCZ1, MAE-SCZ2, and PGC-SCZ. These findings potentially support the endophenotype hypothesis[3], which suggests that intermediate phenotypes (e.g., SCZ subtype MAEs) reside inside the causal pathway from genetics to exo-phenotypes (e.g., SCZ binary diagnosis), making them closer to the underlying etiology[43,44].

Figure 1: **The genetic architecture of the 2024 MAEs**
Three parameters are estimated by SBayesS to delineate the genetic architecture of the 2024 MAEs, including (**a**) the SNP-based heritability (   ), (**b**) the relationship between MAF and effect size (*S*), and (**c**) polygenicity ( ). (**d**) We compared the   and *S* parameters using harmonized GWAS summary data for two AI- and imaging-derived subtypes (MAE-SCZ1 and MAE-SCZ2[1]) from UKBB and the disease endpoint of schizophrenia (PGC-SCZ[2]) from PGC. FinnGen data was not used due to bias stemming from the unavailability of FinnGen-specific linkage disequilibrium data (**Supplementary eMethod 1**). We present the distribution of the estimated parameters for the 2016 brain MAEs using a violin plot; the mean value is denoted by the black horizontal line. These results should be interpreted cautiously for comparative purposes due to limitations stemming from the lack of individual genotype data from FinnGen and PGC, differing linkage disequilibrium structures, and varying sample sizes.

The genetic correlation shows organ-specific and cross-organ associations

We found 132 (P-value < 0.05/2024) and 45 (P-value < 0.05/2024/525) commonly significant positive genetic correlations (r_g) after applying two levels of Bonferroni correction (**Fig. 2**) for the LDSC[28] and GNOVA[34] methods (**Method 3b**, **Supplementary eFile 5**, and **Supplementary eTable 2**). We noted that HDL encountered convergence issues with the models, as detailed in **Method 3b**.

Between these methods, the magnitude of the genetic correlations for the significant signals for both methods differed: mean $\hat{r}_g$=0.24[-0.40~0.52] with 213 significant signals for LDSC, mean $\hat{r}_g$=0.17[-0.30~0.62] for GNOVA with 428 significant signals (**Fig. 2**). The three sets of converged estimates showed a strong correlation: r=0.77 (P-value<1x10^{-10}; N=1,062,577) between LDSC and GNOVA, r=0.81 (P-value<1x10^{-10}; N=59,289) between LDSC and HDL, and r=0.82 (P-value<1x10^{-10}; N=59,289) between GNOVA and HDL. **Supplementary eFigure 4** shows the correlation of the three sets of estimates.

Within the significant signals identified, we observed *i*) organ-specific associations, in which the MAE showed a genetic association with the DE originating from the respective organ system, and *ii*) cross-organ connections, in which the MAE and DE were primarily involved from different organ systems. For example, two brain PSCs showed significant negative genetic correlations with BIP from PGC (C512_368 vs. BIP: -0.16±0.03; C1024_114 vs. BIP: -0.15±0.03). At a less stringent level, the brain MAEs were also genetically associated with DEs from other organ systems, including the positive correlation between C1024_808 and obesity (E4_OBESITY: r_g=0.17±0.13). The cardiovascular BAG was positively correlated with several DEs related to the cardiovascular system, including ischemic heart disease (I9_IHD: r_g=0.26±0.03), coronary heart disease (I9_HEARTFAIL_AND_CHD: r_g=0.26±0.03), angina (I9_ANGINA: r_g=0.25±0.03) and atrial fibrillation (I9_AF: r_g=0.22±0.04). Likewise, the pulmonary BAG was positively associated with multiple DEs related to the lung and respiratory system, including chronic obstructive pulmonary disease (COPD_EARLY: r_g=0.47±0.04) and various forms of asthma (ASTHMA_NAS: r_g=0.43±0.04). Cross-organ connections were established, such as between the pulmonary BAG and substance abuse (KRA_PSY_SUBSTANCE_EXMORE: r_g=0.20±0.03) and hypertension (I9_HYPTENS: r_g=0.17±0.03). Lastly, the metabolic BAG was largely linked to different forms of diabetes (T2D: r_g=0.40±0.04).

Figure 2: **Genetic correlation between the 2024 MAEs and 525 DEs**

The significant genetic correlation estimates (r_g) between 2024 MAEs and 525 DEs are depicted, considering two levels of corrections for multiple comparisons, considering the relatively smaller sample sizes (<40,000) for brain MAEs compared to other organ MAEs (>100,000). Initially, we

reveal significant results shared between LDSC and GNOVA, employing Bonferroni correction
based solely on the number of MAEs (P-value<0.05/2024), uncovering 133 MAE-AE pairs.
Subsequently, a stricter correction based on both the number of MAEs and DEs is applied,
leading to 45 unique MAE-AE pairs marked as red squares; the numeric results are displayed
using results from LDSC. The genetic correlation for non-significant results was set to 0 for
visualization purposes. For the MAEs, readers can explore the BRIDGEPORT portal for a visual
representation of the 2003 brain PSCs (e.g., C256_225: https://labs-
laboratory.com/bridgeport/MuSIC/C256_225) and the other BAGs at the MEDICINE portal:
https://labs-laboratory.com/medicine.
239
The brain, cardiovascular, and pulmonary MAEs are causally linked to DEs of multiple
organ systems
Employing five distinct two-sample Mendelian randomization estimators, we identified 39 (P-
value<0.05/633) and 15 (P-value<0.05/633/524) significant causal relationships, directed from
the MAE to DE, that withstood the Bonferroni correction at two different levels of rigors, as per
the inverse variance weighted (IVW) estimator and at least one of the other four estimators
(**Method 3c** and **Supplementary eTable 3**).
Within the 15 significant causal relationships, the brain MAEs showed causal
connections with DEs from the brain, as well as DEs from other organ systems. For example, the
brain PSC (C1024_598) was causally linked to SCZ from PGC [P-value=9.89×10^{-8}; OR (95%
CI)=0.69 (0.59, 0.79); the number of IVs=7]. C1024_684 was causally linked to Ventral hernia
from FinnGen [K11_VENTHER: P-value=1.09×10^{-7}; OR (95% CI)=1.43 (1.25, 1.63); the
number of IVs=18]. The pulmonary BAG was causally linked to multiple DEs related to the
pulmonary system, including chronic obstructive pulmonary disease (COPD) [J10_COPD: P-
value=2.70×10^{-20}; OR (95% CI)=1.77 (1.56, 2.00); the number of IVs=59] and asthma
[ASTHMA_PNEUMONIA: P-value=1.51×10^{-14}; OR (95% CI)=1.67 (1.41, 1.96); the number of
IVs=59]. The cardiovascular BAG was causally linked to ischemic heart disease (IHD)
[ASTHMA_PNEUMONIA: P-value=1.09×10^{-7}; OR (95% CI)=1.64 (1.36, 1.96); the number of
IVs=37] (**Fig. 3**). The quality check of the significant signals is presented in **Supplementary**
eFolder 1. **Supplementary eFile 6** presents the full set of results for the 521 FinnGen DEs and 4
PGC DEs.
261
262

Figure 3: **Causal relationship from the 2024 MAEs to the 525 DEs**
The causal relationship from the 2024 MAEs to the 525 DEs revealed 39 significant MAE-DE pairs, involving 633 MAEs as effective exposure variables (>8 instrumental variables before harmonization) and 525 DEs as outcomes. Bonferroni correction was applied to identify potential significant causal signals based on *i*) the 633 MAEs (P-value<0.05/633) and *ii*) the 633 MAEs and 525 DEs (P-value<0.05/633/524, denoted by the 15 red rectangles). Furthermore, we verified that the statistical significance attained for the IVW estimator was consistent and persisted across at least one of the other four Mendelian randomization estimators (Egger, weighted median, simple mode, and weighted mode estimators). For visualization purposes, the odds ratios for non-significant results were set to 1 and were left blank. For the MAEs, readers can explore the BRIDGEPORT portal for a visual representation of the 2003 brain PSCs (e.g., C32_4: https://labs-laboratory.com/bridgeport/MuSIC/C32_4) and the other BAGs at the MEDICINE portal: https://labs-laboratory.com/medicine.

The DEs involving Alzheimer's disease, diabetes, asthma, and hypertension exert causal effects on multi-organ MAEs
We then tested the inverse causality by employing the DEs as exposure and MAEs as outcome variables. We identified 47 (P-value<0.05/787) and 23 (P-value<0.05/787/214) significant causal relationships, directed from the DE to MAE, that survived the Bonferroni correction at two different levels of rigors (**Method 3c** and **Supplementary eTable 4**).

Within the 23 significant causal relationships (P-value<0.05/787/214), various forms of Alzheimer's disease were linked to the brain MAEs, including the brain BAG [G6_AD_WIDE: P-value=3.03×10^{-7}; OR (95% CI)=1.10 (1.06, 1.13); the number of IVs=8] and metabolic BAG [G6_AD_WIDE: P-value=3.03×10^{-7}; OR (95% CI)=1.07 (1.04, 1.09); the number of IVs=8]. Type 1 diabetes (E4_DM1NASCOMP) was also causally linked to multiple brain PSCs. In addition, the cardiovascular BAG was causally linked to multiple heart diseases, including hypertension [I9_HYPTENS: P-value=4.67×10^{-31}; OR (95% CI)=1.23 (1.19, 1.27); the number of IVs=110]. Several forms of asthma were causally linked to the pulmonary BAG, such as allergic asthma [ALLERG_ASTHMA: P-value=2.38×10^{-9}; OR (95% CI)=1.09 (1.06, 1.13); the number of IVs=14]. Finally, obesity was also linked to the renal BAG [E4_OBESITY: P-value=2.74×10^{-8}; OR (95% CI)=1.11 (1.07, 1.15); the number of IVs=19] (**Fig. 4**).

295 **Supplementary eFolder 2** presents the quality check results of the significant signals.
296 **Supplementary eFile 7** presents the full set of results for the 521 FinnGen DEs and 4 PGC DEs.
297

298
Figure 4: **Causal relationship from the 525 DEs to the 2024 MAEs**
The causal relationship from the 525 MAEs to the 2024 DEs revealed 47 significant DE-MAE
pairs, involving 214 DEs as effective exposure variables (>8 instrumental variables before
harmonization) and 787 DEs as effective outcomes after quality checks. Bonferroni correction
was applied to identify potential significant causal signals based on *i*) the 787 MAEs (P-
value<0.05/787) and *ii*) the 787 MAEs and 214 DEs (P-value<0.05/787/214, denoted by the 23
red rectangles). Furthermore, we verified that the statistical significance attained for the IVW
estimator was consistent and persisted across at least one of the other four Mendelian
randomization estimators (Egger, weighted median, simple mode, and weighted mode
estimators). For visualization purposes, the odds ratios for non-significant results were set to 1
and were left blank. For the MAEs, readers can explore the BRIDGEPORT portal for a visual
representation of the 2003 brain PSCs (e.g., C128_13: https://labs-
laboratory.com/bridgeport/MuSIC/C128_13) and the other BAGs at the MEDICINE portal:
https://labs-laboratory.com/medicine.
313
The polygenic risk scores of the 2024 MAEs

Using the PRS-CS[45] method, we derived the PRS of the 2024 MAEs. We found that the 1799
MAEs could significantly (P-value<0.05/2024) predict the phenotypic BAGs in the test/target
data (split2 GWAS; detailed in **Method 3d**). Among these, 1791 brain MAEs resulted in
significant incremental R^2 ranging from 0.11% to 10.70% to predict the phenotype of interest.
For example, the PSC (C1024_593 for part of the cerebellum: https://labs-
laboratory.com/bridgeport/MuSIC/C1024_593) showed an incremental of R^2 10.70%. The renal
BAG showed an incremental R^2 of 5.92%, followed by the metabolic ($R^2 = 5.67\%$) and
pulmonary BAG ($R^2 = 3.86\%$) (**Fig. 5a** and (**Supplementary eFile 8**).
We then applied the model to the entire UKBB population and performed a PRS-wide
association study (PWAS), where the 2024 PRS-MAEs were linked to the 59 phenotypes that
were not initially used to compute the respective PRS, to avoid the circular bias[46]
(**Supplementary eTable 5**). Refer to **Method 3d** for details. We found 388 significant
associations (P-value<0.05/2024/59) between 7 PRS-MAEs and 41 phenotypes. Among these,
PSC C32_1 showed the most associations (94%); the lifestyle factor for only fish intake (Field
ID: 16) was highly linked to multiple PRS-MAEs (16%). These results were expected because
the 59 phenotypes (e.g., cognitive and mental traits) are primarily linked to the brain, and
lifestyle factors were largely linked to multiple organ systems (**Fig. 5b** and **Supplementary
eFile 9**). All derived PRS will be returned to UKBB and made available to the community.
333
334

Figure 5: **The polygenic risk score of the 2024 MAEs and PWAS**

(**a**) The incremental R^2 of the PRS derived by PRC-CS to predict the 2024 MAEs in the target/test data (i.e., the split2 GWAS). The y-axis indicates the proportions of phenotypic variation that the PRS can significantly and additionally explain (i.e., incremental R^2). The x-axis lists the 8 organ systems. For the brain, we showed the PRS distribution of the significant results from the 1791 brain PRS-MAEs; the other organ systems only have one PRS-MAE. (**b**) The PWAS links the PRS-MAEs to the 59 additional phenotypes not used to compute the PRS-MAE in the entire UKBB sample (P-value<0.05/2024/59).

Discussion

This study expands previously established genetic atlases[47,32] by integrating AI-derived endophenotypes via the 2024 MAEs within the multi-organ framework solely through GWAS summary statistics. We demonstrate a promising avenue for advancing imaging genetic research in two key aspects: *i*) integrating AI in imaging genetics and *ii*) exploring human aging and disease through a multi-organ perspective.

By comprehensively depicting the genetic architecture of the 2024 MAEs, we showcased that AI endophenotypes supported the endophenotype hypothesis[14,15], in which they showed lower polygenicity and weaker negative selection effects than the disease diagnosis. First, it may suggest that these intermediate phenotypes exist along the causal pathway, bridging the gap between underlying genetics and "exo-phenotypes" like cognitive decline or disease diagnoses in case/control studies, thus positioned closer to the core etiology and pathology. Secondly, many of these 2024 MAEs originated from *in vivo* imaging methodologies like magnetic resonance imaging (MRI). Consequently, they tend to exhibit reduced noise levels (i.e., a higher SNR) in capturing disease-related effects and are less susceptible to biases, such as misclassification[48], case/control-covariate sample bias (e.g., studies matching comorbidities and other factors), and imbalanced case/control ratios, as evidenced in many GWASs in FinnGen. Especially for the former, binary traits have a threshold for disease classification, leading to the dichotomization of individuals into affected and unaffected categories. Thirdly, the 525 DEs often represent complex diseases highly influenced by multiple genetic and environmental factors. Their multifaceted nature, involving numerous genes with modest effects and environmental interactions[49], can lead to a higher vulnerability to disease onset and clinical symptoms. Consistent with this observation, we previously also found that one AI- and imaging-derived subtype of Alzheimer's disease[50] (AD1), but not the binary disease diagnosis, was genetically correlated with brain age (GM- and WM-BAG)[6].

We observed that brain MAEs were overall more polygenic than MAEs from other organ systems. Brain disorders are highly polygenic[51]. First, the brain is a highly complex organ with intricate functions, and disorders affecting it are likely influenced by a larger number of genetic variants[12,52]. Second, many brain disorders are multifaceted, involving various aspects of brain structure, function, and connectivity, which can be influenced by various genetic factors[19]. Additionally, the brain regulates many physiological processes throughout the body, so disruptions in its function can have widespread effects, potentially involving interactions with multiple organ systems[4]. In addition, we found that most of the brain MAEs showed negative selection signatures, including the 9 disease subtype DNEs and 4 brain BAGs; some of the brain PSCs showed a positive *S* estimate (e.g., for the occipital lobe and subcortical structure, *S*=0.31±0.09: https://labs-laboratory.com/bridgeport/MuSIC/C32_18). The anticipated negative selection signatures of biological age across multiple organs and disease subtypes are expected to align with our prior findings, which revealed pervasive signatures of natural selection across a range of complex human traits and functional genomic categories. This negative selection signature prevents mutations with large deleterious effects from becoming frequent in the population[53]. The positive selection signatures identified in certain brain PSCs may suggest that positive selection may also play a role in shaping the genetic architecture of brain structural networks.

The MUTATE atlas uncovered both established and previously undiscovered interactions concerning human systemic diseases within individual organs and across diverse organ systems. For example, within the cardiovascular system, the AI-derived MAE, cardiovascular BAG

showed both substantial genetic correlation (**Fig. 2**) and bi-directional causality (**Fig. 3** and **4**) with multiple heart diseases, such as ischaemic heart disease[54], heart failure[55], and atrial fibrillation[56]. Similarly, pulmonary BAG was also causally linked to multiple diseases related to the lung and respiratory system, including COPD[57] and various forms of asthma[58]. Another organ-specific connection was observed in neurologic diseases, encompassing conditions such as AD[59] and various mental disorders[60] linked to several MAEs associated with the brain, notably several PSCs and WM-BAG. Cross-organ interplay was evidenced for several novel connections. For instance, the brain PSCs exhibited causal connections to conditions extending beyond the brain, such as ventral hernia and vein diseases, as well as systemic conditions, like various forms of diabetes affecting the entire body. In contrast, AD appears to causally impact multiple BAGs across various human organ systems, including the renal, immune, and metabolic systems. It's widely recognized that AD, being a complex condition, triggers detrimental effects that influence several human organ systems[59,61]. Our previous study used imaging genetics to investigate this multi-organ involvement along the disease continuum[62]. These results highlight the clinical relevance and interpretation of these AI endophenotypes to quantify individual-level organ health.

Emphasizing preventative strategies for specific chronic diseases is crucial to enhancing overall multi-organ health. Our MAEs present opportunities as novel instruments for selecting populations in clinical trials and facilitating therapeutic development. AD and various forms of diabetes exemplify disease endpoints significantly impacting multiple human organ systems. AD stands as the leading cause of dementia in older adults, presenting a persistent challenge in medicine despite numerous pharmacotherapeutic clinical trials. These trials have included interventions, such as anti-amyloid drugs[63,64] and anti-tau drugs.[65]. The complexity and multifaceted nature of the underlying neuropathological processes may account for the lack of effective treatments. We call on the scientific community to embrace various mechanistic hypotheses to elucidate AD pathogenesis beyond amyloid and tau[66,67]. Likewise, the complexity of diabetes, with its various contributing factors, renders prevention challenging[68]. Moreover, diabetes often coexists with other chronic conditions affecting multiple organ systems, such as cardiovascular diseases, hypertension, and dyslipidemia[69]. Successful prevention strategies require a holistic approach, encompassing lifestyle adjustments, education, healthcare access, and societal considerations.

Limitation

This study presents several limitations. Primarily, our analyses were centered solely on GWAS summary statistics derived from individuals of European ancestries. Future investigations should extend these findings to diverse ethnic groups, particularly those that are underrepresented, to ascertain broader applicability. This necessitates the research community's commitment to embracing open science in AI and genetics. Secondly, the computational genomics statistical methods utilized in this research rely on several underlying statistical assumptions, which could potentially be violated and introduce bias. We mitigated bias by employing multiple methodologies to compute heritability, genetic correlation, and causality to address this concern. Additionally, we conducted thorough sensitivity checks, and the detailed results are provided accordingly. Additionally, our analysis was limited by the lack of individual-level genotype data from FinnGen and PGC, highlighting the need for future studies utilizing individual-level data to validate our empirical findings. Finally, our study recognizes a tradeoff between clinical interpretability and the detection of genetic associations when using AI-derived phenotypes.

Outlook

In summary, we introduced the MUTATE genetic atlas to comprehensively comprehend the genetic architecture of AI endophenotypes and chronic diseases in multi-organ science. This investigation underscores the potential of integrating AI into genetic research and supports a comprehensive approach to investigating human diseases within a multi-organ paradigm.

Methods

Method 1: The MULTI consortium
The MULTI consortium is an ongoing initiative to integrate and consolidate multi-organ and multi-omics data, including imaging, genetics, and proteomics. Building on existing consortia and studies, such as those listed below, MULTI aims to curate and harmonize the data to model human aging and disease across the lifespan at individual and summary levels. The present study solely utilized GWAS summary statistics; no individual-level data were used. We downloaded the GWAS summary statistics from three web portals for the 2024 MAEs, 521 DEs from FinnGen and 4 DEs from PGC.

UKBB
UKBB is a population-based study of approximately 500,000 people recruited from the United Kingdom between 2006 and 2010. The UKBB study has ethical approval, and the ethics committee is detailed here: https://www.ukbiobank.ac.uk/learn-more-about-uk-biobank/governance/ethics-advisory-committee.
The GWAS summary statistics for all the 2024 MAEs are publicly available at the MEDICINE knowledge portal: https://labs-laboratory.com/medicine, which focuses on disseminating scientific findings on imaging genetics and AI methods in multi-organ science. Specifically, among the 2024 MAEs, 2003 PSCs – at varying scales from C32 to C1024 – were structural covariance networks derived via the sopNMF method[12]. 9 DNEs[1] captured the neuroanatomical heterogeneity of four brain diseases (AD1-2, ASD1-3, LLD1-2, and SCZ1-2) using semi-supervised clustering or representation learning methods[42,62,71,72]. 12 multi-organ BAGs (GM, WM, FC[6], multimodal brain BAGs, cardiovascular BAG, eye BAG, hepatic BAG, immune BAG, musculoskeletal BAG, metabolic BAG, pulmonary BAG, and renal BAG[73]) were derived from various machine learning models to quantify the individual-level deviation from typical brain aging due to various pathological effects. Detailed AI methodologies are presented in **Method 2** for the MAEs, DNEs, and BAGs. All GWASs were performed within European ancestries and using the GRCh37 human genome assembly; the GWAS model (PLINK[74] for linear model and fastGWA[75] for linear mixed-effect model), sample sizes, and covariates included are detailed in the original papers and also in **Supplementary eTable 1a**.
Harmonization of GWAS summary statistics across different models and consortia for various software is crucial, such as aligning the effect allele and the direction of the effect size. There's currently no established standard in the field for this process, although some advice has been proposed[76]. Certain software harmonizes data based on the allele frequency of the effect allele, such as the *TwoSampleMR* package[77] for Mendelian randomization. In our UKBB MAE GWAS summary data, we harmonized the effect allele as the alternative allele from PLINK and A1 from fastGWA and provided its corresponding allele frequency. P-value, effect sizes (e.g., BETA value and SE), and sample sizes are indicated too. The variant identifier is based on the rs ID number, not the chromosome number and position number combination.

FinnGen
The FinnGen[22] study is a research project based in Finland that explores combined genetics and health registry data to understand the underlying causes and mechanisms behind various disease endpoints. It particularly emphasizes the genetic basis of diseases in the Finnish population (>500,000) by conducting extensive GWAS and analyzing large-scale genomic data in collaboration with multiple research institutions and organizations. FinnGen has generously

made their GWAS results publicly available to the community for research purposes
(https://www.finngen.fi/en/access_results).
The present study used the GWAS summary statistics version R9 released to the public
on May 11, 2022, after harmonization by the consortium. In the R9 release, FinnGen analyzed
2269 binary and 3 quantitative endpoints from 377,277 individuals and 20,175,454 variants.
Regenie[78] was used to run the GWAS models, including sex, age, 10 PCs, and genotyping batch
as covariates. Genotype imputation was done with the population-specific SISu v4.0 reference
panel. In our analysis, we concentrated solely on binary DEs with case numbers exceeding 5000
to ensure adequate statistical power, given the highly imbalanced case/control ratios. As the
released data were based on the GRCh38 human genome assembly, we lifted the GWAS
summary statistics to the GRCh37 version for all genetic analyses. **Supplementary eTable 1b**
details the included 521 DEs. More details can be found at the FinnGen website:
https://finngen.gitbook.io/documentation/v/r9/.
The FinnGen team has systematically harmonized the GWAS summary data for the 521
DEs involved. The alternative allele serves as the effect allele. The rsID number represents the
SNP; the chromosome number and position are also shared. The data includes P-values, effect
sizes, and allele frequencies for both the alternative and reference alleles.
506
Psychiatric Genomics Consortium
PGC[23] is an international coalition of researchers exploring the genetic underpinnings of
psychiatric disorders and beyond. This collaborative effort unites scientists globally to examine
and decipher extensive genomic datasets concerning various brain diseases. The primary goal of
PGC involves uncovering and comprehending the genetic elements that contribute to various
psychiatric disorders, such as schizophrenia, bipolar disorder, and major depressive disorder. We
downloaded GWAS summary statistics from the PGC website (https://pgc.unc.edu/for-
researchers/download-results/) and manually harmonized the data to our Mendelian
randomization analyses to replicate the FinnGen findings.
PGC did not harmonize the GWAS summary statistics; the available data information
depends on each study. **Supplementary eTable 1c** details the 4 DEs (AD, ADHD, bipolar
disorder, and schizophrenia) included after the data filtering procedure. First, we ensured that the
study population comprised individuals of European ancestry and, if necessary, lifted the data to
the human genome build assembly GRCh37. Secondly, we excluded two studies where the allele
frequency is unavailable because the *TwoSampleMR* package[77] requires this information to
harmonize the exposure and outcome data (e.g., flip the effect allele and effect size). Thirdly, we
confirmed that the GWAS summary statistics didn't overlap with UKBB data. Specifically, the
AD GWAS summary data[79] explicitly offered a version that excluded participants from UKBB.
In addition, the original dataset lacked a column for the rsID number. To deal with this, we
employed a mapping approach using the chromosome number and position to the dpSNP
database (version 150), which allowed us to obtain the corresponding rsID numbers. All 4 DE
GWAS summary data went through the same harmonization procedure as FinnGen (**Method 3c**)
529
530
Method 2: 2024 multi-organ AI endophenotypes

(a): The 2003 patterns of structural covariance of the brain
In our earlier study[12], we utilized the sopNMF method on an extensive and varied brain imaging
MRI dataset (*N*=50,699, including data from UKBB) to generate the multi-scale brain PSCs. The

scale C ranges from 32 to 1024, progressively increasing by a factor of 2; 11 PSCs vanished
during models.
Biologically, the 2003 PSCs represent data-driven structural networks that co-vary across
brain regions and individuals in a coordinated fashion. Mathematically, the sopNMF method is a
stochastic approximation ("deep learning-analogy") constructed and extended based on
opNMF[80,81]. Consider an imaging dataset comprising n images, each containing d voxels. We
represent the data as a matrix X, where each column corresponds to a flattened image: $X =$
$[x_1, x_2, ..., x_n], X \in \mathbb{R}_{\geq 0}^{d \times n}$. The method factorizes X into two low-rank matrices $W \in \mathbb{R}_{\geq 0}^{d \times r}$ and
$H \in \mathbb{R}_{\geq 0}^{r \times n}$, subject to two important constraints: i) non-negativity and ii) column-wise
orthonormality. More mathematical details can be referred to the original references[12,80,81] and
Supplementary eMethod 2a.
546
(b): The 9 dimensional neuroimaging endophenotypes of the brain
The nine DNEs captured the neuroanatomical heterogeneity of four brain diseases, including
AD1-2 for AD[62], ASD1-3 for autism spectrum disorder[42], LLD1-2 for late-life depression[71], and
SCZ1-2 for schizophrenia[72]. The underlying AI methodologies involved two different semi-
supervised clustering or representation learning algorithms: Surreal-GAN[82] and HYDRA[83]. Refer
to a review for details of the semi-supervised learning[84], which primarily seeks the so-called "*1-*
to-k" mapping patterns or transformations from reference domains (like healthy controls) to
target domains (such as patients).
Surreal-GAN[82] was used to derive AD1-2[62]. It unravels the intrinsic heterogeneity
associated with diseases through a deep representation learning approach. The methodological
innovation, compared to its precentor Smile-GAN[85], lies in how Surreal-GAN models disease
heterogeneity: it interprets it as a continuous dimensional representation, ensures a consistent
increase in disease severity within each dimension, and permits the simultaneous presence of
multiple dimensions within the same participant without exclusivity. More mathematical details
are presented in **Supplementary eMethod 2b.**
HYDRA[83] was employed to derive the other 7 DNEs. It utilizes a widely adopted
discriminative technique, namely support vector machines (SVM), to establish the "*1-to-k*"
mapping. The model extends multiple linear SVMs to the nonlinear domain by piecing them
together. This approach serves the dual purpose of classification and clustering simultaneously.
Specifically, it creates a convex polytope by amalgamating hyperplanes derived from k linear
SVMs. This polytope separates the healthy control group from the k subpopulations within the
patient group. Conceptually, each face of this convex polytope can be likened to encoding each
subtype (categorical trait) or dimension (continuous trait), capturing distinctive disease effects
(Refer to **Supplementary eMethod 2c**).
571
(c): The 12 biological age gaps of nine human organ systems
The nine multi-organ BAGs (brain, cardiovascular, eye, hepatic, immune, musculoskeletal,
metabolic, pulmonary, and renal) were derived from a previous study[5] that used AI to predict the
chronological age of healthy individuals without chronic medical conditions: AI-predicted age –
chronological age. Using a 20-fold cross-validation procedure, we applied the model for each
organ system, employing a linear support vector machine. Before training each model iteration,
standardization was applied to measures (excluding categorical variables) within the training set.
The model was solved using sequential minimal optimization with a gap tolerance of 0.001. The

support vector regression settings were adjusted for optimization, adhering to established
principles in the field[86].
Alongside the nine organ BAGs, we previously derived three multimodal brain BAGs
(GM, WM, and FC-IDP) using features from gray matter (GM), white matter (WM), and
functional connectivity (FC) in MRI scans[6]. We systematically compared four machine learning
models: SVR, LASSO regression, multilayer perceptron, and a five-layer neural network. We
employed nested cross-validation (CV) and included an independent test dataset[87] for a fair
comparison across different models and MRI modalities. This process involved an outer loop CV
with 100 repeated random splits: 80% for training and validation and 20% for testing. Within the
inner loop, a 10-fold CV was utilized for hyperparameter tuning. Furthermore, we reserved an
independent test dataset, which was kept unseen until the fine-tuning of the machine learning
models[88] (e.g., hyperparameters for SVR) was completed.
592

Method 3: Genetic analyses based on GWAS summary statistics

(a): The genetic architecture of the 2024 MAEs and 525 DEs
Primarily, we used SBayesS[39] to estimate three sets of parameters that fully unveil the genetic
architecture of the 2024 MAEs and 525 DEs. SBayesS is an expanded approach capable of
estimating three essential parameters characterizing the genetic architecture of complex traits
through a Bayesian mixed linear model[89]. This method only requires GWAS summary statistics
of the SNPs and LD information from a reference sample. These parameters include SNP-based
heritability (h_{SNP}^2), polygenicity (π), and the relationship between minor allele frequency (MAF)
and effect size (S). We used the software pre-computed sparse LD correlation matrix derived
from the European ancestry by Zeng et al.[39]. More mathematical details can be found in the
original paper from Zeng et al.[39]. We ran the *gctb* command[89] using the argument *--sbayes S*, and
left all other arguments by default. When applying SBayesS to the 2025 MAEs and 525 DEs
summary data, we found that 18 DEs failed to converge in the MCMC sampling, which may be
due to LD differences between FinnGen and UKBB samples (the latter was used as the LD
reference in SBayesS).
To benchmark different methods used in the field for SNP-based heritability estimates,
we also employed two other methods based on GWAS summary data: *i)* LDSC[28] and *ii)*
SumHer[33]. LDSC relies on the principle that the correlation between SNP effect sizes and
linkage disequilibrium with neighboring SNPs can be used to estimate the proportion of
heritability explained by all SNPs using GWAS summary data. For LDSC, we used the
precomputed LD scores from the 1000 Genomes of European ancestry. All other parameters
were set to default in the software. After merging the GWAS summary statistics, we chose the
1000 Genomes reference panel for fair comparisons between the two studies and ensured that
most SNPs were included in the analyses. For example, for the DE
(RX_PARACETAMOL_NSAID), after merging with the reference panel LD, 1,171,361
remained. For the first MAE (C32_1), 1,092,510 SNPs remained after the same merging
procedure. Furthermore, FinnGen didn't provide the original genotype data; they only shared the
LD information via the LDstore software but did not provide the allele information.
Consequently, we cannot generate in-sample LD scores using the LDSC software. Finally, a
prior investigation[90] showcased the robustness of LDSC concerning the selection of LD
reference panels – multi-ethnic European, Finnish-only, non-Finnish European from 1000

Genomes Phase 3 data, and FINRISK Finnish reference panel – regarding heritability estimates
in four lipid traits within a Finnish population.
For SumHer, we used the BLD-LDAK model , as the software suggested. BLD-LDAK
stands for "Bayesian LD-adjusted Kinship," where LD-adjusted kinship refers to the calculation
of genetic relatedness between individuals using information about the correlation of alleles
between nearby SNPs (linkage disequilibrium). We used the software-provided tagging file,
generated from 2000 white British individuals, as a reference penal suggested by the software for
European ancestry groups. The HapMap3 data ([https://www.broadinstitute.org/medical-and-](https://www.broadinstitute.org/medical-and-population-genetics/hapmap-3)
[population-genetics/hapmap-3](https://www.broadinstitute.org/medical-and-population-genetics/hapmap-3)) merged with the tested GWAS summary SNPs. Similarly, we
ensured sufficient SNPs remained after merging with the reference panel. All other parameters
were set to default. SumHer differs from LDSC in several ways: *i*) it models inflation
multiplicatively, whereas LDSC uses an additive approach; *ii*) it accounts for uneven LD patterns
and incorporates MAF on SNP effect; and *iii*) it utilizes a restricted maximum likelihood
solver rather than regression to estimate the h^2_{SNP}.
638
(b): Genetic correlation: We used three different methods to compute the MAE-DE pairwise
(*N*=2024x525=1,062,600) genetic correlations (r_g): *i*) LDSC[28], *ii*) GNOVA[34], and *iii*) HDL[38].
An earlier study[92] highlighted the significance of selecting an appropriate LD score
reference panel for genetic correlation estimates based on summary statistics. We generated the
same reference panel for LD scores across the three software for a fair comparison. For LDSC,
we used the precomputed LD scores from the 1000 Genomes of European ancestry provided by
the software. All other parameters were set by default. To employ GNOVA, we created the LD
scores utilizing the 1000 Genomes of European ancestry using the *--save-ld* argument within the
gnova.py script. For HDL, we used the provided scripts from HDL to generate the LD scores
using the same 1000 Genomes of European ancestry
(https://github.com/zhenin/HDL/wiki/Build-a-reference-panel).
Through our analysis, we found that the three packages have different levels of model
convergence rates, which is critical for future applications as these open-source packages claim
to advance genetic research. In particular, we found that LDSC (1,062,577/1,062,600) and
GNOVA (1,062,600/1,062,600) converged for most of the tested MAE-DE pairs, whereas HDL
failed a substantial proportion of the analyses, leading to only 59,291 out of the 1,062,600 MAE-
DE pairs (refer to the raised issue: https://github.com/zhenin/HDL/issues/30). Therefore, in **Fig.**
2, we presented common significant results after Bonferroni corrections from the LDSC and
GNOVA, resulting in 133 and 45 significant signals corrected on *i*) the number of MAEs and *ii*)
the number of MAEs and DEs.
659
(c): Two-sample bidirectional Mendelian randomization: We employed a bidirectional, two-
sample Mendelian randomization using the *TwoSampleMR* package[77] to infer the causal
relationships between the 2024 MAEs, 521 DEs from FinnGen, and 4 brain DEs from PGC.
The forward Mendelian randomization examined causality from the 2024 MAEs to the
525 DEs, while the inverse analysis investigated causality from the 525 DEs to the 2024 MAEs.
The *TwoSampleMR* package[77] applied five different Mendelian randomization methods. We
presented the significant findings after the Bonferroni correction using the inverse variance
weighted (IVW) estimator, verifying that the correction remained significant in at least one of
the other four estimators (Egger, weighted median, simple mode, and weighted mode
estimators). For the significant signals, we performed several sensitivity analyses. First, a

heterogeneity test was performed to check for violating the IV assumptions. Horizontal
pleiotropy was estimated to navigate the violation of the IV's exclusivity assumption[93] using a
funnel plot, single-SNP Mendelian randomization approaches, and Mendelian randomization
Egger estimator . Moreover, the leave-one-out analysis excluded one instrument (SNP) at a time
and assessed the sensitivity of the results to individual SNP.
Critically, to enhance transparency and reproducibility, we followed a systematic
procedure guided by the STROBE-MR Statement[94] in conducting all causality analyses. This
comprehensive approach encompassed the selection of exposure and outcome variables,
reporting full sets of statistics, and implementing sensitivity checks to identify potential
violations of underlying assumptions. First, we performed an unbiased quality check on the
GWAS summary statistics. Notably, the absence of population overlapping bias[29] was
confirmed, given that FinnGen and UKBB participants largely represent European ancestry
populations without explicit overlap. For the four PGC DEs, we ensured that no UKBB
participants were included in the GWAS summary data. Furthermore, all GWAS summary
statistics were based on or lifted to GRCh37. Subsequently, we selected the effective exposure
variables by assessing the statistical power of the exposure GWAS summary statistics in terms of
instrumental variables (IVs), ensuring that the number of IVs exceeded 8 before harmonizing the
data. Crucially, the function "*clump_data*" was applied to the exposure GWAS data, considering
LD. The function "*harmonise_data*" was then used to harmonize the GWAS summary statistics
of the exposure and outcome variables. This overall resulted in a smaller number (< 525 DEs or
2024 MAEs) of effective exposure/outcome variables in both forward and inverse Mendelian
randomization analyses, as certain GWAS summary data did not have enough IVs.
692
(d): PRS calculation: PRS calculation used the GWAS summary statistics from the split-sample
sensitivity analysis from our previous studies[12,6,4,1]. We established PRS weights using split1
GWAS data as the base/training set, while the split2 GWAS summary statistics were used as the
target/testing data. Details of the quality control (QC) procedures are shown in our previous
studies[12,6,4,1]. Following the QC procedures, PRS for the split2 group was computed using PRS-
CS[45]. PRS-CS infers posterior SNP effect sizes under continuous shrinkage priors using GWAS
summary statistics and an LD reference panel (i.e., UKBB reference). The shrinkage parameter
was not set, and the algorithm learned it via a fully Bayesian approach.
After determining the optimal model, we applied the model to the entire UKBB sample
(~500k individuals). We then performed a PWAS to link the 2024 PRS-MAEs and 59 additional
phenotypes (**Supplementary eTable 5**) not used to compute the PRS-MAE to avoid circular
bias[46]. The 59 phenotypes include cognitive scores (e.g., fluid intelligence score; Field ID:
20016, mental traits (e.g., fed-up feelings; Filed ID: 1960), and lifestyle factors (e.g., tea intake;
Filed ID: 1488). A linear regression was built considering the following covariates: sex (Field
ID: 31), smoking status (Field ID: 20116), weight (Field ID: 21002), standing height (Field ID:
50), waist circumstance (Field ID: 48), age at recruitment (Field ID: 21022), and first 40 genetic
principal components (Field ID: 22009).
710
711

Data Availability

The results of the MUTATE atlas are disseminated at the MUTATE knowledge portal: https://labs-laboratory.com/mutate. The GWAS summary statistics for the 2024 MAEs can be accessed publicly through the MEDICINE knowledge portal: https://labs-laboratory.com/medicine and the BRIDGEPORT knowledge portal: https://labs-laboratory.com/bridgeport. The GWAS summary statistics for the 521 DEs from FinnGen are publicly available at: https://finngen.gitbook.io/documentation/v/r9/. The GWAS summary statistics for the 4 DEs from PGC are publicly available at: https://pgc.unc.edu/for-researchers/download-results/. The study used only GWAS summary statistics rather than individual-level data from the UK Biobank. However, the 2024 MAE GWAS data was initially derived from previous studies conducted under Application Numbers 35148 and 60698 from the UK Biobank.

Code Availability

The software and resources used in this study are all publicly available:

- *GCTB*: https://cnsgenomics.com/software/gctb/#Overview, SNP-based heritability, polygenicity, and MAF/effect size ratio
- *LDSC*: https://github.com/bulik/ldsc, SNP-based heritability and genetic correlation
- *SumHer*: https://dougspeed.com/sumher/, SNP-based heritability
- *GNOVA*: https://github.com/xtonyjiang/GNOVA, genetic correlation
- *HDL*: https://github.com/zhenin/HDL, genetic correlation
- *TwoSampleMR*: https://mrcieu.github.io/TwoSampleMR/index.html, Mendelian randomization
- PRS-CS: https://github.com/getian107/PRScs, PRS
- Surreal-GAN: https://github.com/zhijian-yang/SurrealGAN, to derive AD1 and AD2
- HYDRA: https://github.com/anbai106/mlni, to derive LLD1-2, SCZ1-2, ASD1-3, and GM-, WM-, FC-BAG
- sopNMF: https://github.com/anbai106/SOPNMF, to derive the 2003 brain PSCs
- BioAge: https://github.com/yetianmed/BioAge, to derive the 9 multi-organ BAGs

741 **Competing Interests**

742 None

Authors' contributions

Dr. Wen has full access to all the study data and is responsible for its integrity and accuracy.
Study concept and design: W.J
Acquisition, analysis, or interpretation of data: W.J
Drafting of the manuscript: W.J
Critical revision of the manuscript for important intellectual content: All authors
Statistical analysis: W.J

References

References

1.   Wen, J. *et al.* Neuroimaging-AI Endophenotypes of Brain Diseases in the General

Population: Towards a Dimensional System of Vulnerability. 2023.08.16.23294179

Preprint at https://doi.org/10.1101/2023.08.16.23294179 (2023).

2.   Trubetskoy, V. *et al.* Mapping genomic loci implicates genes and synaptic biology in

schizophrenia. *Nature* **604**, 502–508 (2022).

3.   Kendler, K. & Neale, M. Endophenotype: a conceptual analysis. *Mol Psychiatry* **15**, 789–

797 (2010).

4.   Wen, J. *et al.* The Genetic Architecture of Biological Age in Nine Human Organ Systems.

medRxiv 2023.06.08.23291168 (2023) doi:10.1101/2023.06.08.23291168.

5.   Tian, Y. E. *et al.* Heterogeneous aging across multiple organ systems and prediction of

chronic disease and mortality. *Nat Med* 1–11 (2023) doi:10.1038/s41591-023-02296-6.

6.   Wen, J. *et al.* The genetic architecture of multimodal human brain age. *Nat Commun* **15**,

2604 (2024).

7.   Zhao, B. *et al.* Heart-brain connections: Phenotypic and genetic insights from magnetic

resonance images. *Science* **380**, abn6598 (2023).

8.   McCracken, C. *et al.* Multi-organ imaging demonstrates the heart-brain-liver axis in UK

Biobank participants. *Nat Commun* **13**, 7839 (2022).

9.   Nie, C. *et al.* Distinct biological ages of organs and systems identified from a multi-omics

study. *Cell Reports* **38**, 110459 (2022).

10. Liu, Y. *et al.* Genetic architecture of 11 organ traits derived from abdominal MRI using

deep learning. *eLife* **10**, e65554 (2021).

11. Oh, H. S.-H. *et al.* Organ aging signatures in the plasma proteome track health and disease.

Nature **624**, 164–172 (2023).

12. Wen, J. *et al.* Genomic loci influence patterns of structural covariance in the human brain.

Proceedings of the National Academy of Sciences **120**, e2300842120 (2023).

13. Hodson, R. Precision medicine. *Nature* **537**, S49–S49 (2016).

14. Gottesman, I. I. & Gould, T. D. The endophenotype concept in psychiatry: etymology and

strategic intentions. *Am J Psychiatry* **160**, 636–645 (2003).

15. Cannon, T. D. & Keller, M. C. Endophenotypes in the Genetic Analyses of Mental

Disorders. *Annual Review of Clinical Psychology* **2**, 267–290 (2006).

16. Rajpurkar, P., Chen, E., Banerjee, O. & Topol, E. J. AI in health and medicine. *Nat Med* **28**,

31–38 (2022).

17. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data.

Nature **562**, 203–209 (2018).

18. Miller, K. L. *et al.* Multimodal population brain imaging in the UK Biobank prospective

epidemiological study. *Nature Neuroscience* **19**, 1523–1536 (2016).

19. Elliott, L. T. *et al.* Genome-wide association studies of brain imaging phenotypes in UK

Biobank. *Nature* **562**, 210–216 (2018).

20. Sun, B. B. *et al.* Plasma proteomic associations with genetics and health in the UK

Biobank. *Nature* **622**, 329–338 (2023).

21. Dhindsa, R. S. *et al.* Rare variant associations with plasma protein levels in the UK

Biobank. *Nature* **622**, 339–347 (2023).

22. Kurki, M. I. *et al.* FinnGen provides genetic insights from a well-phenotyped isolated

population. *Nature* **613**, 508–518 (2023).

23. O'Donovan, M. C. What have we learned from the Psychiatric Genomics Consortium.

World Psychiatry **14**, 291–293 (2015).

24. Buniello, A. *et al.* The NHGRI-EBI GWAS Catalog of published genome-wide association

studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res* **47**, D1005–D1012

(2019).

25. Elsworth, B. *et al.* The MRC IEU OpenGWAS data infrastructure. 2020.08.10.244293

Preprint at https://doi.org/10.1101/2020.08.10.244293 (2020).

26. Watanabe, K. *et al.* A global overview of pleiotropy and genetic architecture in complex

traits. *Nat Genet* **51**, 1339–1348 (2019).

27. Shen, L. & Thompson, P. M. Brain Imaging Genomics: Integrated Analysis and Machine

Learning. *Proceedings of the IEEE* **108**, 125–162 (2020).

28. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from

polygenicity in genome-wide association studies. *Nat Genet* **47**, 291–295 (2015).

29. Sanderson, E. *et al.* Mendelian randomization. *Nat Rev Methods Primers* **2**, 1–21 (2022).

30. Bowden, J., Davey Smith, G. & Burgess, S. Mendelian randomization with invalid

instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol*

44, 512–525 (2015).

31. Open science. *Nature* **550**, 7–8 (2017).

32. Bulik-Sullivan, B. *et al.* An atlas of genetic correlations across human diseases and traits.

Nat Genet **47**, 1236–1241 (2015).

33. Speed, D. & Balding, D. J. SumHer better estimates the SNP heritability of complex traits

from summary statistics. *Nat Genet* **51**, 277–284 (2019).

34.   Lu, Q. *et al.* A Powerful Approach to Estimating Annotation-Stratified Genetic Covariance

via GWAS Summary Statistics. *The American Journal of Human Genetics* **101**, 939–964

(2017).

35.   Smith, G. D. & Ebrahim, S. 'Mendelian randomization': can genetic epidemiology

contribute to understanding environmental determinants of disease? *Int J Epidemiol* **32**, 1–

22 (2003).

36.   Davey Smith, G. & Hemani, G. Mendelian randomization: genetic anchors for causal

inference in epidemiological studies. *Hum Mol Genet* **23**, R89-98 (2014).

37.   Pierce, B. L. & Burgess, S. Efficient design for Mendelian randomization studies:

subsample and 2-sample instrumental variable estimators. *Am J Epidemiol* **178**, 1177–1184

(2013).

38.   Ning, Z., Pawitan, Y. & Shen, X. High-definition likelihood inference of genetic

correlations across human complex traits. *Nat Genet* **52**, 859–864 (2020).

39.   Zeng, J. *et al.* Widespread signatures of natural selection across human complex traits and

functional genomic categories. *Nat Commun* **12**, 1164 (2021).

40.   Evans, L. M. *et al.* Comparison of methods that use whole genome data to estimate the

heritability and genetic architecture of complex traits. *Nat Genet* **50**, 737–745 (2018).

41.   Ojavee, S. E., Kutalik, Z. & Robinson, M. R. Liability-scale heritability estimation for

biobank studies of low-prevalence disease. *The American Journal of Human Genetics* **109**,

2009–2017 (2022).

42.   Hwang, G. *et al.* Assessment of Neuroanatomical Endophenotypes of Autism Spectrum

Disorder and Association With Characteristics of Individuals With Schizophrenia and the

General Population. *JAMA Psychiatry* (2023) doi:10.1001/jamapsychiatry.2023.0409.

43. Makowski, C. *et al.* Discovery of genomic loci of the human cerebral cortex using

genetically informed brain atlases. 8 (2022).

44. Matoba, N., Love, M. I. & Stein, J. L. Evaluating brain structure traits as endophenotypes

using polygenicity and discoverability. *Human Brain Mapping* **43**, 329–340 (2022).

45. Ge, T., Chen, C.-Y., Ni, Y., Feng, Y.-C. A. & Smoller, J. W. Polygenic prediction via

Bayesian regression and continuous shrinkage priors. *Nat Commun* **10**, 1776 (2019).

46. Kriegeskorte, N., Simmons, W. K., Bellgowan, P. S. F. & Baker, C. I. Circular analysis in

systems neuroscience: the dangers of double dipping. *Nat. Neurosci.* **12**, 535–540 (2009).

47. Sakaue, S. *et al.* A cross-population atlas of genetic associations for 220 human

phenotypes. *Nat Genet* **53**, 1415–1424 (2021).

48. Lee, S. H. *et al.* Genetic relationship between five psychiatric disorders estimated from

genome-wide SNPs. *Nat Genet* **45**, 984–994 (2013).

49. Hunter, D. J. Gene–environment interactions in human diseases. *Nat Rev Genet* **6**, 287–298

(2005).

50. Wen, J. *et al.* Genetic, clinical underpinnings of subtle early brain change along

Alzheimer's dimensions. 2022.09.16.508329 Preprint at

https://doi.org/10.1101/2022.09.16.508329 (2022).

51. THE BRAINSTORM CONSORTIUM *et al.* Analysis of shared heritability in common

disorders of the brain. *Science* **360**, eaap8757 (2018).

52. Smith, S. M. *et al.* An expanded set of genome-wide association studies of brain imaging

phenotypes in UK Biobank. *Nat Neurosci* **24**, 737–745 (2021).

53. Walsh, B. & Lynch, M. *Evolution and Selection of Quantitative Traits*. (Oxford University

Press, 2018).

54.   Plackett, B. A graphical guide to ischaemic heart disease. *Nature* **594**, S3–S3 (2021).

55.   Peisker, F. *et al.* Mapping the cardiac vascular niche in heart failure. *Nat Commun* **13**, 3027

866   (2022).

867   56.   Brundel, B. J. J. M. *et al.* Atrial fibrillation. *Nat Rev Dis Primers* **8**, 1–23 (2022).

868   57.   Barnes, P. J. *et al.* Chronic obstructive pulmonary disease. *Nat Rev Dis Primers* **1**, 1–21

869   (2015).

870   58.   Holgate, S. T. *et al.* Asthma. *Nat Rev Dis Primers* **1**, 1–22 (2015).

871   59.   Ballard, C. *et al.* Alzheimer's disease. *The Lancet* **377**, 1019–1031 (2011).

872   60.   Marshall, M. The hidden links between mental disorders. *Nature* **581**, 19–21 (2020).

873   61.   Eiser, A. R. & Fulop, T. Alzheimer's Disease Is a Multi-Organ Disorder: It May Already

874   Be Preventable. *J Alzheimers Dis* **91**, 1277–1281 (2023).

875   62.   Wen, J. *et al.* Genetic, clinical underpinnings of subtle early brain change along

876   Alzheimer's dimensions. 2022.09.16.508329 Preprint at

877   https://doi.org/10.1101/2022.09.16.508329 (2022).

878   63.   Guthrie, H. *et al.* Safety, Tolerability, and Pharmacokinetics of Crenezumab in Patients

879   with Mild-to-Moderate Alzheimer's Disease Treated with Escalating Doses for up to

880   133 Weeks. *J Alzheimers Dis* **76**, 967–979 (2020).

881   64.   Sevigny, J. *et al.* The antibody aducanumab reduces Aβ plaques in Alzheimer's disease.

882   *Nature* **537**, 50–56 (2016).

883   65.   Congdon, E. E. & Sigurdsson, E. M. Tau-targeting therapies for Alzheimer disease. *Nat Rev*

884   *Neurol* **14**, 399–415 (2018).

885   66.   Jack, C. R. *et al.* Tracking pathophysiological processes in Alzheimer's disease: an updated

886   hypothetical model of dynamic biomarkers. *Lancet Neurol* **12**, 207–216 (2013).

67.   Frisoni, G. B. *et al.* The probabilistic model of Alzheimer disease: the amyloid hypothesis

revised. *Nat Rev Neurosci* **23**, 53–66 (2022).

68.   Tomic, D., Shaw, J. E. & Magliano, D. J. The burden and risks of emerging complications

of diabetes mellitus. *Nat Rev Endocrinol* **18**, 525–539 (2022).

69.   DeFronzo, R. A. *et al.* Type 2 diabetes mellitus. *Nat Rev Dis Primers* **1**, 1–22 (2015).

70.   Yang, Z. *et al.* Gene-SGAN: discovering disease subtypes with imaging and genetic

signatures via multi-view weakly-supervised deep clustering. *Nat Commun* **15**, 354 (2024).

71.   Wen, J. *et al.* Characterizing Heterogeneity in Neuroimaging, Cognition, Clinical

Symptoms, and Genetics Among Patients With Late-Life Depression. *JAMA Psychiatry*

(2022) doi:10.1001/jamapsychiatry.2022.0020.

72.   Chand, G. B. *et al.* Two distinct neuroanatomical subtypes of schizophrenia revealed using

machine learning. *Brain* **143**, 1027–1038 (2020).

73.   Wen, J. *et al.* The Genetic Architecture of Biological Age in Nine Human Organ Systems.

medRxiv 2023.06.08.23291168 (2023) doi:10.1101/2023.06.08.23291168.

74.   Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based

Linkage Analyses. *Am J Hum Genet* **81**, 559–575 (2007).

75.   Jiang, L. *et al.* A resource-efficient tool for mixed model association analysis of large-scale

data. *Nat Genet* **51**, 1749–1755 (2019).

76.   MacArthur, J. A. L. *et al.* Workshop proceedings: GWAS summary statistics standards and

sharing. *Cell Genomics* **1**, (2021).

77.   Hemani, G. *et al.* The MR-Base platform supports systematic causal inference across the

human phenome. *eLife* **7**, e34408 (2018).

78. Mbatchou, J. *et al.* Computationally efficient whole-genome regression for quantitative and

binary traits. *Nat Genet* **53**, 1097–1103 (2021).

79. Wightman, D. P. *et al.* A genome-wide association study with 1,126,563 individuals

identifies new risk loci for Alzheimer's disease. *Nat Genet* **53**, 1276–1282 (2021).

80. Sotiras, A., Resnick, S. M. & Davatzikos, C. Finding imaging patterns of structural

covariance via Non-Negative Matrix Factorization. *NeuroImage* **108**, 1–16 (2015).

81. Zhirong Yang & Oja, E. Linear and Nonlinear Projective Nonnegative Matrix

Factorization. *IEEE Trans. Neural Netw.* **21**, 734–749 (2010).

82. Yang, Z., Wen, J. & Davatzikos, C. Surreal-GAN:Semi-Supervised Representation

Learning via GAN for uncovering heterogeneous disease-related imaging patterns. *ICLR*

(2021).

83. Varol, E., Sotiras, A. & Davatzikos, C. HYDRA: Revealing heterogeneity of imaging and

genetic patterns through a multiple max-margin discriminative analysis framework.

NeuroImage **145**, 346–364 (2017).

84. Wen, J. *et al.* Subtyping Brain Diseases from Imaging Data. in *Machine Learning for Brain*

Disorders (ed. Colliot, O.) 491–510 (Springer US, New York, NY, 2023). doi:10.1007/978-

1-0716-3195-9_16.

85. Yang, Z. *et al.* A deep learning framework identifies dimensional representations of

Alzheimer's Disease from brain structure. *Nat Commun* **12**, 7065 (2021).

86. Chang, C.-C. & Lin, C.-J. LIBSVM: A library for support vector machines. *ACM Trans.*

Intell. Syst. Technol. **2**, 1–27 (2011).

87.   Samper-González, J. *et al.* Reproducible evaluation of classification methods in

Alzheimer's disease: Framework and application to MRI and PET data. *NeuroImage* **183**,

504–521 (2018).

88.   Wen, J. *et al.* Convolutional neural networks for classification of Alzheimer's disease:

Overview and reproducible evaluation. *Medical Image Analysis* **63**, 101694 (2020).

89.   Zeng, J. *et al.* Signatures of negative selection in the genetic architecture of human complex

traits. *Nat Genet* **50**, 746–753 (2018).

90.   Hautakangas, H. LD Score regression for estimating and partitioning heritability of lipid

levels in the Finnish population. (University of Helsinki, Helsinki, 2018).

91.   Speed, D., Holmes, J. & Balding, D. J. Evaluating and improving heritability models using

summary statistics. *Nat Genet* **52**, 458–462 (2020).

92.   Zhang, Y. *et al.* Comparison of methods for estimating genetic correlation between

complex traits using GWAS summary statistics. *Brief Bioinform* **22**, bbaa442 (2021).

93.   Bowden, J. *et al.* A framework for the investigation of pleiotropy in two-sample summary

data Mendelian randomization. *Stat Med* **36**, 1783–1802 (2017).

94.   Skrivankova, V. W. *et al.* Strengthening the Reporting of Observational Studies in

Epidemiology Using Mendelian Randomization: The STROBE-MR Statement. *JAMA* **326**,

1614–1621 (2021).

Acknowledgment

The MULTI consortium (J.W) aims to integrate multi-organ imaging with multi-omics data to advance our understanding of human aging and disease mechanisms. We sincerely thank the UK Biobank (https://www.ukbiobank.ac.uk/), FinnGen (https://www.finngen.fi/en), and PGC (https://pgc.unc.edu/) team for their invaluable contribution to advancing clinical research in our field.