Towards Maps of Disease Progression: Biomedical Large Language Model Latent Spaces For Representing Disease Phenotypes And Pseudotime

Rafael Zamora-Resendiz[1,*,+]**, Ifrah Khuram**[2,+]**, and Silvia Crivelli PhD**[1,]

[1]San Juan Bautista School of Medicine, USA
[2]Lawrence Berkeley National Laboratory, AMCRD, Berkeley, 94720, USA
[*]corresponding author, rzamoraresendiz@lbl.gov
[+]equal contributors to work

ABSTRACT

In this study, we propose a scientific framework to detect capability among biomedical large language models (LLMs) for representing disease phenotypes and temporal progression. We hypothesize that biomedical LLMs trained on next-token prediction produce latent spaces that implicitly capture "disease states" and disease progression, i.e, the transitions over disease states over time. We describe how foundation models may capture and transfer knowledge from explicit pretraining tasks to specific clinical applications. A scoring function based on Kullback-Leibler divergence was developed to measure the "surprise" seen when subsetting admissions along 13 biomedical LLM latent spaces. By detecting implicit ordering of longitudinal data, we aim to understand how these models self-organize clinical information and support tasks such as phenotypic classification and mortality prediction. We test our hypothesis along a case study for obstructive sleep apnea (OSA) in the publicaly available MIMIC-IV dataset, finding ordering of phenotypic clusters and temporality within latent spaces. Our quantative findings suggest that strong correlation between increased compute, conformance with compute-optimal training, and widening contexts promote better implicit ordering of clinical admissions by disease states, explaining 60.3% of the variance in our proposed implicit task. Preliminary qualitative findings suggest LLMs' latent spaces trace patient trajectories through different phenotypic clusters, terminating at end-of-life phenotypes. This approach highlights the potential of biomedical LLMs in modeling disease progression and evaluating clinical hypotheses related to disease pathways and medical interventions. We underscore the need for larger, high-resolution longitudinal datasets to further validate and enhance understanding of the utility of clinical LLMs in modeling patient trajectories.

Key Points

Question:
Do LLMs order clinical data in a sensible way with respect to applications in predictive medicine?

Findings:
Biomedically-trained LLMs show increasing potential in promoting the organization of patient data to reflect disease progression. In a subcohort of OSA patient, maps derived from LLMs' latent representations reveal traceable disease trajectories.

Meaning:
Maps of disease progression offer an explanation to the utility of LLMs in predictive medicine. Following current pretraining conventions in foundational modeling, scientific inquiry into these maps may help anticipate progress in applications of LLMs for healthcare.

Introduction

Computational medicine has made important strides over the past two decades with the digitization and centralization of patient medical data into electronic health records (EHR)[1,2]. Even so, EHR systems are primarily designed, implemented, and operationalized for administrative function. Common structured medical variables including diagnosis codes, laboratory orders and results, procedure codes, and medication prescriptions typically focus on billable procedures. As a result, relevant patient phenotypes for predictive modeling are under-represented or poorly characterized in structured EHR. This leads to substantial work in order to harmonize clinical concepts and biomarkers using both structured EHR[3] and unstructured text. A means of

integrating health status information locked in clinical text with structured representations of medical knowledge is through the application of natural language processing (NLP)[2, 4]. Recently in clinical medicine, large-language models (LLMs) have been adopted to process clinical text for topic categorization[5], medical concept extraction[6, 7], and medical question-answering[8].

While LLMs continue to show aptitude for general NLP applications as model size increase from millions to trillions of trainable parameters and pretraining scales to tens of terabytes of text[9], clinically pretrained LLMs lag behind current state-of-the-art (SOTA) model pretraining on non-medical text. As the field continues to move towards even larger LLMs[10] and infusion with other socio-economic and biomedical modalities including census tract-level data[11], medical imaging[12] and genetics[13], there remain open questions about the representative capabilities of LLMs in capturing the unique nuances of patient health. In particular, challenges remain in identifying how latent spaces map to structured clinical variables like demographics, disease phenotypes, medical interventions and measures of life expectancy. Additionally, there lacks understanding on the effect of model pretraining in building coherent representations of these common determinants of health even with continued work in interpretability[14] and clinical decision support[15] for medical machine learning (ML) systems.

With the increased accessibility of large-scale EHR for A.I. research[2], there exists greater oppurtunity to observe the effects of LLM scaling in large population-wide studies. We hypothesize that LLMs pretrained on biomedical subject matter capture medically coherent representations of clinical text in their latent spaces, facilitating clinically-relevant subsetting of patient records when embedding population-wide longitudinal corpuses. We propose a mixed-method approach to develop an explanatory framework on the effectiveness of contemporary biomedical LLMs in modelling clinical phenotypes and trajectories. A latent space analysis over the publically available MIMIC-IV clinical dataset[16] was performed across 13 biomedical LLMs of different model size and pre-training strategies including BioBERT, BioBART, ClinicalBert, RadBERT, BioMegatron, Gatortron, BioGPT, and Forge-bio[10, 17–23]. The proposed framework first employs a quantative assessment detecting order in LLM latent spaces with respect to sex, age, race, diagnosis, procedure codes, and death statistics. This is proceeded by qualitative assessment of the LLM latent spaces to interpret the coherence between corpus subsets and structured medical features by visualizing admission-level corpus manifolds, identifying medically-relavent subsetting, and manually reviewing clinical discharge reports for evidence of longitudinal ordering among a case study for obstructive sleep apnea (OSA).

We demonstrate application of the framework for investigating a medical hypothesis related to mechanistic drivers of severe disease along a sub-cohort of patients with diagnosed OSA[24]. OSA is a chronic illness causing interrupted breathing during sleep as result of lack of support of the soft palate due to over relaxation of muscles around the airway. Moreover, OSA is linked to a number of severe illnesses; of which, 6 disease comorbid with OSA are investigated in this study. Previous studies hypothesize obstruction during sleep causes coronary blood flow to increase disproportionately with myocardial work increasing coronary artery vascular resistance. This mechanism may be predictive of increased cardiovascular comorbidities in a dose-dependent relation. Understanding the interaction of OSA with these comorbidities is a necessary step in determining the additive risk of OSA in developing severe illnesses[25]. The case study focuses on discharge reports exhibiting severe OSA and concurrent HF and finds admissions indexed by probability of heart failure diagnosis along LLM latent spaces increases the odd-ratio of mentions of symptoms referencing dyspnea and misadherence of continuous positive airway pressure (CPAP) treatment, surrogates for high doses of respiratory obstruction.

Methods

The proposed framework follows an explantory sequential design splitting analysis of biomedical LLMs' latent spaces into quantitative then qualitative assesment. Quantitative assesment can be summarized as follows:

1. Structured admission-level variables V are mapped to unstructured discharge reports C.

2. For candidate biomedical LLMs $m \in M$, discharge reports C are tokenized then embedded forming corpus manifold E per model m.

3. Per corpus manifold, admissions are clustered into k subsets using K-Means clustering.

4. A Kullback-Liebler Divergence (KLD)-based scoring function is used to measure the amount of information lost per subset when assuming a distribution of variables V expected by random subset assignment for admissions in the collection period.

5. A linear model is used to fit scores for each subsetted latent space against the estimated compute needed for pretraining, the token per parameter ratio between their pretraining data and model size, and the context width of the LLMs.

6. Linear models are used to fit scores against performance on phenotype and mortality classification tasks using cluster distances as input to logistic regression classifiers.

After the subsetted latent spaces are scored, the top performing LLM is selected. To investigate applications of biomedical LLM latent spaces for encoding phenotypes and time-dependant health status variables (referred to as "pseudotime" from here on) as well as representing medical trajectories, the top performing LLM is assessed qualitatively for these properties as follows:

1. UMAP[26] dimensionality reduction is used to plot a 2-dimensional projection of the corpus manifold E. Projections are plotted then shaded using color gradients corresponding to a variable of interest v.

2. Regions of high organization are observed, shading cluster membership by proximity of clusters' centroids in the original high-dimensional space and the informational content over subsets.

3. Temporal ordering is observed, shading projections by time-dependent variables including age at time of admission and time till death at discharge.

4. Phenotypic ordering is observed, shading projections by the prevalence of 6 OSA-related comorbid diseases.

5. Along OSA patients, admissions are plotted then shaded by class probability of comorbid diagnosis and 1-year mortality estimated by logistic regression.

6. Patient admissons are then traced along the corpus manifold and their corresponding discharge reports are reviewed.

Data

The analysis utilized the publicly available MIMIC-IV Clinical Database[27] which includes admissions to the intensive care units (ICU) at Boston's Beth Israel Deaconess Medical Center from 2001 to 2019. EHR for 331,794 admissions, each with a recorded discharge report, were gathered, covering 1.6 million patient-days for a population of 145,915 patients. The unstructured reports were linked to structured medical data including admission-level demographics, billable procedures, billable diagnoses, and mortality outcomes (MIMIC-IV censors time of death past one year after the last admission). The distribution of patient-level demographics is reported in Table 1.

It is important to note that MIMIC-IV contains additional sources of unstructured data, such as radiology reports, but these were not considered in this analysis. The complete corpus of discharge reports totals 3.3 GB of text. Phenotypes for admissions were labeled for the presence of obstructive sleep apnea (OSA), heart failure (HF), stroke (STRK), atrial fibrillation (AFIB), coronary artery disease (CAD), type 2 diabetes mellitus (T2DB), and hypertension (HTN) according to the International Classification of Diseases, 9th and 10th Revision (ICD-9/10). Expert-curated ICD definitions for OSA and comorbid diagnoses are found in Table 2, and the rates of these conditions in the MIMIC-IV patient population are also reported in Table 1.

Large Language Models

LLMs are attention-based deep learning architectures designed to predict the next token or missing tokens in sequentially ordered data[28]. An advantage of transformer-based language models over recurrent architectures like RNNs[29] is the absence of recursive operations for computing next-token probability scores. Instead, transformers utilize parallelizable feed-forward operations. The computational graph for LLM implementations can be naively summarized as stacked blocks of multi-head self-attention followed by feed-forward multilayer perceptrons (MLP) with residual connections between inputs and outputs for both operations.

In essence, multi-head self-attention aggregates information from different sections of a sequence and the MLP projects this aggregated information into a new feature space. Adding more blocks or increasing the number of parameters per block enhances the language model's ability to condition next-token prediction. However, scaling these models requires substantial computational resources and a variety of optimization techniques to manage numerical instability during pretraining[30]. Models of this size heavily rely on data-parallel, model-parallel, and pipeline-parallel training, given that a single GPU cannot accommodate all model parameters or large batches of input sequences in onboard memory[31].

LLMs have been employed previously in biomedical domains. Many LLMs specialized on biomedical domains are trained on publically available datasets such as abstracts and text from scientific literature as well as deidentified EHR datasets like MIMIC-III[32]. Due to limitiations in availabity of public de-identified clinical datasets, many biomedical LLMs are pretrained on the same clinical corpuses and biomedical literature. For example, BioBART, BioMegatron, BioBERT, BioGPT, and Gatortron all contain abstracts and/or full-text from PubMed[33] in their pretraining dataset, and ClinicalBERT and Gatortron have both seen MIMIC-III. While also trained on scientific abstracts, Forge-bio is notable for being trained on a wider set of abstracts sourced from 6 scientific literature databases. Additionally, the Gatortron-line of models and RadBERT have been fit with in-house clinical data from UF Health Integrated Data Repository (IDR) and radiology reports from Veterans Health Administration (VHA) admissions, respectively.

To make comparisons between biomedical LLM architectures, this study gathered 3 characteristics of their pretraining methodolgy including: 1) an estimate of the necessary compute for pretraining (6 floating-point operations per parameter per token) by conventions developed by Yin et al.[31], 2) the ratio between pretraining tokens and model parameter count determining compute optimal training as presented by Hoffman et al.[34], 3) the context width of the model, i.e., the number of input tokens per sequence allowed by an LLM architecture. These characteristics are estimated to the best of our ability from papers and supplementary material, but it is important to note that LLM literature is often opaque with the exact description of pretraining methods, and pretraining conventions have been rapidly changing within recent years. Table 3 reports these characteristics for models covered in the analysis.

Embedding An Admission-Level Corpus

We refer to embedding a complete set of documents C as constructing a corpus manifold E. Embeddings for all 331,794 discharge reports in MIMIC-IV were computed using a set of biomedical LLMs M. This process involved tokenizing the text and then sampling layer activations for each model m.

Without prior knowledge of which specific layer holds the best-compressed representation of the clinical documents, activations were sampled from the last L layers per each model m. The token-wise MLP activations at the end of a transformer block were collected and averaged into a single vector. If the context width for model m was shorter than a full discharge report, the token sequence was chunked into pieces, each of size equal to or less than the maximum context width, without overlap. Considering C as an admission-level corpus of the Boston's Beth Israel Deaconess Medical Center, the transformation $m : C \rightarrow \mathbb{R}^d$ assigns an admission-level feature vector to each report, where d is the number of token-wise features in model m. The construction of an admission-level vector per report can be summarized as

$$e_m^l(c) = \frac{1}{T} \sum_{t=0}^{T} m_t^l(c)$$

where l is the layer being sampled, t is the token index along the sequence, and $c \in C$. The collection of admission-level vectors $e_m^l(c)$ form one corpus manifold E_m^l. Using an allocation of 290.8 node hours on NERSC's Perlmutter supercomputer, we sampled corpus manifolds from the last 3 layers of the 13 models covered in Table 3. Data parallel inference was implemented to accelerate construction of manifolds using 32 Nvidia A100 GPUs. These corpus manifolds were used to ascertain the concentration of medically relevant variables along subregions of the LLMs' latent space as described in the following sections.

Subsetting Latent Spaces Using K-Means

Analysis of healthcare data often requires the subdivision of patient records into smaller, domain-specific cohorts to focus on particular phenotypes or to establish suitable control groups for case-control studies. The exponential increase in records within EHR databases necessitates scalable techniques for grouping patients based on exposures to environmental factors, occurrences of medical events, and longitudinal outcomes[35].

To prepare the latent spaces for evaluation, corpus manifolds were initially partitioned into subregions to assign admission-level representations to different groups. Without prior knowledge of the underlying number of potential subgroups present in the clinical dataset, K-Means clustering[36] was utilized with varied number of expected clusters k. Due to constraints in available computational resources, corpus manifolds were subsetted into 256, 512, 1024, and 2048 clusters. For each corpus manifold E_m^l, admission-level vectors were regularized using max-norm regularization[37] then partitioned into k sets $S = \{S_1, S_2, ..., S_k\}$. For a given set S_i, the cluster's centroid along the corpus manifold is described as

$$\mu_i = \frac{1}{|S_i|} \sum_{e_m^l(c) \in S_i} e_m^l(c)$$

where $|S_i|$ represents the size of the cluster. In addition to assigning each admission to a distinct cluster, the distance relations to each cluster's centroid were computed to be later used as input for classification models described in the subsequent sections. After centroids are identified with K-Means, vectors on corpus manifold E_m^l are mapped to cluster-space $F_m^l \in \mathbb{R}^k$, where each dimension in the cluster-space vector corresponds to the regularized distance from the admission-level vector to each cluster centroid found by K-Means.

Scoring Implicit Ordering in Clinically-Aware Foundation Models

Often referred to as "foundation" models, the capability of LLMs to transfer features learned during pretraining to problems not explicitly defined in their original training policy extends the concept of transfer learning in ML[38]. The two stages of LLM

transfer learning, termed "pretraining" and "finetuning," involve initially training a model on the more general task of token prediction, followed by retraining on a more specific problem with less training data. While transfer learning predates the introduction of LLMs, these models show promise as the first iteration of foundation models due to the apparently tangential yet simple-to-prepare task of pretraining on next-token prediction compared to the domain-specific problems tackled by NLP applications.

Considering auto-regressive architectures exclusively at this stage, the explicit task of language modeling can be expressed using coventions in this paper as follows:

$$m(U) = \sum_{t=0}^{T} log(\mathbb{P}(u_t | u_{t-i}, ..., u_{t-1}, \Theta))$$

where $m(U)$ is the model loss over the full corpus of pretraining text and $\mathbb{P}(u_t | u_{t-i}, ..., u_{t-1}, \Theta)$ is the conditional probability of the token t given i preceding tokens and model parameters Θ[39]. Often referred to as the perplexity of the LLM along the pretraining dataset, its difficult to gather these measures for analysis without access to those datasets used to pretrain each model.

Currently, the field of foundational modeling is considered relatively young, and methods for evaluating foundational capabilities vary widely between projects and domains. By "foundational", we imply a common converging cornerstone representation for a specific domain as opposed to the common catch all "foundation" for LLMs trained at scale. While it is common for LLMs to be evaluated using community NLP benchmarks and human-grade standardized tests, several concerns arise due to the opacity of large-scale LLM research to outside institutions[40]. Particularly, it is challenging to assess the level of bias in benchmark construction (such as cherry-picking benchmarks) and data leakage in the pretraining dataset (where tests and answers may be present in the dataset)[41]. Additionally, the cost of curating gold-standard NLP benchmarks is expensive in terms of experts and time needed for annotation[42].

As opposed to evaluating model capability along a behavioral framework[40], we propose developing more rigorous scientific frameworks to formulate hypotheses about the underlying order found in LLM latent spaces by generating hypotheses of "intelligent designs" solved implicitly during explicit pretraining on next-token prediction, then detecting for the presence of solutions to these hypothetical tasks empirically. Not to be confused with untestable hypotheses on the orgins of life, we use intelligent design as a concept that offers information about how practices in foundation modeling lead to organization along hidden representations of contemporary models. If intermediate solutions for these implicit tasks exist within an LLM's latent space, high degrees of order along these tasks would justify performance seen on downstream finetuning tasks, interpreting high competency on a finetuning problem as indicative of successful encoding of relevant information used in finetuned models along subregions of the latent space.

In the context of clinical finetuning, we hypothesize biomedical LLMs order admission-level reports into collections capturing distinct clinical phenotypes at different stages of disease. We posit a foundational model for clinical text compresses information seen during pretraining into latent representations encoding "disease states" - subsets of admission-level representations with common phenotype and pseudotime. Transitions between disease states would form trajectories detectable along a corpus of longitudinal medical data. We define a Kullback-Liebler-based[43] scoring function to measure the amount of information content or "surprise" captured along subregions on an admission-level corpus manifold as follows

$$\sum_{v \in V} \sum_{S_0}^{S_i} \frac{|S_i|}{|C|} D_{kl}(P_v \| Q_v)$$

where V is a set of medical variables of interest, $|S_i|$ is the size of admission subsets, $|C|$ is total number of admissions in the cohort, P_v is the probability of observing the medical variable of interest v within a subset, and Q_v is the probability of observing the medical variable of interest v along a subsetting which assumes random assignment for admissions occuring during the collection period of the dataset. We consider the demographic determinants of the patient (sex, race, ethnicity), age at time of admission (binned at every 5 years), weeks till death at time of discharge, ICD diagnoses per admission and Current Procedural Terminology (CPT)-coded services as a set of medical variables V necessary to encode a disease state. For ICD and CPT variables, codes were clustered to reduce each set of codes to 100 degrees of freedom for numerical stability. Used as a scoring function for biomedical LLM latent spaces, the measure indicates the amount of information lost per subset when assuming subsetting contains no discernable discrimination between the local statistics of subsetted admissions and the global statistics of admissions in the dataset.

As a consequence, we posit the proposed scoring function will inform progress towards a "clinically-aware" design. Higher scores indicate greater deviation from random assignment of subsets, suggesting a more structured arrangement of

admission-level representations is present within latent spaces. Higher scores may also suggest that models effectively learn to differentiate between distinct clinical observations localized by time, indicating intermediate solutions towards modelling longitudinal outcomes and serving as a strong foundation for downstream clinical tasks in predictive medicine. Conversely, lower scores may indicate challenges in capturing meaningful structure within subregions of the latent space, highlighting areas of high uncertainty where further model refinement should be done through supplementation with additional clinical obeservations.

Admission-Level Classification Using Clusters Distance

To evaluate the hypothesis posited in the previous section, the performance on admission-level classification tasks was assessed for each corpus manifold. If the hypothesis holds true, a positive relationship would be expected between the score of the corpus manifold on the previously described implicit task and the performance of classifiers on explicit phenotypic and pseudotime tasks.

Cluster-space vectors F_m^l for each corpus manifold were utilized as input to classify admissions from patients with history of OSA. Each admission was labeled by the diagnosis of OSA and its six associated comorbid diseases. Furthermore, these admissions were labeled by outcomes for overall mortality used to predict death 1-year, 90-days, and 30-days after discharge. Logistic regression with Elasticnet regularization[44] was employed, using a combination of L1 and L2 penalties to prevent overfitting and improve model interpretability with respect to which regions of the latent space contribute to classification. Without prior knowledge of the needed complexity for each linear model, the regularization coefficient was set to a value of 1.0 and the L2 penalty term was set to 0.5 consistently across all classifers per corpus manifold.

Logistic regression classifiers were fit on the 9 aforementioned binary classification tasks per corpus manifold. K-fold validation on each classifier was conducted, yielding the mean area under the ROC curve (AUC-ROC) over 16 folds. This approach ensured robust evaluation of classifier performance across different subsets of the dataset as per-classifier heuristic optimization is absent at this time. While more optimal classifiers can be constructed with hyperparameter search for each tasks, measuring the difficulty of finding performant classifiers from intermediate solutions on the corpus manifold is the goal of this technique.

Dimensionality Reduction Using UMAP

Given the high-dimensionality of the admission-level embeddings, we explore employing the dimensionality reduction algorithm UMAP to generate human-readable 2-dimensional plots of corpus manifolds for MIMIC-IV. The UMAP algorithm is known for preserving local structure between points while also maintaining more global structure compared to other dimensionality reduction methods like PCA[45] and t-SNE[46]. This property renders UMAP a valuable technique for interpreting encodings or embeddings of high-dimensional data and has been successfully applied in previous work to visualize biological datasets such as transcriptomes of single-cell data[47].

In the context of visualizing a corpus manifold E_m^l, each admission-level vector is mapped to a coordinate (x, y) on a 2-dimensional plane. Each point in the resulting projection corresponds to a single admission within the healthcare system. Through a grid search over UMAP parameters (such as number of nearest neighbors per high-dimensional point and minimum distance between points in the projection), an adequate projection of the corpus manifolds was attained, neither tightly-packing points nor dispersing points uniformly. However, it is important to note that there currently lacks a standardized method for improving the quality of the projection, often relying heavily on trial and error. Additionally, distances along the projection do not necessarily imply meaningful relations between points, but rather denote neighborhood proximity, as is also observed with t-SNE. The cosine distance between admission-level cluster-space vectors F_m^l were used to fit the UMAP projection.

Chart Review For Cases of OSA Patients With Concurrent HF & Controls

As a case study in using biomedical LLMs to characterize severe disease phenotypes, a chart review of discharge reports for patients with history of OSA was performed to assess if comorbid diagnosis probability along the corpus manifold can be used to index severe illness, and if mentions of severe OSA are enriched among regions of the corpus manifold with high probability of HF diagnosed admissions.

Matched stratified sampling was used in this study. OSA patients with concurrent HF were matched with OSA patients with no history of HF along sex, race, and age at time of their first recorded diagnosis of OSA. After cases and their controls were identified, admissions with similar age at time of admission (within a 5 year window) were gathered and 1 admission per case-control match were randomly sampled. This was done as patients potentially have multiple admissions at varied times over the collection period of the dataset. Its important to note that multiple rounds of chart review were conducted to update exclusion criterias for admissions including: 1) removing non-relevant specialties like 'OBSTETRICS/GYNECOLOGY' and 'ORTHOPAEDICS', and 2) removing 'ELECTIVE' admission types.

74 admissions were considered in the final chart review where reviewers were tasked with annotating evidence for severe OSA (prolonged hypoxia or respiratory obstruction) as a mechanistic driver for HF including mentions of: 1) respiratory

distress such as hypoxia, dyspnea, shortness of breath, or related symptoms, 2) history of diagnosed OSA in the patient's medical history preceding current admission, 3) prescribed OSA treatment, such as CPAP devices or specific pharmacological interventions, and 4) adherence/compliance to OSA treatment. After annotation was complete, the HF class probabilities were merged for analysis to measure the odds-ratio of severe OSA or poorly managed OSA over strata of HF probability.

Results

Table 1 reports the cohort characteristics covered by the MIMIC-IV discharge report corpus. The cohort consists of 145,915 patients, with a close to even distribution of sex at 48.7% male and 51.3% female. The majority of patients (68.6%) identify as White, followed by Black (11.98%); 4.8% of patient identifying as Hispanic. Notably, 18.7% of patients in the cohort are marked as deceased with the majority of deaths occuring between the age 60 and 90. The majority of admissions were categorized as emergency (87.8%) with the remaining considered as elective (12.2%). The average number of patient-days summarized by a single discharge report is 4.92 patient-days.

A total of 14,327 patients were diagnosed with OSA with increased rates of males along sex, increased rates of whites and blacks along race, and no discernable change in hispanic ethnicity when compared to the overall population. While overall mortality drops marginally when compared to non-OSA patients, OSA patients have much higher rates of the 6 comorbid diseases than non-OSA patients. For OSA patients, age at time of admission tends towards an older population and shorter lifespans for those patients with reported mortality outcomes. Subsequent sections are divided into results for quanitative and qualitative assements respectively followed by results from the chart review performed on OSA patients with concurrent HF ([PENDING]).

Quanitative Assesment

Subsetting Clinical Admissions By Phenotype and Pseudotime

First, a linear regression analysis was conducted to investigate the potential positive relationship between the three architecture characteristics gathered for each biomedical LLM and the surprise in observing implicit ordering by disease state, as defined in the previous sections. Considering the best layer and subsetting of the corpus manifold per model, the regression model indicates strong evidence that increased compute, compute-optimal training, and a widening context correlate with order among subsetted latent spaces for our proposed implicit task. In other words, the regression model suggests that 60.3% of the variance on the proposed task can be explained by decisions made at the architecture level.

Upper and lower bounds (95% CI) of compute-optimal ratios and context width help explain the underperformance of certain models. Specifically, models with short context widths (512 tokens) underperform even when approaching compute-optimal training, as seen in the publicly-available Gatortron-line models. Interestingly, plotting the trendline for compute-optimal training at the maximum context-width in this sampling of biomedical LLMs retrodicts the performance of the large variant of BioGPT and Forge-bio. Both reported results for their pretraining (October 2022 and November 2023, respectively) after the preprint publication of Hoffman et al.'s paper (March 2022). Gatortron reported results for pretraining in February 2022, one month before Hoffman et al.'s paper was posted on ArXiv. Results for this regression analysis are plotted in Figure 1a.

Table 4 reports more granular measures over the top 10 subsetted corpus manifolds, with Forge-bio containing the highest amount of surprise overall and across age at time of admission, weeks till death after discharge, ICD diagnosis, and CPT services individually. Along demographic determinants, the fully-trained baseline Gatortron model marginally outperformed Forge-bio. This may be a result of observing clinical data, but there are no other models of comparable size trained on clinical data in this work to confirm that pretraining on clinical data allows for better discrimination of demographic determinants at this time.

Second, two linear regression analyses were conducted on the potential positive relationships between the surprise in observing implicit ordering by our proposed definition of disease state and the phenotypic/pseudotime tasks in the OSA case study. Over all subsetted corpus manifolds sampled from the 13 biomedical LLMs, the regression models indicate moderate to strong evidence that implicit ordering by disease state helps provide adequate intermediate solutions for classifiers on the comorbid disease classification and overall mortality prediction tasks. Performance on overall mortality classification by AUC-ROC decreases as the time window widens from 30-day to 1-year mortality, with only weak evidence for correlation between surprise and mortality prediction 30 days out.

Along phenotypic tasks, there is strong evidence of correlation between surprise and improvements in diagnosing admissions for STRK and OSA. Even so, OSA classification only showed marginal improvements as a function of surprise when compared to STRK. Moderate evidence was found for HF, CAD, AFIB, T2DB, and HTN, with HF classifiers performing highest along AUC-ROC. Given that this sampling of classifers are not heuristically optimized, the AUC-ROC scores reported along these results are not conclusive on the limit of performance for any one classification problem, but we expect increasing the computational budget to accommodate for hyperparameter search will help give better assesment on the limits of performance

per task while still indicating a positive correlation between surprise and discriminatory power. Results for these two regression analyses are plotted in Figures 1b and 1c.

Qualitative Assesment

For the best subsetted corpus manifold (last layer of Forge-bio subsetted into 2048 clusters), Figure 2 plots the UMAP projection of the corpus manifold. First, the subset assignment for each admission is shaded by cluster, with the color gradient indicating proximity between the cluster centroids. While no clear separation is found to indicate independent groups of admissions, the UMAP projection does preserve neighborhood proximity found in the higher-dimensional space with bands of admissions expressing similar concentration of colors on the gradient. The overall structure of the projected corpus manifold resembles a curved Y-shaped band centered at $y = 0.55$. A smaller independent patch can be seen towards the bottom left of plot, centered at $(0.25, 0.20)$, and a small thin Y-shape band at $y = 0.7$. Second, plots are shaded by the per-cluster surprise, with the most order following the core of the large Y-shaped band. One point of contention in interpretation of UMAP plots is the difficulty of discriminating what structures in the projections are artifacts of from the algorithms themselves. Even so, we observe enough preservation of the high-dimensional space to proceed with interpretting results for ordering admissions by phenotype and pseudotime.

Ordering By Pseudotime Variables

Figure 3 is divided into two sections, with the first shading the average age at time of admission per subset. Notably, the projected Forge-bio corpus manifold show regions with concentrations of admissions with patients 35 years old and younger centered at $(0.175, 0.175)$. The small thin Y-shaped band referenced in the previous section also concentrates admissions with relatively younger patients (< 40 years). Concentrations of admissions with patients 65 years and older can be seen along the core of the large Y-shaped band. The second section of the figure shades time till death after discharge by prevalence of 1-year, 90-day, and 30-day mortality per admission subsets. As shading moves towards 30-day mortality, we can see recession of prevalence along the large Y-shape band with the strongest concentration remaining around the region at $[(0.1, 0.2), (0.4, 0.5)]$. Interestingly, the thin section within that region, at the top of the independent patch at the bottom left of the plot, has overrepresentation of deaths occuring during admissions to the ICU. In conjunction, these two views demonstrate the temporal ordering of admissions in MIMIC-IV by age and life expectancy. However, its important to note that MIMIC-IV is temporally sparse along longitudinal data and is prodominently acute emergency admission to the ICU. A larger longitudinal dataset is needed to qualitatively assess if these obeservations persist along wider windows of time.

Ordering By Phenotypic Variables

Figure 4 visualizes the prevalence of the 6 comordid diseases of interest for the OSA case study. Visualizations for prevalence of HTN and T2DB show moderate to high prevalence over the large Y-shaped band, the small thin Y-shape band and the independent patch at the bottom left of the plot. Prevalence recedes over subregions that were previously referenced as having higher concentrations of admissions for patients age 40 years and younger. The prevalence of AFIB and CAD concentrate mainly along the large Y-shaped band, with increased concentration on the top of horizontal portion of the band at $y = 0.6125$. Interestingly, this Forge-bio corpus manifold does show clear separation between regions with high concentrations of HF and STRK. STRK admissions are found in abundance following thin bands at $y = 0.675$ and $y = x + 0.225$. For HF, there is heavy overlap with regions concentrating AFIB and OSA. Clear separation of the STRK phenotype may help in interpreting the sharp increase in classification performance seen during quanitative assesment.

Trajectories Along Corpus Manifolds

Admissions for OSA patients with concurrent HF were traced through the corpus manifold to interpret how regions of the projected manifold correspond to narratives found in the discharge reports. Patients whose trajectories terminate as deaths during an admission were selected. Figure 5 visualizes admissions along the Forge-bio corpus manifold for patients with diagnosed OSA. Plots on the left column shade the class probabilities of HF diagnosis from the logistic regression model trained during quanitative assessment. Plots on the right column shade the class probability for 1-year Mortality. Each row in the figure corresponds to a unique OSA patient where admissions with either OSA or HF diagnosis are represented by a cross. Starting at the admission with the earliest recorded diagnosis of OSA, a numbered sequence is traced between admissions to determine if episodes of HF can be inferred by transitioning from regions with low class probability for HF to high probability regions, and if migration towards areas of high class probability for overall mortality occur as admissions get closer to time of death.

Consider the first patient's traced admission history (subject_id=18080257). We see 3 potential episodes of HF along their path through the corpus manifold summarized in discharge reports as follows:

- **First Episode**: Initially, 6.17 years before death ($hadm = 0$), the patient underwent an uneventful orthopedic surgery with no noted complications. 4.86 years before death ($hadm = 1$), the patient underwent another orthopedic procedure,

tolerated it well, and resumed previously prescribed medications. About two months later ($hadm = 2$), the patient began experiencing intermittent chest pain, which occasionally woke him from sleep and was associated with dyspnea.

- **Second Episode**: Approximately 4.19 years before death ($hadm = 3$), the patient experienced significant right shoulder pain following a proximal humerus fracture repair. Two years later ($hadm = 4$), the patient suffered a heart attack necessitating stenting of a coronary artery, complicated by a groin hematoma. A little over four months after that ($hadm = 5$), the patient presented with a fall, hypoxia, and angina, treated with sublingual nitroglycerin and antibiotics for suspected pneumonia. Another 4 months pass ($hadm = 6$), the patient experienced another cardiac event with mild troponin elevation and significant coronary artery blockages. About 1.53 years before death ($hadm = 7$), the patient was hospitalized for severe rectal bleeding and hypotension, complicated by a non-ST elevation myocardial infarction (NSTEMI) requiring intensive care. Only 23 days later ($hadm = 8$), the patient presented with rectal bleeding and unstable angina, with recurrent chest pain indicative of non-ST elevation myocardial infarction amidst ongoing complex medical management.

- **Third Episode**: Just under 1 year before death ($hadm = 9$), the patient was admitted following a fall, complicated by a small subarachnoid hemorrhage and reversal of INR. Thirty-three days before death ($hadm = 10$), the patient required surgery for a periprosthetic joint infection of the right shoulder, managing postoperative pain and progressing well. Four days later ($hadm = 11$), the patient returned with bleeding from the surgical site, complicating severe cardiac issues including a non-ST elevation myocardial infarction (NSTEMI) and acute HF with pulmonary hypertension and reduced ejection fraction. His condition deteriorated further with a subsequent stroke and ongoing fluid management challenges. Six days before death ($hadm = 12$), the patient experienced a life-threatening rectus sheath hematoma requiring embolization, followed by cardiac arrest with successful resuscitation but subsequent multi-system organ failure and decision for palliative care. Cheif cause of death was recored as congestive HF.

Next, consider the second patient's traced admission history (subject_id=16433543). Only 1 potential episode of HF is seen along their path through the corpus manifold, utimately terminating at a different subregion than the previously described patient.

- **First Episode**: Starting 5.82 years before death ($hadm = 0$), the patient underwent a radical nephrectomy without complications. 7 months before death ($hadm = 1$), the patient presented with acute HF with reduced ejection fraction (HFrEF), managed with diuresis and cardiac catheterization showing non-contributory coronary lesions. The patient was discharged to rehabilitation. Two months later ($hadm = 2$), worsening dyspnea, leading to a diagnosis of HF attributed to hypertension and potential sleep disorder or stimulant medications. Ninety days before death ($hadm = 3$), recurrent falls and declining functional status prompted hospitalization. Sixty-four days before death ($hadm = 4$), the patient was readmitted for acute-on-chronic HF exacerbation. The final hospitalization ($hadm = 5$), ending in death from non-traumatic intracerebral hemorrhage, involved a decision to transition to comfort measures only (CMO) due to irreversible neurologic decline and poor prognosis.

Characterizing Drivers of Severe Illness in Obstructive Sleep Apnea [PENDING]
Discussion

A wealth of clinical predictive models apply ML methods on structured data from EHR to help physicians and decision makers integrate large amounts of medical information. Even with the increased push towards automated triaging[48] over the last years, most operationalized models do not fully utilize unstructured clinical text which often hold rich details of patient health status over time. LLMs are trained at scale on broad data with a lower need of data cleaning and curation. Often termed foundation models, we posit LLMs act as central, unifying storage, an impressive hash system that allows to efficiently organize massive amounts of data into highly compressed respresentations. Once pretrained, they can be finetuned for a variety of transfer-learning tasks using significantly smaller amounts of data. We believe clinical foundation models have the potential to capture a full range of associations within clinical corpuses, distill that information into multifaceted dense representations, and integrate diverse sources/modalities of data in medicine. Basic research into the representative capacities of pretrained LLMs for key health indicators is necessary before wide-deployment of models into clinical practice, and insights into how information is stored in LLMs may lead to more informed strategies for solving domain-specific clinical tasks.

Limitations of Analysis

In conjunction, we believe the three linear regression analyses offer researchers in computational medicine a working theory on how the *complexity of LLMs*, influenced by engineers' architectural decisions, determines their capacity for solving downstream tasks. Such a theory would be invaluable for deciding when to reproduce pretraining on larger datasets, justifying the scale of

pretraining over clinical text of healthcare systems, and identifying which classes of clinical tasks are likely to perform well by finetuning.

Several limitations persist with this analysis. First, the architectural characteristics are our best estimations from literature and supplementary materials. Second, there is a lack of understanding regarding the added explanatory power from considering the composition of the pretraining datasets, although Forge-Bio's performance suggests that a more diverse dataset produces better results. Third, the sparsity of MIMIC-IV in terms of longitudinal data makes it difficult to ascertain the consistency of trajectories in latent spaces. More work is needed to determine at what scale LLMs begin to encode trajectories between subregions of the latent space. Lastly, this analysis does not consider any LLMs pretrained on general-domain corpora, though the interest in applying general-domain LLMs in medicine warrants further investigation.

Theoretical vs Empirical Latent Space Analysis

A continued debate persists regarding the relationship between metrics in embedding spaces, concept localization, and semantic relations. Notably, language embedding serves as a demonstration of how vector representations can infer analogies (e.g., king - man = queen - woman), yet the extent of these properties remains an open problem in deep-learning-based NLP[?]. Several attempts have been made to construct a semantic calculus out of token-level representations[?,?], leveraging learned invariants in an embedding space (e.g., (store (not (gross food))) = restaurant). However, it remains uncertain whether these invariants are globally consistent across an embedded manifold and how model pretraining influences them. The introduction of LLMs further complicates the issue, as they demonstrate that many artifacts found in smaller models disappear simply by increasing the number of trainable parameters and the size of the training corpus.

An alternative explanation of how these systems represent language posits that LLMs function as high-capacity locality-sensitive hashes[49]. In essence, LLM parameters serve as a compression of the pretraining corpus. When predicting the next word in a sentence, these models may execute a complex hierarchical lookup in an address space learned through stochastic gradient descent. While the mechanics behind how these address spaces self-organize remain opaque, an empirical framework like the analysis proposed in this work may serve as a sufficient first step in testing LLMs for clinically relevant representations along this line of reasoning. We believe progress in contemporary artificial intelligence research through its persisting and homogenizing coventions[50] warrant the development of new empirical frameworks towards detecting intelligent designs.

The proposed approach is empirical in nature due to its reliance on real-world data and observable outcomes. By scoring the underlying organization of LLM latent spaces with respect to observational data, researchers can empirically evaluate the model's ability to encode clinical phenotypic markers and mortality-related pseudotime. This stands in contrast to theoretical approaches, which may rely more heavily on abstract formulations or theoretical assumptions about the structure of clinical language. Through empirical analysis, researchers can gain practical insights into the capabilities and limitations of LLMs in capturing clinically relevant information within their latent spaces and identify domain-specific areas that need additional refinement.

Scaling Framework to High-Resolution Population-Wide Longitudinal Datasets

Our ultimate goal is to assess the utility of pretrained latent representations for modeling longitudinal trajectories of patients. We demonstrate how measuring excess self-information over subregions of the latent space can effectively detect systematic increase or decrease in organization concerning phenotypic variables such as diagnosis and healthcare services, as well as time-dependent variables like patients' age and life expectancy. However, the analysis over MIMIC-IV provides weak evidence for trajectories due to the sparsity of admissions per patient. We anticipate observing stronger temporal ordering in a larger population study with higher rates of healthcare utilization.

A positive finding would reveal corpus manifolds expressing *attractors*, where patients' embedded clinical histories trace paths through different phenotypic clusters, ultimately terminating at end-of-life phenotypic clusters (e.g., Major depressive disorder -> alcohol abuse -> death by liver failure). If successful, this analysis will serve as a valuable stepping stone toward understanding how clinical LLMs organize information, model disease progression, and how they may be used to evaluate medical hypotheses related to disease progression pathways and the effect of medical interventions on those pathways.

We further hypothesize that the translation of general features learned by clinical LLMs during pretraining to fine-tuned tasks is directly related to the quality of these clinical latent spaces in modeling phenotypes and temporal locality. Specifically, we propose that this implicit task *improves proportionally with the total number and temporal resolution of patient trajectories observed during pretraining*.

However, significant roadblocks exist in confirming the emergence of trajectories along longitudinal datasets, particularly in scaling to populations past tens of millions of patients. To contextualize the engineering feat required, consider the U.S. Department of Veterans Affairs (VA), which houses one of the largest centralized repositories of clinical data in their Corporate Data Warehouse (CDW) . As of 2023, the CDW has archived clinical documents for 4.18 B admissions from nationwide inpatient and outpatient admissions for over 13.8 million patients spanning from January 1, 2000, to January 1, 2022. The urgent need for scalable data analysis is underscored by the VA's estimated growth rate for their clinical corpus, projected to

increase by X00 million documents per year. Even so, the high longitudinal resolution of the VA dataset makes it an important dataset for foundational modeling with the an average of 131.9 admissions per patient and over 4.98 M patient years worth of obeservational data (3 orders of magnitude greater than the 4.47 K patient years covered in MIMIC-IV).

Maps Of Disease Progression

As advancements in computational medicine and artificial intelligence continue to unfold, we anticipate the detection of disease progression will be enhanced through insights derived from large-scale longitudinal datasets and applied foundational modeling, ranging from identifying predisposition for chronic conditions to precipitation of acute illnesses. Mapping disease progression would revolutionize public health strategies enabling policymakers and healthcare professionals to implement targeted interventions and preventive measures tailored to the specific stages of disease development. In identifying early indicators and risk factors associated with disease progression, public health initiatives could be optimized to intervene at critical junctures, thereby reducing the burden of illness on nationwide healthcare systems and improving population health outcomes.

Given sufficient healthcare data, these maps may help understand the rapid collapse of health among frail or high comorbid patients[51]. Additionally, maps may help disentangle the complex interplay between genetic predispositions, environmental factors, and medical interventions. Clinicians can gain insights into the dynamics of disease, allowing for early detection, precise risk stratification, and targeted interventions. Moreover, these maps could serve as invaluable tools for predicting disease trajectories, optimizing treatment strategies, and identifying novel therapeutic targets.

However, with these newfound repositories of clinical knowledge comes ethical considerations and societal responsibilities. Its imperative to ensure accuracy and reduced bias on downstream finetuning applications using foundation models, equitable access to healthcare resources and interventions which employ these systems, and guards against the misuse of personal health data. As tools for understanding the dynamics of disease, maps of disease progression may possess capabilities that extend beyond traditional healthcare applications. The ability to predict disease trajectories, identify vulnerable populations, and anticipate the spread of illnesses may hold implications for shaping societal well-being in positive or negative ways. Methods of ascertaining clinical capability as a function model complexity need to be at the forefront of applied foundational modeling in computational medicine.

References

1. Mesko, B. The role of artificial intelligence in precision medicine (2017).

2. Qiu, J. *et al.* Large ai models in health informatics: Applications, challenges, and the future. *arXiv preprint arXiv:2303.11568* (2023).

3. Hong, C. *et al.* Clinical knowledge extraction via sparse embedding regression (keser) with multi-center large scale electronic health record data. *NPJ digital medicine* **4**, 151 (2021).

4. Koleck, T. A. *et al.* Identifying symptom information in clinical notes using natural language processing. *Nurs. research* **70**, 173 (2021).

5. Liu, F., Shareghi, E., Meng, Z., Basaldella, M. & Collier, N. Self-alignment pretraining for biomedical entity representations. *arXiv preprint arXiv:2010.11784* (2020).

6. Agrawal, M., Hegselmann, S., Lang, H., Kim, Y. & Sontag, D. Large language models are few-shot clinical information extractors. In *Proceedings of the 2022 Conference on Empirical Methods in Natural Language Processing*, 1998–2022 (2022).

7. Alawad, M., Hasan, S. S., Christian, J. B. & Tourassi, G. Retrofitting word embeddings with the umls metathesaurus for clinical information extraction. In *2018 IEEE International Conference on Big Data (Big Data)*, 2838–2846 (IEEE, 2018).

8. Uzuner, Ö., South, B. R., Shen, S. & DuVall, S. L. 2010 i2b2/va challenge on concepts, assertions, and relations in clinical text. *J. Am. Med. Informatics Assoc.* **18**, 552–556 (2011).

9. Nori, H., King, N., McKinney, S. M., Carignan, D. & Horvitz, E. Capabilities of gpt-4 on medical challenge problems (2023). 2303.13375.

10. Yang, X. *et al.* A large language model for electronic health records. *NPJ Digit. Medicine* **5**, 194 (2022).

11. Wang, X. *et al.* An examination of the association between altitude and suicide deaths, suicide attempts, and suicidal ideation among veterans at both the patient and geospatial level. *J. psychiatric research* **153**, 276–283 (2022).

12. Pelka, O., Koitka, S., Rückert, J., Nensa, F. & Friedrich, C. M. Radiology objects in context (roco): a multimodal image dataset. In *Intravascular Imaging and Computer Assisted Stenting and Large-Scale Annotation of Biomedical Data and Expert Label Synthesis: 7th Joint International Workshop, CVII-STENT 2018 and Third International Workshop, LABELS 2018, Held in Conjunction with MICCAI 2018, Granada, Spain, September 16, 2018, Proceedings 3*, 180–189 (Springer, 2018).

13. Kimbrel, N. A. *et al.* Identification of novel, replicable genetic risk loci for suicidal thoughts and behaviors among us military veterans. *JAMA psychiatry* **80**, 135–145 (2023).

14. Martinez, K. M. *et al.* Factors influencing accuracy, interpretability and reproducibility in the use of machine learning in biology. (2024).

15. Dhaubhadel, S. *et al.* High dimensional predictions of suicide risk in 4.2 million us veterans using ensemble transfer learning. *Sci. reports* **14**, 1793 (2024).

16. Johnson, A. *et al.* MIMIC-IV (2020).

17. Lee, J. *et al.* Biobert: a pre-trained biomedical language representation model for biomedical text mining. *Bioinformatics* **36**, 1234–1240 (2020).

18. Yuan, H. *et al.* Biobart: Pretraining and evaluation of a biomedical generative language model. *arXiv preprint arXiv:2204.03905* (2022).

19. Alsentzer, E. *et al.* Publicly available clinical bert embeddings. *arXiv preprint arXiv:1904.03323* (2019).

20. Yan, A. *et al.* Radbert: Adapting transformer-based language models to radiology. *Radiol. Artif. Intell.* **4**, e210258 (2022).

21. Shin, H.-C. *et al.* Biomegatron: Larger biomedical domain language model. *arXiv preprint arXiv:2010.06060* (2020).

22. Luo, R. *et al.* Biogpt: generative pre-trained transformer for biomedical text generation and mining. *Briefings Bioinforma.* **23**, bbac409 (2022).

23. Yin, J., Dash, S., Wang, F. & Shankar, M. Forge: Pre-training open foundation models for science. In *SC23: International Conference for High Performance Computing, Networking, Storage and Analysis*, DOI: 10.1145/3581784.3613215 (2023).

24. Wang, X. *et al.* Obstructive sleep apnea and risk of cardiovascular disease and all-cause mortality: a meta-analysis of prospective cohort studies. *Int. journal cardiology* **169**, 207–214 (2013).

25. Lin, P., Li, X., Zhang, J. & Liang, Z. Association between obstructive sleep apnea and reduced mortality in critically ill patients: A propensity score-based analysis. *Int. J. Gen. Medicine* 4723–4729 (2021).

26. McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and projection for dimension reduction. *arXiv preprint arXiv:1802.03426* (2018).

27. Johnson, A. *et al.* Mimic-iv. *PhysioNet. Available online at: https://physionet. org/content/mimiciv/1.0/(accessed August 23, 2021)* 49–55 (2020).

28. Vaswani, A. *et al.* Attention is all you need. *Adv. neural information processing systems* **30** (2017).

29. Pascanu, R., Gulcehre, C., Cho, K. & Bengio, Y. How to construct deep recurrent neural networks. *arXiv preprint arXiv:1312.6026* (2013).

30. Smith, S. *et al.* Using deepspeed and megatron to train megatron-turing nlg 530b, a large-scale generative language model. *arXiv preprint arXiv:2201.11990* (2022).

31. Yin, J., Dash, S., Gounley, J., Wang, F. & Tourassi, G. Evaluation of pre-training large language models on leadership-class supercomputers. *The J. Supercomput.* **79**, 20747–20768 (2023).

32. Johnson, A. E. *et al.* Mimic-iii, a freely accessible critical care database. *Sci. data* **3**, 1–9 (2016).

33. Canese, K. & Weis, S. Pubmed: the bibliographic database. *The NCBI handbook* **2** (2013).

34. Hoffmann, J. *et al.* Training compute-optimal large language models. *arXiv preprint arXiv:2203.15556* (2022).

35. Mitra, A. *et al.* Associations between natural language processing–enriched social determinants of health and suicide death among us veterans. *JAMA network open* **6**, e233079–e233079 (2023).

36. Dong, Y., Indyk, P., Razenshteyn, I. & Wagner, T. Learning space partitions for nearest neighbor search. *arXiv preprint arXiv:1901.08544* (2019).

37. Lee, J. D., Recht, B., Srebro, N., Tropp, J. & Salakhutdinov, R. R. Practical large-scale optimization for max-norm regularization. *Adv. neural information processing systems* **23** (2010).

38. Bommasani, R. *et al.* On the opportunities and risks of foundation models. *arXiv preprint arXiv:2108.07258* (2021).

39. Li, Y., Huang, Y., Ildiz, M. E., Rawat, A. S. & Oymak, S. Mechanics of next token prediction with self-attention. In *International Conference on Artificial Intelligence and Statistics*, 685–693 (PMLR, 2024).

40. McIntosh, T. R., Susnjak, T., Liu, T., Watters, P. & Halgamuge, M. N. Inadequacies of large language model benchmarks in the era of generative artificial intelligence. *arXiv preprint arXiv:2402.09880* (2024).

41. Sainz, O. *et al.* Nlp evaluation in trouble: On the need to measure llm data contamination for each benchmark. *arXiv preprint arXiv:2310.18018* (2023).

42. Canales, L. *et al.* Assessing the performance of clinical natural language processing systems: development of an evaluation methodology. *JMIR Med. Informatics* **9**, e20492 (2021).

43. Wei, K., Iyer, R. & Bilmes, J. Submodularity in data subset selection and active learning. In *International conference on machine learning*, 1954–1963 (PMLR, 2015).

44. Liu, Q., Sun, Y., Wang, C., Liu, T. & Tao, D. Elastic net hypergraph learning for image clustering and semi-supervised classification. *IEEE Transactions on Image Process.* **26**, 452–463 (2016).

45. Kuhnel, L., Fletcher, T., Joshi, S. & Sommer, S. Latent space non-linear statistics. *arXiv preprint arXiv:1805.07632* (2018).

46. Van der Maaten, L. & Hinton, G. Visualizing data using t-sne. *J. machine learning research* **9** (2008).

47. A single-cell transcriptomic atlas characterizes ageing tissues in the mouse. *Nature* **583**, 590–595 (2020).

48. McCarthy, J. F. *et al.* Evaluation of the recovery engagement and coordination for health–veterans enhanced treatment suicide risk modeling clinical program in the veterans health administration. *JAMA network open* **4**, e2129900–e2129900 (2021).

49. Astefanoaei, M., Cesaretti, P., Katsikouli, P., Goswami, M. & Sarkar, R. Multi-resolution sketches and locality sensitive hashing for fast trajectory processing. In *Proceedings of the 26th ACM SIGSPATIAL International Conference on Advances in Geographic Information Systems*, 279–288 (2018).

50. Huh, M., Cheung, B., Wang, T. & Isola, P. The platonic representation hypothesis. *arXiv preprint arXiv:2405.07987* (2024).

51. Justice, A. C. & Tate, J. P. Strengths and limitations of the veterans aging cohort study index as a measure of physiologic frailty. *AIDS Res. Hum. Retroviruses* **35**, 1023–1033 (2019).

Table 1. Cohort Characteristics For MIMIC-IV Admissions with Discharge Report.

		Overall (n=145,915)	Non-OSA (n=131,588)	OSA (n=14,327)
Sex				
	Male	71,016 (48.7%)	62,469 (47.5%)	8,547 (59.7%)
	Female	74,899 (51.3%)	69,119 (52.5%)	5,780 (40.3%)
Race				
	White	100,085 (68.6%)	89,898 (68.3%)	10,187 (71.1%)
	Black	17,475 (12.0%)	15,456 (11.7%)	2,019 (14.1%)
	Other	20,929 (14.3%)	19,317 (14.7%)	1,612 (11.3%)
	Unknown	7,426 (5.1%)	6,917 (5.3%)	509 (3.6%)
Hispanic Ethnicity				
	Yes	7,006 (4.8%)	6,322 (4.8%)	684 (4.8%)
	No	131,483 (90.1%)	118,349 (89.9%)	13,134 (91.7%)
	Unknown	7,426 (5.1%)	6,917 (5.3%)	509 (3.6%)
Age at Death				
	All	27,247 (18.7%)	24,700 (18.8%)	2,547 (17.8%)
	< 20	9 (0.0%)	9 (0.0%)	0 (0.0%)
	20-29	215 (0.1%)	204 (0.2%)	11 (0.1%)
	30-39	415 (0.3%)	379 (0.3%)	36 (0.3%)
	40-49	1,090 (0.7%)	995 (0.8%)	95 (0.7%)
	50-59	2,960 (2.0%)	2,654 (2.0%)	306 (2.1%)
	60-69	5,014 (3.4%)	4,343 (3.3%)	671 (4.7%)
	70-79	6,128 (4.2%)	5,375 (4.1%)	753 (5.3%)
	80-89	7,329 (5.0%)	6,787 (5.2%)	542 (3.8%)
	> 89	4,087 (2.8%)	3,954 (3.0%)	133 (0.9%)
Diagnosis				
	Stroke (STRK)	18,719 (12.8%)	16,471 (12.5%)	2,248 (15.7%)
	Heart Failure (HF)	23,835 (16.3%)	19,519 (14.8%)	4,316 (30.1%)
	Coronary Artery Disease (CAD)	35,134 (24.1%)	29,891 (22.7%)	5,243 (36.6%)
	Atrial Fibrillation (AFIB)	26,048 (17.9%)	22,179 (16.9%)	3,869 (27.0%)
	Type-2 Diabetes Mellitus (T2DB)	32,752 (22.4%)	26,730 (20.3%)	6,022 (42.0%)
	Hypertension (HTN)	81,640 (55.9%)	70,516 (53.6%)	11,124 (77.6%)
Text Corpus (MB)		3,338	2,781	557
Admitted Patient-Days		1,630,924	1,372,168	258,756
Admissions				
	All	331,794	280,229	51,565
	Emergency	291,255 (87.8%)	247,634 (88.4%)	43,621 (84.6%)
	Elective	40,539 (12.2%)	32,595 (11.6%)	7,944 (15.4%)
Admissions By Age				
	< 20	1,410 (0.4%)	1,402 (0.5%)	8 (0.0%)
	20-29	18,821 (5.7%)	18,115 (6.5%)	706 (1.4%)
	30-39	25,170 (7.6%)	22,999 (8.2%)	2,171 (4.2%)
	40-49	35,634 (10.7%)	30,223 (10.8%)	5,411 (10.5%)
	50-59	59,674 (18.0%)	47,971 (17.1%)	11,703 (22.7%)
	60-69	70,549 (21.3%)	55,336 (19.7%)	15,213 (29.5%)
	70-79	59,461 (17.9%)	48,681 (17.4%)	10,780 (20.9%)
	80-89	45,347 (13.7%)	40,556 (14.5%)	4,791 (9.3%)
	> 89	15,728 (4.7%)	14,946 (5.3%)	782 (1.5%)

Table 2. ICD 9/10 Definitions For OSA and Comorbid Diseases.

Obstructive Sleep Apnea (OSA)	Stroke (STRK)	Heart Failure (HF)	Coronary Artery Disease (CAD)	Atrial Fibrillation (AFIB)	Type-2 Diabetes Mellitus (T2DB)	Hypertension (HTN)
ICD-9 327.20, 327.23, 327.29, 780.51, 780.53, 780.57	- 362.3, 362.3[01234], 362.34, 431, 432.9, 433, 433.0, 433.0[01], 433.1, 433.1[01], 433.2, 433.2[01], 433.3, 433.3[01], 433.8, 433.8[01], 433.9, 433.9[01], 434, 434.0, 434.01, 434.1, 434.1[01], 434.9, 434.9[01], 435, 435.0, 435.1, 435.2, 435.3, 435.8, 435.9, 436, 437.0, 437.1, 437.8, 437.89, 437.9, 438, 438.0, 438.1, 438.1[012], 438.13, 438.14, 438.19, 438.2, 438.2[01], 438.21, 438.22, 438.3, 438.3[012], 438.4, 438.4[012], 438.5, 438.5[0123], 438.6, 438.7, 438.8, 438.8[123459], 438.9, V12.54	- 402.01, 402.11, 402.91, 404.0[13], 404.1[13], 404.9[13], 428, 428.[01], 428.2, 428.2[0123], 428.3, 428.3[0123], 428.4, 428.4[0123], 428.9	- 410, 410.0, 410.0[012], 410.1, 410.1[012], 410.2, 410.2[012], 410.3, 410.3[012], 410.4, 410.4[012], 410.5, 410.5[02], 410.6, 410.6[012], 410.7, 410.7[012], 410.8, 410.8[012], 410.9, 410.9[012], 411, 411.0, 411.1, 411.8, 411.8[19], 412, 413, 413.[019], 414.0, 414.0[1234567], 414.3, 414.[489], 429.7, 429.79, V45.8[12], 36, 36.0, 36.0[34679], 36.1, 36.1[012345679], 36.2, 36.3, 36.3[12349]	- 427.3, 427.3[12]	- 250, 250.0, 250.0[02], 250.1, 250.1[02], 250.2, 250.2[02], 250.3, 250.32, 250.4, 250.4[02], 250.5, 250.5[02], 250.6, 250.6[02], 250.7, 250.72, 250.8, 250.8[02], 250.9, 250.9[02], 357.2, 362.0, 362.0[1234567], 366.41	- 362.11, 437.2, 401.[019], 402, 402.0, 402.00, 402.1, 402.11, 402.9, 402.9[01], 403, 403.0, 403.0[01], 403.1, 403.11, 403.9, 403.9[01], 404, 404.0, 404.0[123], 404.1, 404.1[0123], 404.9, 404.9[0123]
ICD-10 G4730, G4733, G4739	- I61, I61.[012345689], I63, I63.0, I63.00, I63.01, I63.01[1239], I63.02, I63.03, I63.03[1239], I63.09, I63.1, I63.10, I63.11, I63.11[1239], I63.12, I63.13, I63.13[1239], I63.19, I63.2, I63.20, I63.21, I63.21[1239], I63.22, I63.23, I63.23[1239], I63.29, I63.3, I63.30, I63.31, I63.31[1239], I63.32, I63.32[1239], I63.33, I63.33[1239], I63.34, I63.34[1239], I63.4, I63.40, I63.41, I63.41[1239], I63.42, I63.42[1239], I63.43, I63.43[1239], I63.44, I63.44[1239], I63.49, I63.5, I63.50, I63.51, I63.51[1239], I63.52, I63.52[1239], I63.53, I63.53[1239], I63.54, I63.54[1239], I63.59, I63.6, I63.8, I63.8[19], I63.9, I65, I65.0, I65.0[1239], I65.1, I65.2, I65.2[1239], I65.8, I65.9, I66, I66.0, I66.01, I66.0[239], I66.1, I66.1[1239], I66.2, I66.2[1239], I66.3, I66.8, I66.9, I67.2, I67.4, I67.8[129], I67.9, I68.0, I68.8, I69, I69.0, I69.00, I69.01, I69.01[01234589], I69.02, I69.02[1238], I69.03, I69.03[12349], I69.04, I69.04[12349], I69.05, I69.05[12349], I69.06, I69.06[123459], I69.09, I69.09[01238], I69.1, I69.10, I69.11[01234589], I69.12, I69.12[01238], I69.13, I69.13[12349], I69.14, I69.14[12349], I69.15, I69.15[1249], I69.16, I69.16[123459], I69.19, I69.19[01238], I69.2, I69.20, I69.21, I69.21[01234589], I69.22, I69.22[01238], I69.23, I69.23[12349], I69.24, I69.24[12349], I69.25, I69.25[12349], I69.26, I69.26[123459], I69.29, I69.290, I69.29[1238], I69.3, I69.30, I69.31, I69.31[0134589], I69.32, I69.32[01238], I69.33, I69.33[1349], I69.34, I69.34[12349], I69.35, I69.35[12349], I69.36, I69.36[123459], I69.39, I69.39[1238], I69.8, I69.80, I69.81, I69.81[01234589], I69.82, I69.82[01238], I69.83, I69.83[12349], I69.84, I69.84[12349], I69.85, I69.85[12349], I69.86, I69.86[123459], I69.89, I69.89[01238], I69.9, I69.90, I69.91, I69.91[01234589], I69.92, I69.92[01238], I69.93, I69.93[12349], I69.94, I69.94[12349], I69.95, I69.95[12349], I69.96, I69.96[123459], I69.99, I69.99[01238] G45, G45.[01238], G46.[34], H34, H34.0, H34.0[0123], H34.1, H34.1[012], H34.1[23], H34.2, H34.21, H34.21[1239], H34.23, H34.23[1239]	- I11.0, I13.[02], R57.0, I50, I50.1, I50.2, I50.2[0123], I50.3, I50.3[0123], I50.4, I50.4[0123], I50.8, I50.81, I50.81[01234], I50.8[2349], I50.9	- Z95.1, Z95.5, Z98.61, I20, I20.[0129], I21, I21.0, I21.0[129], I21.1, I21.1[19], I21.2, I21.2[19], I21.[349], I21.A, I21.A[19], I22, I22.[01289], I23, I23.[012345678], I24, I24.[0189], I25, I25.1, I25.10, I25.11, I25.11[01289], I25.[2567], I25.70, I25.70[01289], I25.71, I25.71[01289], I25.72, I25.72[01289], I25.73, I25.73[01289], I25.75, I25.75[01289], I25.76, I25.76[01289], I25.79, I25.79[01289], I25.8, I25.81[012], I25.8[2349], I25.9	- I48, I48.0, I48.1, I48.1[19], I48.2, I48.21, I48.3, I48.4, I48.9, I48.9[12]	- E11, E11.0, E11.01, E11.11, E11.2, E11.2[129], E11.3, E11.31, E11.31[19], E11.32, E11.321, E11.321[1239], E11.329, E11.329[1239], E11.33, E11.331, E11.331[239], E11.339, E11.339[1239], E11.34, E11.341, E11.341[1239], E11.349, E11.349[1239], E11.35, E11.351, E11.351[1239], E11.352, E11.352[239], E11.353, E11.353[1239], E11.354, E11.354[1239], E11.355, E11.355[1239], E11.36, E11.37, E11.37X[1239], E11.39, E11.4, E11.4[012349], E11.5, E11.5[129], E11.6, E11.61, E11.61[08], E11.62, E11.62[0128], E11.63, E11.63[08], E11.64, E11.64[19], E11.6[59], E11.[89]	- H35.03[1239], I10, I11, I11.[09], I12, I12.[09], I13, I13.0, I13.1, I13.1[01], I13.2, I16.[019], I67.4

Table 3. Pretraining Characteristics Across Biomedical Large Language Models

	model	parameters	tokens	context width	seen MIMIC	reported datasets
0	RadBERT_2m	109.5 M	466 M	512	No	VHA Radiology Reports
1	Bio_ClinicalBERT	108.3 M	666 M	512	Yes	MIMIC-III
2	BioBART_base	166.4 M	6 B	1024	No	PubMed Abstracts
3	BioBART_large	442.3 M	6 B	1024	No	PubMed Abstracts
4	BioMegatron_base	333.6 M	8.1 B	512	No	PubMed Abstracts, PMC Full Text
5	BioBERT_base	108.3 M	28.4 B	512	No	PubMed Abstracts, PMC Full Text, BooksCorpus US, English Wikipedia
6	BioGPT_base	346.8 M	14 B	1024	No	PubMed Abstracts, PMC Full Text
7	BioBERT_large	364.4 M	28.4 B	512	No	PubMed Abstracts, PMC Full Text, Books Corpus US, English Wikipedia
8	BioGPT_large	1.571 B	14 B	2048	No	PubMed Abstracts, PMC Full Text
9	Gatortron_base	355.3 M	121.33 B	512	Yes	UF Health IRD, MIMIC-III, PubMed Abstracts, PMC Full Text, English Wikipedia
10	Gatortron_s	355.3 M	121.33 B	512	Yes	UF Health IRD, MIMIC-III, PubMed Abstracts, PMC Full Text, English Wikipedia
11	Forge_bio	1.440 B	38 B	2048	No	SciBert Classified Biology/Medicine (Bio/Med) Abstracts From CORE, OAG, MAG, Aminer, ArXiv, and SCOPUS
12	Gatortron_medium	3.913 B	121.33 B	512	Yes	UF Health IRD, MIMIC-III, PubMed Abstracts, PMC Full Text, English Wikipedia

Table 4. Top Subsetted Latent Spaces by Proposed KLD-based Scoring Function

	model	layer	k	sum	age at admission	weeks till death	demographic determinants	dx phenotype	cpt phenotype
0	Forge_bio	0	2048	**9.26**	**0.19**	**0.17**	0.10	**2.07**	**6.73**
1	BioGPT_large	2	2048	8.72	0.18	0.16	0.09	2.00	6.29
2	BioGPT_large	1	2048	8.54	0.18	0.15	0.09	1.94	6.18
3	BioGPT_large	0	2048	8.37	0.17	0.15	0.09	1.91	6.06
4	Gatortron_base	0	2048	8.35	0.17	0.14	**0.11**	1.87	6.06
5	Gatortron_base	1	2048	8.30	0.17	0.13	0.09	1.89	6.01
6	Forge_bio	0	1024	8.29	0.15	0.14	0.07	1.80	6.12
7	Forge_bio	1	2048	8.15	0.17	0.16	0.08	1.91	5.83
8	Gatortron_base	2	2048	8.12	0.17	0.13	0.08	1.83	5.91
9	Gatortron_medium	1	2048	7.97	0.14	0.12	0.07	1.71	5.92

(a)   **(b)** OSA Cohort Phenotyping Tasks   **(c)** OSA Cohort Mortality Prediction Tasks

Figure 1. Quanitative results across biomedical LLMs covered in the study. **(a)** Plots subsetted latent space suprise over the estimated compute needed to pretrain each model with compute optimal conventions reported by Hoffman and Yin . Box plots show the distribution of scores per model over varied clustering hyperparameters. Compute, ratio between training tokens and parameter count, and context length of pretrained models explain 60.3% of variance when regressing against scores. **(b)** Plots area under the ROC curve (ROC-AUC) performance on admission-level phenotyping tasks over subsetted latent space scores. Moderate to strong relationships are found between performance on tasks and scores. **(c)** Plots ROC-AUC performance on admission-level overall mortality prediction over subsetted latent space scores. Moderate to strong relationships are found between 90-day and 1-year mortality prediction and scores.

Figure 2. Qualitative evaluation of best subsetted latent space (Forge-bio, k=2048, penultimate layer). **(a)** UMAP visualization of admission-level corpus manifold. Color gradient corresponds to proximity of K-Means clusters. **(b)** Per cluster KLD scores over corpus manifold. Intensity corresponds to higher amounts of order.

Figure 3. Prevalence of time-dependent variables over clusters on corpus manifold. **(a)** Plots mean age of patients at time of admission. Yellow regions correspond to clusters with patients 35 years-old and younger. Red regions correspond to clusters with patients 65 years-old and older. **(b,c,d)** Plot 1-year, 90-day, and 30-day overall mortality respectively. High prevalence clusters for overall mortality concentrate around center-left region at $[(0.1, 0.2), (0.4, 0.5)]$.

Figure 4. Prevalence of 6 OSA-related commorbid diseases over clusters on corpus manifold. **(a, e)** HTN and T2DB are prevalent across manifold with reduced prevalence observed along clusters with younger average age at time of admission. **(b, f)** AFIB and CAD concentrate on clusters along $y = 0.6125$. **(c, d)** Seperation between HF and STRK is observed. STRK concentrates on clusters following the bands at $y = 0.675$ and $y = x + 0.225$. In contrast, HF overlaps heavily on regions with concentrations of AFIB and CAD.

(a) Heart Failure Dx

(b) 1-Year Mortality

(c) Heart Failure Dx

(d) 1-Year Mortality

Figure 5. Class probabilities for HF diagnosis and 1-year mortality along cases of OSA (51,565 *admissions*) using logistic regression fit on K-Means cluster distances. Paths through corpus manifold are traced for two OSA patients who have been previously diagnosed with HF and died during their last admission to the hospital. Crosses represent unique admissions and thier color gradient correspond the admissions' class probability. **(a, b)** Plots follow admissions with diagnosed HF or OSA for *subject_id* = 18080257. Beginning at the earliest admission with OSA diagnosis (*hadm* = 0), 3 episodes of HF can be inferred at *hadm* = 2, *hadm* = (4, 8), and *hadm* = (11, 12). 1-year mortality probability increases with each episode, ultimately terminating with the patients death at *t* = 12. Cheif cause of death, congestive HF. **(c, d)** For *subject_id* = 16433543, only 1 episode of HF can be inferred at *hadm* = (1, 4). 1-year mortality probability has a sudden increase at *hadm* = 4 with the patient dying in the following admission from non-traumatic intracerebral hemorrhage.