A novel cortical biomarker signature accurately and reliably predicts individual pain sensitivity: The PREDICT longitudinal analytical validation study

Nahian S Chowdhury*[1,2], Chuan Bi*[3, 4], Andrew J Furman[5,6], Alan KI Chiang[1,2], Patrick Skippen[1,7,8], Emily Si[1], Samantha K Millard[1,2], Sarah M Margerison[5,6], Darrah Spies[5,6], Michael L Keaser[5,6], Joyce T Da Silva[5,6], Shuo Chen[3], Siobhan M Schabrun**[1,9,10], David A Seminowicz**[1,5,6,11]

*Co-first author

**Co-senior author

[1]Center for Pain IMPACT, Neuroscience Research Australia, Sydney, New South Wales, Australia
[2]University of New South Wales, Sydney, New South Wales, Australia
[3]Division of Biostatistics and Bioinformatics, Department of Epidemiology and Public Health, University of Maryland School of Medicine
[4]Division of Biostatistics, Center for Devices and Radiological Health, U.S. Food and Drug Administration, USA
[5]Department of Neural and Pain Sciences, University of Maryland School of Dentistry, Baltimore, USA
[6]Center to Advance Chronic Pain Research, University of Maryland Baltimore, USA
[7]Data Sciences, Hunter Medical Research Institute, Newcastle, NSW, Australia
[8]School of Medicine and Public Health, University of Newcastle, Newcastle, NSW, Australia
[9]The Gray Centre for Mobility and Activity, Parkwood Institute, London, Canada
[10]School of Physical Therapy, University of Western Ontario, London, Canada
[11]Department of Medical Biophysics, Schulich School of Medicine & Dentistry, University of Western Ontario, London, Canada

Data availability statement: The data supporting the findings of this study are available from the corresponding author upon reasonable request

Keywords: Pain, Biomarker, Electroencephalography, Transcranial Magnetic Stimulation

Disclosures: This project was funded by the National Institutes of Health (R61 NS113269/NS/NINDS NIH HHS/United States). The authors have no conflicts to declare.

Corresponding Author:
Professor David A Seminowicz
Department of Medical Biophysics
Schulich School of Medicine & Dentistry
University of Western Ontario, London, Canada,
E-mail: dseminow@uwo.ca

Abstract

Background: Biomarkers would greatly assist chronic pain management. The present study aimed to undertake analytical validation of a sensorimotor cortical biomarker signature for pain consisting of two measures: sensorimotor peak alpha frequency (PAF) and corticomotor excitability (CME), using a human model of prolonged temporomandibular pain (masseter intramuscular injection of nerve growth factor [NGF]).

Methods: 150 participants received an injection of NGF to the right masseter muscle on Days 0 and 2, inducing prolonged pain lasting up to 4 weeks. Electroencephalography (EEG) to assess PAF and transcranial magnetic stimulation (TMS) to assess CME were recorded on Days 0, 2 and 5. We determined the predictive accuracy of the PAF/CME biomarker signature using a nested control-test scheme: machine learning models were run on a training set (n = 100), where PAF and CME were predictors and pain sensitivity was the outcome. The winning classifier was assessed on a test set (n = 50) comparing the predicted pain labels against the true labels.

Results: The winning classifier was logistic regression, with an outstanding area under the curve (AUC=1.00). The locked model assessed on the test set had excellent performance (AUC=0.88). Results were reproduced across a range of methodological parameters and inclusion of covariates in the modelling. PAF and CME biomarkers showed good-excellent test-retest reliability.

Conclusions: This study provides evidence for a sensorimotor cortical biomarker signature for an episode of prolonged pain. The combination of accuracy, reproducibility, and reliability, suggests the PAF/CME biomarker signature has substantial potential for clinical translation.

Several objective pain biomarkers have been proposed, including neuroimaging markers of mechanistic/structural abnormalities [1-4] and "multi-omics" metrics of micro RNA [5], proteins [6] , lipids and metabolites [7]. Such biomarkers would greatly assist decision making in the diagnosis, prevention and treatment of chronic pain [8]. However, attempts at establishing pain biomarkers have suffered from either insufficient sample sizes to conduct full-scale analytical validation using machine learning [8-10], failure to use clinically relevant pain models [11-13] or lack of assessment of reproducibility or test-retest reliability [14, 15]. These factors have hindered the clinical translatability of prospective pain biomarkers.

Recent evidence shows promise for a sensorimotor cortical biomarker signature for predicting the severity of a prolonged pain episode. The biomarker signature reflects individual differences in ascending sensory and descending motor processing, comprising two metrics: 1) sensorimotor peak alpha frequency (PAF), defined as the dominant sensorimotor cortical oscillation in the alpha (8-12Hz) range [16], and is related to the efficiency in which the brain can inhibit incoming sensory input [17, 18], and 2) corticomotor excitability (CME), defined as the efficacy at which signals are relayed from primary motor cortex (M1) to peripheral muscles [19]. CME is altered during pain as individuals adopt different movement strategies to cope with pain [20, 21]. Previous work has shown that slower PAF prior to pain onset and reduced CME during prolonged pain ("depression") are associated with more pain, while faster PAF and increased CME ("facilitation") are associated with less pain [21-25]). Given individuals who experience higher pain in the early stages of a prolonged pain episode (e.g. post-surgery) are more likely to develop chronic pain in the future [26], slow PAF prior to an anticipated prolonged pain episode and/or CME depression during the acute stages of pain are potential predictors for the transition to chronic pain.

This paper presents the main outcomes of the PREDICT trial, a pre-registered (NCT04241562, [27]) full-scale analytical validation of the PAF/CME biomarker signature using a human model of prolonged myofascial temporomandibular pain (masseter intramuscular injection of nerve growth factor [NGF]). Repeated NGF injections induce progressively developing prolonged pain lasting up to 4 weeks [25, 28], and has been shown to mimic chronic pain characteristics such as time course (gradual development), type of pain (movement-evoked), functional impairments, hyperalgesia (increased pressure pain thresholds) and mechanism of sensitization [29, 30]. This makes the NGF model a highly

standardised and clinically relevant prolonged pain model with which to undertake biomarker
validation.

The aim of the PREDICT trial was to determine whether individuals could be
accurately classified as high or low pain sensitive based on baseline PAF and early CME
facilitator/depressor classification. We predicted the area under the curve (AUC) of the
receiver operator characteristic (ROC) curve for distinguishing high and low pain sensitive
individuals would be at least 70% (which represents an acceptable AUC) [31].

Methods

Participants

The PREDICT trial enrolled 159 healthy participants (70 females, 89 males, mean age
25.1 ± 6.1), with 150 participants remaining after participant dropouts. Ethical approval was
obtained from the University of New South Wales (HC190206) and the University of
Maryland Baltimore (HP-00085371). Written, informed consent was obtained. The
supplementary appendix contains all additional details regarding participant characteristics
and methodology.

Experimental Protocol

Outcomes were collected over a period of 30 days. Participants attended the laboratory
on Day 0, 2, and 5. Baseline questionnaire data were collected on Day 0. Pressure pain
thresholds, PAF and CME were measured on Day 0, 2 and 5. PAF was obtained via a 5-minute
eyes-closed resting-state EEG recording from 63 scalp electrodes. Sensorimotor PAF was
computed by identifying the component in the signal (transformed by independent component
analysis) that had a clear alpha peak (8-12Hz) upon frequency decomposition and a scalp
topography suggestive of a sensorimotor source. CME was obtained using transcranial
magnetic stimulation (TMS) mapping; single pulses of TMS delivered to the left primary motor
cortex (M1), and motor evoked potentials (MEPs) recorded from the right masseter muscle
using electromyography (EMG) electrodes. TMS was delivered at each site on a 1cm-spaced
grid superimposed over the scalp, and a map of the corticomotor representation of the masseter
muscle was generated. Corticomotor excitability was indexed as map volume, which is
calculated by summing MEP amplitudes from all "active sites" on the grid. NGF was injected
into the right masseter muscle at the end of the Day 0 and 2 laboratory sessions. Electronic pain

diaries were collected from Days 1 to 30 at 10am and 7pm each day, where participant rated their pain (0-10) during various activities. Pain upon functional jaw movement is a key criterion for the diagnosis of TMD [32]. Moreover, previous research has shown that, after an NGF injection to the masseter muscle, pain during chewing and yawning are higher compared to other activities [30, 33]. As such, the primary outcomes were pain upon chewing and yawning. The protocol and methodology are shown in Figure 1A and 1B.

Analytical Validation Plan

Division of the Data. Analysis was conducted in R, MATLAB and Python, with code publicly available https://github.com/DrNahianC/PREDICT_Scripts. Figure 1C details the analysis plan. We adopted a nested-control-testing scheme by partitioning 150 participants into a training and test set of 100 and 50 participants respectively.

Growth Mixture Modelling. We used growth mixture modelling (GMM) in R [34-36] to form two participant classes: high and low pain sensitive. For this categorization, we used the sum of pain upon chewing and yawning data, and pain diary trajectories from Days 1-7 for the classification, as this was the timeframe when pain was most prominent (Supplementary Figure 3). As such, participants would more reliably fall into high and low pain sensitive classes during this timeframe. The first and last 40 participants (80 in total) in the training set, based on the ordering of probabilities of the pain intensity trajectory belonging to one of the classes, were then labelled as high and low pain sensitive. The trained GMM model, once established, was locked and utilized to label the test set. Consequently, 38 out of 50 test set participants (24 high and 14 low pain) were labelled. These labels were recorded for subsequent comparison with the predicted labels produced by the trained machine learning model.

Machine Learning Model Selection and Fine Tuning. We utilized five machine learning models on the labelled training set —logistic regression, random forest, gradient boosting, support vector machine, and neural network. The dependent variable was pain sensitivity label (high or low) identified from the GMM and independent variables were sensorimotor PAF and CME: the latter was typified as facilitator and depressor, depending on whether they showed an increase or decrease in map volume on Day 5 relative to Day 0, respectively. For each model, we identified optimized parameters through 5-fold cross-validation: we randomly divided the 80 participants into an internal training set of 64 participants (consisting of four equal folds of 16) and a validation set of 16. The optimized models in the internal training set were then employed to predict labels in the validation set to

facilitate model selection. The model with the best performance (area under the curve) on the

validation set was then locked in.

Test Set Prediction. The locked machine learning model was assessed on the test set.

The participant IDs in this set did not coincide with those in the pain diary data, thereby

preserving the double-blind nature of the analysis. By using the ground truth labels

(shuffled), predicted labels (unshuffled), and the shuffling order for the test set, we were able

to evaluate the model's performance by comparing the reordered predicted labels against the

ground truth labels established by the GMM. Performance was assessed via receiver

operating characteristic (ROC) area under the curve (AUC), with 95% confidence intervals

reported. AUC values between 0.7-0.8, 0.8-0.9 and 0.9-1 were considered "acceptable",

"excellent", and "outstanding" respectively [31].

178   **Results**

PAF/CME demonstrated good-excellent test-retest reliability

PAF and ΔCME showed good to excellent test-retest reliability across sessions

(Supplementary Figures 5 and 7).

Outstanding performance on the training validation set

Figure 2A shows the pain scores for participants in the training and test set classified

as high and low pain sensitive based on GMM. Figure 2B (upper) shows the performances of

the machine learning models across the internal training and validation sets. Logistic

regression was chosen as the optimal classifier based on its outstanding performance

(AUC=1.00[1.00-1.00]) when applied to the validation set (Figure 2B lower), with slower

PAF and CME depression predicting higher pain.

Excellent performance on the test set

When the locked logistic regression model was applied to the test set, performance

(Figure 2C upper) was excellent (AUC=0.88[0.78-0.99]). Figure 2C (lower) shows the

differences in pain scores between participants predicted to have high or low pain. Visually

one can observe slower peak alpha frequency in those predicted to have high vs. low pain

sensitivity (Figure 2D), which was further confirmed with a two-sample t-test ($t(48)=5.8$,

$p<.001$). Moreover, one can observe a decrease in CME within the masseter motor maps in

those predicted to have high pain (Figure 2E), whereas those predicted to have low pain exhibit an increase in CME. The differences in the change in CME relative to Day 0 between these groups was further confirmed with a two-sample t-test ($t(48)$=2.81, p=.007).

A benefit for a *combined* signature

We reran the models to determine whether the combined PAF/CME signature out-performed each measure individually (Supplementary Figure 9). For PAF alone, the performance of the logistic regression model on the training validation and test set were respectively excellent (AUC=0.95[0.84-1.00]) and outstanding (AUC=0.83[0.70-0.96]). For CME alone, the performance of the logistic regression model for the training validation and test set were respectively excellent (AUC=0.88[0.69-1.00]) and acceptable (AUC=0.75[0.60-0.91]).

Results were reproducible when including covariates

We evaluated the performance of the biomarker combined with demographic and clinical attributes. As we collected a large amount of this data, we applied feature selection, i.e. filtering features by inspecting p-values when associating predictors and labels, and using parameter tuning to optimize the coefficients associated with the filtered features. Five features were subsequently selected and optimized – Sensorimotor PAF, CME, Sex, Pain Catastrophizing Scale (PCS) Total and PCS Helplessness. The associations between labels and biomarkers/covariates in the training vs. test set, and performance of the models are shown in Figure 3A and 3B. When including these five features, the performance of the logistic regression model was for the training validation and test set were respectively outstanding (AUC=1.00[1.00-1.00]) and excellent (AUC=0.81[0.67-0.95]).

Results were reproducible across methodological choices

To determine whether our results were robust across different methodological choices, we repeated the analysis using PAF calculated using component level data (with the sensorimotor component chosen manually or using an automated script) vs. sensor level data (with a sensorimotor region of interest), using different frequency windows (8-12Hz vs. 9-11Hz) and using different CME calculation methods (map volume vs. map area). We found that, regardless of the choices, logistic regression was the best or equal-best performing model when applied to the validation set (Figure 3C), with AUCs varying from acceptable (0.77) to outstanding (1.00). When the locked models were applied to the test set,

performance varied from acceptable (AUC=0.73) to excellent (AUC=0.88) (Figure 3D).

Lastly, excellent performance was demonstrated when the data was analysed two other ways

(Supplementary Figure 10 and 11): where GMM pain labels were established using the whole

30 days rather than the first 7 days (training validation AUC=0.84[0.64-1], test AUC

=0.89[0.79-0.99]), and when missing pain diary data was not imputed (training validation

AUC=0.81[0.6-1], test AUC=0.89[0.79-0.99]).

Discussion

A full-scale analytical validation of the PAF and CME biomarker signature was

conducted using a clinically relevant prolonged pain model. In an initial training set (n=100),

we found that a logistic regression was the optimal classifier based on its outstanding

performance (AUC=100%), with slower PAF and CME depression predicting higher pain.

When this model was applied to an independent test set, the AUC was excellent (88%).

PAF/CME showed good-excellent test-retest reliability, and results were reproduced across a

range of methodological parameters and consideration of covariates. Overall, the combination

of sample size, pain model validity, and biomarker accuracy, reproducibility and reliability

suggest the PAF/CME biomarker signature has substantial potential for clinical translation.

Our results suggest that individuals who have slow PAF prior to an anticipated

prolonged pain episode and show corticomotor depression during a prolonged pain episode,

are more likely to experience higher pain. Model performance was higher combining the two,

suggesting consideration of both ascending sensory and descending motor pain processing

mechanisms provides more information regarding pain sensitivity. Overall, we believe this

biomarker could be particularly useful in contexts such as predicting post-operative pain. For

example, a recent study showed that individuals with slower PAF experienced more pain

following a thoracotomy [25]. Given that higher acute pain post-surgery predicts the

development of chronic pain [26], our findings suggest individuals with slow PAF/reduced

CME could be more likely transition to chronic pain. Indeed, individuals who show lower

CME during the acute stages of low back pain were more likely to develop chronic pain at 6-

months follow-up [37]. These preliminary findings, along with our analytical validation

study, suggest PAF and CME could be susceptibility biomarkers for the transition from acute

to chronic pain.

There are several aspects of our study which stand out within the field. The first is sample size: with recent advancements in machine learning, it has become possible to conduct analytical validation of pain biomarkers. However, deep learning requires a large amount of labelled samples to conduct rigorous training on validation and test sets [8]. Unfortunately, many pain susceptibility biomarker studies have not been sufficiently sampled to adopt such approaches [9, 10], and the ones that have used machine learning failed to reach the sample sizes similar to that of the present study [1, 2].

Another strength of our findings is reproducibility. The majority of work has shown similar associations between higher pain and slower PAF [16, 22] and CME depression [21, 25] in models of upper limb pain. The present study replicated these results in a model of prolonged jaw pain, suggesting these associations hold across pain locations. It is important to note that some studies have not shown a negative relationship between PAF and pain sensitivity [38, 39] or a positive relationship between CME depression and pain sensitivity [33]. However, these studies were not sufficiently sampled to conduct analytical validation of the kind presented in this study. Nonetheless, the mixed findings could also arise from differences in methodological choices in the estimation of PAF e.g. frequency windows [39] and use of sensor vs. component space data [40] and estimation of CME e.g. map volume [21] vs. area [33]. For this reason, we repeated the main analysis using different methodological choices and found at least acceptable AUCs. In addition, we found that the inclusion of covariates such as pain catastrophizing and sex did not alter our results, further supporting the reproducibility of our results.

The PAF/CME measures demonstrated good-excellent reliability. Reliability is a highly desirable characteristic which assists in the widespread application of pain biomarkers [8]. We found in the present analysis, and previously [14], that participants exhibit stable PAF across days despite the presence of pain, and even when considering different methodological factors that may influence the reliability of PAF such as pre-processing pipeline, recording length and frequency window. Indeed, we found reliable PAF with a recording length as short as 2 minutes and minimal data pre-processing. We also showed that those who show CME depression on Day 2 are also likely to show CME depression on Day 5 (and vice versa for those who show CME facilitation). This finding was shown even when an automated method of determining MEP amplitude on each trial was applied. Thus, our work not only shows that PAF and CME can predict pain, but the relative ease with which reliable PAF/CME data can be obtained is promising for subsequent clinical translation.

Another strength of this study is the clinical relevance of our pain model, making

clinical translation of the current findings highly feasible. While other pain biomarker studies

have shown promising results, these studies were restricted to pain models utilizing transient

painful stimuli lasting seconds to minutes [11-13]. The brief nature of the painful stimuli

questions the external validity of these findings and limits generalizability to clinical

populations. In contrast, the present study used a prolonged pain model lasting weeks.

Several other studies have shown that injections of NGF to the neck, elbow or masseter

muscles can mimic symptoms of clinical neck pain [41], chronic lateral epicondylalgia [29]

and TMD [30] respectively. Thus, the observed relationships between PAF/CME and pain in

the present study show promise in terms of clinical applicability.

Lastly, the PAF/CME biomarker demonstrated high performance. A previous study

found that connectivity between medial prefrontal cortex and nucleus accumbens in 39 sub-

acute low back pain patients (pain duration 6-12 weeks) could predict future pain persistence

at ~7, 29 and 54 weeks, with AUCs of 67-83% [1]. Another study on 24 sub-acute low back

pain patients showed that white matter fractional anisotropy measures in the superior

longitudinal fasciculus and internal capsule predicted pain persistence over the next year,

with an AUC of 81% [2]. Though the present did not directly assess the transition to chronic

pain, our AUCs of 100% (validation set) and 88% (test set) appear comparatively high. We

therefore encourage future clinical studies to determine whether PAF/CME can predict the

transition from acute to chronic pain.

Conclusions

312   A novel biomarker signature comprised of PAF and CME accurately and reliably

313   distinguishes high and low pain sensitive individuals during prolonged jaw pain with an

314   excellent AUC of 88% in an independent test set. No other pain biomarker study has shown

315   this combination of biomarker accuracy, reproducibility, reliability and pain model validity,

316   suggesting high potential for clinical translation.

References

1. Baliki, M.N., et al., *Corticostriatal functional connectivity predicts transition to chronic back pain.* Nature neuroscience, 2012. **15**(8): p. 1117-1119.

2. Mansour, A.R., et al., *Brain white matter structural properties predict transition to chronic pain.* PAIN®, 2013. **154**(10): p. 2160-2168.

3. Musto, A.E., et al., *Hippocampal neuro-networks and dendritic spine perturbations in epileptogenesis are attenuated by neuroprotectin d1.* PLoS One, 2015. **10**(1): p. e0116543.

4. Vachon-Presseau, E., et al., *Corticolimbic anatomical characteristics predetermine risk for chronic pain.* Brain, 2016. **139**(7): p. 1958-1970.

5. Wang, W.-T., et al., *Circulating microRNAs identified in a genome-wide serum microRNA expression analysis as noninvasive biomarkers for endometriosis.* The Journal of Clinical Endocrinology & Metabolism, 2013. **98**(1): p. 281-289.

6. Singh, J.A., S. Noorbaloochi, and K.L. Knutson, *Cytokine and neuropeptide levels are associated with pain relief in patients with chronically painful total knee arthroplasty: a pilot study.* BMC Musculoskeletal Disorders, 2017. **18**(1): p. 1-6.

7. Sasamoto, N., et al., *Presurgical blood metabolites and risk of postsurgical pelvic pain in young patients with endometriosis.* Fertility and Sterility, 2022. **117**(6): p. 1235-1245.

8. Zhang, Z., J.S. Gewandter, and P. Geha, *Brain imaging biomarkers for chronic pain.* Frontiers in Neurology, 2022. **12**: p. 734821.

9. Sluka, K.A., et al., *Predicting chronic postsurgical pain: current evidence and a novel program to develop predictive biomarker signatures.* Pain, 2023. **164**(9): p. 1912-1926.

10. Eldabe, S., et al., *Biomarkers for chronic pain: significance and summary of recent advances.* Pain Research and Management, 2022. **2022**.

11. Lee, J.-J., et al., *A neuroimaging biomarker for sustained experimental and clinical pain.* Nature medicine, 2021. **27**(1): p. 174-182.

12. Woo, C.-W., et al., *Quantifying cerebral contributions to pain beyond nociception.* Nature communications, 2017. **8**(1): p. 14211.

13. Wager, T.D., et al., *An fMRI-based neurologic signature of physical pain.* New England Journal of Medicine, 2013. **368**(15): p. 1388-1397.

14. Chowdhury, N.S., et al., *The reliability of two prospective cortical biomarkers for pain: EEG peak alpha frequency and TMS corticomotor excitability.* Journal of Neuroscience Methods, 2023. **385**: p. 109766.

15. Davis, K.D., et al., *Discovery and validation of biomarkers to aid the development of safe and effective pain therapeutics: challenges and opportunities.* Nature Reviews Neurology, 2020. **16**(7): p. 381-400.

16. Furman, A.J., et al., *Cerebral peak alpha frequency predicts individual differences in pain sensitivity.* NeuroImage, 2018. **167**: p. 203-210.

17. Mazaheri, A., D.A. Seminowicz, and A.J. Furman, *Peak alpha frequency as a candidate biomarker of pain sensitivity: the importance of distinguishing slow from slowing.* NeuroImage, 2022. **262**: p. 119560.

18. Van Diepen, R.M., J.J. Foxe, and A. Mazaheri, *The functional role of alpha-band activity in attentional processing: the current zeitgeist and future outlook.* Current opinion in psychology, 2019. **29**: p. 229-238.

19. Weavil, J.C. and M. Amann, *Corticospinal excitability during fatiguing whole body exercise.* Progress in brain research, 2018. **240**: p. 219-246.

20. Summers, S.J., et al., *Motor adaptation varies between individuals in the transition to sustained pain.* Pain, 2019. **160**(9): p. 2115-2125.

21. Seminowicz, D.A., T. Thapa, and S.M. Schabrun, *Corticomotor depression is associated with higher pain severity in the transition to sustained pain: A longitudinal exploratory study of individual differences.* The Journal of Pain, 2019. **20**(12): p. 1498-1506.

22. Furman, A.J., et al., *Sensorimotor peak alpha frequency is a reliable biomarker of prolonged*
pain sensitivity. Cerebral Cortex, 2020. **30**(12): p. 6069-6082.
23. Millard, S.K., et al., *Predicting postoperative pain in lung cancer patients using preoperative*
peak alpha frequency. British Journal of Anaesthesia, 2022. **128**(6): p. e346-e348.
24. Seminowicz, D.A., et al., *Slow peak alpha frequency and corticomotor depression linked to*
high pain susceptibility in transition to sustained pain. bioRxiv, 2018: p. 278598.
25. Chowdhury, N.S., et al., *The Effect of Acute and Sustained Pain on Corticomotor Excitability:*
A Systematic Review and Meta-analysis of group-and individual-level data. The Journal of
Pain, 2022. **23**(10): p. 1680-1696.
26. Hah, J.M., et al., *Factors associated with acute pain estimation, postoperative pain*
resolution, opioid cessation, and recovery: secondary analysis of a randomized clinical trial.
JAMA network open, 2019. **2**(3): p. e190168-e190168.
27. Seminowicz, D.A., et al., *A novel cortical biomarker signature for predicting pain sensitivity:*
protocol for the PREDICT longitudinal analytical validation study. Pain Reports, 2020. **5**(4).
28. Svensson, P., et al., *Injection of nerve growth factor into human masseter muscle evokes*
long-lasting mechanical allodynia and hyperalgesia. Pain, 2003. **104**(1-2): p. 241-247.
29. Bergin, M.J.G., et al., *Movement evoked pain and mechanical hyperalgesia after*
intramuscular injection of nerve growth factor: a model of sustained elbow pain. Pain
Medicine, 2015. **16**(11): p. 2180-2191.
30. Schabrun, S.M., et al., *Intramuscular injection of nerve growth factor as a model of*
temporomandibular disorder: nature, time-course, and sex differences characterising the
pain experience. Neurobiology of Pain, 2023. **13**: p. 100117.
31. Hosmer Jr, D.W., S. Lemeshow, and R.X. Sturdivant, *Applied logistic regression*. Vol. 398.
2013: John Wiley & Sons.
32. Schiffman, E., et al., *Diagnostic criteria for temporomandibular disorders (DC/TMD) for*
clinical and research applications: recommendations of the International RDC/TMD
Consortium Network and Orofacial Pain Special Interest Group. Journal of oral & facial pain
and headache, 2014. **28**(1): p. 6.
33. Costa, Y.M., et al., *Masseter corticomotor excitability is decreased after intramuscular*
administration of nerve growth factor. Eur J Pain, 2019. **23**(9): p. 1619-1630.
34. Proust-Lima, C., V. Philipps, and B. Liquet, *Estimation of extended mixed models using latent*
classes and latent processes: the R package lcmm. arXiv preprint arXiv:1503.00890, 2015.
35. Li, F., et al., *Latent growth modeling of longitudinal data: A finite growth mixture modeling*
approach. Structural Equation Modeling, 2001. **8**(4): p. 493-530.
36. Ram, N. and K.J. Grimm, *Methods and measures: Growth mixture modeling: A method for*
identifying differences in longitudinal change among unobserved groups. International
journal of behavioral development, 2009. **33**(6): p. 565-576.
37. Jenkins, L.C., et al., *Cortical function and sensorimotor plasticity are prognostic factors*
associated with future low back pain after an acute episode: the UPWaRD prospective cohort
study. Pain, 2022: p. 10.1097.
38. Nir, R.-R., et al., *Pain assessment by continuous EEG: association between subjective*
perception of tonic pain and peak frequency of alpha oscillations during stimulation and at
rest. Brain research, 2010. **1344**: p. 77-86.
39. De Martino, E., et al., *Slowing in peak-alpha frequency recorded after experimentally-induced*
muscle pain is not significantly different between high and low pain-sensitive subjects. The
Journal of Pain, 2021. **22**(12): p. 1722-1732.
40. Millard, S.K., et al., *Effects of nicotine compared to placebo gum on sensitivity to pain and*
mediating effects of peak alpha frequency. 2023, eLife Sciences Publications, Ltd.
41. Christensen, S.W.M., et al., *Effect of prolonged experimental neck pain on exercise-induced*
hypoalgesia. Pain, 2022. **163**(12): p. 2411-2420.

Figure 1. **(A)** Experimental protocol showing timeline of data collection procedures. On Day 0, we measured peak alpha frequency (PAF) and corticomotor excitability (CME). At the end of the session, an injection of nerve growth factor (NGF) was administered to the right masseter muscle. On Day 2, PAF and CME were measured, followed by a second NGF injection. On Day 5, PAF and CME were measured. From Days 1-30, electronic diaries measuring jaw pain were sent to participants at 10AM and 7PM each day. **(B)** Details of the methodology. Sensorimotor PAF was measured using a 5 minutes eyes closed resting state EEG recording. Sensorimotor PAF was computed by identifying the component in the signal (transformed by independent component analysis) that had a clear alpha peak in the 8–12 Hz range upon frequency decomposition and a scalp topography suggestive of a source predominately over the sensorimotor cortex. TMS mapping was conducted by stimulating the scalp area over left M1 to obtain a map of the representation of the right masseter muscle. The map consists of the motor-evoked potential (MEP) amplitude at each stimulated location, with CME corresponding to the map volume (sum of all MEPs from active sites). **(C)** Details of the analysis plan. We adopted a nested-control-test scheme by partitioning the 150 subjects into a training set consisting of 100 subjects and an independent test set of 50 subjects. We labelled a subset of participants in the training (n = 80) and test set (n = 38) as high or low pain sensitive using growth mixture modelling (GMM) to establish "ground-truth" labels. We then ran various machine learning models on the labelled training set (with PAF/CME as predictors, and pain severity labels as outcome), and determined optimized parameters through 5-fold cross-validation i.e. randomly dividing the 80 subjects into an internal training set of 64 subjects (with 4 equal folds of 16) and a validation set of 16. The optimized models in the internal training set were employed to predict labels in the validation set to facilitate model selection. The model with the best performance on the validation set was then locked in, and applied to the labelled test set, comparing the predicted labels of high/low pain sensitive with the ground-truth labels of high/low pain sensitive.

Figure 2. **(A)** Results of the growth mixture modelling which categorized 80 participants in the training set (left) and 38 participants in the test (right) as high or low pain sensitive. Data shows mean pain score (chew + yawn pain rating) for each timepoint, while the shaded area shows 95% confidence intervals. **(B)** The upper panel shows performances (AUC [95% confidence intervals]) of various machine learning models for the internal training set and validation set. Logistic regression (LR) was chosen as the optimal classifier based on outstanding AUC of 100% as shown in the lower panel. **(C)** The upper panel shows the performance of the locked logistic regression model when applied to the test set, which was in the excellent range (AUC of 88%). The lower panel shows the pain trajectories (mean chew + yawn pain and 95% confidence intervals) of participants predicted to have high or low pain sensitivity based on the locked logistic regression model. **(D)** Individual and mean z-transformed spectral plots and topography of the sensorimotor alpha component on Day 0 for participants predicted to have high or pain sensitivity based on the locked logistic regression model. **(E)** The mean motor cortex maps on Day 0 and Day 5 showing normalized motor evoked potential (MEP) amplitude (expressed as a proportion of the maximal MEP amplitude) for participants predicted to have high or low pain sensitivity based on the locked logistic regression model.

Figure 3. **(A)** Visualisation of biomarkers and covariates for the training and test sets across high (red) and low (blue) pain labels identified from the GMM. Data on PAF, PCS total and PCS helplessness are plotted as boxplots, while data on CME and Sex are plotted according facilitator: depressor (Fac: dep) and female: male (fem: mal) split respectively, including odd ratios. A lower odds ratio means a lower probability of high pain sensitive individuals belonging to the facilitator or female categories. For PAF and CME, low pain was associated with fast PAF and CME facilitation for both training and test sets. In contrast, the relationship between covariates and labels were in the opposite direction for the training and test set, suggesting the relationship between biomarkers and labels was consistent. **(B)** The left panel shows the performance of the locked logistic regression model on the test set when including covariates in the model. The right panel shows pain trajectories (mean chew + yawn score and 95% confidence intervals) of participants predicted to have high or low pain sensitivity based on the locked logistic regression model including covariates. **(C)** The performance of each machine learning model (AUC [95% confidence intervals]) on the training validation set across different PAF/CME calculation methods. This includes the sensorimotor component chosen manually after an independent component analysis, component identified using an automated script after an independent component analysis, or using a sensorimotor region of interest (ROI, mean of Cz, C3 and C4) in electrode space, to calculate PAF. We also looked at different frequency windows for computing PAF (8-12Hz vs. 9-11Hz) or CME calculated using map area or map volume. **(D)** The performance of the locked logistic regression model (AUC [95% confidence intervals]) when applied to the test set, across different PAF/CME calculation methods.

Figure 1

508   **Figure 2**

515   **Figure 3**