Directed exploration is elevated in affective disorders but reduced by an aversive

interoceptive state induction

Ning Li[1*], Claire A. Lavalley[1*], Ko-Ping Chou[1], Anne E. Chuning[1], Samuel Taylor[1], Carter M.

Goldman[1], Taylor Torres[1], Rowan Hodson[1], Robert C. Wilson[3,4], Jennifer L. Stewart[1], Sahib S.

Khalsa[1,2], Martin P. Paulus[1,2] & Ryan Smith[1,2]

*Co-first authorship

[1]Laureate Institute for Brain Research, Tulsa, OK.

[2]Oxley College of Health and Natural Sciences, University of Tulsa, Tulsa, OK.

[3]Department of Psychology, University of Arizona, Tucson, AZ.

[4]Cognitive Science Program, University of Arizona, Tucson, AZ.

Corresponding author: Ryan Smith, rsmith@laureateinstitute.org

Laureate Institute for Brain Research. 6655 S. Yale Ave. Tulsa, Oklahoma, 74136, United States

Abstract

Elevated anxiety and uncertainty avoidance are known to exacerbate maladaptive choice in

individuals with affective disorders. However, the differential roles of state vs. trait anxiety remain

unclear, and underlying computational mechanisms have not been thoroughly characterized. In the

present study, we investigated how a somatic (interoceptive) state anxiety induction influences

learning and decision-making under uncertainty in individuals with clinically significant levels of

trait anxiety. A sample of 58 healthy comparisons (HCs) and 61 individuals with affective

disorders (iADs; i.e., depression and/or anxiety) completed a previously validated explore-exploit

decision task, with and without an added breathing resistance manipulation designed to induce

state anxiety. Computational modeling revealed a pattern in which iADs showed greater

information-seeking (i.e., directed exploration; Cohen's d=.39, p=.039) in resting conditions, but

that this was reduced by the anxiety induction. The affective disorders group also showed slower

learning rates across conditions (Cohen's d=.52, p=.003), suggesting more persistent uncertainty.

These findings highlight a complex interplay between trait anxiety and state anxiety. Specifically,

while elevated trait anxiety is associated with persistent uncertainty, acute somatic anxiety can

paradoxically curtail exploratory behaviors, potentially reinforcing maladaptive decision-making

patterns in affective disorders.

Keywords: decision-making; computational modeling; exploration; anxiety; depression

Introduction

Persistent uncertainty and maladaptive avoidance are key maintenance factors in anxiety disorders and a major focus of psychotherapy (1, 2). While considerable progress has been made in understanding the neural and cognitive mechanisms associated with anxiety, underlying computational processes have only been examined in a limited number of studies to date (3, 4). For example, subclinical levels of trait anxiety have been associated with reduced flexibility in learning rates when there is a change in the stability of environmental statistics (5), and individuals with anxiety disorders show elevated learning rates in general (6), suggesting a belief that action-outcome contingencies often change unexpectedly. Both depression and anxiety have also been associated with elevated learning from punishment in particular (7). More recently, trait anxiety has been linked to a greater tendency to infer changes in the underlying causes of aversive outcomes during extinction learning – which may facilitate return of fear and reduce the long-term efficacy of behavioral therapies (8).

While such studies provide important insights into the learning processes that may contribute to depression and anxiety disorders, they do not fully account for avoidance or other behaviors driven by intolerance of uncertainty. There could also be multiple computational mechanisms underlying avoidance behavior, each representing distinct hypotheses and possible treatment targets. For example, individuals must often seek information to learn that a feared situation is tolerable. Avoidance prevents such learning, which can in turn maintain avoidance. This speaks to the explore-exploit dilemma (9, 10), which has begun to receive attention in psychiatry and substance use research (11-13). This dilemma reflects the need to judge whether one has sufficient information to maximize reward, or whether one should first seek more information. In avoidance, an individual may hold the confident belief that avoided situations are dangerous, while, in fact, exploration would allow them to learn otherwise. Yet, there are multiple types of exploration, and factors that could deter exploration, which have not been thoroughly evaluated.

To date, only a few studies have examined the relationship between affective disorders and information-seeking. For example, one study found that 'directed' exploration (DE) in a community sample was lower in those with higher anxiety and depression symptoms (14), while another study associated lower DE with greater somatic anxiety in particular (15). This type of exploration is strategically directed toward situations and actions for which an individual has had

fewer past experiences. Further studies examining effects of traumatic/unpredictable childhood

environments (16, 17) – which often correlate with affective disorders (18-20) – also suggest

negative effects on DE. Other supportive work has shown that: higher stress/anxiety is associated

with less exploration in a virtual-reality plus-maze (21), agoraphobia and anxiety sensitivity are

associated with less exploratory behavior (22), and increases in cortisol in response to an acute

stressor and scores on a chronic stress questionnaire are both associated with under-exploration

in foraging tasks (23). Aversive arousal states are also generally known to reduce cognitive

control network activity in neuroscience studies (24-28), consistent with reduced cognitive

reflection tendencies seen in those who display lower DE (14). Notably, a distinct exploratory

strategy – 'random' exploration (i.e., where choices become less reward-driven as a means of

gaining information) – has not shown such associations.

Past research therefore suggests that anxiety may selectively reduce DE, consistent with

maintained avoidance. Yet, these studies are largely correlational and have tended to focus on

trait anxiety, and they have primarily investigated sub-clinical symptoms in community samples.

There are also some reasons to expect relationships in the oppositive direction. Namely, the

cognitive aspects of anxiety involve elevated uncertainty and worry, which could promote over-

exploration. Indeed, intolerance of uncertainty could drive over-exploration as a means of

continually attempting to reduce it. This is consistent with some work instead suggesting greater

DE in those with higher trait/cognitive aspects of anxiety (29, 30) and increased exploratory

behavior in depression (30, 31). Combined with the work reviewed above, this suggests the

possibility that somatic anxiety might reduce exploration, while uncertainty- or worry-related

cognitive anxiety could instead promote it.

Here we sought to build on this previous work by manipulating state anxiety with a somatic

anxiety induction while individuals performed an explore-exploit task. By comparing exploration

and learning with vs. without this induction, while also gathering information about trait anxiety

and other clinical, cognitive, and affective dimensions, we aimed to disentangle the role of state-

and trait-related anxiety and relationships with cognitive vs. somatic aspects of anxiety. By doing

this in both healthy comparison (i.e., low-anxiety) and clinical (i.e., high-anxiety) groups, we

also sought to clarify potential differences between subclinical variation in anxiety and that

associated with psychopathology. We hypothesized that anxiety induction would reduce DE and

that somatic vs. cognitive components of anxiety sensitivity might show negative and positive

relationships with exploration, respectively. Given the mixed results described above with

respect to trait anxiety, we did not have a confident directional hypothesis regarding group

differences. As secondary aims, we also sought to reproduce the aforementioned relationships

found between DE and both cognitive reflectiveness (14) and early adversity (16, 17), as these

associations could offer additional explanatory power if present in our sample.

<div align="center">

Methods

</div>

Participants

Data were collected as part of a larger, multi-visit study to investigate the cognitive and neural

correlates of psychiatric disorders at the Laureate Institute for Brain Research (LIBR), with

participants recruited from the community in and surrounding Tulsa, OK, USA. Clinical

diagnoses were assessed by a licensed clinician according to the Mini International

Neuropsychiatric Interview 7 (MINI; (32)). Healthy comparisons (HCs) were not previously

diagnosed with, or treated for, any mental health disorder and had a score of < 8 on the Overall

Anxiety Severity and Impairment Scale (OASIS; (33)). Individuals within the high-anxiety

affective disorders group (iADs) had a current diagnosis of one or more anxiety or depressive

disorders and scored $\geq$ 8 on the OASIS in a pre-visit screening. Any iADs who were taking

psychiatric medications (e.g., SSRIs, SNRIs, etc.) were required to have a stable dosage for at

least 6 weeks prior to study entry. No other neurological medications (e.g., antipsychotics,

stimulants, or anticonvulsive medications) were allowed; nor were the following diagnoses:

bipolar disorder, personality disorders, substance use disorders, eating disorders, schizophrenia,

or obsessive-compulsive disorder. Recruitment aimed to match participants in the two groups by

age, sex, and education level. Participants were asked not to consume any drugs (e.g.,

benzodiazepines, marijuana) or alcohol in the 48 hours preceding participation and were required

to pass a drug panel during their study visit.

Seven individuals withdrew from participation due to discomfort with the anxiety induction

paradigm (2 HCs, 5 iADs, all female; details in **Supplementary Materials**), leading to a final

sample of 119 participants: 58 HCs, 61 iADs (see **Table 1** below). A power analysis (using the

wp.rmanova function in the *WebPower* package in R (34)) indicated that this sample size would

afford 80% power to detect a medium effect size of $\eta_p^2 = .063$ for the effect of the anxiety

induction.

It is important to note here that data from the healthy sample in this study was previously used

for comparison to a different clinical population (methamphetamine use disorders) (45). Here,

we instead focus on differences between this group and iADs.

Measures

Primary measures were chosen to assess state and trait anxiety (State-Trait Anxiety Inventory

[STAI]; (35)) and to account for clinical symptoms of depression (Patient-Health Questionnaire

[PHQ]; (36)). Our secondary aim of replicating prior work motivated inclusion of measures to

test links between DE and both childhood adversity (Childhood Trauma Questionnaire [CTQ];

(37)) and cognitive reflectiveness (Cognitive Reflection Test [CRT]; (14, 16, 17)) – the tendency

to 'think things through' before responding based on intuition. Detailed descriptions of each

measure are included in **Supplementary Methods**. Descriptive information for all participants is

shown in **Table 1**, as well as preliminary group comparisons. Additional sample demographics

are included in **Supplementary Table S1** and a diagnostic breakdown for iADs is shown in

Supplementary Table S2.

Table 1. Demographic and clinical characteristics (mean and SD).

Measure	Healthy Comparisons (*N*=58)	Affective Disorders (*N*=61)	Statistic
Age	35.41 (13.08)	33.44 (10.97)	$t(117) = 0.89, p = .374$
Sex	72% Female	75% Female	$\chi^2(1) = 0.03, p = .870$
PHQ	1.66 (2.27)	9.46 (5.15)	**$t(117) = 10.60, p < .001$**
OASIS	1.07 (1.52)	8.41 (3.02)	**$t(117) = 16.62, p < .001$**
QIDS	2.17 (1.77)	8.77 (4.01)	**$t(117) = 11.52, p < .001$**
STAI Trait	29.17 (7.4)	51.87 (9.06)	**$t(117) = 14.92, p < .001$**
CTQ Total*	34.44 (10.36)	48.07 (16.19)	**$t(112) = 5.31, p < .001$**
Physical Neglect	6.18 (2.14)	8.41 (3.79)	**$t(112) = 3.82, p < .001$**
Physical Abuse	6.67 (2.33)	7.95 (3.26)	**$t(112) = 2.39, p = .019$**
Emotional Neglect	8.76 (3.88)	13.03 (5.02)	**$t(112) = 5.05, p < .001$**
Emotional Abuse	6.89 (2.98)	10.59 (4.59)	**$t(112) = 5.07, p < .001$**
Sexual Abuse	5.93 (3.05)	8.08 (5.01)	**$t(112) = 2.75, p = .007$**
CRT	3.33 (2.18)	2.18 (1.93)	**$t(117) = 3.05, p = .003$**

Note. While OASIS scores at screening were required to be ≥ 8 for inclusion in the Affective Disorders
group, values shown here are at the later date of study participation. As OASIS scores are rated with
respect to the prior week, some scores represented here were < 8 in this group. Legend: PHQ = Patient
Health Questionnaire-9 (depressive symptoms); OASIS = Overall Anxiety Severity and Impairment

Scale; STAI = State-Trait Anxiety Inventory, CTQ = Childhood Trauma Questionnaire; CRT = Cognitive
Reflection Test (scores correspond to number of correct answers).
*This measure was collected as part of the larger study at an earlier date (35 days prior to participation on
average).
154

Somatic Anxiety Induction and Sensitivity Assessment

To manipulate state anxiety, we utilized a previously established interoceptive (breathing-based)

anxiety induction paradigm (38-40). In this paradigm, participants are asked to breathe through a

silicon mask attached to a valve (**Figure 1a**) that allows application of different levels of

inspiratory resistance, creating air hunger-related sensations that induce anxiety. To provide

participants a chance to familiarize themselves with the paradigm, and assess baseline sensitivity,

participants were first exposed to a series of resistances in ascending order (0, 10, 20, 40, 60, and

80 cmH2O/L/sec) applied for 60 seconds each. Participants were asked to rate their anxiety level

after each exposure from 0 ("no anxiety") to 10 ("maximum possible anxiety"). Other questions

relating to subjective breathing difficulty, arousal, and other affective states were also presented

(see **Supplementary Methods**).

This induction was later applied using a moderate resistance level (40 cmH2O/L/sec) during

performance of one run of the behavioral task. This level of resistance was chosen because of

prior work demonstrating its effectiveness and feasibility (41); we also confirmed that it induced

moderate (roughly 5 out of 10) levels of anxiety within our clinical sample (**Supplementary**

Figure S1).

Figure 1. A) Equipment used for anxiety induction including silicon mask with adjustable straps and single breathing port, and an example resistor used to create resistance on inhalation. B) Graphical depiction of the Horizon Task. Shown are the two horizon conditions (H1 and H6) at the last forced choice (red box) and the first free choice (green boxes). The H1 example also shows an equal information trial (2 outcomes for each choice), while the H6 example shows an unequal information trial (3 forced choices on the right, 1 on the left. C) Table outlining each of the trial type combinations, counterbalanced across the task.

Behavioral Task

Participants completed two runs of the Horizon Task (42), a previously validated task used to measure explore-exploit decision-making. During each task run, players repeatedly choose between two slot machines in 80 games. Each game involves either 5 or 10 sequential choices. The first four choices in all games are "forced", meaning the player is told which option to choose (see **Figure 1b**). The remaining choices are "free", where either option can be chosen. Trials with one free choice (H1) and those with six free choices (H6) appear an equal number of times throughout the task.

Forced-choice patterns create two different information conditions: *equal*, in which each option is chosen twice; and *unequal,* in which one option is chosen three times and the other is chosen only one time. The goal of the task is to maximize the number of points won by repeatedly

choosing the option expected to provide higher point values on average. Point values were

sampled (and rounded to the nearest integer) from a Gaussian distribution with a mean of either

of 40 or 60 on one option (standard deviation of 8). Then, for each game, the other option had a

mean point value difference of +/- 4, 8, 12, 20, or 30 from the first. The same pseudorandom

sequences of forced-choice outcomes were shown to each participant, consistent with the

generative means. All game dimensions (length, information type, mean difference, and

better/worse side) were counterbalanced across the task (**Figure 1c**).

Participants wore the mask for each task run. In the *no-resistance run*, resistance was not added

(0 cmH2O/L/sec). In the *resistance run*, a moderate resistance (40 cmH2O/L/sec) was

consistently applied throughout the run. The order was counterbalanced across participants in

each group. The first six participants played a version of the task with different outcome values

than the rest in a small number of games (but still sampled from the same underlying

distributions). To ensure this did not confound results below, this difference in task version was

evaluated as a potential covariate.

Computational Model and Model Fitting

The computational model used here is outlined in Zajkowski, Kossut (43). We present this in

detail in **Supplementary Methods**. Brief definitions of each computational parameter are

provided in **Table 2**.

Table 2. Descriptions of model parameters.

Model Parameter	General Description
Information Bonus (IB)	IB parameters (separated by horizon condition to yield IB1 and IB6) quantify an individual's tendency to choose the option that maximizes information gain. This parameter only affects decisions in the unequal information condition, where selecting the option that was chosen only once during the forced choices provides more information about its average reward value. Directed exploration (DE) is derived from these parameters (IB6 minus IB1). This is because information gain in H1 cannot guide future choices, while information gain in H6 can guide future choices. Thus, DE can be seen as the tendency to increase information-seeking when it would be useful.
Decision Noise (DN)	The DN parameters (separated by horizon condition to yield DN1 and DN6) quantify an individual's tendency to choose the option with a lower observed mean reward. Random exploration (RE) is derived by subtracting DN1 from DN6

	in the equal information condition. Thus, RE can be seen as the tendency to make less reward-sensitive choices when information gain would be useful.
Spatial Bias	These parameters (one per horizon/information condition) account for the possibility that an individual prefers to choose one option over the other simply because of which side it is on.
Initial Learning Rate (α_0)	The initial value of the learning rate before making the first forced choice.
Asymptotic Learning Rate (α_∞)	The learning rate an individual would converge to if the game were played indefinitely (i.e., when uncertainty about mean reward values would be minimal).

Statistical Analyses

Resistance sensitivity

214   To verify that the anxiety induction was effective, we examined affective responses to each

215   resistance level in the initial exposure protocol. To this end, we estimated linear mixed effects

216   models (LMEs) with self-reported anxiety (i.e., the single anxiety question on a scale from 0 to

217   10) as the outcome variable and resistance level and group as predictors (and their interaction).

Primary model-based analyses

Before addressing our primary hypotheses, we checked for outliers using an iterative Grubbs

method (threshold: *p<*.01; using *grubbs.test* from the *outliers* package in R (44)). This resulted in

one data point being removed for IB1 (in the resistance condition), two for DN6 in the equal

information condition (one in each resistance condition), and two for RE (one in each resistance

condition). Then, we estimated LMEs that included group, resistance condition, and their

interaction as sum-coded predictors for each model parameter (HCs=-1, iADs=1; no-resistance=-

1, resistance=1). All models predicting $\boldsymbol{\alpha_\infty}$ also accounted for $\boldsymbol{\alpha_0}$ values. To rule out possible

alternative explanations, we also re-ran these models including covariates to confirm that any

group or resistance effects could not be explained by age, sex (Male=-1, Female=1), or task

version (new=-1, old=1).

We then assessed continuous relationships between computational parameters and affective

symptoms. We first tested separate models using PHQ scores or STAI Trait scores. These LMEs

contained interactions with resistance condition and any covariates found to be significant

predictors in the group models. Then, we included both symptom measures in the same models

(and their interactions with resistance) to establish potentially independent effects.

Model-free task performance

To better interpret observed effects on task behavior and assess relationships between task

performance and model parameters, we also performed additional supportive analyses examining

overall accuracy on free choices (as measured by the number of times participants chose the

option with the higher underlying mean reward). We first restricted analyses to first free choice

and estimated an LME predicting accuracy based on horizon, information condition, resistance,

and group as predictors. We also included possible three-way interactions between group,

horizon, and information condition, and between group, horizon, and resistance (and associated

two-way interactions). This allowed us to explore whether groups might show greater differences

depending on horizon, information condition, or anxiety induction. As commonly observed in

this task, we expected accuracy in H1 trials would be higher than in H6 trials (i.e., reflecting

random exploration).

A subsequent model was run predicting accuracy across free choice trials in H6 (choices 5-10).

We also tested three-way interactions between group, free choice number, and information

condition, and between group, free choice number, and resistance (and associated two-way

interactions) to explore whether groups might differ in the slope of accuracy improvement over

time depending on information condition or anxiety induction.

To evaluate whether some parameter values might be considered more optimal than others, we

then tested LMEs with parameters predicting accuracy on H6 free choice trials, excluding the

first free choice to which model parameters were fit. As information bonus (IB) is only

calculated in the unequal information condition, accuracy on the unequal information trials alone

was predicted in these models, with resistance condition, group, free choice number, IB for H6

(IB6), and the interaction between free choice number and IB6 as predictors. Analogous models

were used to test relationships between accuracy and the other model parameters within the

relevant trial types (i.e., decision noise for H6 with equal information condition accuracy;

learning rates and accuracy in both information conditions).

Secondary replication analyses

Additional LMEs were run to accomplish our secondary aim of replicating prior relationships

found between DE and both early adversity (16, 17) and cognitive reflectiveness (14). First,

separate LMEs were run predicting DE based on each subscale of the CTQ, while also including

group, resistance condition, and their interaction as predictors (while controlling for known

effects of age). Analogous LMEs were run instead using CRT scores as predictors.

266   **Results**

The Breathing Resistance Protocol Successfully Induced Anxiety During Task Performance

Anxiety ratings during the initial breathing resistance sensitivity protocol are shown in

Supplementary Figure S1. Briefly, anxiety increased as the resistance level increased, anxiety

ratings were higher in iADs than HCs, and iADs showed greater increases in anxiety compared

to HCs as resistance levels increased. LMEs predicting self-reported anxiety scores at baseline

and during both task runs showed the same pattern (**Figure 2**). See **Supplementary Materials**

for full results.

Figure 2. Self-reported anxiety and STAI State at baseline and during each run of the Horizon
Task (i.e., with and without the added breathing resistance of 40 cmH2O/L/sec). Results showed
greater anxiety in iADs, greater anxiety with higher resistance level, and greater increases in
anxiety in iADs than HCs when resistance was added. Boxplots show median and quartile values
along with individual datapoints.

Directed Exploration was Elevated in Affective Disorders Patients, but Reduced by State Anxiety Induction

Plots depicting parameter values by group and condition are shown in **Figure 3**. In an initial LME predicting DE, there was a significant *Group x Resistance Condition* interaction ($F(1,117)=3.98$, $p=.048$, $\eta_p^2=.03$), such that iADs showed higher DE in the no-resistance condition (estimated marginal mean [EMM]=5.69) than in the resistance condition (EMM=4.89; $t(117)=-2.09$, $p=.039$), while DE in HCs remained at lower values and did not change between conditions ($t(117)=0.75$, $p=.456$). Further exploration suggested this was driven primarily by greater IB values in iADs compared to HCs in H6 games within the no-resistance condition (see **Supplementary Figure S2**). Main effects of group or resistance were nonsignificant. In a subsequent model accounting for age, sex, and task version, this interaction remained significant ($F(1,117)=3.98$, $p=.048$, $\eta_p^2=.03$). There was also a negative association with age (b=-0.06; $F(1,114)=5.65$, $p=.019$, $\eta_p^2=.05$).

Figure 3. Individual data points and density plots by group and resistance condition for each parameter. HCs showed stable DE values in both conditions. Those with ADs instead showed elevated DE in the no-resistance condition, which then decreased after application of the resistance. For illustrative purposes, the individual information bonus (IB) values underlying DE for each horizon condition (H1 and H6) are shown in the top-right (recall that DE corresponds to the difference in information bonuses between horizon conditions; DE = IB6 – IB1). As can be seen, the H6 information bonus is higher in the no-resistance condition than in the resistance condition for iADs only. RE did not differ by group or condition (bottom left). Learning rates tended to show lower values in iADs than HCs across conditions. As visible in the lower middle panel, α_0 showed a bimodal distribution, which motivated the cluster-based analysis approach described in the text. Asymptotic learning rates (α_∞) were significantly lower in iADs than HCs in both conditions (bottom right). Note that stars and Cohen's *d* effect sizes correspond to the interaction between group and resistance for DE (upper left) and for the group difference in α_∞ across resistance conditions (bottom right) in post-hoc comparisons.

No significant effects were found in LMEs predicting RE ($Fs \leq 3.45$, $ps \geq .067$).

Because of bimodality in the distributions for α_0 (see **Figure 3**), we used *k*-means clustering (46)

across resistance conditions to divide participants into groups with high vs. low values and

treated this as a 2-level categorical variable (low=-1; high=1). In logistic regression models

predicting high vs. low α_0 values (using *glmer* in the *lme4* package; (47)), there were no

significant effects before or after adding covariates (zs≤|1.60|, *ps*≥.110).

The LME predicting α_∞ (accounting for α_0) revealed a significant main effect of group

($F(1,117)=7.73$, $p=.006$, $\eta_p^2=.06$; see **Figure 3**), such that HCs (EMM=.33) had higher learning

rates than iADs (EMM=.28, $t(117)=-2.79$, $p=.006$). The expected positive association with α_0

was also observed ($F(1,200)=15.64$, $p<.001$, $\eta_p^2=.07$; b=.11). Both effects remained significant

when accounting for covariates, and no additional effects were observed ($Fs≤1.73$, $ps≥.192$).

We also evaluated within-group associations with symptom severity. This was done to explore

potentially differential effects of depression and anxiety, noting that associated measures (PHQ

and STAI Trait) were highly correlated (*rs*=.83-.86; see **Supplementary Figure S3**). In models

predicting DE in iADs alone (based on PHQ, STAI Trait, or both measures in a single model) no

significant effects were found. However, marginal effects of resistance condition recapitulated

the *Group x Resistance* interaction effects observed in our primary analyses above ($Fs≥3.76$,

.05<*p*≤.058).

Within HCs, models predicting DE with either PHQ or STAI Trait showed no significant effects.

However, when both symptom measures were included in the same model, there was a

significant interaction between resistance condition and STAI Trait ($F(1,55)=5.48$, $p=.023$,

$\eta_p^2=.09$). This reflected a greater positive association between anxiety and DE in the no-

resistance condition than in the resistance condition (no-resistance: estimated trend [ET]=.236;

resistance: ET=.072; $t(55)=2.34$, $p=.023$), with a nonsignificant positive association with STAI

Trait overall ($F(1,54)=2.85$, $p=.097$, $\eta_p^2=.05$). PHQ instead showed a marginal negative

association with DE that interacted with resistance ($F(1,55)=3.43$, $p=.069$, $\eta_p^2=.06$), suggesting

this negative relationship was stronger in the no-resistance condition (no-resistance: ET=-.484;

resistance: ET=-.060; $t(55)=-1.85$, $p=.070$).

Equivalent analyses were also carried out for other model parameters, which suggested potential

opposing associations between PHQ, STAI Trait, and RE when included in the same model

(detailed in **Supplementary Results**).

Higher Levels of Directed and Random Exploration are Associated with Greater Task

Performance

In LMEs predicting H6 accuracy (on choices 6-10) using each model parameter, we found main

effects of learning rates (**Table 3**). Specifically, α_0 predicted accuracy on equal information

trials, while α_∞ predicted accuracy on unequal information trials. Additionally, there was an

interaction between IB6 and free choice number indicating that, in unequal information trials,

those with higher IB6 showed steeper increases in accuracy from early to later choices. Similar

results were found for decision noise when predicting accuracy on equal information trials.

We also tested an analogous model using DE as a predictor of free choice accuracy and saw

comparable results to those seen for IB6 (i.e., main effect of free choice number and a significant

interaction with DE; $Fs \geq 9.69$, $ps \leq .002$). Similarly, results when using RE to predict equal

information accuracy corroborated those found for decision noise (main effect of free choice:

$F(1,1057)=114.74$, $p<.001$; *RE x free choice*: $F(1,1057)=15.78$, $p<.001$; main effect of group:

$F(1,117)=5.67$, $p=.019$). Finally, motivated by the effect of resistance on DE observed above in

iADs, we tested a model including a *DE x Resistance Condition* interaction term. This revealed a

significant interaction ($F(1,1089)=4.46$, $p=.035$), such that greater DE was positively predictive

of accuracy in the resistance condition (ET=0.002) but not in the no-resistance condition (ET=-

0.001; $t(1089)=2.11$, $p=.035$). Jointly, these results suggest higher values for DE were more

optimal.

Table 3. Free choice accuracy (separated by information condition) predicted by model parameters.

	Parameter	Resistance	Group	Free Choice Number	Parameter*Free Choice
IB6 (unequal IC)	$F(1,992)=0.01$, $p=.913$	$F(1,1071)=0.03$, $p=.853$	$F(1,117)=1.82$, $p=.181$	**$F(1,1067)=91.96$, $p<.001$; b=0.012**	**$F(1,1067)=9.89$, $p=.002$; b=0.001**
DN6 (equal IC)	$F(1,1174)=1.54$, $p=.215$	$F(1,1059)=2.66$, $p=.103$	**$F(1,117)=5.65$, $p=.019$**	**$F(1,1057)=113.83$, $p<.001$; b=0.007**	**$F(1,1057)=9.93$, $p=.002$; b=0.03**
α_0 Cluster (equal IC)	**$F(1,1167)=4.54$, $p=.033$; b=0.021**	$F(1,1068)=2.84$, $p=.092$	**$F(1,117)=5.65$, $p=.019$**	**$F(1,1067)=114.37$, $p<.001$; b=0.022**	$F(1,1067)=1.37$, $p=.242$
α_0 Cluster (unequal IC)	$F(1,1165)=0.94$, $p=.332$	$F(1,1068)=0.01$, $p=.909$	$F(1,117)=1.77$, $p=.186$	**$F(1,1067)=91.36$, $p<.001$;**	**$F(1,1067)=6.26$, $p=.013$; b= -0.009**

				b=0.021	
α_{∞} (equal IC)	$F(1,1301)=1.18$, $p=.278$	$F(1,1309)=2.21$, $p=.137$	**$F(1,119)=4.90$, $p=.029$**	**$F(1,1304)=86.60$, $p<.001$; b=0.015**	$F(1,1304)=0.20$, $p=.656$
α_{∞} (unequal IC)	**$F(1,1194)=18.88$, $p<.001$; b=0.145**	$F(1,1310)=0.01$, $p=.928$	$F(1,120)=1.06$, $p=.305$	**$F(1,1304)=128.71$, $p<.001$; b=0.025**	**$F(1,1304)=12.67$, $p<.001$; b= -0.038**

361 **Note.** For each significant effect of group, contrasts indicated that accuracy in the iADs (EMM=0.80) was
362 lower than accuracy in the HCs (EMM=0.85; $p<.05$). The models predicting accuracy including α_{∞} also
363 saw positive effects of α_0 ($Fs \geq 6.77$, $ps<.01$, bs$\geq$0.046). IC=information condition.

Task Performance was Reduced in Affective Disorders in Specific Conditions

Secondary analyses of free choice accuracy indicated high performance across both groups

(HCs: M=.84, SD=.13; iADs: M=.80, SD=.16; see **Supplementary Figure S4**). LMEs

predicting first free choice accuracy (**Supplementary Table S3**) revealed significant effects of

horizon type (greater accuracy in H1; $F(1,823)=278.81$, $p<.001$) and information condition

(greater accuracy in equal information trials; $F(1,823)=72.08$, $p<.001$). The *Group x Horizon x*

Information Condition interaction was also significant ($F(1,823)=4.18$, $p=.041$), indicating that

HCs had significantly greater accuracy than iADs in H1/equal information trials and in

H6/unequal information trials only (see post-hoc contrasts). Subsequent LMEs predicting

accuracy across free choices in H6 showed significant effects of choice number (greater accuracy

in later choices; $F(1,2727)=201.30$, $p<.001$), information condition (greater accuracy in equal

information trials; $F(1,2727)=51.37$, $p<.001$), and a *Group x Information Condition* interaction

(greater accuracy in HCs only in the equal information condition; $F(1,2727)=8.93$, $p=.003$).

Replication Analyses Confirm Associations with Early Adversity and Cognitive Reflectiveness

CTQ scores were greater in iADs than HCs ($t(112)=5.31$, $p<.001$). In separate LMEs predicting

DE with each subscale of the CTQ, group, resistance condition, and their interaction as

predictors (also controlling for age), and after correcting for multiple comparisons ($p \leq .01$), there

was a significant negative association with physical abuse (b=-.26; $F(1,109)=6.96$, $p=.010$,

$\eta_p^2=.06$) such that exploration was lower in those who experienced greater physical abuse. This

remained significant in iADs alone (b=-.32; $F(1,56)=7.27$, $p=.009$, $\eta_p^2=.11$). No effects were

observed for any other parameter.

CRT scores were lower in iADs than HCs ($t(117)=3.05$, $p=.003$). In an LME assessing the effect

of CRT on DE across all participants, there was a significant positive association (b=.32;

$F(1,116)=5.06$, $p=.026$, $\eta_p^2=.04$). The effect of CRT was directionally the same in iADs alone

(and stronger in effect size), but only marginally significant due to smaller sample size (b=.40;

$F(1,58)=3.89$, $p=.053$, $\eta_p^2=.06$). There were no significant effects of CRT on RE or α_∞ in either

the full sample or in the clinical group alone ($ps>.054$). CRT scores positively predicted α_0

cluster membership in the full sample (z=-2.91, $p=.004$; log-odds=-.41, CI=[-0.68,-0.13]), but

not in iADs alone (z=-1.57, $p=.115$; log-odds=-.38, CI=[-0.86,0.09])).

396   **Discussion**

In this study, we compared decision-making behavior on an explore-exploit task in participants

with and without clinically elevated levels of trait anxiety and depression. A somatic

(interoceptive) anxiety induction was also used to allow dissociation of the influences of state vs.

trait anxiety. We hypothesized that state anxiety induction would reduce direction exploration

(DE) as a potential mechanism promoting avoidance behavior. Results were mixed, offering only

partial support for our primary hypotheses. In particular, there was a significant interaction in

which elevated DE in iADs at baseline was reduced by the anxiety induction, while DE in HCs

remained stable between conditions. The elevated DE at baseline suggests more persistent and

exaggerated uncertainty in iADs, similar to that observed in some previous work in non-clinical

samples (29). The reduction in DE after anxiety induction also appears consistent with

correlational work suggesting anxiety and depression are associated with less exploration (14), as

well as with work showing that stress reduces exploration (21-23). It is also somewhat in line

with a previous study showing lower DE in those with higher trait somatic anxiety (15). Thus,

our results show partial consistency with findings of multiple previous studies and suggest a

general differentiation between state and trait effects.

While the reduction in DE with state anxiety induction in iADs is consistent with our initial

hypothesis, the expected decrease in HCs was not observed. This suggests the change in iADs

could be viewed as a normalization toward healthy levels. On the other hand, higher levels of DE

were also associated with steeper improvements in task performance over time, suggesting the

anxiety induction had a maladaptive effect. This interpretation is also not entirely supported,

however, as choice accuracy was not significantly different between resistance conditions, and

task performance was marginally worse in iADs overall. This would therefore suggest greater

exploratory behavior in the no-resistance condition did not in fact improve performance, or that

there were other counteracting factors (e.g., differences in learning rates). Thus, differences in

DE did not relate to optimality in task performance in a clear way.

The finding that iADs showed slower asymptotic learning rates across conditions may offer

some additional explanatory insights. First, depression and anxiety symptoms (and particularly

trait anxiety levels in the no-resistance condition) were negatively associated with asymptotic

learning rates, while these learning rates were positively associated with task performance. This

suggested that slower learning may have contributed to worse performance in iADs. Here,

slower learning implies that beliefs remained closer to uninformative prior values; thus,

confidence in the better choice would increase more slowly (similar to less flexible learning rates

previously associated with higher anxiety; (5)). Within this task, slower learning rates might thus

support a type of persistent uncertainty about the best course of action, with suboptimal effects

on decision-making. Interpreting this result in light of previous findings is subtle, however, as

faster learning rates are theoretically linked to greater uncertainty, and prior work has linked

anxiety to both faster learning (6) and a greater tendency to infer changes in context (8). On the

other hand, the type of uncertainty in these studies pertains to volatility, or how frequently

environmental contingencies are expected to change. Thus, if anxious individuals believe the

world is ever-changing, then learning rates should be high.

However, if "uncertainty" instead pertains to the estimated stochasticity of (i.e., noise in) the

mapping from underlying states to observations, learning rate should instead be low, so as not to

overfit beliefs to random outcomes (48). Thus, in the present task, slower learning in iADs could

represent greater uncertainty about the informativeness of each outcome when inferring the

underlying reward mean. This could offer a complementary means by which uncertainty is

maintained in anxious individuals, consistent with the previous findings reviewed above. It could

also relate to another recent study showing that individuals who experienced greater early

adversity, itself a predictor of subsequent affective disorders (49-51), also showed a slower

learning rate (16). It should be kept in mind, however, that our sample includes individuals with

affective disorders, while most prior work has examined sub-clinical anxiety levels. Thus, some

results between studies may not be fully comparable. This was suggested by our follow-up

results in which, for HCs, DE showed potentially opposing positive associations with trait

anxiety and negative associations with depression in the no-resistance condition (i.e., when both

symptom measures were included in the same models), while no such relationships were found

in the clinical sample.

In line with our secondary aims, we were also able to successfully replicate prior results showing

higher DE in those with greater cognitive reflectiveness (14) and lower DE in those who

experienced greater childhood adversity (physical abuse) and environmental unpredictability (16,

17). These findings suggest that less reflection on uncertainty and greater exposure to

unpredictable early environments could each contribute to clinical differences seen here in

exploration and learning. It is also worth noting that cognitive reflectiveness has been shown to

improve with training (52-54). Thus, future studies could examine whether improving

reflectiveness also optimizes exploratory behavior or reduces affective symptoms. Another

consideration is that multiple previous studies have shown strong relationships between physical

abuse in childhood and later avoidance behavior (55, 56). Results here suggest associated

differences in DE could contribute to this avoidance behavior, perhaps especially in relation to

somatic anxiety and physical abuse, potentially highlighting DE as a possible treatment target for

those recovering from childhood trauma.

There are important limitations to consider. First, the breathing resistance only induced mild

anxiety in HCs and moderate anxiety in iADs. While we intentionally chose a tolerable level of

resistance for patients, it is possible that effects would be larger with a stronger induction

approach. Additionally, the sample size was only moderate and may not have afforded sufficient

power to detect some effects within the clinical group alone. Future research will also be needed

to see whether results generalize to other explore-exploit tasks as well as tasks designed to

distinguish learning rates in relation to volatility vs. stochasticity (48).

Conclusions

The results of this study suggest elevated directed exploration and slower learning rates in

individuals with affective disorders, as well as a causal role of somatically focused state anxiety

in reducing directed exploration in this population. It also replicates potential links to cognitive

reflection and early adversity that may be of clinical relevance. These findings highlight potential

computational mechanisms underlying both persistent uncertainty and maladaptive avoidance. If

confirmed in future work, this could suggest potential benefits of treatments aimed at optimizing

levels of information-seeking and belief testing in relation to current affective states as well as

adjusting beliefs about the reliability of new experiences in revising expectations.

Conflict of interest or competing financial interests

482   The authors have no competing interests to disclose.

Funding

484   This project was funded by National Institute of General Medical Sciences (P20GM121312 [R.S.

485   and M.P.P.]) and the Laureate Institute for Brain Research.

Author contributions

487   NL and CAL took the lead in data analysis and writing the manuscript. RS designed the study,

488   planned and oversaw analyses, and played a major role in writing the manuscript. ST, AEC,

489   CMG, and TT assisted analyses and in data collection. All authors contributed to the manuscript.

References

1.   Barlow DH, Allen LB, Choate ML. Toward a unified treatment for emotional disorders - republished article. Behav Ther. 2016;47(6):838-53.

2.   Hayes S, Strosahl K, Wilson K. Acceptance and commitment therapy: An experiential approach to behaviour change. 2003.

3.   Sharp PB, Eldar E. Computational models of anxiety: nascent efforts and future directions. Current Directions in Psychological Science. 2019;28(2):170-6.

4.   Lloyd A, Roiser JP, Skeen S, Freeman Z, Badalova A, Agunbiade A, et al. Reviewing explore/exploit decision-making as a transdiagnostic target for psychosis, depression, and anxiety. Cogn Affect Behav Neurosci. 2024.

5.   Browning M, Behrens TE, Jocham G, O'Reilly JX, Bishop SJ. Anxious individuals have difficulty learning the causal statistics of aversive environments. Nat Neurosci. 2015;18(4):590-6.

6.   Huang H, Thompson W, Paulus MP. Computational dysfunctions in anxiety: failure to differentiate signal from noise. Biol Psychiatry. 2017;82(6):440-6.

7.   Pike AC, Robinson OJ. Reinforcement learning in patients with mood and anxiety disorders vs control individuals: A systematic review and meta-analysis. JAMA psychiatry. 2022;79(4):313-22.

8.   Zika O, Wiech K, Reinecke A, Browning M, Schuck NW. Trait anxiety is associated with hidden state inference during aversive reversal learning. Nat Commun. 2023;14(1):4203.

9.   Wilson RC, Bonawitz E, Costa VD, Ebitz RB. Balancing exploration and exploitation with information and randomization. Curr Opin Behav Sci. 2021;38:49-56.

10.   Schulz E, Gershman SJ. The algorithmic architecture of exploration in the human brain. Current opinion in neurobiology. 2019;55:7-14.

11.   Addicott MA, Pearson JM, Wilson J, Platt ML, McClernon FJ. Smoking and the bandit: a preliminary study of smoker and nonsmoker differences in exploratory behavior measured with a multiarmed bandit task. Exp Clin Psychopharmacol. 2013;21(1):66-73.

12.   Addicott MA, Pearson JM, Froeliger B, Platt ML, McClernon FJ. Smoking automaticity and tolerance moderate brain activation during explore-exploit behavior. Psychiatry Res. 2014;224(3):254-61.

13.   Addicott MA, Pearson JM, Sweitzer MM, Barack DL, Platt ML. A Primer on Foraging and the Explore/Exploit Trade-Off for Psychiatry Research. Neuropsychopharmacology. 2017;42(10):1931-9.

14.   Smith R, Taylor S, Wilson RC, Chuning AE, Persich MR, Wang S, et al. Lower levels of directed exploration and reflective thinking are associated with greater anxiety and depression. Frontiers in Psychiatry. 2022;12:782136.

15.   Fan H, Gershman SJ, Phelps EA. Trait somatic anxiety is associated with reduced directed exploration and underestimation of uncertainty. Nature Human Behaviour. 2023;7(1):102-13.

16.   Lloyd A, McKay RT, Furl N. Individuals with adverse childhood experiences explore less and underweight reward feedback. Proceedings of the National Academy of Sciences. 2022;119(4):e2109373119.

17.   Xu Y, Harms MB, Green CS, Wilson RC, Pollak SD. Childhood unpredictability and the development of exploration. Proceedings of the National Academy of Sciences. 2023;120(49):e2303869120.

18.   Chorpita BF, Barlow DH. The development of anxiety: The role of control in the early environment. The Neurotic Paradox, Vol 2. 2018:227-64.

19.   Nelson CA, Gabard-Durnam LJ. Early adversity and critical periods: neurodevelopmental consequences of violating the expectable environment. Trends in neurosciences. 2020;43(3):133-43.

20.   Spadoni AD, Vinograd M, Cuccurazzu B, Torres K, Glynn LM, Davis EP, et al. Contribution of early‐life unpredictability to neuropsychiatric symptom patterns in adulthood. Depression and anxiety. 2022;39(10-11):706-17.

21.   Biedermann SV, Biedermann DG, Wenzlaff F, Kurjak T, Nouri S, Auer MK, et al. An elevated plus-
maze in mixed reality for studying human anxiety-related behavior. BMC Biol. 2017;15(1):125.
22.   Walz N, Muhlberger A, Pauli P. A Human Open Field Test Reveals Thigmotaxis Related to
Agoraphobic Fear. Biol Psychiatry. 2016;80(5):390-7.
23.   Lenow JK, Constantino SM, Daw ND, Phelps EA. Chronic and Acute Stress Promote
Overexploitation in Serial Decision Making. J Neurosci. 2017;37(23):5681-9.
24.   Teigen K. Yerkes-Dodson: A law for all seasons. Theory and Psychology. 1994;4:525-47.
25.   Arnsten AF. Stress weakens prefrontal networks: molecular insults to higher cognition. Nat
Neurosci. 2015;18(10):1376-85.
26.   Robbins TW, Arnsten AF. The neuropsychopharmacology of fronto-executive function:
monoaminergic modulation. Annu Rev Neurosci. 2009;32:267-87.
27.   Arnsten A, Robbins T. Neurochemical modulation of prefrontal cortical function in humans and
animals. Principles of frontal lobe function2002. p. 51-84.
28.   Arnsten AF. Catecholamine modulation of prefrontal cortical cognitive function. Trends Cogn Sci.
1998;2(11):436-47.
29.   Aberg KC, Toren I, Paz R. A neural and behavioral tradeoff underlies exploratory decisions in
normative anxiety. Molecular Psychiatry. 2022;27:1573–87.
30.   Witte K, Wise T, Huys QJ, Schulz E. Exploring the Unexplored: Worry as a Catalyst for Exploratory
Behavior in Anxiety and Depression. 2024.
31.   Blanco NJ, Otto AR, Maddox WT, Beevers CG, Love BC. The influence of depression symptoms on
exploratory decision-making. Cognition. 2013;129(3):563-8.
32.   Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-International
Neuropsychiatric Interview (MINI): the development and validation of a structured diagnostic psychiatric
interview for DSM-IV and ICD-10. Journal of clinical psychiatry. 1998;59(20):22-33.
33.   Campbell-Sills L, Norman SB, Craske MG, Sullivan G, Lang AJ, Chavira DA, et al. Validation of a
brief measure of anxiety-related severity and impairment: the Overall Anxiety Severity and Impairment
Scale (OASIS). Journal of affective disorders. 2009;112(1-3):92-101.
34.   Zhang Z, Mai Y. WebPower: Basic and Advanced Statistical Power Analysis. R package version
0.9.4 ed2023.
35.   Spielberger C, Gorsuch R, Lushene R, Vagg P, Jacobs G. Manual for the State-Trait Anxiety
Inventory; Palo Alto, CA, Ed. Palo Alto: Spielberger. 1983.
36.   Sun Y, Fu Z, Bo Q, Mao Z, Ma X, Wang C. The reliability and validity of PHQ-9 in patients with
major depressive disorder in psychiatric hospital. BMC Psychiatry. 2020;20:1-7.
37.   Bernstein DP, Stein JA, Newcomb MD, Walker E, Pogge D, Ahluvalia T, et al. Development and
validation of a brief screening version of the Childhood Trauma Questionnaire. Child abuse & neglect.
2003;27(2):169-90.
38.   Stewart JL, May AC, Poppa T, Davenport PW, Tapert SF, Paulus MP. You are the danger:
attenuated insula response in methamphetamine users during aversive interoceptive decision-making.
Drug Alcohol Depend. 2014;142:110-9.
39.   Kruschwitz JD, Kausch A, Brovkin A, Keshmirian A, Paulus MP, Goschke T, et al. Self-control is
linked to interoceptive inference: Craving regulation and the prediction of aversive interoceptive states
induced with inspiratory breathing load. Cognition. 2019;193:104028.
40.   Berk L, Stewart JL, May AC, Wiers RW, Davenport PW, Paulus MP, et al. Under pressure:
adolescent substance users show exaggerated neural processing of aversive interoceptive stimuli.
Addiction. 2015;110(12):2025-36.
41.   Paulus MP, Flagan T, Simmons AN, Gillis K, Kotturi S, Thom N, et al. Subjecting elite athletes to
inspiratory breathing load reveals behavioral and neural signatures of optimal performers in extreme
environments. PLoS one. 2012;7(1):e29394.

42.   Wilson RC, Geana A, White JM, Ludvig EA, Cohen JD. Humans use directed and random
exploration to solve the explore–exploit dilemma. Journal of Experimental Psychology: General.
2014;143(6):2074.
43.   Zajkowski WK, Kossut M, Wilson RC. A causal role for right frontopolar cortex in directed, but not
random, exploration. Elife. 2017;6:e27430.
44.   Komsta L. outliers: Tests for Outliers. R package version 0.15 ed2022.
45.   Goldman CM, Takahashi T, Lavalley CA, Li N, Taylor S, Chuning AE, et al. Individuals with
Methamphetamine Use Disorder Show Reduced Directed Exploration and Learning Rates Independent of
an Aversive Interoceptive State Induction. medRxiv. 2024:2024.05.17.24307491.
46.   Jain AK, Dubes RC. Algorithms for clustering data: Prentice-Hall, Inc.; 1988.
47.   Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest package: tests in linear mixed effects
models. Journal of statistical software. 2017;82(13).
48.   Piray P, Daw ND. A model for learning based on the joint estimation of stochasticity and volatility.
Nat Commun. 2021;12(1):6587.
49.   Heim C, Newport DJ, Mletzko T, Miller AH, Nemeroff CB. The link between childhood trauma and
depression: insights from HPA axis studies in humans. Psychoneuroendocrinology. 2008;33(6):693-710.
50.   Heim C, Binder EB. Current research trends in early life stress and depression: review of human
studies on sensitive periods, gene-environment interactions, and epigenetics. Exp Neurol.
2012;233(1):102-11.
51.   Springer KW, Sheridan J, Kuo D, Carnes M. Long-term physical and mental health consequences
of childhood physical abuse: results from a large population-based sample of men and women. Child
Abuse Negl. 2007;31(5):517-30.
52.   Lachaud L, Jacquet B, Bourlier M, Baratgin J. Mindfulness-based stress reduction is linked with an
improved Cognitive Reflection Test score. Frontiers in Psychology. 2023;14:1272324.
53.   Calvillo DP, Bratton J, Velazquez V, Smelter TJ, Crum D. Elaborative feedback and instruction
improve cognitive reflection but do not transfer to related tasks. Thinking & Reasoning. 2023;29(2):276-
304.
54.   Orona GA, Trautwein U. Thinking Disposition Education Improves Cognitive Reflection:
Experimental Results from Two Intervention Studies. 2023.
55.   Fiorillo D, Papa A, Follette VM. The relationship between child physical abuse and victimization in
dating relationships: The role of experiential avoidance. Psychological trauma: theory, research, practice,
and policy. 2013;5(6):562.
56.   Unger JAM, De Luca RV. The relationship between childhood physical abuse and adult
attachment styles. Journal of family violence. 2014;29:223-34.
619
620

621 # Supplementary Materials

622 ## **Supplementary Methods**

623 **Table S1. Sample Demographics**

Demographics	Healthy Comparisons (n = 58)	Affective Disorders (n = 61)	Sample Comparison
Age (mean (SD))	35.41 (13.08)	33.44 (10.97)	$t(117) = 0.89, p = 0.374$
Gender Identity = Woman (counts)	42 (72.4%)	46 (75.4%)	$\chi^2(1) = 0.03, p = .870$
Sexual Orientation (counts)			$\chi^2(3) = 4.07, p = 0.254$
Bisexual	3 (5.2%)	5 (8.5%)	
Lesbian	0 (0.0%)	3 (5.1%)	
None of these describe me	1 (1.7%)	2 (3.4%)	
Straight, that is, not gay or lesbian, etc.	54 (93.1%)	49 (83.1%)	
Education (counts)			$\chi^2(7) = 4.47, p = 0.724$
GED or equivalent	0 (0.0%)	1 (1.6%)	
High school graduate	1 (1.7%)	4 (6.6%)	
Some college, no degree	6 (10.3%)	8 (13.1%)	
Associate's degree: occupational, technical, or vocational program	4 (6.8%)	5 (8.2%)	
Associate's degree: academic program	3 (5.2%)	4 (6.6%)	
Bachelor's degree	25 (43.1%)	18 (29.5%)	
Master's degree	7 (12.1%)	7 (11.5%)	
Doctoral degree	1 (1.7%)	1 (1.6%)	
Information Unavailable	11 (19.0%)	13 (21.3%)	
Ethnicity = Not Hispanic/Latino (counts)	52 (89.7%)	54 (88.5%)	$\chi^2(1) = 0, p = 1$
Race (counts)			$\chi^2(5) = 1.327, p = 0.932$
American Indian or Alaska Native	8 (13.8%)	8 (13.6%)	
Asian	4 (6.9%)	3 (5.1%)	
Black or African American	3 (5.2%)	4 (6.8%)	

Native Hawaiian or Pacific Islander	1 (1.7)	0 (0.0)
Other	3 (5.2)	3 (5.1)
White	39 (67.2)	41 (69.5)

Table S2. Diagnostic breakdown of individuals with high anxiety (N=61).

iADs (*N*=61)	MDD	GAD	Social	PTSD	Panic
MDD	**58***				
GAD	20	**22**			
Social	12	6	**13**		
PTSD	12	4	1	**12**	
Panic	1	0	0	1	**1**

Note. MINI diagnoses were given up to 12 months prior to study participation. Participants in this group were required to have a lifetime diagnosis of one or more anxiety/depressive disorder(s) and OASIS $\geq$ 8 for eligibility. MDD=major depressive disorder, GAD=generalized anxiety disorder, Social=social anxiety disorder, PTSD=post-traumatic stress disorder, Panic=panic disorder. Numbers on the diagonal indicate the total number of participants with a lifetime occurrence of a given diagnosis. Numbers below the diagonal indicate participants with a given comorbidity.
*Of those with a lifetime diagnosis of MDD, 20 had current MDD, 30 were in partial remission, and 8 were in full remission.

Participant Withdrawals

The sensitivity assessment served not only as a way to obtain individual anxiety ratings to the breathing resistances, but also to allow initial exposure to the mask before participants began the Horizon Task. Seven participants who consented for the study withdrew either during or after the sensitivity assessment because of discomfort with the breathing mask. Of the 7 individuals who consented for this project and withdrew before completion due to tolerability of the breathing resistance, 5 were iADs and 2 were HCs. The iADs who withdrew were numerically older (M=44, SD=14.09) than those who did not and had comparable OASIS scores (M=8.5, SD=1.29) and numerically lower PHQ scores (M=8.25, SD=4.79). All 5 were diagnosed with MDD, 4 GAD, 1 PTSD, and 1 social anxiety disorder. Two were taking psychiatric medications. All 7 participants who withdrew were female. The small number of individuals who withdrew did not allow for formal statistical comparison to those who remained in the study.

Measure Details

State-Trait Anxiety Inventory (STAI-State/Trait). The STAI consists of two sets of 20 items

designed to assess state and trait anxiety, respectively. Respondents were asked to rate each item

on a scale of 1 ("not at all" for state anxiety; "almost never" for trait anxiety) to 4 ("very much

so" for state anxiety; "almost always" for trait anxiety) based on how much they agree or

disagree with the associated statement. The STAI has shown good to excellent internal

consistency (Cronbach's alphas ranging from .86 to .96; (38)).

Overall Anxiety Severity and Impairment Scale (OASIS). The OASIS is a 5-item self-report

measure designed to assess anxiety-related impairment across various areas of life over the past

week. Total scores are obtained by summing scores of individual items (0-4), ranging from 0 to

20, where higher scores indicate greater impairment. The OASIS has shown a high internal

consistency, with a Cronbach's alpha of .80 (39).

Patient Health Questionnaire (PHQ-9). The PHQ is a screening tool for depression and

includes 9 self-report items, which correspond to the diagnostic criteria of major depressive

disorder. Each item statement is rated on a 4-point Likert Scale, ranging from 0 (not at all) to 3

(nearly every day) to describe the extent to which a depressive symptom has bothered them in

the past 2 weeks. The total score reflects level of depression severity. The PHQ-9 has been found

to show good internal consistency ($\alpha = .89$; (40)).

Childhood Trauma Questionnaire (CTQ). The CTQ short form (41) includes 28 self-report

items that measure childhood abuse and neglect through five subscales: physical abuse,

emotional abuse, sexual abuse, physical neglect, and emotional neglect. Each item is scored

using a Likert scale, ranging from 1 ("never true") to 5 ("very often true"), with relevant items

summed to provide each subscale score. The total score is the mean of the subscale scores.

Higher scores indicate higher levels of childhood abuse and neglect.

Cognitive Reflection Test (CRT-7). The CRT-7 (42) is comprised of 7 short questions that are

designed to assess trait cognitive reflectiveness. Lower reflectiveness is operationalized as an

individual's tendency to provide an immediate intuitive answer that is incorrect, when more

detailed reflection on the content of the question would lead to a less intuitive but correct answer.

Example item: "Jerry received both the 15th highest and the 15th lowest mark in the class. How

many students are in the class?" (intuitive answer: 30; correct answer: 29). Scores are based on

the number of correct answers. The CRT-7 has shown acceptable reliability (α=.72; (42)).

Sensitivity Assessment

In addition to self-reported anxiety levels, participants also gave responses to a series of other

related questions immediately after exposure to each breathing resistance during the sensitivity

protocol (exact wording listed below). Self-Assessment Manikin items assessed happiness and

excitement (1) rated on a 5-point scale, while the others were rated similarly to self-reported

anxiety (11-point scale). See **Figure S1** below for associated bar plots demonstrating ratings

across resistance levels.

"How calm or excited did you feel during loaded breathing?" (on the scale from 1 to 5)

"How happy or unhappy did you feel during loaded breathing?" (on the scale from 1 to

5)

"How difficult did it feel to breathe?" (0-no difficulty; 10-maximal difficulty you could

tolerate)

"How much fear did you feel while breathing?" (0-no fear at all; 10-maximal fear you

could tolerate)

"How unpleasant did it feel to breathe?" (0-not at all unpleasant; 10-maximal

unpleasantness you could tolerate)

Computation Model and Model Fitting

Learning Model

The learning model is designed to reflect how the participant estimates the mean reward of both

bandits during forced-choice trials. In order to estimate the mean reward values, the model, based

on a Kalman filter (Kalman, 1960), assumes that rewards r are generated from a Gaussian

distribution with a fixed standard deviation σ_r and a mean μ that differs between bandits and can

change over time. Changes in the mean estimate are determined by a Gaussian random walk with

mean 0 and standard deviation σ_d. Importantly, the dynamic mean assumptions made by this

model do not reflect the true generative model of the Horizon Task (where means are constant

throughout a game; i.e., $\sigma_d = 0$) allowing for the possibility of suboptimal learning as well as a

recency bias in mean estimates.

For bandit i on trial t, estimation of mean reward R_t^i is updated based on the reward prediction

error (i.e., difference between the observed reward and prior mean), which is scaled by a time

varying learning rate α_t^i. This gives the following:

709
$$R_{t+1}^i = R_t^i + \alpha_t^i \left(r_t - R_t^i\right)$$

710 and

711
$$\frac{1}{\alpha_t^i} = \frac{1}{\alpha_{t-1}^i + \alpha_d} + 1$$

712 where

713
$$\alpha_t^i = \frac{(\sigma_{t+1}^i)^2}{\sigma_r^2} \quad \text{and} \quad \frac{1}{(\sigma_{t+1}^i)^2} = \frac{1}{(\sigma_t^i)^2 + \sigma_d^2} + \frac{1}{\sigma_r^2}.$$

714 The learning rate, as shown here, is defined by uncertainty in: 1) the estimated mean of a bandit

715 (σ_t^i), 2) the drift in this mean over time (σ_d), and 3) the assumed standard deviation of the

716 generative distribution (σ_r). If bandit i is not played on trial t, the estimated mean is assumed to

717 stay the same, while the uncertainty in that estimate grows in accordance with σ_d:

718
$$\left(\sigma_{t+1}^i\right)^2 = \left(\sigma_t^i\right)^2 + \sigma_d^2.$$

719 The dynamic learning rate can be decomposed into two free parameters that are fit to individual

720 behavior. The starting learning rate is defined by:

721
$$\alpha_0 = \frac{\sigma_0^2 + \sigma_d^2}{\sigma_r^2}.$$

722 The "asymptotic" learning rate, or the value of α_t^i if bandit i is chosen ad infinitum, is instead:

723
$$\alpha_\infty = \frac{1}{2}\left(-\alpha_d + \sqrt{\alpha_d^2 + 4\alpha_d}\right),$$

724 where

$$725 \qquad \alpha_d = \frac{\sigma_d^2}{\sigma_r^2}.$$

726   This reparameterization then constrains both learning rate values to [0,1].

Decision Model

728   After estimating the mean reward of each bandit, R_t^i, the participant uses it to choose between

729   two bandits probabilistically according to a logistic choice rule:

$$730 \qquad p(choose\ right) = \frac{1}{1 + exp\left(\frac{\Delta R + A\Delta I + B}{\sigma}\right)},$$

where ΔR is denoted as the differences in expected reward between two bandits and ΔI is denoted

as the differences in information between two bandits. Here, +1 is specified when the left bandit

is more informative, -1 when the right bandit is more informative, and 0 when both bandits

convey equal information. The decision process is characterized by three parameters: *A* (i.e., the

information bonus), *B* (i.e., the spatial bias), and (i.e., the decision noise). Separate values of each

parameter are estimated for each horizon (i.e., H1 and H6) and the spatial bias and decision noise

parameters are further separated by uncertainty conditions (i.e., equal and unequal information),

while information bonus is only calculated in the unequal information condition.

Model Fitting

A previously used hierarchical Bayesian model was used to fit each parameter to the behavior of

each participant. In this approach to model fitting, it is assumed that each parameter for each

participant is taken from the same group-level prior distribution, and is estimated using a Markov

Chain Monte Carlo (MCMC) sampling procedure. Group-level prior distributions are

parameterized by hyperpriors set to have broad, minimally informative distributions.

Model fitting was performed using the JAGS package (Plummer, 2003) to run Markov Chain

Monte Carlo simulations in MATLAB (psiexp.ss.uci.edu/research/programs_data/ jags/). This

package draws samples from the posterior distribution given the observed behavioral data to

approximate the posterior distribution over model parameters. Specifically, 1000 samples were

generated from the posterior distribution over model parameters for each of 4 independent

Markov chains (i.e., 4000 samples in total). We discarded the first 500 samples of each chain

(i.e., burn-in = 500) that might impact the posterior distribution and generated posterior samples

at a thin rate of 1.

<u>Supplementary Results</u>

Breathing Resistance Sensitivity

Anxiety ratings during the breathing resistance sensitivity protocol are shown in **Figure S1**. In

the LME predicting anxiety including resistance level, group, and their interaction as predictors,

all effects were significant ($Fs \geq 5.77$, $ps < .001$). Post-hoc contrasts indicated that 1) iADs had

higher self-reported anxiety levels than HCs at each resistance level ($ps < .013$), 2) that anxiety at

resistance levels 40, 60, and 80 were not significantly different from each other in HCs

($ps \geq .076$), and 3) that each increase in resistance level resulted in a significant increase in

anxiety for the iADs, but anxiety in HCs did not change from 40 through 80. These results

indicate that anxiety induction was successful across participants, and that iADs showed greater

sensitivity than HCs.

LMEs predicting self-reported anxiety scores at baseline and during both task runs (**Figure 2**)

showed significant effects of group ($F(1,117)=49.44$, $p<.001$), resistance condition

($F(2,234)=105.89$, $p<.001$), and their interaction ($F(2,234)=8.04$, $p<.001$). Post-hoc analysis

revealed that: 1) iADs (estimated marginal mean [EMM]=3.06) reported higher anxiety ratings

than HCs (EMM=1.03; $t(117)=7.03$, $p<.001$); 2), anxiety ratings were highest during the task

condition with the added resistance (EMM=3.20), followed by the task condition without the

resistance (EMM=1.63) and baseline ratings (EMM=1.31; $ps<.022$); and 3) self-reported anxiety

ratings did not increase from baseline to the no-resistance task run for HCs ($EMM_{baseline}=0.53$,

$EMM_{no\text{-}resistance}=0.69$; $p=.440$), but did for iADs ($EMM_{baseline}=2.08$, $EMM_{no\text{-}resistance}=2.57$;

$p=.013$).

Similar results were found for complementary STAI-State ratings. Here there were main effects

of group ($F(1,117)=54.93$, $p<.001$) and resistance ($F(2,234)=85.51$, $p<.001$), but no interaction

($F(2,234)=0.56$, $p=.570$). Post-hoc contrasts indicated that iADs reported higher anxiety

(EMM=44.9) than HCs (EMM=33.1; $t(117)=7.41$, $p<.001$) and that anxiety during the run with

the added breathing resistance (EMM=44.3) was higher than anxiety at baseline (EMM=35.7)

and when no resistance was added ($ps<.001$), but baseline was not significantly different from

the no resistance condition ($t(234)=1.73$, $p=.085$).

Disentangling Depression and Anxiety for Other Parameters

In models predicting RE in iADs based on effects of depression (PHQ) alone or anxiety (STAI

Trait) alone, only a negative effect of age was found ($Fs\geq3.92$, $ps\leq.052$). With PHQ and STAI in

the same model, there was a significant interaction between PHQ and resistance condition

($F(1,58)=4.40$, $p=.040$). Post-hoc analysis indicated that this was driven by a stronger negative

association between PHQ and RE in the resistance condition than in the no-resistance condition

(resistance condition: ET=-.178; no-resistance condition: ET=-.016; $t(58)=2.10$, $p=.040$). The

interaction between resistance condition and STAI Trait was also significant in this model

($F(1,58)=4.67$, $p=.035$). This reflected a positive association between anxiety and RE in the

resistance condition and no relationship between these variables in the no-resistance condition

(resistance condition: ET=.102; no-resistance condition: ET=.007; $t(58)=-2.16$, $p=.035$). There

were also a marginal overall positive effect of STAI ($F(1,57)=3.38$, $p=.071$, b=.055) and negative

effect of PHQ ($F(1,57)=3.46$, $p=.068$, b=-.098) driven by the above interactions.

When analogous models of RE were run in HCs, those containing only effects of depression

(PHQ) or anxiety (STAI Trait) separately showed no significant effects. However, when both

PHQ and STAI were included in the same model, results showed a marginal effect of STAI Trait

($F(1,53)=3.30$, $p=.075$) suggesting a possible positive association (b=.043).

In logistic regression models predicting α_0 in iADs, no significant effects were found in separate

or combined models. Analogous models in HCs found no significant effects when PHQ and

STAI were included alone. However, in a model containing both symptom measures together,

there were a couple noteworthy marginal effects. Namely, higher STAI Trait was suggestively

associated with higher α_0 (z=-1.93, $p=.054$; log-odds=-.23, CI=[-.47, .00])), and there was an

interaction between STAI Trait and resistance condition (z=1.769, $p=.077$; log-odds=.22, CI=[-

.02, .46]). This interaction suggested higher STAI scores predicted higher α_0, but only in the no-

resistance condition.

All analogous models predicting α_∞ in iADs showed no significant effects. In HCs, there were

similarly no significant effects in models including either PHQ or STAI alone. However, in a

model predicting α_∞ with both PHQ and STAI together, the interaction between PHQ and

resistance condition was marginal ($F(1,55)$=2.85, p=.097), suggesting a positive association

between PHQ and α_∞ in the no-resistance condition (ET=.009), but a negative association in the

resistance condition (ET=-.010; $t(55)$=1.69, p=.097). Additionally, the interaction between STAI

Trait and resistance condition was marginal ($F(1,55)$=3.23, p=.078), suggesting a negative

association between STAI and α_∞ in the no-resistance condition (ET=-.001), but a positive

association in the resistance condition (ET=.005; $t(55)$=-1.80, p=.078).

Finally, while main results of models predicting DE based on symptoms are reported in the main

text, we provide some additional statistical details here in relation to secondary resistance effects.

Namely, when predicting DE in iADs, specific statistics included marginal effects of resistance

condition in the models with PHQ alone ($F(1,59)$=3.94, p=.052, η_p^2=.06, b=-.525), STAI Trait

alone ($F(1,59)$=3.90, p=.053, η_p^2=.06, b=-.627), and both together ($F(1,58)$=3.89, p=.053,

η_p^2=.06, b=-.716).

Supplementary Figures

Figure S1. Participant ratings during the breathing resistance sensitivity protocol across 6 levels

of breathing resistance (0 to 80 cmH2O/L/sec). Boxplots show medians and quartiles as well as

individual datapoints.

Figure S2. Decision noise parameters separated by horizon, group, and resistance condition.

Figure S3. Post-hoc correlations between model parameters and self-report measures for the interested reader. Statistics below the diagonal indicate correlations with parameters in the resistance condition, while those above the diagonal are from the no-resistance condition. Note that the correlation between α_0 and other variables were conducted using Spearman correlations due to non-normality.

Figure S4. Fraction of correct responses (i.e., option with higher generative mean) for each free choice in H1 and H6 trials. Color shades indicate task performance with (light) and without (dark) breathing resistance separately in the HCs (left/blue) and iADs (right/pink). Plots are also shown separately for equal (top) or unequal (bottom) information conditions.