Comparison of Fine-Scale Malaria Strata Derived from Population Survey Data Collected Using mRDTs, Microscopy and qPCR in South-Eastern Tanzania

Issa H. Mshani[1, 2]*, Frank M. Jackson[1], Elihaika G Minja[1,3,4], Said Abbas[1], Nasoro S. Lilolime[5], Faraji E. Makala[1], Alfred B. Lazaro[1], Idrisa S. Mchola[1], Linda N. Mukabana[1,6], Najat Kahamba[1,2], Alex Limwagu[1], Rukia. M. Njalambaha[1], Halfan S. Ngowo[1], Donal Bisanzio[1, 8], Francesco Baldini[1, 2†], Simon A. Babayan[2†] and Fredros Okumu[1, 2, 7, 9†]*

1. Ifakara Health Institute, Environmental Health, and Ecological Sciences Department, Morogoro, United Republic of Tanzania.
2. The University of Glasgow, School of Biodiversity, One Health and Veterinary Medicine, Glasgow, UK.
3. University of Basel, Petersplatz, 4001 Basel, Switzerland.
4. Swiss Tropical and Public Health Institute, Kreuzstrasse 2, CH-4123 Allschwil, Switzerland.
5. Ifakara Health Institute, Interventions and Clinical Trials Department, Pwani, Bagamoyo, United Republic of Tanzania.
6. Jomo Kenyatta University of Agriculture and Technology, School of Public Health, Juja, Kenya.
7. The University of the Witwatersrand, School of Public Health, Park Town, South Africa.
8. RTI International, Washington, District of Columbia, USA
9. Nelson Mandela African Institution of Science and Technology, School of Life Sciences and Biotechnology, Arusha, United Republic of Tanzania.

†Fredros Okumu, Simon A. Babayan and Francesco Baldini are equally co-supervised this work.

*Correspondence:

1. Issa H. Mshani: imshani@ihi.or.tz & fredros@ihi.or.tz

Abstract

Introduction: Malaria-endemic countries are increasingly adopting data-driven risk stratification, often at district or higher regional levels, to guide their intervention strategies. The data typically comes from population-level surveys collected by rapid diagnostic tests (RDTs), which unfortunately perform poorly in low transmission settings. Here, we conducted a high-resolution survey of *Plasmodium falciparum* prevalence rate (PfPR) in two Tanzanian districts and compared the fine-scale strata obtained using data from RDTs, microscopy and quantitative polymerase chain reaction (qPCR) assays.

Methods: A cross-sectional survey was conducted in 35 villages in Ulanga and Kilombero districts, south-eastern Tanzania between 2022 and 2023. We screened 7,628 individuals using RDTs (SD-BIOLINE) and microscopy, with two thirds of the samples further analyzed by qPCR. The data was used to categorize each district and village as having very low (PfPR<1%), low (1%≤PfPR<5%), moderate (5%≤PfPR<30%), or high (PfPR≥30%) parasite prevalence. A generalized linear model was used to analyse infection risk factors. Other metrics, including positive predictive value (PPV), sensitivity, specificity, parasite densities, and Kappa statistics were computed for RDTs or microscopy using qPCR as reference.

Results: Significant fine-scale variations in malaria risk were observed within and between districts, with village prevalence ranging from 0% to >50%. Prevalence varied by testing method: Kilombero was low risk by RDTs (PfPR=3%) and microscopy (PfPR=2%) but moderate by qPCR (PfPR=9%); Ulanga was high risk by RDTs (PfPR=39%) and qPCR (PfPR=54%) but moderate by microscopy (PfPR=26%). RDTs and microscopy classified majority of the 35 villages as very low to low risk (18 - 21 villages). In contrast, qPCR classified most villages as moderate to high risk (29 villages). Using qPCR as the reference, PPV for RDTs and microscopy ranged from <20% in very low transmission villages to >80% in moderate to high transmission villages. Sensitivity was 62% for RDTs and 41% for microscopy; specificity was 93% and 96%, respectively. Kappa values were 0.58 for RDTs and 0.42 for microscopy. School-age children (5-15years) had higher malaria prevalence and parasite densities

than adults (P<0.001). High-prevalence villages also had higher parasite densities (Spearman r=0.77, P<0.001 for qPCR; r=0.55, P=0.003 for microscopy).

Conclusion: This study highlights significant fine-scale variability in malaria risk within and between districts and emphasizes the variable performance of the testing methods when stratifying risk. While RDTs and microscopy were effective in high-transmission areas, they performed poorly in low-transmission settings; and classified most villages as very low or low risk. In contrast, qPCR classified most villages as moderate or high risk. While we cannot conclude on which public health decisions would be subject to change because of these differences, the findings suggest the need for improved testing approaches that are operationally feasible and sufficiently sensitive, to enable precise mapping and effective targeting of malaria in such local contexts. Moreover, public health authorities should recognize the strengths and limitations of their available data when planning local stratification or making decisions.

Key words: Malaria; Fine scale stratifications; Prevalence rate, Rapid diagnostic tests (RDTs), Polymerase chain reaction (PCR); Microscopy; Population surveys; Micro-stratification; Malaria screening.

Background

Precise mapping of malaria prevalence is crucial for the eventual elimination of the disease from different localities. In line with World Health Organization (WHO) guidelines, National Malaria Control Programs (NMCPs) in Africa are increasingly adopting data-driven stratification of malaria risk, in most cases at either district or higher regional levels [1–3]. These stratifications involve assessing risk levels in geographical areas at the subnational level (e.g. zones, regions, and districts) [2,4,5], and can include fine scale mapping (down to wards and villages levels) as countries progress towards elimination [6–8]. The data for such stratification may come from health facilities, active malaria screening during population surveys, or proxy data sources such as antenatal care clinic visits [6,9,10].

When developing country-level malaria strategies, the prevalence of malaria, representing the proportion of confirmed positive cases of *Plasmodium falciparum* (or other *Plasmodium* sp.) among all individuals tested [11,12], can be classified into various transmission categories. The WHO has previously used the following cutoff points for malaria endemicities: below 1% as very low, 1-10% as low, 10-35% as moderate, and above 35% as high risk malaria stratum [12]. Different NMCPs may adapt these criteria with slight adjustments based on local epidemiological insights. For instance, some countries, including Tanzania and Kenya, have used the parasite prevalence data to categorize their geographic zones as either very low risk (PfPR < 1%), low risk (1% ≤ PfPR < 5%), moderate risk (5% ≤ PfPR < 30%), or high risk strata (PfPR ≥ 30%) [2,13]. Another measure that can be used for generating these strata is the annual parasite incidence (API), which is the number of diagnostically confirmed malaria cases per 1000 individuals per year and is usually obtained from health facilities data [12,14]. API offers precise malaria stratifications, though it fails to account for sub-clinical malaria infections, which can contribute to transmission and impede malaria elimination efforts [15].

National malaria programs usually rely on different actively and passively collected data to inform malaria burden and monitor the effectiveness of control measures [16–18]. For instance, Tanzania employs multiple platforms, including the District Health Information

software (DHIS2) populated with data from routine health facility visits, the Malaria

Indicator Surveys (MIS) and Tanzania Demographic and Health Surveys (TDHS), which

are done every 4-5 years through household surveys, and the school malaria parasite

surveillance targeting kids aged 5-16 years during (SMPS) [2,19–21]. A common

feature of these established systems is that most rely primarily on rapid diagnostic tests

(RDTs) and microscopy [13,22,23], though samples are sometime also preserved for

PCR assays.

Microscopy, long used in malaria diagnosis, can quantify parasite loads and identify

different *Plasmodium* species, which are essential for precise treatment choices [24,25].

However, its effectiveness depends significantly on the skill and experience of the

microscopist, making it unreliable in some contexts, and it can miss a substantial

number of true infections due to sub-optimal accuracy [26–28]. In contrast, Rapid

Diagnostic Tests (RDTs) offer a consistent and user-friendly option, enabling quick, on-

site diagnosis without specialized skills or equipment. RDTs have become widely used

in both point-of-care settings and population surveys due to their operational simplicity

and cost-effectiveness [29–33]. While the technique enhances access to diagnostics,

especially in remote areas, RDTs have lower sensitivity for detecting low-level infections

and cannot quantify parasite density or distinguish between *Plasmodium* species.

Additionally, current RDTs may detect antigens for over three weeks post-treatment,

leading to poor specificity and potential overestimation of malaria cases in high

transmission areas [30,33,34].

In contrast, polymerase chain reaction (PCR) assays are known for their high sensitivity

and specificity [35]. While conventional PCR assays typically provide qualitative

information on malaria infections, quantitative PCR (qPCR) can offer additional

quantitative measures of malaria parasite density [36,37]. Unfortunately, the widespread

use of PCR assays for population surveys is hindered by cost constraints and the need

for specialized expertise and infrastructure for implementation [35,38,39].

The increased focus on evidence-based strategies in malaria control also includes a

transition from broad subnational stratifications to more granular, fine-scale approaches

[6,10]. However, although current methods like RDTs and microscopy are favored for

their operational simplicity, their effectiveness in detailed risk stratification, which are

critical for targeting both clinical and sub-clinical infections for malaria elimination,

remains poorly understood. Some authors have also suggested that RDTs may have

vastly reduced performance in settings where the malaria burden has been significantly

reduced [40]. This calls for a rigorous evaluation and comparison of these methods

against highly sensitive techniques such as qPCR to refine malaria stratification

approaches for malaria elimination. Indeed, available evidence, including data from

Kenya and Tanzania, suggest that PCR assays are generally better at pinpointing main

malaria hotspots in communities than RDTs and microscopy [41,42]. The study from

Tanzania further showed that in subsequent treatment campaigns relying on RDT-

based screening, ~45% of infections remain untreated, even if treatment is offered to all

members of households with an infected individual [42]. In the Kenyan study, the

authors went further to suggest that since detection of hotspots depends on the

sensitivity of diagnostic tools, health authorities working in malaria elimination settings

should consider using PCR to guide detection of the residual hotspots, as this provides

greatest opportunities to find asymptomatic individuals and sub-patent parasite

reservoirs in the communities [41].

All these studies clearly show that while sub-national stratification may be the most

effective approach to decide on how to allocate resources, the type of data used for

such epidemiological profiling matters significantly; especially when the stratification is

done at local-sub-district levels. In places like southeastern Tanzania, which has

experienced decades of sustained malaria interventions and progress, and where

robust entomological surveillance already exists [43], addition of detailed parasite

prevalence data from population-level surveys is required to enable more precise, fine-

scale stratifications at both district and sub-district levels.

The aim of this study was therefore to generate a high-resolution population-level

survey map of *P. falciparum* prevalence in two districts in south-eastern Tanzania, and

compare the fine scale malaria risk strata obtained when using data from different test

methods, namely RDTs, microscopy and qPCR. The study also examined the

association between population-level parasite prevalence and the parasite densities as

determined by both qPCR and microscopy in the different study villages. Lastly, we

investigated the malaria detection capacity of RDTs and microscopy across villages with
varying parasite prevalences and densities.

Methods

Study site

The study was conducted in Morogoro region, in south-eastern Tanzania (Figure 1), in
the two districts of Kilombero (population: ~583,000; 8.2414°S, 36.3349°E; elevation:
~270m) and Ulanga (population: ~233,000; 8.9889°S, 36.6133°E; elevation: ~800m).
The average malaria prevalence in the Morogoro region has previously been estimated
to exceed 10%, with *P. falciparum* as the dominant malaria species [19,20,44]. The
main economic activities for residents include rice farming, sugarcane farming and
maize farming, though the area also has other food crops and large commercial tree
plantations (teak). The known annual rainfall range is 1200-1400 mm in the lower-lying
plains of Kilombero district, and1400-2100 mm in the higher areas in Ulanga district
[45]. Approximately 90% of the rainfall occurs during the wet seasons between
December to April, with dry seasons typically lasting from June through September [45].
The annual mean daily temperature is around 27°C in the lowlands and approximately
23°C in the highlands. Relative humidity averages from 75% in the lowlands to 80% in
the highlands.

Figure 1: Study villages in Kilombero and Ulanga districts, south-eastern Tanzania.

186
187
188

Study design, procedures and survey tools

The cross-sectional surveys were conducted over two consecutive years, from 2022 to 2023, covering the transition from the end of rainy seasons to the peak of dry seasons. Villages were randomly selected from each district, and sample sizes were proportionately determined based on the population of each village using the Cochran formula adjusted for finite populations [46–48]. The expected prevalence varied depending on the village and was derived from previous surveys and health centers within each village. Based on the estimated sample size mentioned above, representative households were selected by systematically randomizing the names of all households obtained from the respective village administrations.

The screening criteria included individuals aged 5-60 years who had not taken malaria
medications in the preceding two weeks. This precaution aimed to prevent potential
overestimation by RDTs, as they may detect residual traces post-treatment [49].
Individuals needing special medical attention, such as pregnant women, were excluded
from the study. Each consenting individual undergoing malaria screening was assigned
a unique identity number that was also linked to their corresponding household ID. On-
site malaria diagnosis was conducted using: 1) RDTs via finger prick, 2) creating thick
and thin blood smears, and 3) collecting 3-5 dried blood spots (cycles) on Whatman^{TM}
filter paper cards. Subsequently, these samples were transported to the reference
laboratory for microscopy and qPCR analysis. For children under five years, only
screening with RDTs was done for prevalence estimations and not used for strata
comparisons.

Ethical considerations, survey team, and trainings

Permission to conduct this study was obtained from the Ifakara Health Institute Review
Board (Ref: IHI/IRB/No: 1/2021) and the National Institute for Medical Research-NIMR
(NIMR/HQ/R.8a/Vol. 1X/3735). Additionally, approvals were obtained from regional,
district, ward, and respective selected village authorities before commencing the
surveys, given the screening was done at centralized location in each village. Written
informed consent was obtained from individual adult participants (and parents or
guardians of those aged below 18) on the day before the actual testing. The study team
consisted of 11 members, including three molecular laboratory technologists, four
licensed medical laboratory microscopists, two licensed clinical officers, and two social
scientists. Prior to the survey commencement, a five-day training session was
conducted at the Ifakara Health Institute laboratory. This training covered explanations
of the study protocols, pilot implementations, procedures for protecting human
participants, quality assurance and training on data collection tools.

Tests using malaria rapid diagnostic tests (RDTs)

A small blood drop obtained through a finger prick was collected onto the RDTs (SD
Bioline Ag *Pf*/Pan), following the manufacturer's instructions. The buffer solution was
applied according to standard RDTs procedures and left on the bench surface for up to

20 minutes. The type RDT used were capable of detecting *P. falciparum* infections by

targeting histidine-rich proteins 2, which react on the *Pf*-line. Additionally, they could

detect *P. malariae* and *P. ovale* by targeting glycolytic lactate dehydrogenase,

expressed by the Pan-line on RDTs [30]. The RDT results were recorded on a paper

form, and any individuals who tested positive for malaria were promptly treated with

Arthemether Lumefantrine (ALu), following Tanzania's national malaria treatment

guidelines [50].

Tests using microscopy

Thick and thin blood smears were created in the field, stained with 10% Giemsa for 15

minutes then examined for the presence of malaria parasites under oil immersion at

100X magnification [47,51,52]. Two experienced microscopists independently read the

slides, and discrepancies between them were resolved by a third, more experienced

microscopist. They read the thick smear first, and if an infection was detected, the thin

smear was read to identify parasite species. The presence of both asexual and sexual

malaria parasite stages discriminating *P. falciparum*, *P. malariae*, and *P. ovale* was

recorded. Asexual stage parasites were counted per 200 white blood cells and

assuming 8000 WBC/µL [53]. The mean count of malaria parasite by microscopy

between the two readers was calculated and confirmed by the third reader.

Tests using real-time qPCR assays

A representative sample of approximately two thirds of all samples was randomly

selected from each village and screened further by quantitative polymerase chain

reaction (qPCR) i.e. 4905 samples out of the total 7628 samples. Out of five spots

created on the Whatman^{TM} filter paper card, three spots of DBS were used for DNA

extraction using Quick-DNA^{TM} Miniprep plus kit (Zymo Research,USA) [54], and eluted

with50 µL of elution buffer, stored at -20°C for further detection and quantification of *P.*

falciparum infections using probe-level allele-specific quantification (PlasQ)-multiplex

qPCR assays protocols [37,55,56]. The detection and quantification of *P. falciparum*

parasites were performed using the Bio-Rad CFX96 real-time PCR system (Bio-Rad

Laboratories, USA) [55] and analyzed with Bio-Rad CFX maestro software. The qPCR

reaction, PlasQ primers and probes mix, are summarized in supplementary online table

1& 2. DNA amplification processes included: activation at 95°C for 1 min, denaturation

at 95°C for 15 seconds, and annealing and elongation at 57°C for 45 seconds for 45

cycles, followed by melting [55].

The qPCR assays were run with positive controls (samples with confirmed *P.*

falciparum) and a non-template control (samples with no *P. falciparum* as negative

control). For absolute parasite quantification, the WHO international standard for *P.*

falciparum nucleic acid amplification techniques were used (WHO reference from

NIBSC#04/176) [37]. The standard was reconstituted following the manufacturer's

instructions and serially diluted in the range of 100,000 parasites/µL to 0.01 and

analyzed in triplicates.

During the qPCR assay, the prepared standards were run together with unknown

samples, and at the end of the assay, the standard curve and samples were normalized

and analyzed with Bio-Rad CFX maestro software. The obtained normalized Ct values

of the samples and the linear regression equation derived from the standard curve were

used to calculate the parasites density of the unknown samples, expressed as parasites

per microlitre (parasites/µL) of blood.

Data analysis

All results from RDTs, microscopy, and qPCR were entered into the Open Data Kit

(ODK) system [57], and subsequently downloaded as an excel file for further cleaning.

The datasets for RDTs, microscopy, and qPCR results were merged based on the

participant's ID using the Pandas Python package [58].

Generalized linear models (GLM) were used to evaluate the association between

malaria infection risk and the variables of age and gender in a univariate analysis, run in

Python using the stats model package [59]. Additionally, to evaluate the performance of

RDTs and microscopy in fine-scale malaria stratifications compared to qPCR, their

agreement was tested using Kappa statistic [60], and the resulting Kappa values

interpreted as follows: $\kappa < 0.20$ as poor agreement, 0.21–0.40 as fair, 0.41–0.60 as

moderate, 0.61–0.80 as substantially good and 0.81–1 as almost perfect agreement

[61]. In addition, the positive predictive value (PPV) for RDTs and microscopy was

computed, using qPCR results as the reference, per village, as (proportion of positive

test results that are actually true positives, estimated as PPV = True Positives / (True

Positives + False Positives)). Fine-scale stratification by villages using data from qPCR,

RDT, and microscopy were used to generate the prevalence maps using quantitative

Geographic Information System (qGIS) software, allowing visualization of malaria

prevalence across villages. Additionally, the risk maps were created using inverse

distance weighting (IDW) interpolation techniques. This method helped to estimate

malaria risk by spatially interpolating the prevalence data, which provided a continuous

surface of malaria risk across the study area.

The geometric mean of parasite density estimated by microscopy and qPCR was

computed, and statistically compared the parasite densities between gender and age

groups. The non-parametric Mann-Whitney statistics were used to compare the parasite

densities between two categorical groups, while Kruskal-Wallis statistical tests were

used to compare more than two categorical groups [62,63]. For example, differences in

parasite densities between age groups were tested using Kruskal-Wallis statistics, and if

statistically significant, the Mann-Whitney statistics were applied for pairwise statistical

significance tests. All analyses that compared parasite densities excluded the negative

class and focused solely on investigating parasite density distribution among malaria-

positive patients. Lastly, to test for statistical correlations between parasite prevalence

and parasite densities estimated by both qPCR and Microscopy, non-parametric

Spearman's rank correlation tests were employed [64]. Additionally, logistic regression

model was also used to assess the probability of diagnostic tools to detect malaria

infections at various parasite densities.

Results

Baseline study population.

This survey covered 35 villages and 93 sub-villages across Ulanga and Kilombero

districts. A total of 7,628 participants (>5 years) were recruited upon consent and tested

for malaria using RDTs and microscopy, with 64.3% (4,905) of them also tested by

qPCR (Figure 2). Males comprised 38% of the study population, while females made up

62%. Among the participants, 35% were school-aged children (5-15 years), and 65%

were aged 16 years and above (Table 1). For children under five years, for whom

screening was conducted with only RDTs to estimate the burden, the results are not

reported here.

Figure 2: *Schematic representation of the study sampling procedures*

Table 1: Baseline characteristics of the study populations

	Kilombero District n (%)	Ulanga District n (%)	Total N (%)
Villages	19 (54.3)	16 (45.7)	35.0
Sub Villages	48 (51.6)	45 (48.4)	93.0
Gender			
Female	2164 (67.4)	2573 (58.3)	4737 (62.1)
Male	1047 (32.6)	1844 (41.7)	2891 (37.9)
Total	**3211**	**4417**	**7628**
Age Group			
5-10 years	519 (16.2)	935 (21.2)	1454 (19.1)
11-15 years	400 (12.5)	802 (18.2)	1202 (15.8)
16-20 years	198 (6.2)	333 (7.5)	531 (7.0)
>20 years	2094 (65.2)	2347 (53.1)	4441 (58.2)
Total	**3211**	**4417**	**7628**

Malaria prevalence by RDTs, Microscopy and qPCR

In the Ulanga district, malaria transmission was found to be high by both qPCR and

RDTs, with *P. falciparum* prevalence rates of 53.89% [95% CI, 52.06-55.72] and

38.35% [95% CI, 36.92-39.79] respectively. However, microscopy categorized it as

moderate, with a prevalence rate of 26.07% [95% CI: 24.77-27.36] (Table 2). Within this

moderate to high transmission strata in Ulanga, males had a significantly higher

prevalence of malaria compared to females. The odds of malaria infection in males

compared to females were estimated as 1.54 [95% CI: 1.36-1.74] (P < 0.001) by RDTs,

1.45 [95% CI: 1.27-1.66] (P < 0.001) by microscopy, and 1.54 [95% CI: 1.36-1.74] (P <

0.001) by qPCR (Table 2). All tests - RDTs, microscopy, and qPCR - indicated that

school-age children (5-15 years) had a significantly higher prevalence of malaria

infections than the other age groups (Table 2).

In Ifakara council, within the Kilombero district, both RDTs and microscopy categorized

the area as a low risk stratum, with observed prevalence rates of 2.68 [95% CI: 2.12-

3.24] and 1.84 [95% CI: 1.37 – 2.30], respectively (Table 3). However, qPCR classified

Kilombero district as a moderate risk stratum with a prevalence rate of 8.77 [95% CI:

7.55-9.99] (Table 3). Notably, there were no statistically significant differences in

malaria infection risk between males and females in this low to moderate transmission

setting, as indicated by both RDTs (Odds prevalence 1.51% [95% CI; 0.97, 2.33],

P=0.065) and microscopy (Odds prevalence 1.33% [95% CI; 0.78, 2.25], P=0.293), as

well as qPCR (Odds prevalence 1.09% [95% CI: [0.80-5.67], P = 0.567). Additionally,

school age children (5-15 years) exhibited a significantly higher risk of malaria infections

compared to those above 15 years old, as demonstrated by both RDTs, microscopy,

and PCR (Table 3).

Micro-stratification of malaria risk using data collected by qPCR, RDTs, and

microscopy

Significant variability in malaria infections was observed at the individual village level,

with prevalence rates ranging from 0% to over 50% across the study area (Figure 3 &

Table 4). Additionally, the method used to test for malaria significantly impacted the risk

categorization of villages. Among the 35 villages surveyed, qPCR data indicated that

only one village (1.2% of all villages) had very low malaria prevalence (PfPR < 1%). In

contrast, RDTs identified 12 villages (34.3% of all villages) and microscopy identified 11

villages (31.4% of all villages) as having very low prevalence. For moderate

transmission, qPCR, RDTs, and microscopy categorized 15, 9, and 8 villages,

respectively. For high transmission, qPCR identified 14 villages, RDTs identified 8, and

microscopy identified 6. Notably, qPCR detected more malaria infections than RDTs

and microscopy, resulting in many villages being classified into higher transmission

categories.

Overall, using qPCR data, over 80% of the villages were classified as moderate to high

risk, significantly higher than the 48% classified by RDTs and 40% by microscopy.

Conversely, while only 17% of the villages were classifiable as having low or very low

malaria risk based on qPCR data, as high as 51% and 60% of the villages were

classified into these same categories based on RDT and microscopy data (Table 4 and

Figure 4)

Table 2: Malaria prevalence in Ulanga district by sex and age groups, as measured using RDTs, microscopy and qPCR

Attribute	RDTs				Microscopy				qPCR			
	Positive/ Total tested	Prevalence (%) [95% CI]	Odds of infection [95% CI]	P-value	Positive/ Total tested	Prevalence (%) [95% CI]	Odds of infection [95% CI]	P-value	Positive/ Total tested	Prevalence (%) [95% CI]	Odds of infection [95% CI]	P-value
Overall prevalence	1689/4404	38.4 [36.9-39.8]	-	-	1148/4404	26.1 [24.8 – 27.4]	-	-	1531/2841	53.9 [52.1-55.7]	-	-
Sex												
Female	874/2566	34.1 [32.2-35.9]	Ref		593/2566	23.1 [21.5 – 24.7]	Ref	-	830/1659	50.0 [47.6-52.4]	Ref	-
Male	815/1838	44.3 [42.1-46.6]	1.5 [1.4-1.7]	<0.001	555/1838	30.2 [28.1-32.3]	1.4 [1.3-1.6]	<0.001	701/1182	59.3 [56.5-62.1]	1.5 [1.4-1.8]	<0.001
Age group												
5-10 years	497/933	53.3 [50.1-56.5]	Ref		341/933	36.6 [33.5-39.6]	Ref	-	338/622	54.3 [50.4-58.3]	Ref	-
11-15 years	479/801	59.8 [56.4-63.2]	1.3 [1.1-1.6]	0.006	358/801	44.7 [41.3-48.1]	1.4 [1.2-1.7]	<0.001	355/534	66.5 [62.5-70.5]	1.3 [1.1-1.6]	0.006
16-20 years	154/332	46.4 [41.0-51.8]	0.8 [0.6-1.0]	0.031	109/332	32.8 [27.9-37.9]	0.9 [0.7-1.1]	0.224	127/210	60.5 [53.9-67.1]	0.8 [0.6-1.0]	0.031
>20 years	559/2338	23.9 [22.2-25.6]	0.3 [0.2-0.4]	<0.001	340/2338	14.5 [13.1-16.0]	0.3 [0.3-0.4]	<0.001	711/1475	48.2 [45.7-50.8]	0.3 [0.2-0.3]	<0.001

371
372

373   **Table 3**: Malaria prevalence in the Kilombero district by sex and age groups, as measured using RDTs, microscopy and
374   qPCR

	RDTs				Microscopy				qPCR			
Attribute	**Positive/ Total tested**	**Prevalence (%) [95% CI]**	**Odds of infection [95% CI]**	**P-value**	**Positive/ Total tested**	**Prevalence (%) [95% CI]**	**Odds of infection [95% CI]**	**P-value**	**Positive/ Total tested**	**Prevalence (%) [95% CI]**	**Odds of infection [95% CI]**	**P-value**
Overall prevalence	86/3211	2.7 [2.1-3.2]	-	-	59/3211	1.8 [1.4 – 2.3]	**-**	-	181/2064	8.8 [7.6-10.0]	**-**	**-**
Sex												
Female	50/2164	2.3 [1.7-2.9]	**Ref**	-	36/2164	1.2 [1.1-2.2]	**Ref**		118/1385	8.5 [7.1-10.0]	**Ref**	-
Male	36/1047	3.4 [2.3-4.5]	1.5 [0.9-2.3]	0.065	23/1047	2.2 [1.3 – 3.1]	1.3 [0.8-2.3]	0.293	63/679	9.3 [7.1-11.5]	1.1 [0.8-1.5]	0.567
Age group												
5-10 years	21/519	4.1 [2.4-5.7]	**Ref**	-	16/519	3.1 [1.6-4.6]	**Ref**	-	33/343	9.6 [6.5-12.7]	**Ref**	-
11-15 years	17/400	4.3 [2.3-6.2]	1.1 [0.6-2.0]	0.877	10/400	2.50 [1.0-4.03]	0.8 [0.4-1.8]	0.597	28/256	10.9 [7.1-14.8]	1.2 [0.7-2.0]	0.59
16-20 years	9/198	4.6 [1.6-7.5]	1.1 [0.5-2.5]	0.765	4/198	2.02 [0.1-3.4]	0.6 [0.2-2.0]	0.443	16/123	13.0 [7.1-19.0]	1.4 [0.7-2.7]	0.295
>20 years	39/2094	1.9 [1.3-2.4]	0.5 [0.3-0.8]	0.003	29/2094	1.38 [0.88-1.89]	0.4 [0.2-0.8]	0.009	104/1342	7.8 [6.3-9.2]	0.8 [0.5-1.2]	0.259

Figure 3: *Fine-scale malaria mapping of 35 surveyed villages in the Ulanga and Kilombero districts using qPCR, RDTs, and microscopy data is shown in the top panel. The bottom panel indicates malaria risk generated by interpolating prevalence data obtained for each surveyed village by qPCR, RDTs, and Microscopy.*

385 ***Table 4****: Number of villages categorized into different risk strata based on the* P.
386 falciparum *prevalence rate (PfPR) data from qPCR, RDTs and microscopy.*

Risk strata	Prevalence	Number of villages					
		qPCR		RDTs		Microscopy	
		Count	%	No.	%	No.	%
Very low	PfPR < 1%	1	2.9	12	34.3	11	31.4
Low	1% ≤ PfPR < 5%	5	14.3	6	17.1	10	28.6
Moderate	5% ≤ PfPR < 30%	15	42.9	9	25.7	8	22.9
High	PfPR ≥ 30%	14	40.0	8	22.9	6	17.1
Total		35	100.0	35	100.0	35	100.0

Figure 4: *Percentage of villages categorized by different testing methods as either very*
low risk, low risk, moderate risk or high risk (total number of villages = 35).

Comparison of the perfomance of RDTs and microscopy relative to qPCR.

In this comparative analysis, only samples tested by all three methods—PCR, RDTs,

and microscopy—were included, totaling 64.3% (n = 4905) of the samples (Table 5).

Among these, qPCR identified 1712 (34.9%) as positive, whereas RDTs and

microscopy classified 1289 (26.3%) and 843 (17.2%) positives, respectively (Table 5).

A substantial number of positive samples, otherwise detectable by qPCR, were missed

by both RDTs and microscopy, suggesting the existence of subpatent infections (table 5

and Figure 5). Of all samples analyzed by qPCR, RDTs, and microscopy, 647 out of

4,905 (13.2%) were categorized as positive by all three methods (Figure 5).

Interestingly, qPCR classified 56 out of 4,905 (1.1%) samples as negative, while both

RDTs and microscopy agreed they were positive (Figure 5). Similarly, the same

proportion was agreed upon by qPCR and microscopy as positive, while RDTs

classified them as negative. Microscopy classified 415 out of 4,905 (8.5%) samples as

negative, while both RDTs and qPCR agreed they were positive.

Table 5: Overall prevalence in the 35 surveyed villages estimated using qPCR, RDTs,
and microscopy. Additionally, the table summarizes the proportion of malaria positive
samples missed by RDTs and microscopy when qPCR is used as the reference.

	No. tested	*Pf*. Positive	Prevalence
qPCR	4905	1712	34.9%
RDTs	4905	1289	26.3%
Microscopy	4905	843	17.2%
Proportion of missed positives when qPCR is the reference			
	Total Positive by PCR	**Sample missed**	**(%) of missed Positive**
RDTs	1712	650	37.9%
Microscopy	1712	1009	58.9%

Both RDT and microscopy missed several infections otherwise identified by qPCR. This

category of false negatives included cases where qPCR identified a sample as positive,

but microscopy identified it as negative, cases classified as positive by qPCR but

negative by RDTs, and cases where RDTs indicated positive results while microscopy

indicated negative result. Out of the 1712 positives detected by qPCR, RDTs missed

650 (37.97%) and microscopy missed 1009 (58.9%) (Table 5). Additionally, when

comparing microscopy to RDTs, microscopy failed to detect 45.46% (586/1289) of

malaria infections detected by RDTs. RDTs correctly identified 1062 (62.03%) samples

as true positives, while microscopy identified 703 (41.06%) as true positives (Table 6).

Furthermore, RDTs misclassified 227 (7.10%) samples as false positives, while

microscopy misclassified 140 (4.38%) (Table 6).

Figure 5: *The Venn diagram illustrates positive samples detected exclusively by a specific tool while the other two missed them (qPCR only: 594 positive, RDT only: 171 positive, Microscopy only: 84 positive). Additionally, it shows intersections indicating positive detection by two tools when one detects negative (qPCR & RDT: 415 positive; qPCR & Microscopy: 56 positive; RDT & Microscopy: 56 positive). It also indicates intersections where all tools detect positive samples (qPCR, RDT, & Microscopy: 647 positive).*

Table 6: Evaluation metrics for assessing the performance of RDTs and microscopy relative to qPCR during the fine-scale stratification of malaria risk in Ulanga and Kilombero districts, south-eastern Tanzania.

Test characteristics	RDTs		Microscopy (Thick smear)	
	Evidence from Published studies	Evidence from this study	Evidence from Published studies	Evidence from this study
True Positives (PCR positive=1712)	-	1062	-	703
False Positives (PCR negative)	-	227	-	140
True Negatives (PCR negative=3193)	-	2966	-	3057
False Negatives (PCR positive)	-	650	-	1009
Sensitivity [95% CI]	56 - 78% [55,65–68]	62.0% [95 CI: 60.0-64.2]	31-63% [55,65–68]	41.0% [95 CI: 38.8-43.4]
Specificity [95% CI]	91-99% [55,65–68]	92.9% [95 CI: 92.0-93.7]	82-100% [55,65–68]	95.6% [95 CI: 94.9-96.3]
Positive Predictive Value [95% CI]	75 -100 [69,70]	82.4% [95 CI: 80.3 -84.4]	70 -100 [66,69]	83.4% [95 CI: 80.8-86.0)
Negative Predictive Value [95% CI]	75-99 [66,68,71]	82.0% [95 CI: 80.7-83.2]	66-99 [66,68]	75.2% [95 CI: 73.8-76.4)
Kappa value [95% CI]	50-77 [55,65–68]	0.6 [95 CI: 0.6-0.6]	54% [48-60] [55,65–68]	0.4 [95 CI: 0.4-0.40)
Accuracy	-	82.1%	-	76.57%

Positive predictive values (PPVs), sensitivity, specificity, and agreement of RDTs and microscopy when compared to qPCR

Considering qPCR as the benchmark, the sensitivity (the proportion of actual positives which were correctly identified as such) of RDTs was 62.0% [95% CI: 60.0-64.2], while that of microscopy was 41.0% [95% CI: 38.8 - 43.4]. The specificity (proportion of actual negatives which were correctly identified as such) was 92.9% [95% CI: 92.00-93.7] for RDTs and 95.6% [95% CI: 94.9- 96.3] for microscopy (Table 6). Overall, the positive predictive value (PPV), i.e. the probability that individuals with a positive test result actually have the disease, was 82.4% [95% CI: 80.3 -84.4] for RDTs and 83.4% [95% CI: 80.8-86.0] for microscopy (Table 6). Importantly however, the PPV for both RDTs and microscopy varied with malaria endemicity, generally increasing with prevalence, ranging from less than 20% in very low transmission areas to over 80% in high transmission areas (Figure 06).

When considering the micro-strata generated using qPCR data, the PPV of RDTs and microscopy started at 0% in very low risk strata and gradually increased to >80% as villages shifted towards high risk strata (Figure 6A). However, when referring to the strata generated using RDTs data (Figure 6B), the PPV of both RDTs and microscopy started at 20% in very low risk strata and gradually increased to >80% in high-risk strata. The agreement between RDTs and qPCR was good (Kappa value = 0.58 [95% CI: 0.56-0.61]), while the agreement between microscopy and qPCR showed fair agreement (Kappa value = 0.42 [95%CI: 0.39-0.44]), (Table 6).

The sensitivity of both RDTs and microscopy varied by age, where RDTs sensitivity was higher for school-aged children (>80%) and dropped to 75% and 60% for 16-20 years and >20 years, respectively (Figure 6C). A similar trend of sensitivity was observed for microscopy (Figure 6C), indicating that RDTs and microscopy perform better in detecting malaria in school-aged children compared to adults.

The relationship between parasite density and the malaria detection probability by RDTs and microscopy was also examined. In this analysis, the probability of RDTs detecting

positive malaria infections was maximized reaching 1 at 100 parasites/µL, where the

logistic regression (logit (p)) model saturated (Figure 6D). At this density in contrast, the

probability of microscopy to detect malaria infections was only 0.85% (Figure 6D),

suggesting higher sensitivity of RDTs vs microscopy.

Figure 6: *Estimates of the positive predictive values (PPVs) of RDTs and microscopy at*
different malaria endemicities across the study villages, defined based on either qPCR
data (A) or RDT data (B). Panel C) illustrates the trend of sensitivity of both RDTs and
microscopy relative to age groups. Panel D) displays the detection probability of both
RDTs and microscopy relative to the parasite density estimated by qPCR.

Parasites density estimates and their correlations with *Plasmodium* prevalence

Further, asexual parasite densities estimated by both microscopy and qPCR were investigated and compared across different sex and age groups using Mann-Whitney statistics for two categories and the Kruskal-Wallis statistical test for more than two categories. Overall, PCR was capable of detecting approximately 100 fold lower parasite densities compared to microscopy. The geometric mean asexual parasite density estimated by microscopy was 2206.4 parasites/µL (95% CI: 1976.7-2462.8), while that estimated by PCR was 27.07 parasites/µL (95% CI: 23.23-31.54) (Table 7).

The asexual parasite density of infected individuals significantly differed between males and females as estimated by qPCR ($P < 0.001$), with males harboring a higher parasite density compared to females, though similar trend was observed by microscopy, this sex difference was not statistically detectable by microscopy ($P = 0.11$). Importantly, the geometric mean parasite density estimated by both microscopy and qPCR per village demonstrated a significant positive correlation with the parasite prevalence of the respective village. Thus, villages with high malaria prevalence also had high malaria parasite densities compared to villages with lower prevalence (Figure 7C-F). Considering qPCR-estimated geometric mean parasite densities, the Spearman rank correlation score was 0.77 ($P < 0.001$) and 0.76 ($P < 0.001$) when the malaria prevalence of the villages was estimated by qPCR and RDTs, respectively (Figure 7C & 7E). On the other hand, the Spearman rank correlation for the parasite density estimated by microscopy was 0.55 ($P < 0.003$) and 0.48 ($P < 0.012$) for qPCR and RDTs estimated prevalence of the villages, respectively (Figure 7D & 7F).

When parasite density by age groups were analyzed, both microscopy and qPCR revealed a significant difference in estimated malaria parasite densities between age groups based on Kruskal-Wallis statistics ($P < 0.001$) (Table 7). Pairwise tests by Mann-Whitney statistics revealed that school-aged children (5-15 years old) harbored a higher parasite density than those 16 years old and above ($P < 0.001$) as indicated by both microscopy and qPCR (Figure 7A & 7B).

499 **Table 7:** Parasite densities by gender and age as estimated using Microscopy slides and quantitative PCR assays
500

	Microscopy			qPCR		
	Geometric mean (/µL)	95% CI (/µL)	P-value	Geometric mean (/µL)	95% CI (/µL)	P-value
Overall parasite density	2206.39	1976.68-2462.81	-	27.07	23.23-31.54	-
Sex						
Female	2085.06	1775.94-2447.98	**0.11** [a]	20.48	16.76-25.02	**<0.001** [a]
Male	2341.22	2014.74-2720.61		38.32	30.33-48.41	
Age group						
5-10 years	3715.39	3040.63-4534.91	**<0.001** [b]	108.32	74.55-157.42	**<0.001** [b]
11-15 years	2398.92	2017.28—2852.75		76.83	55.26-106.83	
16-20 years	2117.02	1443.62-3104.53		30.22	17.47-52.25	
>20 years	1182	968.15-1444.34		9.05	7.59-10.79	

501 **a=** Mann-Whitney Test
502 **b=** Kruskal-Wallis Test

Figure 7: *Geometric mean parasite densities per age group estimated by A) qPCR and B) microscopy, and correlations with parasite prevalence (C-F).*

Discussion

In malaria-endemic countries, data-driven risk stratification is increasingly used at district or higher regional levels to guide intervention strategies and optimize resource allocation. Additionally, the geographical variations in levels of endemicities and the shift towards elimination in some settings necessitates finer resolution for optimal resource allocation [6,10,72]. In most settings in Africa, the data used for epidemiological stratification of malaria typically comes from rapid diagnostic tests (RDTs) or microscopy-based testing, which despite wide scale availability and low operational costs, often perform poorly in low transmission settings [41,73,74]. While direct comparisons of these diagnostic tools for fine-scale stratification are currently limited, selecting the most appropriate data sources and testing methods is crucial, as different methods can yield significantly different results depending on endemicity, particularly in elimination settings. Even without alternative testing methods, data users and decision-makers need to understand the limitations of their selected approaches, especially the weaknesses of current dominant data sources like RDTs or microscopy. In this study, we conducted a high-resolution survey of *P. falciparum* malaria in two Tanzanian districts, comparing fine-scale strata obtained using RDTs, microscopy, and qPCR assays. The findings will provide the necessary evidence to the malaria programs at national and district levels on how RDTs, microscopy, and qPCR compare when used at local levels. Moreover, the findings demonstrate that selection of appropriate test methods is essential for more precise resource allocation and improved understanding of disease patterns.

The findings highlight significant variability in malaria risk at a fine scale. Within less than 150 kilometers, malaria prevalence estimates ranged from 0% to over 50% across contiguous villages in an area broadly classified as moderate risk (~17% PfPR) by recent government stratification [44]. Such fine-scale variability is not uncommon and has been observed in several other settings [75]. In one study in Madagascar, there was a tenfold difference of malaria prevalence within a radius of less than 50 kilometers [76]. For precise micro-stratifications, this study emphasizes the importance of carefully selecting diagnostic tools, especially for local malaria elimination efforts. Our findings indicate that RDTs and microscopy have poor positive predictive values, which can be

even less than 20% in villages with very low and low transmission as the proportion of truly infected individuals is very small compared to non-infected persons. There were also significant discrepancies in the resulting micro-strata depending on the test method used. For instance, among the 35 surveyed villages, RDTs and microscopy classified 12 and 11 as very low and 6 and 10 as low risk strata, respectively, while qPCR identified only 1 village as very low and 5 as low transmission. This means RDTs and microscopy classified majority of the villages as very low to low risk while qPCR classified most villages as moderate to high risk (Table 4).

Clear demarcation of areas with very low to low risk versus those with moderate to high risk is essential, particularly in the push towards elimination. As countries increasingly adopt data-driven decision-making for malaria control, there is a risk of improper resource allocation or premature withdrawal of effective interventions from localities erroneously deemed as nearing elimination. Local authorities need to decide which data to use for local-level micro-stratification and whether RDTs, commonly used for broader-scale sub-national stratification, suffice for fine-scale local decision-making. Previous evidence has shown that hotspots identified by RDTs are less stable than those identified by microscopy and PCR [41]. Hotspots of febrile malaria infections are also generally unstable and variable over geographical spaces, while hotspots of asymptomatic cases tend to be more permanent and can be more practically targeted for transmission control [77]. In this current study, we also found significant positive correlations between malaria parasite densities and malaria prevalence in southeastern Tanzania, emphasizing the need to incorporate tests that depict sub-microscopic infections into malaria stratification and decision making to better target the hotspots. As already reported in other studies, villages in our study area, which were classified as low transmission areas had lower geometric mean parasite density compared to those with higher transmission rates.

Our findings, benchmarked against qPCR, reveal limited detection capabilities of RDTs and microscopy in overall fine-scale stratifications, especially in low transmission settings. Previous studies have emphasized the usefulness of routine hospital data for micro-stratifications [8,10,78,79]. However, evidence indicates that both microscopy and RDTs are less effective in identifying stable febrile malaria hotspots, except for

asymptomatic hotspots, which are reliably identified by microscopy [77], however still
not stable when transmission is low [80]. Our research underscores the importance of
identifying subclinical infections using sensitive tools to advance malaria elimination,
particularly through fine-scale population surveys. Additionally, there is evidence
suggesting potential benefits from integrating hospital and school-age children survey
data or even antenatal care centers [2,9]. Nevertheless, these approaches heavily rely
on rapid diagnostic tests (RDTs) as the primary tool for malaria detection. While the
World Health Organization (WHO) recommends monitoring RDT performance alongside
microscopy, our study is particularly relevant as we have directly compared the fine-
scale stratification capabilities of RDTs, microscopy, and qPCR at a fine scale.

This study demonstrated overall good agreement between RDTs and qPCR, while
microscopy showed fair agreement. However, RDTs missed over 38% of malaria
infections, particularly among adults over twenty years old, who were found to harbor
lower parasite densities compared to those under twenty years old. However, RDTs
remain useful in testing fever-positive malaria cases in hospitals and are widely
employed in population surveys due to their cost-effectiveness and ease of
implementation [32]. As evidenced in this study, carefully re-consideration of using
RDTs for finer-scale mapping and intervention planning at sub-district level should be a
priority. Similarly, microscopy missed >50% of the malaria infections detected by qPCR,
which is consistent with previous studies, including a meta-analysis of 42 studies, which
showed that microscopy misses over 50% of malaria infections [26,66]. Operational
challenges, such as the level of expertise required for accurate detection and the need
for electricity and precise sample handling procedures, contribute to these limitations.
Interestingly, microscopy underestimated malaria risk by classifying more villages as
low strata compared to qPCR. Nonetheless, microscopy still plays a crucial role when
used in conjunction with tools like RDTs, providing valuable information about malaria
parasite densities [81–83]. Here, we estimate a 100-fold higher parasite density when
measured by microscopy compared to densities measured by qPCR, consistent with
similar trends observed in previous studies [55]. The findings of this study also indicate
that the false-negative rate of microscopy decreases with increasing parasite density, a
pattern observed in other studies too [81].

The analysis also revealed variations in parasite densities across different age groups,

with school-age children (5-15 years old) exhibiting higher parasite densities compared

to individuals aged 16 and above. Notably, our study identified a reduced sensitivity of

both RDTs and microscopy among adults aged over 16 years, consistent with findings

from prior studies conducted before 2015 in various regions [84–91]. It is possible this

pattern is driven by the age-related differences in malaria parasite prevalence, which

was also observed (Table 3). Furthermore, our findings suggest that this trend may be

attributed to the lower parasite density estimates observed in adults within the study

(Figure 7A & 7B). Significantly, our research provides valuable insights, highlighting the

potential implications of these trends, particularly in fine-scale mapping scenarios,

where RDTs and microscopy may underestimate stratifications at very low and low

transmission strata, with qPCR serving as the reference standard in this study.

Ultimately, this study underscores the potential need to expand the use of molecular

approaches such as qPCR to achieve more precise mapping of malaria infections,

which could reduce false negatives and facilitate more precise resource allocations.

However, qPCR has operational challenges, including the need for well-designed

infrastructure, high costs, and expertise, and it is not portable for remote areas [82].

Efforts are underway to develop portable qPCR technologies, but cost and expertise

remain significant barriers. To address these gaps, NMCPs should develop innovative

plans, which might include: a) centralized facilities receiving and processing qPCR

samples and then doing such surveys only infrequently, say every 3 years; b) partnering

with local research organizations to support the high accuracy evaluations with nucleic

acid based tests. A related point to emphasize is the overall need for highly-sensitive,

cost-effective, and potentially reagent-free tools that align with the economic context of

malaria-endemic settings. Recent innovations such as the saliva-based tests [92] or the

use of Infrared spectroscopy (IR) and machine learning (ML) [82,93] have shown

promise in detecting malaria infections at sensitivities equivalent to PCR, but further

research are needed before these technologies can be routinely deployed. Such

reagent-free assays like the IR-AI based approaches would be particularly

transformative for scaling up effective micro stratification of malaria risk in Africa.

This study also raised some important new questions. For example, it is baffling to

observe that areas with low transmission also have persistently low parasite densities

despite individuals having low immunity compared to those in higher transmission

settings who acquire immunity and become protective. Studies have demonstrated that

in low transmission areas, highly virulent parasites are more exposed to facilitate

malaria transmission by mosquitoes compared to low virulent ones [26,94–96].

Consequently, high virulent parasites are detected and treated, leading to their removal

from the population [26,94–96]. This leaves behind low virulent parasites that are less

exposed and maintain low densities, becoming symptomatic, undetectable, and

untreated [26,94–96]. This phenomenon may contribute to long-term parasite

transmission strategies, highlighting the importance of using highly sensitive tools for

screening [55,97,98].

One limitation of this study is its failure to consider factors that may contribute to the

broader heterogeneity of malaria infections in southeastern Tanzania. Future

investigations should delve into potential environmental, geographical, immunological or

genetic diversity of the parasite influences underlying this variability. Additionally, the

biological significance of missed infections by both RDTs and microscopy was not

explored. Consequently, the study did not estimate the transmission burden associated

with these undetected positive samples, nor assess the parasite densities necessary to

sustain transmission in the population.

Conclusion

As countries progress towards malaria elimination, fine-scale mapping of malaria risk

becomes increasingly important. This study highlights significant variability in village-

level malaria risk within and between districts in southeastern Tanzania, an area where

the scale-up of effective interventions has led to substantial progress, yet cases persist

despite high intervention coverage. Secondly, the study underscores the variable

performance of different testing methods in stratifying risk. While RDTs and microscopy,

the primary test methods used in low-income endemic settings and the main sources of

data for ongoing epidemiological stratification efforts, were effective in high-transmission

areas, they performed poorly in low-transmission settings, often classifying most

villages as very low or low risk. In contrast, qPCR classified most villages as moderate

or high risk. These findings demonstrate the importance of using appropriate testing

methods for data-driven, fine-scale risk stratification to enhance targeted interventions

aimed at reducing and eliminating malaria. The study underscores the need for

improved testing approaches that are both operationally feasible and sufficiently

sensitive to enable precise mapping and effective targeting of malaria in local contexts.

More importantly, public health authorities must recognize the strengths and limitations

of their available data when planning local stratification or making decisions. While

innovation for more effective strategies is ongoing, sensitive molecular tools like qPCR,

despite their operational challenges, will be crucial for accurate malaria risk mapping

and intervention planning, especially in settings with significantly reduced risk. Going

forward, developing new tools that balance operational costs and sensitivity, particularly

in low transmission settings, will be essential for effective malaria control and eventual

elimination.

Acknowledgements

We extend our sincere gratitude to the dedicated team of the Deep Diagnostics Project

at Ifakara Health Institute for their invaluable support throughout this survey. Special

recognition is due to Abigael W. Magesa, Tumpe G. Mwandyala, Magreth C. Magoda,

Scolastica N. Hangu, Alex T. Ngonyani, Magreth I. Henry, Jiddah A. Ally, Alfred J.

Simfukwe, Amisa N. Rajabu, Scolastica A. Ndonde, Rahma S. Kikweo, Benson D.

Makua, Jacob J. Nyooka, Festo I. Tangaliola, Stephen J. Magasa, Joyce Y. Jackson,

Fidelis D. Mbena, Slyakus V, Mlembe, and Agnes Kimaro for their invaluable

contributions to conducting the surveys, administering questionnaires, screenings, and

laboratory procedures. We also express our gratitude to Judith Mshumbusi and the

procurement team for their assistance in procuring materials, often on short notice.

Additionally, we extend our thanks to Faraji Abilahi, the Ifakara laboratory manager, and

fellow scientists, students and staff at Ifakara Health institute and University of Glasgow,

who contributed direct or indirectly to this study.

Author contributions

Conceptualization: I.H.M., F.O., F.B., and S.A.B. Data curation: I.H.M., F.M., E.G.M.,

I.M., and F.E.M. Formal analysis: I.H.M., S.A.B., F.O.O., D.B.,H.S.N., F.B. Funding

acquisition: F.O.O, S.A.B., F.B., and I.H.M. Investigation: I.H.M., F.M., E.G.M., I.M.,

N.S.L, S.A, and F.E.M. Methodology: I.H.M., A.B.L., S.A.B., F.O.O., F.B. Project

administration: I.H.M., R.M.N., F.M., Resources: I.H.M., F.M.,H.S.N S.A.B., F.O.O., and

F.B. Supervision: S.A.B., F.O.O., and F.B. Validation: S.A.B., F.O.O., and F.B.

Visualization: I.H.M., S.A.B., F.O.O., F.B., N.F.K, A.L. D.B. Writing original draft: I.H.M.,

S.A.B., F.O.O., and F.B. Writing review & editing: All authors.

Funding

This work was supported in part by the Bill & Melinda Gates Foundation (Grant number

Grant No. OPP 1217647 and Grant No. INV-002138 to Ifakara Health Institute). Under

the grant conditions of the Foundation, a Creative Commons Attribution 4.0 Generic

License has already been assigned to the Author Accepted Manuscript version that

might arise from this submission. Support was also received from Rudolf Geigy

Foundation through Swiss Tropical & Public Health Institute (to Ifakara Health Institute),

Howard Hughes Medical Institute (HHMI) and Gates Foundation (Grants: OPP1099295)

awarded to FOO and HMF, Royal Society (Grant No ICA/R1/191238 to SAB University

of Glasgow and Ifakara Health Institute). Academy Medical Science Springboard Award

(ref: SBF007\100094) to FB.

Ethics approvals

708   This work was approved under Ifakara Health Institute Review Board (Ref: IHI/IRB/No:

709   1 2021) and the National Institute for Medical Research-NIMR (NIMR/HQ/R.8a/Vol.

710   1X/3735). Permission to publish this work has been granted with reference No: BD.

711   242/437/01C/6.

Data availability

713   The dataset supporting the findings is available upon a reasonable request, which can

714   be directed to either the corresponding author, the Ifakara Health Institute ethical review

715   board in Tanzania or National Health Research Ethics Committee in Tanzania with

716   reference to ethical clearance certificate of NIMR/HQ/R.8a/Vol. 1X/3735.

Reference

1. World Health Organization (WHO). Malaria policy advisory group (MPAG) meeting report, 18–20 April 2023. Geneva PP - Geneva: World Health Organization;

2. Thawer SG, Chacky F, Runge M, Reaves E, Mandike R, Lazaro S, et al. Sub-national stratification of malaria risk in mainland Tanzania: a simplified assembly of survey and routine data. Malar J. 2020;19: 1–12.

3. Mategula D, Mitambo C, Sheahan W, Masingi Mbeye N, Gumbo A, Kwizombe C, et al. Malaria Burden Stratification in Malawi- A report of a consultative workshop to inform the 2023-2030 Malawi Malaria Strategic Plan. Wellcome open Res. 2023;8: 178. doi:10.12688/wellcomeopenres.19110.1

4. Djaskano MI, Cissoko M, Diar MSI, Israel DK, Clément KH, Ali AM, et al. Stratification and adaptation of malaria control interventions in Chad. Trop Med Infect Dis. 2023;8: 450.

5. Cissoko M, Magassa M, Sanogo V, Ouologuem A, Sangaré L, Diarra M, et al. Stratification at the health district level for targeting malaria control interventions in Mali. Sci Rep. 2022;12: 8271. doi:10.1038/s41598-022-11974-3

6. Thawer SG, Golumbeanu M, Munisi K, Aaron S, Chacky F, Lazaro S, et al. The use of routine health facility data for micro-stratification of malaria risk in mainland Tanzania. Malar J. 2022;21: 345. doi:10.1186/s12936-022-04364-7

7. Rijal KR, Adhikari B, Adhikari N, Dumre SP, Banjara MS, Shrestha UT, et al. Micro-stratification of malaria risk in Nepal: implications for malaria control and elimination. Trop Med Health. 2019;47: 21. doi:10.1186/s41182-019-0148-7

8. Thawer SG, Golumbeanu M, Lazaro S, Chacky F, Munisi K, Aaron S, et al. Spatio-temporal modelling of routine health facility data for malaria risk micro-stratification in mainland Tanzania. Sci Rep. 2023;13: 10600. doi:10.1038/s41598-023-37669-x

9. Kitojo C, Gutman JR, Chacky F, Kigadye E, Mkude S, Mandike R, et al. Estimating malaria burden among pregnant women using data from antenatal care centres in Tanzania: a population-based study. Lancet Glob Heal. 2019;7: e1695–e1705.

10. Alegana VA, Suiyanka L, Macharia PM, Ikahu-Muchangi G, Snow RW. Malaria micro-stratification using routine surveillance data in Western Kenya. Malar J. 2021;20: 1–9.

11. Buring JE. Epidemiology in medicine. Lippincott Williams & Wilkins; 1987.

12. World Health Organization (WHO). A framework for malaria elimination. World Health Organization; 2017.

13. Alegana VA, Macharia PM, Muchiri S, Mumo E, Oyugi E, Kamau A, et al. Plasmodium falciparum parasite prevalence in East Africa: Updating data for

malaria stratification. PLOS Glob Public Heal. 2021;1: e0000014.

14. Ssempiira J, Kissa J, Nambuusi B, Kyozira C, Rutazaana D, Mukooyo E, et al. The effect of case management and vector-control interventions on space–time patterns of malaria incidence in Uganda. Malar J. 2018;17: 1–11.

15. Yekutiel P. Problems of epidemiology in malaria eradication. Bull World Health Organ. 1960;22: 669–683.

16. Brooker S, Kolaczinski JH, Gitonga CW, Noor AM, Snow RW. The use of schools for malaria surveillance and programme evaluation in Africa. Malar J. 2009;8: 1–9.

17. Smith Gueye C, Newby G, Tulloch J, Slutsker L, Tanner M, Gosling RD. The central role of national programme management for the achievement of malaria elimination: a cross case-study analysis of nine malaria programmes. Malar J. 2016;15: 1–21.

18. Bakar S, Holzschuh A, Ross A, Stuck L, Abdul R, Al-Mafazy A-WH, et al. Risk of imported malaria infections in Zanzibar: a cross-sectional study. Infect Dis Poverty. 2023;12: 80. doi:10.1186/s40249-023-01129-5

19. ICF. Tanzania demographic and health survey 2022 - final report . Rockville, Maryland, USA : ICF ; 2023.

20. Chacky F, Runge M, Rumisha SF, Machafuko P, Chaki P, Massaga JJ, et al. Nationwide school malaria parasitaemia survey in public primary schools, the United Republic of Tanzania. Malar J. 2018;17: 452. doi:10.1186/s12936-018-2601-1

21. Ashton RA, Bennett A, Al-Mafazy A-W, Abass AK, Msellem MI, McElroy P, et al. Use of routine health information system data to evaluate impact of malaria control interventions in Zanzibar, Tanzania from 2000 to 2015. EClinicalMedicine. 2019;12: 11–19.

22. Bastiaens GJH, Bousema T, Leslie T. Scale-up of malaria rapid diagnostic tests and artemisinin-based combination therapy: challenges and perspectives in sub-Saharan Africa. PLoS Med. 2014;11: e1001590.

23. Noor AM, Gething PW, Alegana VA, Patil AP, Hay SI, Muchiri E, et al. The risks of malaria infection in Kenya in 2009. BMC Infect Dis. 2009;9: 1–14.

24. World Health Organization (WHO), Regional Office for the Western Pacific. Giemsa Staining of Malaria Blood Films. Malaria Microscopy Standard Operating Procedure – Mm-Sop-07a 1. 2016.

25. Tangpukdee N, Duangdee C, Wilairatana P, Krudsood S. Malaria diagnosis: a brief review. Korean J Parasitol. 2009;47: 93. doi:10.3347/kjp.2009.47.2.93

26. Okell LC, Bousema T, Griffin JT, Ouédraogo AL, Ghani AC, Drakeley CJ. Factors determining the occurrence of submicroscopic malaria infections and their relevance for control. Nat Commun. 2012;3: 1–9. doi:10.1038/ncomms2241

27.	Manjurano A, Okell L, Lukindo T, Reyburn H, Olomi R, Roper C, et al. Association of sub-microscopic malaria parasite carriage with transmission intensity in north-eastern Tanzania. 2011; 1–8.

28.	Payne D. Use and limitations of light microscopy for diagnosing malaria at the primary health care level. Bull World Health Organ. 1988;66: 621.

29.	UNITAID. 2015 Malaria Diagnostics Landscape Update. 2015.

30.	Moody A. Rapid diagnostic tests for malaria parasites. Clin Microbiol Rev. 2002;15: 66–78.

31.	Shillcutt S, Morel C, Goodman C, Coleman P, Bell D, Whitty CJM, et al. Cost-effectiveness of malaria diagnostic methods in sub-Saharan Africa in an era of combination therapy. Bull World Health Organ. 2008;86: 101–110. doi:10.2471/BLT.07.042259

32.	Cunningham J, Jones S, Gatton ML, Barnwell JW, Cheng Q, Chiodini PL, et al. A review of the WHO malaria rapid diagnostic test product testing programme (2008-2018): Performance, procurement and policy. Malar J. 2019;18. doi:10.1186/s12936-019-3028-z

33.	Drakeley C, Reyburn H. Out with the old, in with the new: the utility of rapid diagnostic tests for malaria diagnosis in Africa. Trans R Soc Trop Med Hyg. 2009;103: 333–337.

34.	World Health Organization (WHO). Universal access to malaria diagnostic testing: An operational manual. WHO Press. 2011;12: 31A.

35.	Roth JM, Korevaar DA, Leeflang MMG, Mens PF. Molecular malaria diagnostics: a systematic review and meta-analysis. Crit Rev Clin Lab Sci. 2016;53: 87–105.

36.	Snounou G, Viriyakosol S, Jarra W, Thaithong S, Brown KN. Identification of the four human malaria parasite species in field samples by the polymerase chain reaction and detection of a high prevalence of mixed infections. Mol Biochem Parasitol. 1993;58: 283–292. doi:https://doi.org/10.1016/0166-6851(93)90050-8

37.	Kamau E, Alemayehu S, Feghali KC, Saunders D, Ockenhouse CF. Multiplex qPCR for detection and absolute quantification of malaria. PLoS One. 2013;8: e71539.

38.	Snounou G. Detection and identification of the four malaria parasite species infecting humans by PCR amplification. Methods Mol Biol. 1996;50: 263–291. doi:10.1385/0-89603-323-6:263

39.	Perandin F, Manca N, Calderaro A, Piccolo G, Galati L, Ricci L, et al. Development of a real-time PCR assay for detection of Plasmodium falciparum, Plasmodium vivax, and Plasmodium ovale for routine clinical diagnosis. J Clin Microbiol. 2004;42: 1214–1219.

40.	Moonen B, Cohen JM, Snow RW, Slutsker L, Drakeley C, Smith DL, et al. Operational strategies to achieve and maintain malaria elimination. Lancet

(London, England). 2010;376: 1592–1603. doi:10.1016/S0140-6736(10)61269-X

41.   Mogeni P, Williams TN, Omedo I, Kimani D, Ngoi JM, Mwacharo J, et al. Detecting Malaria Hotspots: A Comparison of Rapid Diagnostic Test, Microscopy, and Polymerase Chain Reaction. J Infect Dis. 2017;216: 1091–1098. doi:10.1093/infdis/jix321

42.   Mosha JF, Sturrock HJW, Greenhouse B, Greenwood B, Sutherland CJ, Gadalla N, et al. Epidemiology of subpatent Plasmodium falciparum infection: implications for detection of hotspots with imperfect diagnostics. Malar J. 2013;12: 1–9.

43.   Finda MF, Limwagu AJ, Ngowo HS, Matowo NS, Swai JK, Kaindoa E, et al. Dramatic decreases of malaria transmission intensities in Ifakara, south-eastern Tanzania since early 2000s. Malar J. 2018;17: 1–18.

44.   MoHCDGEC. Ministry of Health, Community Development, Gender, Elderly and Children (MoHCDGEC). The 2021 school malaria and nutrition survey (SMNS) report Mainland Tanzania. 2022.

45.   Ruth EP, Kashaigili JJ, Majule AE. Availability, Access and Use of Weather and Climate Information by Smallholder Farmers in the Kilombero River Catchment, Tanzania. Clim Impacts Agric Nat Resour Sustain Africa. 2020; 489–506.

46.   Kasiulevičius V, Šapoka V, Filipavičiūtė R. Sample size calculation in epidemiological studies. Gerontologija. 2006;7: 225–231.

47.   Arya R, Antonisamy B, Kumar S. Sample size estimation in prevalence studies. Indian J Pediatr. 2012;79: 1482–1488.

48.   Cochran WG. Sampling Techniques. Wiley; 1977.

49.   Mouatcho JC, Goldring JPD. Malaria rapid diagnostic tests: challenges and prospects. J Med Microbiol. 2013;62: 1491–1505.

50.   National guidelines for malaria diagnosis and treatment, 2006 / United Republic of Tanzania, Ministry of Health and Social Welfare. [Dar es Salaam]□: National Malaria Control Programme, [2006].

51.   Bailey JW, Williams J, Bain BJ, Parker-Williams J, Chiodini PL, Haematology GHTF of the BC for S in. Guideline: the laboratory diagnosis of malaria. Br J Haematol. 2013;163: 573–580.

52.   World Health Organization (WHO). Basic malaria microscopy. World Health Organization; 2010.

53.   World Health Organization (WHO). Malaria parasite counting. World Health Organization; 2016.

54.   Quick-DNA^{TM} Miniprep Kit, Rapid and simple isolation of ultra-pure DNA from biological liquids and cells.

55.   Hofer LM, Kweyamba PA, Sayi RM, Chabo MS, Maitra SL, Moore SJ, et al. Malaria rapid diagnostic tests reliably detect asymptomatic Plasmodium

falciparum infections in school-aged children that are infectious to mosquitoes. Parasit Vectors. 2023;16: 217.

56. Hofmann N, Mwingira F, Shekalaghe S, Robinson LJ, Mueller I, Felger I. Ultra-Sensitive Detection of Plasmodium falciparum by Amplification of Multi-Copy Subtelomeric Targets. PLOS Med. 2015;12: e1001788.

57. Hartung C, Lerer A, Anokwa Y, Tseng C, Brunette W, Borriello G. Open data kit: tools to build information services for developing regions. Proceedings of the 4th ACM/IEEE international conference on information and communication technologies and development. 2010. pp. 1–12.

58. McKinney W. Python for data analysis: Data wrangling with Pandas, NumPy, and IPython. " O'Reilly Media, Inc."; 2012.

59. Seabold S, Perktold J. statsmodels: Econometric and statistical modeling with python. 9th Python in Science Conference. 2010.

60. Landis JR, Koch GG. The Measurement of Observer Agreement for Categorical Data Published by: International Biometric Society Stable URL: https://www.jstor.org/stable/2529310. 1977;33: 159–174.

61. Munoz SR, Bangdiwala SI. Interpretation of Kappa and B statistics measures of agreement. J Appl Stat. 1997;24: 105–112.

62. McKight PE, Najab J. Kruskal-wallis test. corsini Encycl Psychol. 2010; 1.

63. McKnight PE, Najab J. Mann-Whitney U Test. Corsini Encycl Psychol. 2010; 1.

64. Sedgwick P. Spearman's rank correlation coefficient. Bmj. 2014;349.

65. Berzosa P, de Lucio A, Romay-Barja M, Herrador Z, González V, García L, Fernández-Martínez A, Santana-Morales M, Ncogo P, Valladares B, Riloha M BA. Comparison of three diagnostic methods (microscopy, RDT, and PCR) for the detection of malaria parasites in representative samples from Equatorial Guinea. Malar J. 2018;17: 1–12.

66. Mfuh KO, Achonduh-Atijegbe OA, Bekindaka ON, Esemu LF, Mbakop CD, Gandhi K, et al. A comparison of thick-film microscopy, rapid diagnostic test, and polymerase chain reaction for accurate diagnosis of Plasmodium falciparum malaria. Malar J. 2019;18: 1–8.

67. Ngasala B, Mutemi DD, Mwaiswelo RO. Diagnostic performance of malaria rapid diagnostic test and microscopy compared with PCR for detection of Plasmodium falciparum infections among primary schoolchildren in Kibiti District, Eastern Tanzania: an area with moderate malaria transmission. Am J Trop Med Hyg. 2019;101: 809.

68. Opoku Afriyie S, Addison TK, Gebre Y, Mutala A-H, Antwi KB, Abbas DA, et al. Accuracy of diagnosis among clinical malaria patients: comparing microscopy, RDT and a highly sensitive quantitative PCR looking at the implications for submicroscopic infections. Malar J. 2023;22: 76. doi:10.1186/s12936-023-04506-

69. Abdalla ZA, Rahma NA, Hassan EE, Abdallah TM, Hamad HE, Omer SA, et al. The diagnostic performance of rapid diagnostic tests and microscopy for malaria diagnosis in eastern Sudan using a nested polymerase chain reaction assay as a reference standard. Trans R Soc Trop Med Hyg. 2019;113: 701–705.

70. Galatas B, Mayor A, Gupta H, Balanza N, Jang IK, Nhamussua L, et al. Field performance of ultrasensitive and conventional malaria rapid diagnostic tests in southern Mozambique. Malar J. 2020;19: 1–15.

71. Fransisca L, Kusnanto JH, Satoto TBT, Sebayang B, Supriyanto , Andriyan E, et al. Comparison of rapid diagnostic test Plasmotec Malaria-3, microscopy, and quantitative real-time PCR for diagnoses of Plasmodium falciparum and Plasmodium vivax infections in Mimika Regency, Papua, Indonesia. Malar J. 2015;14: 1–11.

72. Ozodiegwu ID, Ogunwale AO, Surakat O, Akinyemi JO, Bamgboye EA, Fagbamigbe AF, et al. Description of the design of a mixed-methods study to assess the burden and determinants of malaria transmission for tailoring of interventions (microstratification) in Ibadan and Kano metropolis. Malar J. 2023;22: 255. doi:10.1186/s12936-023-04684-2

73. Sumari D, Mwingira F, Selemani M, Mugasa J, Mugittu K, Gwakisa P. Malaria prevalence in asymptomatic and symptomatic children in Kiwangwa, Bagamoyo district, Tanzania. Malar J. 2017;16: 1–7.

74. Jiram AI, Ooi CH, Rubio JM, Hisam S, Karnan G, Sukor NM, et al. Evidence of asymptomatic submicroscopic malaria in low transmission areas in Belaga district, Kapit division, Sarawak, Malaysia. Malar J. 2019;18: 1–12.

75. Amratia P, Psychas P, Abuaku B, Ahorlu C, Millar J, Oppong S, et al. Characterizing local-scale heterogeneity of malaria risk: a case study in Bunkpurugu-Yunyoo district in northern Ghana. Malar J. 2019;18: 1–14.

76. Rice BL, Golden CD, Randriamady HJ, Rakotomalala AANA, Vonona MA, Anjaranirina EJG, et al. Fine-scale variation in malaria prevalence across ecological regions in Madagascar: a cross-sectional study. BMC Public Health. 2021;21: 1018.

77. Bejon P, Williams TN, Liljander A, Noor AM, Wambua J, Ogada E, et al. Stable and unstable malaria hotspots in longitudinal cohort studies in Kenya. PLoS Med. 2010;7: e1000304.

78. Alegana VA, Okiro EA, Snow RW. Routine data for malaria morbidity estimation in Africa: challenges and prospects. BMC Med. 2020;18: 1–13.

79. Oduro AR, Bojang KA, Conway DJ, Corrah T, Greenwood BM, Schellenberg D. Health centre surveys as a potential tool for monitoring malaria epidemiology by area and over time. PLoS One. 2011;6: e26305.

80. Kangoye DT, Noor A, Midega J, Mwongeli J, Mkabili D, Mogeni P, et al. Malaria hotspots defined by clinical malaria, asymptomatic carriage, PCR and vector numbers in a low transmission area on the Kenyan Coast. Malar J. 2016;15: 1–13.

81. Oyibo W, Latham V, Oladipo O, Ntadom G, Uhomoibhi P, Ogbulafor N, et al. Malaria parasite density and detailed qualitative microscopy enhances large-scale profiling of infection endemicity in Nigeria. Sci Rep. 2023;13: 1599. doi:10.1038/s41598-023-27535-1

82. Mshani IH, Siria DJ, Mwanga EP, Sow BBD, Sanou R, Opiyo M, et al. Key considerations, target product profiles, and research gaps in the application of infrared spectroscopy and artificial intelligence for malaria surveillance and diagnosis. Malar J. 2023;22: 346. doi:10.1186/s12936-023-04780-3

83. Bejon P, Andrews L, Hunt-Cooke A, Sanderson F, Gilbert SC, Hill AVS. Thick blood film examination for Plasmodium falciparum malaria has reduced sensitivity and underestimates parasite density. Malar J. 2006;5: 1–4.

84. Swarthout TD, Counihan H, Senga RKK, Van den Broek I. Paracheck-Pf® accuracy and recently treated Plasmodium falciparum infections: is there a risk of over-diagnosis? Malar J. 2007;6: 1–6.

85. Reyburn H, Mbakilwa H, Mwangi R, Mwerinde O, Olomi R, Drakeley C, et al. Rapid diagnostic tests compared with malaria microscopy for guiding outpatient treatment of febrile illness in Tanzania: randomised trial. Bmj. 2007;334: 403.

86. Abeku TA, Kristan M, Jones C, Beard J, Mueller DH, Okia M, et al. Determinants of the accuracy of rapid diagnostic tests in malaria case management: evidence from low and moderate transmission settings in the East African highlands. Malar J. 2008;7: 1–10.

87. Hopkins H, Kambale W, Kamya MR, Staedke SG, Dorsey G, Rosenthal PJ. Comparison of HRP2-and pLDH-based rapid diagnostic tests for malaria with longitudinal follow-up in Kampala, Uganda. Am J Trop Med Hyg. 2007;76: 1092–1097.

88. Harris I, Sharrock WW, Bain LM, Gray K-A, Bobogare A, Boaz L, et al. A large proportion of asymptomatic Plasmodium infections with low and sub-microscopic parasite densities in the low transmission setting of Temotu Province, Solomon Islands: challenges for malaria diagnostics in an elimination setting. Malar J. 2010;9: 1–8.

89. Laurent A, Schellenberg J, Shirima K, Ketende SC, Alonso PL, Mshinda H, et al. Performance of HRP-2 based rapid diagnostic test for malaria and its variation with age in an area of intense malaria transmission in southern Tanzania. Malar J. 2010;9: 1–9.

90. Mtove G, Nadjm B, Amos B, Hendriksen ICE, Muro F, Reyburn H. Use of an HRP2-based rapid diagnostic test to guide treatment of children admitted to hospital in a malaria-endemic area of north-east Tanzania. Trop Med Int Heal.

2011;16: 545–550.

91.   Fryauff DJ, Gomez-Saladin E, Sumawinata I, Sutamihardja MA, Tuti S, Subianto B, et al. Comparative performance of the ParaSight F test for detection of Plasmodium falciparum in malaria-immune and nonimmune populations in Irian Jaya, Indonesia. Bull World Health Organ. 1997;75: 547.

92.   Tao D, McGill B, Hamerly T, Kobayashi T, Khare P, Dziedzic A, et al. A saliva-based rapid test to quantify the infectious subclinical malaria parasite reservoir. Sci Transl Med. 2019;11: eaan4479.

93.   Mshani I, Jackson F, Mwanga R, Kweyamba P, Mwanga E, Tambwe M, et al. Using AI-Powered Mid-Infrared Spectroscopy for Malaria Screening Under Varying Parasite Densities and Anaemic Conditions. 2024.

94.   Okell LC, Ghani AC, Lyons E, Drakeley CJ. Submicroscopic infection in plasmodium falciparum-endemic populations: A systematic review and meta-analysis. J Infect Dis. 2009;200: 1509–1517. doi:10.1086/644781

95.   Bousema T, Okell L, Felger I, Drakeley C. Asymptomatic malaria infections: detectability, transmissibility and public health relevance. Nat Rev Microbiol. 2014;12: 833–840. doi:10.1038/nrmicro3364

96.   Slater HC, Ross A, Felger I, Hofmann NE, Robinson L, Cook J, et al. The temporal dynamics and infectiousness of subpatent Plasmodium falciparum infections in relation to parasite density. Nat Commun. 2019;10: 1433. doi:10.1038/s41467-019-09441-1

97.   Nguyen T-N, von Seidlein L, Nguyen T-V, Truong P-N, Do Hung S, Pham H-T, et al. The persistence and oscillations of submicroscopic Plasmodium falciparum and Plasmodium vivax infections over time in Vietnam: an open cohort study. Lancet Infect Dis. 2018;18: 565–572.

98.   Björkman AB. Asymptomatic low-density malaria infections: a parasite survival strategy? Lancet Infect Dis. 2018;18: 485–486.