Complement 3a Receptor 1 on Macrophages and Kupffer cells is not required for the

Pathogenesis of Metabolic Dysfunction-Associated Steatotic Liver Disease

Edwin A. Homan[1], Ankit Gilani[1], Alfonso Rubio-Navarro[1], Maya A. Johnson[1], Odin M.

Schaepkens[1], Eric Cortada[1], Renan Pereira de Lima[1], Lisa Stoll[1], James C. Lo[1*].

Affiliations:

[1] Division of Cardiology, Department of Medicine, Cardiovascular Research Institute, Weill

Center for Metabolic Health, Weill Cornell Medicine, New York, New York, 10021

*Corresponding author. Email: jlo@med.cornell.edu.

Abstract

Together with obesity and type 2 diabetes, metabolic dysfunction-associated steatotic liver

disease (MASLD) is a growing global epidemic. Activation of the complement system and

infiltration of macrophages has been linked to progression of metabolic liver disease. The role of

complement receptors in macrophage activation and recruitment in MASLD remains poorly

understood. In human and mouse, *C3AR1* in the liver is expressed primarily in Kupffer cells, but

is downregulated in humans with MASLD compared to obese controls. To test the role of

complement 3a receptor (C3aR1) on macrophages and liver resident macrophages in MASLD,

we generated mice deficient in C3aR1 on all macrophages (C3aR1-MφKO) or specifically in

liver Kupffer cells (C3aR1-KpKO) and subjected them to a model of metabolic steatotic liver

disease. We show that macrophages account for the vast majority of *C3ar1* expression in the

liver. Overall, C3aR1-MφKO and C3aR1-KpKO mice have similar body weight gain without

significant alterations in glucose homeostasis, hepatic steatosis and fibrosis, compared to

controls on a MASLD-inducing diet. This study demonstrates that C3aR1 deletion in

macrophages or Kupffer cells, the predominant liver cell type expressing *C3aR1*, has no

significant effect on liver steatosis, inflammation or fibrosis in a dietary MASLD model.

Keywords: obesity, hepatic steatosis, steatohepatitis, C3aR1, macrophage, Kupffer cell

Introduction

Obesity and related metabolic diseases such as type 2 diabetes (T2D) and metabolic dysfunction-associated steatotic liver disease (MASLD) remain a worldwide epidemic with increasing prevalence[1,2]. MASLD describes the constellation of hepatic lipid deposition, inflammation, and fibrosis associated with obesity and T2D that ultimately leads to MASH cirrhosis, which has become the leading cause of liver transplantation in the United States[3-6]. Notably, MASLD is increasingly recognized as an important risk-enhancing factor for atherosclerotic cardiovascular disease[7,8].

Liver macrophages help to maintain hepatic homeostasis and consist of embryo-derived resident macrophages called Kupffer cells, which self-renew and do not migrate, or peripheral monocyte-derived macrophages, which infiltrate into liver tissue upon metabolic or toxic liver injury and under certain circumstances can take on Kupffer cell-like identity[9-13]. In obesity, bone marrow-derived myeloid cells migrate to the steatotic liver, and pro-inflammatory recruited macrophages are postulated to drive the progression of MASLD to MASH[14]. Spatial proteogenomics reveals a population of lipid-associated macrophages near bile canaliculi that is induced by local lipid exposure and drives fibrosis in steatotic regions of murine and human liver[15]. In addition, deep transcriptomic profiling in human MASLD has identified candidate gene signatures for steatohepatitis and fibrosis with possible therapeutic implications[16].

Activation of the body's complement system leads to increased cell lysis, phagocytosis, and inflammation[17], and it is increasingly recognized as an important contributor to regulation of metabolic disorders such as T2D and MASLD[18,19]. In human liver biopsies, higher lobular inflammation scores correlate with activation of the complement alternative pathway[20], which can signal *via* the C3a receptor 1 (C3aR1), a G_i-coupled G protein-coupled receptor[21]. The complement 3 polypeptide (C3) is cleaved by C3 convertase to the activated fragment, C3a, which then binds C3aR1[22]. Complement factor D (CFD), also known as the adipokine adipsin, is the rate-limiting step in the alternative pathway of complement activation[23,24].

Several studies have reported opposing roles of adipsin and C3aR1 on hepatic steatosis

in diet-induced obesity[25-27]. Our lab has found that adipsin/CFD is critical for maintaining

pancreatic beta cell mass and function[28,29]. Murine obese and diabetic models such as *db/db*

mice and high fat diet (HFD) feeding result in very low circulating adipsin[23]. Replenishing adipsin

in *db/db* mice raises levels of C3a and insulin, lowers blood glucose levels, and inhibits hepatic

gluconeogenesis[28]. However, whole-body deletion of C3aR1 decreases macrophage infiltration

and activation in adipose tissue, protects from HFD-induced obesity and glucose intolerance,

and decreases hepatic steatosis and inflammation[30]. In a model of fibrosing steatohepatitis,

bone marrow-derived macrophages were found to activate hepatic stellate cells, which was

blunted in whole-body C3aR1 KO mice[31].

In the present study we aim to explore the macrophage-specific effect of complement

receptor signaling in MASLD pathogenesis. To determine the consequences of macrophage and

Kupffer cell ablation of C3aR1, we use a murine dietary model of MALFD/MASH, the Gubra

Amylin Nash (GAN) diet, which has macronutrient similarities to the Western diet and produces

similar histologic and transcriptomic changes to human MASLD/MASH[32-34].

Results

C3AR1 is expressed in human and mouse liver, primarily in Kupffer cells.

In the scRNA-Seq database, Human Protein Atlas, *C3AR1* is broadly expressed

throughout the body, with increased abundance in tissues rich in immunologic cell types, such

as bone marrow and appendix (Fig. 1A)[35]. In a single-cell transcriptomic database of healthy

human liver, *C3AR1* expression predominates in the macrophage and Kupffer cell population,

with minimal-to-undetectable *C3AR1* expression in hepatocytes or hepatic stellate cells by

scRNA-Seq (Fig. 1B)[36]. In the mouse liver scRNA-Seq database, Tabula Muris, *C3ar1* is

similarly expressed primarily in Kupffer cells (Fig. S1)[37].

Hepatic CFD and C3AR1 are downregulated in human MASLD/MASH.

We also examined data from Suppli and coworkers, who performed bulk transcriptomic

analysis of human liver samples from an age-matched cohort of healthy controls and obese

controls without MASLD, as well as MASLD and MASH patients without cirrhosis[38]. Both *CFD*

and C3AR1 were unchanged in obese subjects without MASLD compared to healthy controls,

but both *CFD* and *C3AR1* were significantly downregulated in liver biopsies from both MASLD

and MASH patients compared to both healthy controls and obese subjects without MASLD (Fig.

1C). Interestingly, both *CFD* and *C3AR1* levels were slightly higher in MASH individuals

compared to those with MASLD only.

Murine MASH model recapitulates key features of human MASH

At 5 weeks of age, we subjected *C3ar1* flox/flox control mice to standard regular diet

(RD) or GAN diet[32,33]. After 28 weeks of GAN diet, male mice gained body weight compared to

RD (Fig. 1D), primarily as fat mass (Fig. S2-3), but weight gain in female GAN-fed mice was

attenuated. Histologic signs of MASLD were present in GAN-fed mice (Fig. 1E), most notably

hepatic steatosis and hepatocyte ballooning (Fig. 1F), and liver fibrosis measured by collagen

deposition nearly doubled with GAN compared to RD (Fig. 1G). Both hepatic *C3ar1* and *Cfd*

gene expression were robustly increased on GAN compared to RD, as were markers of

macrophage infiltration, hepatic inflammation, and fibrosis, including collagen gene expression,

indicating progression to fibrotic MASH (Fig. 1H). In female control mice on GAN diet, there

were no significant differences in *C3ar1* expression or other gene markers, though there was a

nonsignificant trend toward increased inflammation and fibrosis compared to regular diet (Fig.

S4).

Macrophage-specific C3aR1 deletion does not alter glucose homeostasis.

Owing to higher levels of *C3ar1* in murine MASLD and the differential regulation of

C3AR1 gene in MASLD humans, this motivated us to interrogate the role of pathophysiological

role of *C3ar1* in macrophages in MASLD. We generated transgenic mice with macrophage-

specific deletion of C3aR1 by crossing *C3ar1* floxed mice with LysM-Cre transgenic mice

(C3aR1-MφKO) to target both liver resident macrophages and recruited monocytes. *C3ar1*

floxed mice were used as controls. Successful deletion of *C3ar1* in macrophages from the

C3aR1-MφKO mouse was confirmed by quantitative RT-PCR of isolated peritoneal

macrophages that were F4/80+ and CD68+ by fluorescence-activated cell sorting (Fig. 2A). In

liver tissue, *C3ar1* expression was reduced by ~88% in both male and female C3aR1-MφKO

(Fig. 2B). These results indicate that macrophages account for the vast majority of *C3ar1*

expression in the liver.

When placed on GAN diet, there was no significant difference in weight gain between

control and C3aR1-MφKO mice (Fig. 2C). There was similarly no difference in percent lean or

fat mass between these mice (Fig. 2D). Glucose tolerance tests performed in fasted mice after

27 weeks GAN diet found no significant differences between control and C3aR1-MφKO mice

(Fig. 2E). There was also no difference in insulin sensitivity as measured by insulin tolerance

tests in male mice (Fig. S5). Insulin resistance as measured by comparing the ratio of fasting

glucose level to fasting insulin level (HOMA-IR) was also unchanged between controls and

C3aR1-MφKO mice (Fig. S6). Circulating serum ALT levels were unchanged in male control and

C3aR1-MφKO mice on GAN diet (Fig. S7).

Macrophage-specific C3aR1 deletion does not significantly impact hepatic steatosis or fibrosis.

Liver samples collected after 28-30 weeks of GAN or regular diet did not show significant

differences in liver mass between control and C3aR1-MφKO mice (Fig. 2F). Male mice on GAN

diet developed similar qualitative appearance on histology (Fig. 2G), and slide image analysis

showed similar proportions of lipid droplet area and collagen area (Figs. 2H, 2I). This indicates

that there were no significant differences in steatosis or fibrosis between GAN-fed control and

C3aR1-MφKO male mice. While *C3ar1* expression was markedly reduced in the C3aR1-MφKO

liver tissue (Fig. 2B), there were no detectable gene expression changes in markers of fibrosis,

inflammation, or lipid handling on either GAN or regular diet (Fig. 2J-K). Similarly, in female mice

there were also no significant differences between control and C3aR1-MφKO mouse liver on

either GAN or regular diet in a subset of key gene markers of fibrosis or inflammation (Fig. S8).

Kupffer cell-specific C3aR1 deletion does not alter weight gain or glucose homeostasis.

To explore whether there may be competing effects between recruited monocytes and

liver resident macrophages (Kupffer cells), we next generated Kupffer cell-specific C3aR1

knockout mice (C3aR1-KpKO) by crossing *C3ar1* floxed mice to Clec4f-Cre transgenic mice and

fed them GAN diet. *C3ar1* floxed mice were used as controls. Body weight gain was similar

between genotypes for both male and female mice (Fig. 3A), and there was no difference in

body composition between control and C3aR1-KpKO mice on GAN diet (Fig. 3B). There was

similarly no significant difference in glucose homeostasis between the genotypes during a

glucose tolerance test (Fig. 3C).

Kupffer cell-specific C3aR1 deletion does not significantly impact hepatic steatosis or fibrosis.

Liver mass was not significantly different between control and C3aR1-KpKO mice on

GAN diet (Fig. 3D). Liver sections appeared qualitatively similar by histology stained with

Masson's trichrome (Fig. 3E). There were similar levels of hepatic steatosis in these mice as

measured by percent lipid droplet area (Fig. 3F). When measured by collagen proportional area,

there was no significant differences in liver fibrosis between C3aR1-KpKO and control mice

(Fig. 3G). While *C3ar1* expression was reduced by 73% in liver tissue of C3aR1-KpKO mice,

there were no significant differences in expression of inflammatory, fibrotic, or lipid handling

gene markers (Fig. 3H). *C3ar1* expression similarly decreased by ~90% in liver tissue of female

C3aR1-KpKO mice fed regular diet compared to control mice (Fig. S9). These data also indicate

that Kupffer cells account for ~80% of hepatic *C3ar1* gene expression in our mouse model of

MASLD/MASH.

Discussion

Overall, we found that macrophage or Kupffer cell expression of *C3ar1* does not impact

body weight gain or histologic/transcriptomic features of MASLD/MASH in a murine dietary

model. Deletion of C3aR1 in the macrophage population throughout the body, or specifically in

Kupffer cells, did not affect weight gain, glucose homeostasis, or extent of hepatic

steatosis/fibrosis. With long term GAN diet feeding that has been previously shown to model

human MASLD/MASH, we did not observe significant differences in liver abnormalities with the

KO mice.

Our findings in macrophage-specific C3aR1 KO mice contrast with prior observations in

whole-body C3aR1 KO mice[30], which are protected from diet-induced obesity, have improved

glucose tolerance, and exhibit decreased hepatic steatosis. In both our macrophage- and

Kupffer cell-specific C3aR1 KO mice, which had similar degrees of obesity compared to

controls, there was no detectable effect on liver steatosis or fibrosis despite the near abrogation

of *C3ar1* expression. This raises the possibility that the lower levels of hepatic steatosis and

insulin resistance previously observed in the whole body C3aR1 KO mice may be secondary to

protection from obesity. Protection from diet-induced obesity in whole-body C3aR1 KO mice

may be mediated by a non-macrophage cell type, since our macrophage-specific C3aR1 KO

mice were not afforded this protection. The *C3ar1*-expressing cell types that promote obesity

and MASLD remains to be determined.

Our laboratory recently reported sex-dependent regulation of thermogenic adipose

tissue mediated by adipocyte-derived C3aR1[39]. However, no such sexual dimorphism was

observed in hepatic expression of key MASH genes in response to GAN diet in our

macrophage- or Kupffer cell-specific C3aR1-deficient mice. Other work has suggested possible

compensatory effects from its sister anaphylatoxin receptor C5aR1, with increased cold-induced

adipocyte browning and attenuated diet-induced obesity seen in C3aR1/C5aR1 double KO

mice[40].

The strengths of our study include careful metabolic and transcriptomic phenotyping of

cell type-specific transgenic mice. Some limitations were our use of a single MASLD dietary

model and our focus on the C3aR1 pathway. While the GAN diet recapitulates many features of

human MASH due to its similarity to Western diet[34], relatively low levels of fibrosis were seen in

our study, potentially related to initiating the diet at young age; more rapid fibrosis induction has

been seen when GAN diet is initiated at older ages[41]. It is possible that in other models of liver

injury that we did not test (e.g., short-term treatment with a hepatotoxin such as carbon

tetrachloride),[42] there may be differences in liver injury in mice lacking *C3ar1* in macrophages.

However, the GAN diet model has been shown to better parallel the gene expression changes

in human MAFLD/MASH.[33] Lastly, while *C3AR1/C3ar1* expression is very low in non-

macrophage cells (Fig. B, S1), C3aR1 signaling on other hepatic cell types not explored in this

study, such as hepatic stellate cells, could mediate the observed effect in the whole-body

C3aR1 KO mouse.

Deletion of C3aR1 in macrophages generally, or in liver resident macrophages

specifically, had no major effect on systemic glucose homeostasis and hepatic steatosis,

inflammation, and fibrosis in this murine dietary model of MASLD/MASH. The complement

system is a complex entity directing an important part of the body's inflammatory and tissue

repair response in MASLD. Further work is needed to elucidate the mechanisms of the role of

C3aR1 in the pathogenesis of MASH and cirrhosis.

Materials and Methods

Animals

C3ar1 flox/flox mice were on the C57BL/6J background as described[43]. Homozygous LysM-Cre mice on the C57BL/6J background (Strain #004781) as well as homozygous Clec4f-Cre mice on the C57BL/6J background (Strain #003296) were purchased from Jackson Laboratories. *C3ar1 flox/flox* homozygous mice on C57BL/6J background were used in the experiments as controls from the same backcross generation[39]. All mice were maintained in plastic cages under a 12h/12h light/dark cycle at constant temperature (22°C) with free access to water and food. Mice were fed regular diet containing 4.5%kcal fat PicoLab Rodent diet 20 (LabDiet) or GAN diet containing 40%kcal HFD (mostly palm oil) with 20% fructose and 2% cholesterol (D09100310, Research Diets) for 28-30 weeks. Fat mass and lean mass were determined via noninvasive 3-in-1 body composition analyzer (EchoMRI). Mice were humanely euthanized with CO_2 inhalation followed by exsanguination by cardiac puncture.

Blood chemistry and serum insulin analysis

Mice were fasted overnight (14-16 hours) for glucose tolerance tests and injected intraperitoneally with syringe-filtered D-glucose solution (2g/kg). For insulin tolerance test, mice were fasted for 6 hours and injected with 0.5 mIU/kg insulin. Blood glucose levels were assayed by commercial glucometer (OneTouch) by tail vein blood samples. Plasma insulin levels were measured from mice fasted for 6 hours. Tail vein blood was collected into lithium heparin-coated tubes, centrifuged at 2000xg at 4ºC, and plasma insulin levels were determined by ELISA using a standard curve (Mercodia). Serum alanine aminotransferase levels were measured in serum from blood collected via cardiac puncture using a commercially available colorimetric assay (TR71121, ThermoFisher Scientific).

Peritoneal macrophage isolation and flow cytometry

Peritoneal macrophages were isolated from as previously described[44]. Briefly, mice were euthanized then immediately injected intraperitoneally with 10 mL phosphate-buffered

saline (PBS, pH 7.4) at room temperature. After a 3-5 minute incubation period, peritoneal fluid

was removed with sterile needle and syringe and placed on ice. After centrifugation at 300xg,

the pellet was resuspended in PBS containing 2% fetal bovine serum and 0.1% sodium azide.

Cells were stained with phycoerythrin-conjugated anti-F4/80 (clone BM8, cat. #123110) and

fluorescein isothiocyanate-conjugated anti-CD11b (clone M1/70, cat. #101206) fluorescent

antibodies (Biolegend). Stained cells were loaded on MA900 fluorescence-activated cell sorter

(Sony), and dual-positive F480+/CD11b+ cells were sorted for subsequent RNA extraction.

Histological studies

A mid-distal portion of the left liver lobe was fixed with 10% buffered formalin and

transferred to 70% ethanol. Samples were embedded in paraffin, sectioned at ~5μm thickness,

and stained with Masson's trichrome. Slides were imaged using Zeiss Axioscan7 at 20x

magnification. Histologic analyses were performed using ImageJ software (version 1.53t). Lipid

droplet area was quantified by subtracting non-droplet area in the green channel from total

section area of 2-3 independent sections. Collagen proportionate area was quantified by

measuring total area in the red channel after reducing intensity threshold to 60-70.

RNA extraction and real-time quantitative PCR analysis

Total RNA from liver tissue lysates was extracted using Trizol reagent (Invitrogen)

followed by RNAeasy Mini kit (Qiagen) as per manufacturer's protocol. RNA was reverse-

transcribed using the High Capacity cDNA RT kit (Thermo). Quantitative PCR was performed

using SYBR Green Master Mix (Quanta) and specific gene primers on QuantStudio6 Flex Real-

Time PCR Systems (Thermo Fisher Scientific) using the delta-delta Ct method. Expression

levels were normalized to Ribosomal protein S18 (*Rps18*). Primer sequences are listed in

Supplementary Table A.

Statistical analyses

All statistical analyses were performed using GraphPad Prism10. Unpaired two-tailed

Student's *t* test with Welch correction for most analyses, with Holm-Šídák correction for multiple

comparisons where applicable, and $p < 0.05$ was considered statistically significant.

Funding

E.A.H. was supported by NIH T32 5T32HL160520-02. A.G. was supported by ADA 9-22-

PDFPM-01. R.P.L was supported by AHA 23DIVSUP1074485. L.S. was supported by AHA

908952 and an Ehrenkranz Young Scientist Award. J.C.L. was supported by NIH R01

DK121140, R01 DK121844, and R01 DK132879. The views expressed in this manuscript are

those of the authors and do not necessarily represent the official views of the American

Diabetes Association, the American Heart Association, the National Institute of Diabetes and

Digestive and Kidney Diseases, or the National Institutes of Health.

Acknowledgments

We would like to thank Dr. Baran Ersoy, Dr. Robert Schwartz, and Dr. Saloni Sinha for their

technical advice and assistance.

Declaration of Competing Interest

None

Data Availability

Data will be made available upon reasonable request.

Figure Legends

Figure 1. C3AR1 is found in macrophages, is modulated by MASLD/MASH in humans,

and is induced by a murine dietary model of MASH.

A) Relative *C3AR1* human tissue expression level by tissue, derived from deep sequencing

of the mRNA combined dataset (HPA and GTEx) in the Human Protein Atlas, shown as

normalized transcripts per million (nTPM). Liver is highlighted in purple and immunologic

tissues are highlighted in red.

B) Single-cell RNA sequencing distribution of *C3AR1* expression in human liver (tSNE, t-

distributed Stochastic Neighbor Embedding).

C) Analysis of *CFD* and *C3AR1* expression from liver biopsy samples in patients with

MASH, MASLD, obesity without MASLD, and age-matched healthy controls (n = 12-16

per group, Welch *t* test with Holm-Šídák correction for multiple comparisons).

D) Weight curve in male and female flox/flox control mice placed on GAN high-fat diet

compared to regular diet (RD) controls (males, n = 7; females, n = 6).

E) Representative liver section staining by Masson's Trichrome in male control mice on RD

or GAN diet for 28 weeks (scale bar = 100 mm).

F) Lipid droplet area quantification in liver sections from male control mice, excluding

vessel lumens (RD, n = 3; GAN, n = 7).

G) Collagen area quantification in liver sections of male control mice (RD, n = 3; GAN, n =

7).

H) Gene expression of key macrophage or fibrosis genes in male control mice on GAN or

RD (n = 6 per group).

Unpaired two-tailed Student's *t* test (Except 1C as above). Annotations: *, $p < 0.05$; **, $p < 0.01$;

***, $p < 0.001$

Figure 2. C3aR1 deletion in all macrophages does not affect weight gain, glucose homeostasis, liver steatosis or fibrosis.

A) Expression of *C3ar1* in isolated peritoneal F4/80+/CD68+ cells from flox/flox control mice (n = 6) or C3aR1-MφKO male mice (n = 3).

B) Expression of *C3ar1* in whole liver from control or C3aR1-MφKO mice (n = 11-12 per male group, n = 13-14 per female group).

C) Body mass curve of control or C3aR1-MφKO mice on GAN high-fat diet starting at 5 weeks of age (n = 11-12 per male group, n = 14 per female group).

D) Body composition analysis by EchoMRI in control or C3aR1-MφKO mice after 30 weeks GAN diet (n = 6-9 per male group, n = 9-13 per female group).

E) Glucose tolerance test in control or C3aR1-MφKO mice with 14h fast after 28 weeks GAN diet (n = 6-9 per male group, n = 9-14 per female group).

F) Liver mass in control or C3ar1-MφKO male mice at time of euthanasia after 30 weeks GAN diet (n = 6-9 per male group, n = 9-14 per female group).

G) Representative liver section staining by Masson's Trichrome in male control or C3ar1-MφKO mice (scale bar = 100 mm).

H) Lipid droplet area in liver sections from male control or C3ar1-MφKO mice, excluding vessel lumens (n = 6-7 per group).

I) Collagen area in liver sections from male control or C3ar1-MφKO mice (n = 6-7 per group).

J,K) Relative mRNA expression of key markers for inflammation, fibrosis, and liver metabolism in liver from male control or C3ar1-MφKO mice after 30 weeks of either GAN (J) diet (n = 11-12 per group) or regular (K) diet (n = 3-5 per group).

Unpaired two-tailed Student's *t* test: Student's *t* test: *, $p < 0.05$.

Figure 3. C3aR1 deletion in Kupffer cells does not affect weight gain, glucose homeostasis, liver steatosis or fibrosis.

A) Body mass curve on GAN diet in flox/flox control or C3aR1-KpKO mice beginning at 5 weeks of age (n = 8-10 per group).

B) Body composition analysis by EchoMRI in control or C3aR1-KpKO mice after 28 weeks GAN diet (n = 8-10).

C) Glucose tolerance test in control or C3aR1-KpKO mice with 14h fast after 26 weeks GAN diet (n = 8-10).

D) Liver mass in control or C3aR1-KpKO male mice at time of euthanasia after 30 weeks GAN diet (n = 8-10).

E) Representative liver section staining by Masson's Trichrome in control or C3aR1-KpKO male mice (scale bar = 100 mm).

F) Lipid droplet area quantified on liver sections of control or C3aR1-KpKO male mice, excluding vessel lumens (n = 8-9).

G) Collagen area quantified on whole liver section of control or C3aR1-KpKO male mice (n= 8-9).

H) Relative gene expression in male control or C3aR1-KpKO mice after 30 weeks GAN diet (n = 5-6).

Unpaired two-tailed Student's *t* test: **, $p < 0.01$.

Supplementary figures.

S1)   Single cell RNA sequencing analysis of *C3ar1* expression in mouse liver tissue (see text).

S2)   Percent lean and fat mass of flox/flox control mice after 20 weeks of GAN or RD (n = 6-7 per group).

S3)   Absolute lean and fat mass of flox/flox control mice after 20 weeks of GAN or RD (n = 6-7 per group).

S4)   Relative gene expression in control female mice after 33 weeks on RD (n = 4-6 per group).

S5)   Insulin tolerance test in control or C3aR1-MφKO male mice with 14h fast after 29 weeks GAN diet (n = 6-9 per group).

S6)   HOMA-IR measurement of insulin resistance in control or C3aR1-MφKO mice with 6h fast after 27 weeks GAN diet (n = 6-9 per male group, n = 9-13 per female group).

S7)   Serum alanine aminotransferase levels in control or C3aR1-MφKO male mice after 30 weeks GAN diet (n = 4 per group).

S8)   Relative gene expression in control or C3aR1-MφKO female mice after 30 weeks of either GAN (n = 13-14 per group) or RD (n = 5-6 per group).

S9)   Relative *C3ar1* expression in control or C3aR1-KpKO female mice after 30 weeks RD (n = 2-3 per group).

Unpaired two-tailed Student's *t* test: **, p < 0.01; ***, p < 0.001.

367 **Supplementary Table S1.**

Mus musculus gene name	Forward qPCR primer	Reverse qPCR primer
Acc2	GCCTCCACTCACATTGGTTT	ATTGAAGAAAGCTGGGCTGA
Acta2	GGCTCTGGGCTCTGTAAGG	CTCTTGCTCTGGGCTTCATC
Adgre1	TGCATCTAGCAATGGACAGC	GCCTTCTGGATCCATTTGAA
C3ar1	TGACAGGTCAGCTCCTTCCT	CATTAGGAGGCTTTCCACCA
Ccr2	ATCCACGGCATACTATCAACATC	CAAGGCTCACCATCATCGTAG
Cd163	TCCACACGTCCAGAACAGTC	CCTTGGAAACAGAGACAGGC
Cfd	CGTACCATGACGGGGTAGTC	ATCCGGTAGGATGACACTCG
Col1a1	GTGCTCCTGGTATTGCTGGT	GGCTCCTCGTTTTCCTTCTT
Col1a2	GCCACCATTGATAGTCTCTCC	CACCCCAGCGAAGAACTCATA
Col3a1	GGGTTTCCCTGGTCCTAAAG	CCTGGTTTCCCATTTTCTCC
Fasn	TTGCTGGCACTACAGAATGC	AACAGCCTCAGAGCGACAAT
Fgf21	CTGCTGGGGGTCTACCAAG	CTGCGCCTACCACTGTTCC
Hnf1a	GACCTGACCGAGTTGCCTAAT	CCGGCTCTTTCAGAATGGGT
Il1b	CTGGTGTGTGACGTTCCCATTA	CCGACAGCACGAGGCTTT
Il6	ACAACCACGGCCTTCCCTACTT	CACGATTTCCCAGAGAACATGTG
Pepck1	TCATCATCACCCAAGAGCAG	CACATAGGGCGAGTCTGTCA
Rps18	CATGCAGAACCCACGACAGTA	CCTCACGCAGCTTGTTGTCTA
Scd1	CGCCCAAGCTGGAGTACGTC	CGCCCAAGCTGGAGTACGTC
Srebp1c	CTGGCAGTTCCATTGACAAG	ACTGAAGCTGGTGACTGCTG
Tgfb1	TGCGCTTGCAGAGATTAAAA	AGCCCTGTATTCCGTCTCCT
Tlr4	TGTCATCAGGGACTTTGCTG	GGACTCTGATCATGGCACTG
Tnfa	ACGGCATGGATCTCAAAGAC	AGATAGCAAATCGGCTGACG
Trem2	CTACCAGTGTCAGAGTCTCCGA	CCTCGAAACTCGATGACTCCTC

References

1   Ge, X., Zheng, L., Wang, M., Du, Y. & Jiang, J. Prevalence trends in non-alcoholic fatty
liver disease at the global, regional and national levels, 1990-2017: a population-based
observational study. *BMJ Open* **10**, e036663 (2020). https://doi.org:10.1136/bmjopen-
2019-036663
2   Younossi, Z. *et al.* Global burden of NAFLD and NASH: trends, predictions, risk factors
and   prevention.   *Nat   Rev   Gastroenterol   Hepatol*   **15**,   11-20   (2018).
https://doi.org:10.1038/nrgastro.2017.109
3   Ferguson, D. & Finck, B. N. Emerging therapeutic approaches for the treatment of
NAFLD   and   type   2   diabetes   mellitus.   *Nat   Rev   Endocrinol*   **17**,   484-495   (2021).
https://doi.org:10.1038/s41574-021-00507-z
4   Friedman, S. L., Neuschwander-Tetri, B. A., Rinella, M. & Sanyal, A. J. Mechanisms of
NAFLD   development   and   therapeutic   strategies.   *Nat   Med*   **24**,   908-922   (2018).
https://doi.org:10.1038/s41591-018-0104-9
5   Stefan, N., Haring, H. U. & Cusi, K. Non-alcoholic fatty liver disease: causes, diagnosis,
cardiometabolic consequences, and treatment strategies. *Lancet Diabetes Endocrinol* **7**,
313-324 (2019). https://doi.org:10.1016/S2213-8587(18)30154-2
6   Kim, H. *et al.* Metabolic Spectrum of Liver Failure in Type 2 Diabetes and Obesity: From
NAFLD to NASH to HCC. *Int J Mol Sci* **22** (2021). https://doi.org:10.3390/ijms22094495
7   Duell, P. B. *et al.* Nonalcoholic Fatty Liver Disease and Cardiovascular Risk: A Scientific
Statement From the American Heart Association. *Arterioscler Thromb Vasc Biol* **42**,
e168-e185 (2022). https://doi.org:10.1161/ATV.0000000000000153
8   Kasper, P. *et al.* NAFLD and cardiovascular diseases: a clinical review. *Clin Res Cardiol*
110, 921-937 (2021). https://doi.org:10.1007/s00392-020-01709-7
9   Barreby, E., Chen, P. & Aouadi, M. Macrophage functional diversity in NAFLD - more
than   inflammation.   *Nat   Rev   Endocrinol*   **18**,   461-472   (2022).
https://doi.org:10.1038/s41574-022-00675-6
10   Cai, J., Zhang, X. J. & Li, H. The Role of Innate Immune Cells in Nonalcoholic
Steatohepatitis. *Hepatology* **70**, 1026-1037 (2019). https://doi.org:10.1002/hep.30506
11   Guilliams, M. & Scott, C. L. Liver macrophages in health and disease. *Immunity* **55**,
1515-1529 (2022). https://doi.org:10.1016/j.immuni.2022.08.002
12   Park, S. J., Garcia Diaz, J., Um, E. & Hahn, Y. S. Major roles of kupffer cells and
macrophages in NAFLD development. *Front Endocrinol (Lausanne)* **14**, 1150118 (2023).
https://doi.org:10.3389/fendo.2023.1150118
13   Sakai, M. *et al.* Liver-Derived Signals Sequentially Reprogram Myeloid Enhancers to
Initiate   and   Maintain   Kupffer   Cell   Identity.   *Immunity*   **51**,   655-670   e658   (2019).
https://doi.org:10.1016/j.immuni.2019.09.002
14   Krenkel, O. *et al.* Myeloid cells in liver and bone marrow acquire a functionally distinct
inflammatory phenotype during obesity-related steatohepatitis. *Gut* **69**, 551-563 (2020).
https://doi.org:10.1136/gutjnl-2019-318382
15   Guilliams, M. *et al.* Spatial proteogenomics reveals distinct and evolutionarily conserved
hepatic   macrophage   niches.   *Cell*   **185**,   379-396   e338   (2022).
https://doi.org:10.1016/j.cell.2021.12.018
16   Govaere, O. *et al.* Transcriptomic profiling across the nonalcoholic fatty liver disease
spectrum reveals gene signatures for steatohepatitis and fibrosis. *Sci Transl Med* **12**
(2020). https://doi.org:10.1126/scitranslmed.aba4448
17   Merle, N. S., Church, S. E., Fremeaux-Bacchi, V. & Roumenina, L. T. Complement
System Part I - Molecular Mechanisms of Activation and Regulation. *Front Immunol* **6**,
262 (2015). https://doi.org:10.3389/fimmu.2015.00262

Kolev, M. & Kemper, C. Keeping It All Going-Complement Meets Metabolism. *Front Immunol* **8**, 1 (2017). https://doi.org:10.3389/fimmu.2017.00001

Zhao, J. *et al.* Association of complement components with the risk and severity of NAFLD: A systematic review and meta-analysis. *Front Immunol* **13**, 1054159 (2022). https://doi.org:10.3389/fimmu.2022.1054159

Segers, F. M. *et al.* Complement alternative pathway activation in human nonalcoholic steatohepatitis. *PLoS One* **9**, e110053 (2014). https://doi.org:10.1371/journal.pone.0110053

Markiewski, M. M. & Lambris, J. D. The role of complement in inflammatory diseases from behind the scenes into the spotlight. *Am J Pathol* **171**, 715-727 (2007). https://doi.org:10.2353/ajpath.2007.070166

Yadav, M. K. *et al.* Molecular basis of anaphylatoxin binding, activation, and signaling bias at complement receptors. *Cell* **186**, 4956-4973 e4921 (2023). https://doi.org:10.1016/j.cell.2023.09.020

Flier, J. S., Cook, K. S., Usher, P. & Spiegelman, B. M. Severely impaired adipsin expression in genetic and acquired obesity. *Science* **237**, 405-408 (1987). https://doi.org:10.1126/science.3299706

Xu, Y. *et al.* Complement activation in factor D-deficient mice. *Proc Natl Acad Sci U S A* **98**, 14577-14582 (2001). https://doi.org:10.1073/pnas.261428398

Lim, J. *et al.* C5aR and C3aR antagonists each inhibit diet-induced obesity, metabolic dysfunction, and adipocyte and macrophage signaling. *FASEB J* **27**, 822-831 (2013). https://doi.org:10.1096/fj.12-220582

Polyzos, S. A., Kountouras, J. & Mantzoros, C. S. Adipokines in nonalcoholic fatty liver disease. *Metabolism* **65**, 1062-1079 (2016). https://doi.org:10.1016/j.metabol.2015.11.006

Han, J. & Zhang, X. Complement Component C3: A Novel Biomarker Participating in the Pathogenesis of Non-alcoholic Fatty Liver Disease. *Front Med (Lausanne)* **8**, 653293 (2021). https://doi.org:10.3389/fmed.2021.653293

Lo, J. C. *et al.* Adipsin is an adipokine that improves beta cell function in diabetes. *Cell* **158**, 41-53 (2014). https://doi.org:10.1016/j.cell.2014.06.005

Gomez-Banoy, N. *et al.* Adipsin preserves beta cells in diabetic mice and associates with protection from type 2 diabetes in humans. *Nat Med* **25**, 1739-1747 (2019). https://doi.org:10.1038/s41591-019-0610-4

Mamane, Y. *et al.* The C3a anaphylatoxin receptor is a key mediator of insulin resistance and functions by modulating adipose tissue macrophage infiltration and activation. *Diabetes* **58**, 2006-2017 (2009). https://doi.org:10.2337/db09-0323

Han, J. *et al.* Bone marrow-derived macrophage contributes to fibrosing steatohepatitis through activating hepatic stellate cells. *J Pathol* **248**, 488-500 (2019). https://doi.org:10.1002/path.5275

Boland, M. L. *et al.* Towards a standard diet-induced and biopsy-confirmed mouse model of non-alcoholic steatohepatitis: Impact of dietary fat source. *World J Gastroenterol* **25**, 4904-4920 (2019). https://doi.org:10.3748/wjg.v25.i33.4904

Hansen, H. H. *et al.* Human translatability of the GAN diet-induced obese mouse model of non-alcoholic steatohepatitis. *BMC Gastroenterol* **20**, 210 (2020). https://doi.org:10.1186/s12876-020-01356-2

Vacca, M. *et al.* An unbiased ranking of murine dietary models based on their proximity to human metabolic dysfunction-associated steatotic liver disease (MASLD). *Nat Metab* (2024). https://doi.org:10.1038/s42255-024-01043-6

Uhlen, M. *et al.* Proteomics. Tissue-based map of the human proteome. *Science* **347**, 1260419 (2015). https://doi.org:10.1126/science.1260419

36   MacParland, S. A. *et al.* Single cell RNA sequencing of human liver reveals distinct
intrahepatic macrophage populations. *Nat Commun* **9**, 4383 (2018).
https://doi.org:10.1038/s41467-018-06318-7
37   Tabula Muris, C. *et al.* Single-cell transcriptomics of 20 mouse organs creates a Tabula
Muris. *Nature* **562**, 367-372 (2018). https://doi.org:10.1038/s41586-018-0590-4
38   Suppli, M. P. *et al.* Hepatic transcriptome signatures in patients with varying degrees of
nonalcoholic fatty liver disease compared with healthy normal-weight individuals. *Am J*
Physiol Gastrointest Liver Physiol **316**, G462-G472 (2019).
https://doi.org:10.1152/ajpgi.00358.2018
39   Ma, L. *et al.* Adipsin and Adipocyte-derived C3aR1 Regulate Thermogenic Fat in a Sex-
dependent Fashion. *JCI Insight* (2024). https://doi.org:10.1172/jci.insight.178925
40   Kong, L. R. *et al.* Loss of C3a and C5a receptors promotes adipocyte browning and
attenuates diet-induced obesity via activating inosine/A2aR pathway. *Cell Rep* **42**,
112078 (2023). https://doi.org:10.1016/j.celrep.2023.112078
41   Li, X. *et al.* A new NASH model in aged mice with rapid progression of steatohepatitis
and fibrosis. *PLoS One* **18**, e0286257 (2023).
https://doi.org:10.1371/journal.pone.0286257
42   Tsuchida, T. *et al.* A simple diet- and chemical-induced murine NASH model with rapid
progression of steatohepatitis, fibrosis and liver cancer. *J Hepatol* **69**, 385-395 (2018).
https://doi.org:10.1016/j.jhep.2018.03.011
43   Cumpelik, A. *et al.* Dynamic regulation of B cell complement signaling is integral to
germinal center responses. *Nat Immunol* **22**, 757-768 (2021).
https://doi.org:10.1038/s41590-021-00926-0
44   Zhang, X., Goncalves, R. & Mosser, D. M. The isolation and characterization of murine
macrophages. *Curr Protoc Immunol* **Chapter 14**, 14 11 11-14 11 14 (2008).
https://doi.org:10.1002/0471142735.im1401s83
494

Figure 1.

Figure 1. C3AR1 is found in macrophages, is modulated by MASLD/MASH in humans, and is induced by a murine dietary model of MASH.

A) Relative *C3AR1* human tissue expression level by tissue, derived from deep sequencing of the mRNA combined dataset (HPA and GTEx) in the Human Protein Atlas, shown as normalized transcripts per million (nTPM). Liver is highlighted in purple and immunologic tissues are highlighted in red.

B) Single-cell RNA sequencing distribution of *C3AR1* expression in human liver (tSNE, t-distributed Stochastic Neighbor Embedding).

C) Analysis of *CFD* and *C3AR1* expression from liver biopsy samples in patients with MASH, MASLD, obesity without MASLD, and age-matched healthy controls (n = 12-16 per group, Welch *t* test with Holm-Šídák correction for multiple comparisons).

D) Weight curve in male and female flox/flox control mice placed on GAN high-fat diet compared to regular diet (RD) controls (males, n = 7; females, n = 6).

E) Representative liver section staining by Masson's Trichrome in male control mice on RD or GAN diet for 28 weeks (scale bar = 100 μm).

F) Lipid droplet area quantification in liver sections from male control mice, excluding vessel lumens (RD, n = 3; GAN, n = 7).

G) Collagen area quantification in liver sections of male control mice (RD, n = 3; GAN, n = 7).

H) Gene expression of key macrophage or fibrosis genes in male control mice on GAN or RD (n = 6 per group).

Unpaired two-tailed Student's *t* test (Except 1C as above). Annotations: *, p < 0.05; **, p < 0.01; ***, p < 0.001.

Figure 2.

Figure 2. C3aR1 deletion in all macrophages does not affect weight gain, glucose homeostasis, liver steatosis or fibrosis.

A) Expression of *C3ar1* in isolated peritoneal F4/80+/CD68+ cells from control mice (n = 6) or C3aR1-MφKO male mice (n = 3).

B) Expression of *C3ar1* in whole liver from control or C3aR1-MφKO mice (n = 11-12 per male group, n = 13-14 per female group).

C) Body mass curve of control or C3aR1-MφKO mice on GAN high-fat diet starting at 5 weeks of age (n = 11-12 per male group, n = 14 per female group).

D) Body composition analysis by EchoMRI in control or C3aR1-MφKO mice after 30 weeks GAN diet (n = 6-9 per male group, n = 9-13 per female group).

E) Glucose tolerance test in control or C3aR1-MφKO mice with 14h fast after 28 weeks GAN diet (n = 6-9 per male group, n = 9-14 per female group).

F) Liver mass in control or C3ar1-MφKO male mice at time of euthanasia after 30 weeks GAN diet (n = 6-9 per male group, n = 9-14 per female group).

G) Representative liver section staining by Masson's Trichrome in male control or C3ar1-MφKO mice (scale bar = 100 µm).

H) Lipid droplet area in liver sections from male control or C3ar1-MφKO mice, excluding vessel lumens (n = 6-7 per group).

I) Collagen area in liver sections from male control or C3ar1-MφKO mice (n = 6-7 per group).

J,K) Relative mRNA expression of key markers for inflammation, fibrosis, and liver metabolism in liver from male control or C3ar1-MφKO mice after 30 weeks of either GAN (J) diet (n = 11-12 per group) or regular (K) diet (n = 3-5 per group).

Unpaired two-tailed Student's *t* test: Student's *t* test: *, p < 0.05.

Figure 3.

Figure 3. C3aR1 deletion in Kupffer cells does not affect weight gain, glucose homeostasis, liver steatosis or fibrosis.

A) Body mass on GAN diet in flox/flox control or C3aR1-KpKO mice beginning at 5 weeks of age (n = 8-10 per group).

B) Body composition analysis by EchoMRI in control or C3aR1-KpKO mice after 28 weeks GAN diet (n = 8-10).

C) Glucose tolerance test in control or C3aR1-KpKO mice with 14h fast after 26 weeks GAN diet (n = 8-10).

D) Liver mass in control or C3aR1-KpKO male mice at time of euthanasia after 28 weeks GAN diet (n = 8-10).

E) Representative liver section staining by Masson's Trichrome in control or C3aR1-KpKO male mice (scale bar = 100 μm).

F) Lipid droplet area quantified on liver sections of control or C3aR1-KpKO male mice, excluding vessel lumens (n = 8-9).

G) Collagen area quantified on whole liver section of control or C3aR1-KpKO male mice (n= 8-9).

H) Relative gene expression in control or C3aR1-KpKO male mice after 30 weeks GAN diet (n = 5-6).

Unpaired two-tailed Student's t test: **, $p < 0.01$.

Supplementary Figures.

Supplementary figures.

S1)   Single cell RNA sequencing analysis of *C3ar1* expression in mouse liver tissue (see text).

S2)   Percent lean and fat mass of flox/flox control mice after 20 weeks of GAN or RD (n = 6-7 per group).

S3)   Absolute lean and fat mass of flox/flox control mice after 20 weeks of GAN or RD (n = 6-7 per group).

S4)   Relative gene expression in control female mice after 30 weeks on RD (n = 4-6 per group).

S5)   Insulin tolerance test in control or C3aR1-MφKO male mice with 14h fast after 29 weeks GAN diet (n = 6-9 per group).

S6)   HOMA-IR measurement of insulin resistance in control or C3aR1-MφKO mice with 6h fast after 27 weeks GAN diet (n = 6-9 per male group, n = 9-13 per female group).

S7)   Serum alanine aminotransferase levels in control or C3aR1-MφKO male mice after 30 weeks GAN diet (n = 4 per group).

S8)   Relative gene expression in control or C3aR1-MφKO female mice after 30 weeks of either GAN (n = 13-14 per group) or RD (n = 5-6 per group).

S9)   Relative *C3ar1* expression in control or C3aR1-KpKO female mice after 30 weeks RD (n = 2-3 per group).

Unpaired two-tailed Student's *t* test: **, p < 0.01; ***, p < 0.001.