**Facilitating the Virtual Exercise Games for Youth with Type 1 Diabetes (ExerT1D) Peer Intervention: Protocol Development and Feasibility**

Garrett I. Ash[1,2], Soohyun Nam[3], Selene S. Mak[4], Matthew Stults-Kolehmainen[5,6], Adrian D. Haughton[1,7], Carolyn Turek[8], Julien S. Baker[9], Kimberly Hieftje[10], Asher Marks[10], Annette Chmielewski[1], Michael Shelver[1], Elizabeth Considine[10], James L. Lukasik[10], Stuart A. Weinzimer[3,10], Laura M. Nally[10]

[1]Department of Internal Medicine and Department of Biomedical Informatics and Data Science. Yale School of Medicine, New Haven, CT, United States
[2]Center for Pain, Research, Informatics, Medical Comorbidities and Education Center (PRIME), VA Connecticut Healthcare System, West Haven, CT, United States
[3]Yale School of Nursing, Orange, CT, United States
[4]Center for the Study of Healthcare Innovation, Implementation and Policy (CSHIIP), VA Greater Los Angeles Healthcare System, Los Angeles, US.
[5]Center for Weight Management, Yale New Haven Hospital, North Haven, CT, United States
[6]Department of Biobehavioral Sciences, Teachers College – Columbia University, New York, NY, United States
[7]Department of Health and Movement Sciences Programs, Southern Connecticut State University, New Haven, CT, United States
[8]Child Study Center, Yale School of Medicine, New Haven, CT, United States
[9]Centre for Health and Exercise Science Research, Population Health and Medical Informatics, Hong Kong Baptist University, Kowloon Tong, Hong Kong SAR, China.
[10]Department of Pediatrics, Yale School of Medicine, New Haven, CT, United States

**Correspondence:**
Garrett I. Ash, PhD
Yale School of Medicine
100 Church St South, Rm F201c
New Haven, CT 06519
Phone: 203-444-3079
Fax: 203-937-3829 (attn: James Lukasik)
Email: garrett.ash@yale.edu

**Running Title:** Virtual Peer Group Physical Activity

**Keywords:** Diabetes Mellitus, Type 1; Adolescent; Motivation; Exercise

**Other Publication:** This work was presented as a podium presentation and a poster at the Society of Behavioral Medicine Annual Meeting, March 13-16, Philadelphia, PA, USA

**Text Word Count**

4,283

**Table & Figure Count**

4 Tables + 4 Figures = 8

## ABSTRACT

**Aims:** Adolescents with type 1 diabetes (T1D) face several unique challenges to engaging in moderate-to-vigorous physical activity (MVPA), including uncertainty with diabetes self-management, risk of hypoglycemia, limited access to safe or supportive environments, and diabetes-specific stigma. Opportunities for T1D peer activities with T1D role model support are limited. To address this need, we tested iterative refinements of pilot Virtual Exercise Games for Youth with T1D (ExerT1D) for feasibility and acceptability.

**Materials and Methods:** Youth with T1D (ages 14-19) were recruited at a pediatric diabetes clinic and through social media. The program included 6 versions: study #1 (versions 1.1-1.4) included an active videogame, while study #2 (versions 2.1-2.2) included a virtual reality active videogame. All versions of the program included education about managing T1D with MVPA and objective habitual MVPA goal-setting guided by clinicians and young adult coaches living with T1D.

**Results:** Seventeen adolescents (median age 15.4 [IQR 14.6 - 16.4] years, 7 non-Hispanic white, 8 male, median HbA1c 8.1% [IQR 7.4%-11.1%]) enrolled. In total there were 260 person-sessions offered over 130 person-weeks. 15 participants attended 83% of sessions. Sessions began with pre-MVPA safety briefing with glucose checks, followed by the MVPA session, and ended with debriefing, goal-setting, and role-playing skits involving educational points related to T1D management. Participants rated the program, comfort, clinicians, coaches, and group cohesion high/very high (4.2±0.5 to 4.8±0.3 out of 5). They also rated motivation for the videogame high (4.1±0.4 out of 5). Building T1D and MVPA self-management skills achieved an excellent rating at a majority of sessions in both versions (76%), as did peer interactions (69%) and enriched communication (56%) after the addition of immersive virtual reality in version #2. Transitions between

virtual reality apps forced group delays of 19±6min per session and some

individuals to exercise separately. Compared to baseline, glycemic metrics

appeared to decrease over time (d= -1.12, 90% CI [-1.78, -0.48]). No participants

experienced diabetic ketoacidosis, severe hypoglycemia, or injuries during the

study period.

**Conclusions:** Overall, ExerT1D facilitated excellent peer support by engaging

diverse youth with T1D in an MVPA program led by role models living with T1D.

**Abstract word count:** 346

## INTRODUCTION

The adolescent years present unique challenges for youth with type 1 diabetes

(T1D). Only ~20% of youth meet the recommended A1c level below 7% shown to

correlate with long-term health.[1,2] Moderate-to-vigorous physical activity (MVPA)

can be leveraged to improve glycemia, weight, and cardiovascular risk factors[3]. In

addition, MVPA has been demonstrated to improve overall body composition,

muscle strength, cardiopulmonary fitness, mental health, diabetes-specific risks of

glycemic outcomes and insulin resistance.[3] MVPA sustained in adulthood, is

associated with less fewer microvascular complications and mortality.[4] Managing

glycemia during MVPA requires more frequent glucose monitoring and

adjustments to diet and insulin dosing, that are further complicated in adolescence

by impaired executive function[5] and puberty-specific hormones.[3,6-8] Perceived

barriers to physical activity reported by adolescents include the fear of glycemic

destabilization[9] and social stigma (i.e., "negative social judgments, stereotypes,

prejudices"[10]) associated with disclosing an individual's diagnosis when engaging

in MVPA.[11] Thus, it is warranted to pursue multifaceted interventions targeted at

addressing these barriers.

Current pediatric T1D MVPA guidelines are available but not actively

understood or prescribed by most clinicians.[12] Additionally, there are limited data

addressing feasibility and acceptability of studies supporting physical activity for

youth with low baseline MVPA.[3] Active videogames and virtual reality (VR) are

potential strategies to promote MVPA among youth.[13,14] Group activities

incorporating active videogame and VR technology with T1D mentors provide an

exciting new avenue to deliver components of a successful T1D self-management

intervention:[15] 1) behavior modification (goal setting, diabetes self-management);

2) psychosocial support (engagement with T1D peers and role models); and 3)

medical guidance (real-time diabetes self-management advice). Herein we present

six cohorts of 3-5 participants evaluating the feasibility, acceptability, and

preliminary safety of iterative refinements of a home-based virtual intervention to

promote MVPA among adolescents with T1D.

**METHODS**

**Enrollment**

Adolescents (14-19yr old, T1D ≥6mo) not achieving recommended MVPA

targets (i.e., <4days/wk with 60+min self-reported[16]) were recruited from the Yale

Children's Diabetes Center, clinicaltrials.gov (NCT05163912, NCT05662826),

and diabetes social media groups between December 23, 2021 and June 15,

2022 (study #1; versions #1.1-1.4) and January 12th – March 2nd, 2023 (study #2;

versions #2.1-2.2). Screening was conducted by telephone using the Physical

Activity Readiness Questionnaire verified against medical record[17]. For any

health conditions presenting special considerations for MVPA, additional

documentation was requested from the primary care physician or sub-specialist

for MVPA clearance (n=1, arthritis). Informed consent or parental

permission/assent (<18yr) was completed and signed using REDCap eConsent

over a HIPAA-compliant tele-video call (Zoom Video Communications, San Jose,

CA). Data on demographics, insulin therapy, anthropometrics, MVPA,[16] and

sedentary time[18] were collected by parental REDCap survey and verified against

medical record. Screening and intake data for study #2 were also used as long-

term follow-up for study #1. The research team mailed supplies to the participant

and held a second tele-video call to confirm proper setup.

**Intervention Development**

The intervention, based upon Social Cognitive Theory, integrated peer

and role model interactions into home-based MVPA[19]: networked active

videogame sessions and virtual discussions about experiences managing

diabetes and MVPA. In study #2, we added VR as a delivery tool to enhance the

drivers of adolescent development and motivation according to self-

determination theory, which are: competence, autonomy, and relatedness[20-22].

VR enhances competence by providing a safe space to learn competence in

specific activities like MVPA and self-monitoring with a controlled degree of

distractions and barriers. VR enhances autonomy by allowing users to embody a

self-designed avatar. Finally, VR enhances relatedness by allowing users to

experience new unique environments for the first time as a group.

Study #1 participants fitted their home television monitor with a Nintendo

Switch Entertainment System (Nintendo of America, Inc., Redmond, WA)

including Ring Fit Adventure which alternates low- (stationary walking or jogging)

with 60+ higher-intensity exercises that are dynamic (e.g., knee-lifts) or

resistance-based (e.g., abdominal press) tracked by a leg-strap accelerometer

and handheld ring with gyroscopic/accelerometer sensors. Success with

exercises unlocks more advanced adaptations (e.g., Russian twist). They were

also provided a yoga mat, poster with Borg rating of perceived exertion scale,

Precision Xtra Blood Ketone meter and strips (Abbott Diabetes Care, Alameda,

CA), and Fitbit Inspire 2 (Google, Menlo Park, CA). Sessions were livestreamed

over Zoom (Figure 1).

**Figure 1.** Study #1 virtual exercise session livestreamed over Zoom. Sessions were guided by a coach, and a clinician with expertise in T1D exercise management was present to advise diabetes-specific adjustments for exercise management.

Study #2 participants engaged in physical activity sessions using a virtual reality headset (Meta Quest 2 headset, Menlo Park, CA). Participants completed headset tutorials to adjust the lenses for proper focus, draw a guardian boundary to demarcate an area without obstacles or tripping hazards, and learn how to use the controllers. Participants also designed a unique Avatar to represent themselves during activity using Meta's features (clothing, accessories, general appearance [e.g., skin tone, eye color, body shape], hairstyle, and voice modifications) and use two applications within the headsets. The first application, FitXR (Meta, Menlo Park, CA) facilitated group exercise classes using the avatars side-by-side to other participants to perform exercises guided by verbal and visual instructions from an automated avatar coach. Participants received individual and group scores within each activity. Following exercise activities, participants used VR headsets to meet in a virtual environment (Foretell Reality,

Glimpse Group, New York, NY). Environments included a campfire with roasted

marshmallow props, hot chocolate, and a guitar or a tropical beach with sea

animals. These environments contain presence-enhancing features (spatial

audio, hand and face tracking).

Four coaches living with T1D aged 20 to 26 years old (2 non-Hispanic

White male, 1 non-Hispanic Black female, 1 non-Hispanic White female) with

experience managing MVPA with T1D facilitated the sessions. The first coach was

trained by reading a prototype manual and lesson guides prepared by the principal

investigator with input from the multidisciplinary investigative team (pediatric

endocrinologists, a pediatric psychologist, an exercise physiologist, an exercise

telemedicine instructor, and behavioral interventionists).  Then, the coach

conducted two mock sessions with investigative team members as participants.

The subsequent coaches were trained by reading the manual, observing

recordings of prior sessions, and running mock sessions with the principal

investigator and research assistants (RAs) as participants. All mock and live

sessions were followed by feedback and refinements (c.f. Assessments below).

An RA facilitated intervention sessions and followed a source document to

record the completion of each activity along with related diabetes-specific data

(Appendix 1). The RA also helped participants troubleshoot technical difficulties

with using the video games and equipment. A clinician (pediatric endocrinologist,

certified diabetes care and education specialist) provided guidance about

diabetes-specific exercise management according to published guidelines[3,23] and

an exercise physiologist monitored adherence to the intervention protocol

(Appendix 1).

Participants were recruited in sequential cohorts for the 6-week (version 1)

and 4 week (version 2) intervention with meetings twice weekly (Table S1) over

Zoom. At the first session in each cohort, the clinician gave an overview of safe

exercise practices related to T1D management based on national and international

guidelines.[6,7,24] Prior to activity, coaches confirmed that the participants had water,

fast-acting carbohydrates to treat hypoglycemia, proper exercise shoes and

clothing, and a 2.4m-by-2.4m unobstructed space for exercise. Participants

communicated to the coach if they needed to leave the session for any reason,

and the RA maintained a list of each participant's physical location in case

emergency services were needed.

**Diabetes Management**

At each session, before and after exercise, teens reported their blood

glucose or sensor glucose value, trend arrow, and insulin administration history

(time and amount of last bolus and/or insulin-on-board from pump). Exercise was

paused, and clinical guidance was provided if a glucose level was less than

70mg/dL and/or symptoms of hypoglycemia were reported. Ketone levels were

checked if a glucose level exceeded 250mg/dL or there were clinical concerns for

elevated ketones. Clinical guidance was provided when blood ketone levels were

elevated ≥0.6mmol/L or urine ketones were moderate or large. On a weekly basis,

the clinician reviewed continuous glucose monitor (CGM) and ketone data

collected by questionnaire or upload (Keto-Mojo GK+, Napa, CA), identified any

safety concerns (>4% time with hypoglycemia), and communicated these

concerns with participants and parents.

For the purposes of coaching exercise, we tracked the proportion of each

session spent in active movement, repetitions of each Ring Fit exercise or total

FitXR target hits, Borg 6-20 rating of perceived exertion of each Ring Fit exercise,

Borg 1-10 rating of the total session, and average Fitbit heart rate.

Following the 20-40min exercise activity, the last 20-30min of each session

was spent on discussion over Zoom (version 1) or in virtual environments. Week 1

sessions were dedicated to creating individual fitness-related goals following the

SMART (Specific, Measurable, Attainable, Relevant, Timely) principle. In addition,

participants in each cohort formulated a group goal of cumulative achievements

(e.g., group steps) to promote a cooperative rather than competitive approach.[25]

Goal progress was checked and discussed each week, including the provision of

MVPA and T1D management information that helped support the specific goals.

During middle week sessions, coaches engaged teens to create role-playing skits

on Zoom that involved educational points related to T1D management. During the

final sessions, the coaches led a discussion and painting exercise allowing

participants to express the relevance of the intervention to building and sustaining

an overall active lifestyle (Figure 2).

**Figure 2.** The version 2 virtual reality discussion environment included 3-dimensional painting where adolescents could express where their MVPA goals fit with their overall personal perspectives on MVPA, T1D, peers, family, other life topics, and their intersections. In the above example an adolescent represents themselves as a star that shines and dims with mood and life events, surrounded by diabetes (syringe), food managed with diabetes (hamburger), school, homework and projects (book), sports managed with diabetes (football), and being outside (runner). In version #1, similar discussions occurred verbally without visual expression.

**Assessments**

Session recordings were double-coded by exercise physiologists or psychology

RA to identify missed opportunities for practicing each core competency of a role

model and peer intervention[26-31] which were then discussed with instructors weekly

to iteratively improve the intervention. The competencies were:

**1) Facilitating Peer Interactions.** We encouraged peer support with group

discussions led by coaches, inviting each participant to share general life updates,

discuss diabetes management challenges and successes, and physical activity

goals.

**2) Practicing enriching communication.** Instructors and clinicians modeled

empathy, resilience, and realistic optimism, validating challenges, empowering

youth to manage their condition with confidence and self-compassion.  The team

encouraged conversations that were supportive, meaningful, and growth-oriented.

**3) Building T1D and MVPA Management Skills.** Clinicians led T1D self-

management skill-building exercises, reviewing participants' glucose changes

during exercise shared in the physical activity session to augment glucose pattern

recognition and problem solving, emphasizing that glucose levels during physical

activity can vary based on recent insulin doses ("insulin on board"), carbohydrate

intake and time of day.  We also reviewed strategies to mitigate hypoglycemia and

hyperglycemia during activity (e.g., reducing insulin prior to activity or setting an

"activity" or "exercise" mode or "temp target" prior to and after physical activity).

At the end of the study, participants completed a follow-up retrospective

survey of acceptability including the motivation for exergame play inventory

($\alpha=0.79$),[32] perceived cohesion scale ($\alpha=0.78$-$0.83$),[33] and a satisfaction survey

from our past work[27] to evaluate program components/strategies (5 items, $\alpha=0.72$),

personal comfort level (6 items, $\alpha=0.81$), and interactions with instructors (12

items, $\alpha=0.95$). Participants also rated the top 3 factors by importance.

**Feasibility Outcomes and Analytic Methods**

Study outcomes and pre-specified thresholds included recruitment uptake

(≥35%),[34] attendance (≥75% attending ≥75% of sessions),[35] biosensor wear-time

(≥70%),[36] survey completion (≥85%), fidelity ratings (≥70% excellent, ≥90% at

least fair), and psychosocial perceptions of acceptability (≥3.0 out of 5). Variable

distributions were described using visual inspection, Shapiro-Wilks test of

normality, Levene's test of equal variances, and appropriate effect size formulas

to quantify temporal trends (SPSSv28, Chicago, IL). Inferential statistics such as

ANOVA or comprehensive pairwise testing were not performed due to the small

sample size. We refined the curriculum based on data after each cohort, ending

the trial when no further refinements were found. This sample size (n=15) had 83%

statistical power to detect any feasibility problems having ≥10% incidence

according to Viecthbauer's formula $n = (\ln[1-\gamma])/(\ln[1-\pi])$ where $\gamma$ is power and $\pi$ is

minimum incidence of a feasibility problem to be detected.[37]

The intervention was unique in that it included both short-term behavioral

reinforcement (e.g., peer encouragement of MVPA) and psychoeducation

targeting long-term habit change (e.g., role-playing skits involving educational

points). As such, we observed metrics not only during the 4-6-week interventions

(e.g., CGM data) but also up to 1 year later (e.g., medical record HbA1c) in accord

with standard practices for brief psychoeducational interventions (e.g.,[38,39]).

250                                **RESULTS**

**Participants**

Seventeen participants referred by local clinicians, Facebook posts, and T1D

community organizations were enrolled and completed follow-up (Figure S1),

representing diverse geographic (8 states) and demographic characteristics: 8

female sex, 6 female gender, 2 non-binary gender, 4 public insurance, 2 with

household incomes below the federal poverty line, and 7 household income

<$60,000 (Table 1). Most participants had A1c levels above American Diabetes

Association targets: 12 had an A1c of greater than 7.0%, and 5 had an A1c level

of greater than 10.0%.

**Table 1.** Baseline characteristics of the study participants.

| | Study #1 | Study #2* |
|---|---|---|
| N | 15 | 10 |
| Age (yr)[1] | 15.4 (14.5 - 16.4) | 15.6 (15.0 – 16.0) |
| Gender | 6 female (40%)<br>2 non-binary (13%)<br>7 male (47%) | 4 female (40%)<br>1 non-binary (10%)<br>5 male (50%) |
| Sex | 8 female (53%)<br>7 male (47%) | 5 female (50%)<br>5 male (50%) |
| Race | 3 Black (20%)<br>10 White (67%)<br>1 More than 1 race (7%)<br>1 Other (7%) | 3 Black (30%)<br>5 White (50%)<br>2 More than 1 race (20%)<br>0 Other (0%) |
| Ethnicity | 4 Hispanic (27%)<br>11 Non-Hispanic (73%) | 3 Hispanic (30%)<br>7 Non-Hispanic (70%) |
| A1c (%)† | 7.8 (7.4 – 11.2) | 8.4 (7.4 – 10.3) |
| Duration of type 1 diabetes (yr) | 8.2±3.8 | 7.3±3.3 |
| Household income annual | 2 <$20,000 (13%)<br>0 $20,000-$39,999 (0%)<br>5 $40,000-$59,999 (33%)<br>0 $60,000-$79,999 (0%)<br>3 $80,000-$99,999 (20%)<br>4 >$100,000 (27%)<br>1 declined to respond (7%) | 1 <$20,000 (10%)<br>0 $20,000-$39,999 (0%)<br>3 $40,000-$59,999 (30%)<br>0 $60,000-$79,999 (0%)<br>2 $80,000-$99,999 (20%)<br>4 >$100,000 (40%)<br>0 declined to respond (0%) |
| Insurance (N public, %) | 4 (27%) | 1 (13%) |
| Therapy | 1 multiple daily injections (7%)<br>11 sensor-augmented pump (73%)<br>3 automated insulin delivery (20%) | 0 multiple daily injections (0%)<br>4 sensor-augmented pump (40%)<br>6 automated insulin delivery (60%) |
| Prior CGM use (N, %) | 15 (100%) | 10 (100%) |
| Total daily insulin dose (U/kg) | 0.9±0.3 | 0.9±0.2 |
| Body mass index (%)† | 67 (63 – 87) | 71 (51 – 84) |
| Physical activity (days/wk with 60+ min MVPA) | 1.5±1.1 | 1.4±1.1 |
| Screen time (hours/day) | 8.8±4.6 | 10.3±5.4 |
| Data are presented as mean±SD for normally distributed data and median (interquartile range) otherwise.<br>*Included 8 participants that also participated in study #1.<br>†Skewed distribution (age and HbA1c positive-skewed, body mass index negative-skewed). | | |

**<u>Intervention Feasibility</u>**

<u>Attendance and Adherence</u>

Fifteen participants attended ≥80% of sessions, while the remaining two attended

50%-60% of the sessions. Absences were due to family activities (7), school

activities (7), negative mood (6), oversleeping (6), acute illness (4), or no answer

was given (2). Most of these absences (27/32, 84%) occurred on Saturdays, the

few absences on Wednesdays being due to school activities (1), negative mood

(3), or no answer was given (1).

All participants were using a CGM in their clinical care before the study (14

Dexcom G6, 3 Abbott Libre 2). Overall, weekly audits found CGM adherence was

73.6% in study #1 and 75.0% in study #2 of the possible 288 daily readings were

captured across all study participants. Data completeness at the participant level

was 79% (SD 25%) with 12 participants at ≥70%, 4 participants at 24%-54%, and

1 participant only sharing visual CGM data without a csv file (excluded from Tables

5-6).

Fitbit wear time on average met literature standards (70.5%, or 16.9 hours

per day suggesting an average of 10-15 hours during waketime).[36] Version 1.1

participants stopped wearing the Fitbit 2-4 weeks into the program, so we inserted

a biweekly check of wear-time and troubleshooting after which this pattern did not

repeat.

<u>Technical Barriers</u>

In study 1, one participant was unable to participate in one session due to

"dead" RingFit controller batteries (rated as moderate-high disruption). This

participant was also unable to communicate verbally in two other sessions due to

phone audio interruption (though could communicate by text chat). Two other

participants had brief Zoom connectivity interruptions on 3 occasions respectively

(reported as mild disruption).

In study 2, there were 15 malfunctions of the Oculus system (8 microphone,

3 speakers, 2 guardian boundary misplacement, 2 passcode), 9 with FitXR (5

times full class did not launch, 4 times individual participant ejected from class),

and 5 with Foretell Reality Rooms (4 times individual participant locked out or

ejected, 1 time no audio in room). FitXR was used in 12 of the 16 sessions and

Foretell Reality was used in all 16 sessions.  All issues resolved after restarting the

app and/or system. FitXR classes were not started until all individuals could join

the class without malfunctions, causing a delay of 19 (SD=6) minutes per session.

When malfunctions occurred for individual participants mid-session, activities were

paused unless rejoining was impossible.  In these cases, a backup coach joined

the participant in a separate FitXR class (4 instances) or group discussion (1

instance). Most malfunctions (80%) occurred during weeks 3-4 of study 2 before

Meta updated its firmware (v50), when other users nationally reported similar

issues.

<u>Fidelity</u>

In version #1, intervention evaluators rated coach competencies highest for

building T1D self-management and MVPA management skills followed by

enriching communication and facilitating peer interactions at the group sessions

(Table 2). Most competencies trended upward for each successive cohort except

for cohort #4. Cohort #3 was the largest size and most balanced with respect to

race & ethnicity, income, HbA1c levels, and recruitment source (Table 3). Cohort

#4 had lowest size plus lowest attendance among those enrolled. One coach

covered the first two cohorts and another coach covered the last two, both

achieved similar average scores on most competencies. In version #2, facilitating

peer interaction ratings increased and reached similar rating as building skills.

Practicing enriching communication also increased, to a lesser extent and building

skills remained high.

**Table 2.** Fidelity, Satisfaction, Perceived Cohesion, and Exergaming Motivation Overall and by Cohort

| Version | Instructor Competencies (% of sessions rated excellent) | | | Psychosocial Perceptions mean±SD (% rated ≥4 out of 5) | | | | Exergaming Motivation mean±SD (% rated ≥4 out of 5) |
|---|---|---|---|---|---|---|---|---|
| | Facilitating Peer Interactions at the Group Sessions | Enriching Communication at the Group Sessions | Building T1D and MVPA Management Skills | Interactions with Clinicians and Coaches | Group Cohesion (Perceived) | Program Components and Strategies | Comfort | Exergaming Motivation |
| #1.1 (n=3) | 0 | 25 | 75 | 4.6±0.7 (67%) | 4.2±0.9 (67%) | 4.1±0.2 (100%) | 3.9±0.9 (67%) | 3.8±0.4 (33%) |
| #1.2 (n=4) | 27 | 46 | 100 | 4.9±0.0 (100%) | 4.6±0.4 (100%) | 4.5±0.3 (100%) | 4.5±0.1 (100%) | 4.1±0.2 (75%) |
| #1.3 (n=5) | 0 | 50 | 100 | 4.9±0.2 (100%) | 4.2±0.4 (80%) | 4.3±0.6 (60%) | 4.2±0.4 (80%) | 3.9±0.2 (60%) |
| #1.4 (n=3) | 0 | 0 | 38 | 4.8±0.3 (67%) | 4.7±0.2 (100%) | 4.3±0.7 (33%) | 4.3±0.6 (33%) | 4.6±0.2 (100%) |
| **Study #1 Total (n=15)** | **8** | **29** | **78** | **4.8±0.3 (93%)** | **4.4±0.5 (87%)** | **4.3±0.5 (73%)** | **4.2±0.5 (80%)** | **4.1±0.4 (60%)** |
| #2.1 (n=5) | 75 | 50 | 63 | 4.7±0.4 (100%) | 4.3±0.5 (80%) | 4.6±0.3 (!00%) | 4.7±0.4 (100%) | 3.8±0.5 (40%) |
| #2.2 (n=5) | 63 | 63 | 88 | 4.5±0.5 (100%) | 4.4±0.8 (80%) | 3.8±0.2 (40%) | 3.7±0.6 (40%) | 3.9±0.4 (60%) |
| **Study #2 Total (n=10)** | **69** | **56** | **75** | **4.6±0.4 (100%)** | **4.3±0.6 (80%)** | **4.2±0.5 (70%)** | **4.1±0.7 (60%)** | **3.8±0.4 (50%)** |

Study #1 (versions 1.1-1.4) is 2-dimensional Ring Fit game and discussions by Zoom shown in methods Figure 1. Study #2 (versions 2.1-2.2) is virtual reality immersive environment shown in methods Figure 2.T1D, type 1 diabetes. MVPA, moderate-to-vigorous physical activity.

Version 1.4 took one week longer than expected to recruit the minimum 3 participants. In fairness to the 2 participants who signed up on the preset schedule, we offered them 2 sessions during the week of delay which were credited as extra sessions.

**Figure 3.** Competency ratings. Dots represent cohorts and are color-coded by study. Purple is study #1 (versions 1.1-1.4; 2-dimensional Ring Fit game and

discussions by Zoom shown in methods Figure 1). Blue is study #2 (versions 2.1-2.2; virtual reality immersive environment shown in methods Figure 2).

**Table 3.** Cohort Characteristics

| Cohort | Race/ Ethnicity | Gender | Age | Household Income (annual) | A1c | Recruitment | Attendance (Mean %)[1] |
|---|---|---|---|---|---|---|---|
| #1.1 (n=3) | 2 HispWhite<br>1 NHispWhite | 1 NB<br>2 M | 1, 14-15yr<br>2, 16-17yr | 1 under $20k<br>2 under $60k | 3 over 7.0%<br>1 over 10.0% | 3 from clinic<br>0 from social media | 75 |
| #1.2 (n=4) | 4 NHispWhite | 4 M | 3, 14-15yr<br>1, 16-17yr | 0 under $20k<br>0 under $60k | 2 over 7.0% | 0 from clinic<br>4 from social media | 92 |
| #1.3 (n=5) | 3 NHispBlack<br>1 More than 1 race<br>1 NHispWhite | 1 NB<br>3 F<br>1 M | 5, 14-15yr | 0 under $20k<br>3 under $60k<br>1 not reported | 5 over 7.0%<br>2 over 10.0% | 2 from clinic<br>3 from social media | 95 |
| #1.4 (n=3) | 2 HispWhite<br>1 NHispWhite | 3 F | 1 14-15yr<br>1, 16-17yr<br>1, 18-19yr | 1 under $20k<br>2 under $60k | 2 over 10.0% | 2 from clinic<br>1 from social media | 72 |
| #2.1 | 3 NHispBlack<br>2 More than 1 race | 3 F<br>2 M | 4, 14-15yr<br>1, 16-17yr | 0 under $20k<br>3 under $60k | 5 over 7.0%<br>2 over 10.0% | 2 from clinic<br>3 from social media | 90 |
| #2.2 | 2 HispWhite<br>3 NHispWhite | 1 F<br>3 M<br>1 NB | 4, 14-15yr<br>1, 16-17yr | 1 under $20k<br>1 under $60k | 4 over 7.0%<br>1 over 10.0% | 2 from clinic<br>3 from social media | 83 |

HispWhite, Hispanic White. NHispWhite, Non-Hispanic White. NHispBlack, Non-Hispanic Black. NB, non-binary. M, male. F, Female.

[1]Cohort #4 took one week longer than expected to recruit the minimum 3 participants. In fairness to the 2 participants who signed up on the preset schedule, we offered them 2 sessions during the week of delay which were credited as extra sessions.

**Intervention Acceptability**

Participant satisfaction ratings in both versions were highest for interactions with

clinicians and coaches, followed by perceived group cohesion, program

components and strategies, comfort, and lastly, active videogame motivation

(Table 2). The clinicians and coaches component was also most frequently

selected as a top-1 or top-3 (72%, 64%).

**Intervention Safety and Monitoring**

No participants experienced diabetic ketoacidosis, severe hypoglycemia, or other

study-related adverse events during the study period. Two participants developed

elevated ketone levels prior to exercise (1.1, 1.4 mmol/L) due to an overnight pump

malfunction restricting their prior insulin, that resolved after insulin administration.

One participant deferred exercise due to feeling unwell as a result of having a high

sensor glucose level (400mg/dL) without elevated ketones. Three participants

reported hypoglycemia symptoms during a session that resolved with

carbohydrate treatment and three had sensor glucose levels <70mg/dL less than

1 hour after a session that participants resolved independently without staff

guidance.

Weekly clinician reviews of CGM data resulted in 5 participants being

referred to their healthcare providers for insulin dose adjustments to address

elevated percentages of low sensor glucose values. Topics related to avoiding

hypoglycemia after exercise were discussed in the group lesson, with participants

directly.

Heart rate data were captured for 129/156 person-sessions (83%), with 15

out of 17 participants having at least one session. Among them, one averaged in

the vigorous range of 70%-90% age-predicted maximum, 13 averaged in the

moderate range of 55%-70%, and one averaged below the moderate range. The

average participant heart rate across sessions was 61.4% ± 5.2% of maximum

playing Ring Fit and 57.7% ± 6.0% playing FitXR.

**Intervention Acceptability**

Ratings of specific components were the highest for activities as follows: group 3-

D painting, seeing personal and peers' scores for hitting boxing targets during

MVPA, freely exploring the virtual reality environments, and exercising with other

teens with T1D. The lowest-rated activities were those either administered in the

non-avatar Zoom audio-chat (icebreaker questions, blood sugar checks) or

automated videogame characters not part of the peer group (FitXR automated

coach) (Table 4).

Participants set individual and group goals related to achieving Fitbit step

counts, playing a sport, recreational exercise, or playing Ring Fit outside of group

sessions, or managing diabetes (Table S2) with 28% success on individual goals

and 50% success on group goals.

Average steps in study #1 for those with Fitbit step goals were 8274±3734

per day and for the total sample were 7178±4013 per day. Average steps in study

#2 for the participant with a Fitbit step goal was 6,034 per day and for the total

sample was 6467±2998 per day. Each skit successfully integrated 3-5 educational

points (Table S3).

**Table 4.** Ranking of satisfaction of specific study #2 activities (n=10). Not done for study #1.

| Component | Mean±SD (% rated ≥4 out of 5) |
|---|---|
| Group 3D painting | 4.9±0.3 (100%) |
| Seeing individual score | 4.6±0.5 (100%) |
| Seeing everyone's scores | 4.4±0.7 (90%) |
| Freely exploring virtual environments | 4.4±0.7 (90%) |
| Exercising with other teens with type 1 diabetes | 4.3±0.7 (90%) |
| Exercise Movements | 4.2±0.6 (90%) |
| Exercising with coach that has type 1 diabetes | 4.1±0.6 (90%) |
| Diabetes skit | 4.1±1.0 (80%) |
| Talking about individual physical goals | 4.1±1.0 (80%) |
| Discussing exercise management strategies with clinicians and coaches | 4.1±0.7 (80%) |
| Talking about group physical activity goals | 4.1±0.7 (80%) |
| Icebreaker questions | 4.0±0.9 (80%) |
| Checking blood sugars and reporting them to clinicians and coaches | 4.0±0.9 (67%)† |
| Automated Exercise Coach | 3.7±0.7 (60%) |

†One participant declined to answer.

**<u>Ancillary Analysis of Glycemic Outcomes</u>**

Between the two timepoints that HbA1c was available for the same participants -

Study #1 baseline and follow-up (n=15) - it was on average lowered (8.9% vs 8.3%)

but equally positive-skewed (Levene's $F_{1,2}=0.458$) (Figure 4). Such effect is best

standardized as the Wilcoxon Signed Rank test index W= -1.905, equal to biserial

r= -0.49 or Cohen's d= -1.12 (90% CI [-1.78, -0.48]) (i.e., probably a medium or

large effect).[40] Total daily insulin showed no indication of increasing (0.88±0.27 to

0.83±0.24 U/kg/day) and participants upgrading to a closed-loop system during

follow-up (n=7) had the same average HbA1c change as the full cohort. Body mass

index percentile remained negative-skewed and showed no indication of change

(median 67 vs 66, mean 75 vs 75, IQR 63-87 vs 53-86).

During the interventions, CGM clinical targets were met for time in

hypoglycemia level 1 (1.8%-2.2% vs target ≤4%) and level 2 (0.5%-0.7% vs target

≤1%), but not hyperglycemia (42%-51% vs target ≤25%) or in target range (46%-

55% vs target ≥70%) (Figure 4). Full CGM summary metrics are given in tables S4

and S5.

**Figure 4.** Glycemic metrics over time. **Top.** Metrics indicating average glucose. Version #1 baseline is laboratory HbA1c taken an average of 7 weeks (range 1-14) before intervention start, extracted from medical record. Version #1 follow-up is laboratory HbA1c taken an average of 45 weeks (range 19-77) after intervention start, extracted from medical record. Missing values (1 participant at weeks 1-6 due to not sharing downloadable CGM data, 2 participants at follow-up due to not sharing long-term access to medical record) were imputed by intention-to-treat, last observation carried forward. Other boxes are glucose management indicator (estimation of HbA1c based on CGM). Dash through box indicates median, plus sign indicates mean, box indicates interquartile range, whiskers indicate minimum and maximum. **Bottom.** Metrics indicating time in glucose ranges. Data completeness at the participant level was 79% (SD 25%) with 12 participants at ≥70%, 4 participants at 24%-54%, and 1 participant only sharing visual CGM data without a csv file (excluded). The 1 participant who did not share downloadable CGM data was excluded. Bars are omitted from timepoints where ≥5 participants had no shared data.

**DISCUSSION**

419    The primary finding of the study was that the virtual, compared to in-person

420 protocol,[26] greatly increased feasibility while maintaining excellent safety

421 standards and opportunities to interact with peers, clinicians, and young adult role

422 model coaches. Compared to our in-person group, we doubled recruitment uptake

423 proportion from 16% to 35%, which is comparable to 37% seen in personalized

424 MVPA prescription for T1D that did not require fitting a group schedule.[34] We also

425 expanded from clinic to social media venues. In addition, we improved the

426 attendance from 56% to 85%-90%, while also doubling the frequency of sessions

427 from once to twice per week. A glucose monitoring protocol was followed with no

428 adverse events and occurrence of MVPA-induced hypoglycemia no more

429 frequently than typical daily living. The program was also highly cost-efficient.

430 Social media advertisements utilized community postings rather than paid

431 advertisements, and virtual technology was all within the scope of devices normally

432 owned by diverse families of teens with T1D. Videogame technology is popular

433 among teens and substantially more so among those from lower-income families.[41]

434    The high acceptability of the in-person protocol[27] was also maintained

435 based on mean evaluation scores, though unbalanced across subscales. Building

436 T1D and MVPA self-management skills achieved an excellent rating at a majority

437 of sessions in both versions (76%), as did peer interactions (69%) and practicing

438 enriching communication (56%) after the addition of immersive virtual reality in

439 version #2. Perceived group cohesion was also rated highly by the teens, reflecting

440 some combination of perceived cohesion to the instructors and other teens.

441 Program structure and comfort were rated less highly than clinician and coach

442 interactions and perceived group cohesion, though nonetheless received ratings

443 of 4+ out of 5 and were likely essential to the program's feasibility.

Prior studies supporting MVPA for adolescents with T1D provided self-

management education and decision guidance[34,42-44] and only one included

personal support, which came from key family members such as a parent.[34] In

real-world contexts, 91% of adolescents with T1D report that their parents

discourage MVPA[9] and that teachers and coaches have limited T1D

knowledge.[45] Therefore, interventions aimed at improving health outcomes need

to be introduced during the transition period of adolescence when youth are

developing the skills they will need as adults to manage T1D.[26,27]

Support from peers and peer mentors is rated as important by

adolescents with T1D[46,47] and associated with increased diabetes self-care.[48]

However, the incorporation of peers and role models is in nascent stages[28,49] and

mostly focused on T1D-specific behaviors rather than those important for general

health such as MVPA. Such restriction of peer support to T1D-specific behaviors

has been negatively associated with diabetes self-care,[48] similar to the "nagging"

perceived from parents.[50] It may also be less engaging, as these prior peer

interventions have been almost entirely restricted to asynchronous

communications without physical activities.[29,30] When synchronous chats have

been added, just 39%-48% of participants attended one or more sessions.[30,49]

Overall, the present study shows the promise of peer group MVPA as a novel

strategy both for supporting MVPA and generally engaging adolescents with

T1D.

A major strength of the study was the enrollment of a diverse sample of

adolescents in greatest need of support for MVPA and T1D self-management

(lower income, people of color, elevated HbA1c).[51] This study had no A1c upper

limit, thus incorporating a group of youth with T1D at high risk of dysglycemia that

has been largely excluded from research studies. Intervention attendance was

excellent and overcame logistical barriers related to alignment of group

schedules, availability of MVPA supplies, timing MVPA with insulin and diet, and

provision of real-time guidance. Given the focus on feasible uptake of technology

and inclusion of youth with all HbA1c levels, we enrolled participants even if they

were wearing their CGM inconsistently at enrollment, which meant glycemic

outcomes could only be assessed by combining HbA1c and GMI values in an

ancillary analysis. HbA1c and GMI can be discordant due to the timeframe they

reflect and non-glycemic factors that can impact HbA1c.[52] Nonetheless, glycemic

reductions were seen during and after the intervention. These findings were

encouraging, especially in light of the elevated HbA1c levels in the cohort at

baseline, as these youth are largely excluded from research.

We acknowledge some limitations of the intervention. Coaches were

trained in group facilitation strategies derived from our expert planning group

discussions and two pilot sessions, but for future studies we will utilize an

accredited group facilitation course. Second, Fitbit measurements of heart rate

may be less accurate for those with darker skin tone.[53] However, Fitbit was mainly

used for step-counting which does not have this limitation, and heart rate was only

used for an assessment of convergent validity with the validated videogame.[54] That

the study was limited to those who were able to afford Internet access at home

and time to attend the sessions does limit some generalizability and

implementation. Finally, as a small single-group study design, we had limited ability

to robustly determine efficacy outcomes.

492                                    **CONCLUSIONS**

In summary, we report protocol development and feasibility of a virtual

home intervention to promote MVPA among peer groups of adolescents with T1D.

Successes included attraction of a diverse group, high attendance at the sessions,

T1D self-management guidance that promoted safe MVPA, and meaningful

connections to including shared experiences and skills in T1D self-management

and coping. The challenge was technical delays especially transitions between

virtual reality apps. Therefore, future trials testing efficacy and moderators warrant

financial investment to integrate and interoperate the virtual reality apps. The

advantages of our program were effectiveness at connecting teens to peers,

clinicians, and young adult coaches in a way that was logistically feasible, time

efficiency for the clinicians and coaches since they could interact with a group all

at once, and sustainment of favorable MVPA and clinical metrics.

**DATA AVAILABILITY**

Individual de-identified participant data (including data dictionaries) were

shared. This includes individual participant data that underlies the results reported

in any aspect of a published article (text, tables, figures, and appendices). Other

documents that will be available include the study protocol, statistical analysis plan,

informed consent form, and analysis code. The data will be available immediately

following publication, with no end dates. The data will be shared with researchers

who provide a methodologically sound proposal to achieve the aims of the

approved proposal. The proposals should be directed to Dr. Garrett Ash at Yale

University (garrett.ash@yale.edu). To gain access, data requesters must sign a

data access agreement.

**AUTHORSHIP CONFIRMATION/CONTRIBUTION STATEMENT**

**GIA:** Conceptualization (lead), data curation (lead), formal analysis (lead), funding

acquisition (lead), investigation (equal), methodology (equal), project

administration (lead), resources (lead), software (support), supervision (lead),

validation (equal), visualization (lead), writing – original draft (lead). **SN:**

Conceptualization (equal); investigation (equal); methodology (equal); writing –

review and editing (equal). **MSK:** Methodology (supporting); Writing – review and editing (equal). **ADH:** Investigation (equal); methodology (equal); project administration (equal); software (equal); writing, review, and editing (supporting). **CT:** Investigation (equal); methodology (equal); writing, review, and editing (equal). **AC:** Investigation (equal); methodology (equal); writing – review and editing (equal). **MS:** Investigation (equal); methodology (equal); writing, review, and editing (equal). **JSB:** Methodology (supporting); Writing – review and editing (equal). **SAW:** Conceptualization (equal); investigation (equal); methodology (equal); writing – review and editing (equal); writing – review and editing (equal). **LMN:** Conceptualization (equal), formal analysis (equal), investigation (equal), methodology (lead), software (lead), visualization (equal), writing – review and editing (lead).

## FUNDING SOURCE

The study and G.I.A. were supported by American Heart Association Grant #852679 (G.I.A., 2021–2024) and a Robert E. Leet and Clara Guthrie Patterson Trust Mentored Research Award, Bank of America, N.A., Trustee. G.I.A. and L.M.N. were supported by the National Institute of Diabetes, Digestive, and Kidney Diseases of the National Institutes of Health under mentored research scientist development awards (K01DK129441, K23DK128560). None of these entities were involved in the manuscript writing, editing, approval, or decision to publish.

## CONFLICT-OF-INTEREST DISCLOSURE

G.I.A. scientifically advises Behavioral Tech Health Innovations. S.A.W. receives grant support to his university from Abbott Laboratories. M.S. is employed as a product manager by Abbott Laboratories and receives restricted stock units. L.M.N. receives funding for research from the National Institutes of Health and Medtronic Diabetes. She is also a consultant for Medtronic, WebMD and Calm.

None of these entities supported the above study, and the authors conducted the

research outside of their responsibilities and affiliations with these entities.

**ACKNOWLEDGMENTS**

The authors thank the families and participants that help made this research

possible. We also acknowledge Sa'Ra Skipper and Juanita Montoya for their

assistance with conducting the study. Study data were collected and managed

using REDCap electronic data capture tools hosted at Yale University.

555
556 **REFERENCES**
557
1.    Foster NC, Beck RW, Miller KM, et al. State of Type 1 Diabetes
Management and Outcomes from the T1D Exchange in 2016-2018. Diabetes
technology & therapeutics 2019;21(2):66-72, doi:10.1089/dia.2018.0384 [doi]
2.    Nathan DM, Group DER. The diabetes control and complications
trial/epidemiology of diabetes interventions and complications study at 30 years:
overview. Diabetes care 2014;37(1):9-16, doi:10.2337/dc13-2112 [doi]
3.    Adolfsson P, Taplin CE, Zaharieva DP, et al. ISPAD Clinical Practice
Consensus Guidelines 2022: Exercise in children and adolescents with diabetes.
Pediatr Diabetes 2022;23(8):1341-1372, doi:10.1111/pedi.13452
4.    Pongrac Barlovic D, Harjutsalo V, Groop PH. Exercise and nutrition in type
1 diabetes: Insights from the FinnDiane cohort. Front Endocrinol (Lausanne)
2022;13(1064185, doi:10.3389/fendo.2022.1064185
5.    Perez KM, Patel NJ, Lord JH, et al. Executive function in adolescents with
type 1 diabetes: relationship to adherence, glycemic control, and psychosocial
outcomes. Journal of Pediatric Psychology 2017;42(6):636-646
6.    Moser O, Riddell MC, Eckstein ML, et al. Glucose management for
exercise using continuous glucose monitoring (CGM) and intermittently scanned
CGM (isCGM) systems in type 1 diabetes: position statement of the European
Association for the Study of Diabetes (EASD) and of the International Society for
Pediatric and Adolescent Diabetes (ISPAD) endorsed by JDRF and supported by
the American Diabetes Association (ADA). Pediatric diabetes 2020;21(8):1375-
1393, doi:10.1111/pedi.13105 [doi]
7.    Riddell MC, Gallen IW, Smart CE, et al. Exercise management in type 1
diabetes: a consensus statement. The lancetDiabetes & endocrinology
2017;5(5):377-390, doi:S2213-8587(17)30014-1 [pii]
8.    Amiel SA, Sherwin RS, Simonson DC, et al. Impaired insulin action in
puberty. A contributing factor to poor glycemic control in adolescents with diabetes.
The New England journal of medicine 1986;315(4):215-219,
doi:10.1056/NEJM198607243150402 [doi]
9.    Jabbour G, Henderson M, Mathieu ME. Barriers to Active Lifestyles in
Children with Type 1 Diabetes. Canadian journal of diabetes 2016;40(2):170-172,
doi:10.1016/j.jcjd.2015.12.001 [doi]
10.    Speight J, Holmes-Truscott E, Garza M, et al. Bringing an end to diabetes
stigma and discrimination: an international consensus statement on evidence and

recommendations. Lancet Diabetes Endocrinol 2024;12(1):61-82,
doi:10.1016/S2213-8587(23)00347-9
11.    Lawton K. People with type 1 diabetes held back from sport due to stigma
and fear of hypos. Juvenile Diabetes Research Foundation: 2023.
12.    Chinchilla P, Dovc K, Braune K, et al. Perceived Knowledge and
Confidence for Providing Youth-Specific Type 1 Diabetes Exercise
Recommendations amongst Pediatric Diabetes Healthcare Professionals: An
International, Cross-Sectional, Online Survey. Pediatric Diabetes
2023;2023(8462291, doi:10.1155/2023/8462291
13.    Benzing V, Schmidt M. Exergaming for Children and Adolescents:
Strengths, Weaknesses, Opportunities and Threats. Journal of clinical medicine
2018;7(11):422. doi: 10.3390/jcm7110422, doi:10.3390/jcm7110422 [doi]
14.    Council On C, Media. Media Use in School-Aged Children and
Adolescents. Pediatrics 2016;138(5), doi:10.1542/peds.2016-2592
15.    Akhter K, Turnbull T, Simmons D. A systematic review of parent/peer-
based group interventions for adolescents with type 1 diabetes: interventions
based on theoretical/therapeutic frameworks. Br J Diabetes 2018;18(51-65
16.    Prochaska JJ, Sallis JF, Long B. A physical activity screening measure for
use with adolescents in primary care. Archives of Pediatrics & Adolescent
Medicine 2001;155(5):554-559
17.    Warburton DE, Gledhill N, Jamnik VK, et al. Evidence-based risk
assessment and recommendations for physical activity clearance: Consensus
Document 2011. Applied physiology, nutrition, and metabolism = Physiologie
appliquee, nutrition et metabolisme 2011;36 Suppl 1(S266-98, doi:10.1139/h11-
062 [doi]
18.    Currie C IJ, Molcho M, Lenzi M, Veselska Z & Wild F (ed). Health Behaviour
in School-aged Children (HBSC) Study Protocol: Background, Methodology and
Mandatory items for the 2013/14 Survey. CAHRU: St Andrew;
19.    Allen NA. Social cognitive theory in diabetes exercise research: an
integrative literature review. The Diabetes educator 2004;30(5):805-819,
doi:10.1177/014572170403000516 [doi]
20.    Ryan RM, Deci EL. Self-determination theory and the facilitation of intrinsic
motivation, social development, and well-being. Am Psychol 2000;55(1):68-78,
doi:10.1037//0003-066x.55.1.68
21.    Ng JY, Ntoumanis N, Thogersen-Ntoumani C, et al. Self-Determination
Theory Applied to Health Contexts: A Meta-Analysis. Perspect Psychol Sci
2012;7(4):325-40, doi:10.1177/1745691612447309
22.    Huang Y-C, Backman SJ, Backman KF, et al. An investigation of motivation
and experience in virtual learning environments: A self-determination theory.
Education and Information Technologies 2019;24(591-611
23.    Riddell MC, Gallen IW, Smart CE, et al. Exercise management in type 1
diabetes: a consensus statement. Lancet Diabetes Endocrinol 2017;5(5):377-390,
doi:10.1016/s2213-8587(17)30014-1
24.    Adolfsson P, Riddell MC, Taplin CE, et al. ISPAD Clinical Practice
Consensus Guidelines 2018: Exercise in children and adolescents with diabetes.
Pediatric diabetes 2018;19 Suppl 27(205-226, doi:10.1111/pedi.12755 [doi]
25.    Staiano AE, Abraham AA, Calvert SL. Motivating effects of cooperative
exergame play for overweight and obese adolescents. Journal of diabetes science
and technology 2012;6(4):812-819, doi:dst.6.4.0812 [pii]
26.    Ash GI, Joiner KL, Savoye M, et al. Feasibility and safety of a group
physical activity program for youth with type 1 diabetes. Pediatric diabetes 2019,

doi:10.1111/pedi.12841 [doi]

27. DeJonckheere M, Joiner KL, Ash GI, et al. Youth and Parent Perspectives
on the Acceptability of a Group Physical Activity and Coping Intervention for
Adolescents With Type 1 Diabetes. Sci Diabetes Self Manag Care 2021;47(5):367-
381, doi:10.1177/26350106211040429

28. Titoria R, Amed S, Tang TS. Peer Support Interventions on Digital
Platforms for Children With Type 1 Diabetes and Their Caregivers. Diabetes
Spectr 2022;35(1):26-32, doi:10.2337/ds21-0081

29. Walker AF, Haller MJ, Gurka MJ, et al. Addressing health disparities in type
1 diabetes through peer mentorship. Pediatr Diabetes 2020;21(1):120-127,
doi:10.1111/pedi.12935

30. Troncone A, Cascella C, Chianese A, et al. Psychological support for
adolescents with type 1 diabetes provided by adolescents with type 1 diabetes:
The chat line experience. Pediatric diabetes 2019;20(6):800-810,
doi:10.1111/pedi.12873 [doi]

31. Floyd BD, Block JM, Buckingham BB, et al. Stabilization of glycemic control
and improved quality of life using a shared medical appointment model in
adolescents with type 1 diabetes in suboptimal control. Pediatric diabetes
2017;18(3):204-212, doi:10.1111/pedi.12373 [doi]

32. Staiano AE, Adams MA, Norman GJ. Motivation for Exergame Play
Inventory: Construct validity and relationship to game play. Cyberpsychology:
Journal of Psychosocial Research on Cyberspace 2019;13(3):Article 7,
doi:10.5817/CP2019-3-7

33. Chin WW, Salisbury WD, Pearson AW, et al. Perceived Cohesion in Small
Groups: Adapting and Testing the Perceived Cohesion Scale in a Small-Group
Setting. Small Group Research 1999;30(6):751-766,
doi:10.1177/104649649903000605

34. Faulkner MS, Michaliszyn SF, Hepworth JT, et al. Personalized exercise
for adolescents with diabetes or obesity. Biological research for nursing
2014;16(1):46-54, doi:10.1177/1099800413500064 [doi]

35. Quirk H, Blake H, Tennyson R, et al. Physical activity interventions in
children and young people with Type 1 diabetes mellitus: a systematic review with
meta-analysis. Diabetic medicine : a journal of the British Diabetic Association
2014;31(10):1163-1173, doi:10.1111/dme.12531 [doi]

36. Troiano RP, Berrigan D, Dodd KW, et al. Physical activity in the United
States measured by accelerometer. Medicine and science in sports and exercise
2008;40(1):181-188, doi:10.1249/mss.0b013e31815a51b3

37. Viechtbauer W, Smits L, Kotz D, et al. A simple formula for the calculation
of sample size in pilot studies. Journal of clinical epidemiology 2015;68(11):1375-
1379, doi:10.1016/j.jclinepi.2015.04.014 [doi]

38. Weissberg-Benchell J, Shapiro JB, Bryant FB, et al. Supporting Teen
Problem-Solving (STEPS) 3 year outcomes: Preventing diabetes-specific
emotional distress and depressive symptoms in adolescents with type 1 diabetes.
J Consult Clin Psychol 2020;88(11):1019-1031, doi:10.1037/ccp0000608

39. Grey M, Whittemore R, Jeon S, et al. Internet psycho-education programs
improve outcomes in youth with type 1 diabetes. Diabetes care 2013;36(9):2475-
2482, doi:10.2337/dc12-2199 [doi]

40. Fritz CO, Morris PE, Richler JJ. Effect size estimates: current use,
calculations, and interpretation. J Exp Psychol Gen 2012;141(1):2-18,
doi:10.1037/a0024338

41. Gottfried JS, O. Teens and Video Games Today. 2024. Available from:

https://www.pewresearch.org/internet/2024/05/09/teens-and-video-games-today/.
42.    Wong CH, Chiang YC, Wai JP, et al. Effects of a home-based aerobic
exercise programme in children with type 1 diabetes mellitus. Journal of Clinical
Nursing 2011;20(5-6):681-691, doi:10.1111/j.1365-2702.2010.03533.x [doi]
43.    Marrero DG, Fremion AS, Golden MP. Improving compliance with exercise
in adolescents with insulin-dependent diabetes mellitus: results of a self-motivated
home exercise program. Pediatrics 1988;81(4):519-525
44.    Shetty VB, Soon WHK, Roberts AG, et al. A Novel Mobile Health App to
Educate and Empower Young People With Type 1 Diabetes to Exercise Safely:
Prospective    Single-Arm    Mixed    Methods    Pilot    Study.    JMIR    Diabetes
2021;6(4):e29739, doi:10.2196/29739
45.    Ryninks K, Sutton E, Thomas E, et al. Attitudes to Exercise and Diabetes
in Young People with Type 1 Diabetes Mellitus: A Qualitative Analysis. PloS one
2015;10(10):e0137562, doi:10.1371/journal.pone.0137562 [doi]
46.    Ye CY, Jeppson TC, Kleinmaus EM, et al. Outcomes That Matter to Teens
With    Type    1    Diabetes.    The    Diabetes    educator    2017;43(3):251-259,
doi:10.1177/0145721717699891 [doi]
47.    Xie LF, Housni A, Nakhla M, et al. Adaptation of an Adult Web Application
for Type 1 Diabetes Self-management to Youth Using the Behavior Change Wheel
to Tailor the Needs of Health Care Transition: Qualitative Interview Study. JMIR
Diabetes 2023;8(e42564, doi:10.2196/42564
48.    Doe E. An analysis of the relationships between peer support and diabetes
outcomes in adolescents with type 1 diabetes. J Health Psychol 2018;23(10):1356-
1366, doi:10.1177/1359105316656228
49.    Bisno DI, Reid MW, Pyatak EA, et al. Virtual Peer Groups Reduce HbA1c
and Increase Continuous Glucose Monitor Use in Adolescents and Young Adults
with    Type    1    Diabetes.    Diabetes    Technol    Ther    2023;25(9):589-601,
doi:10.1089/dia.2023.0199
50.    Luyckx K, Seiffge-Krenke I, Missotten L, et al. Parent-adolescent conflict,
treatment adherence and glycemic control in Type 1 diabetes: the importance of
adolescent    externalising    symptoms.    Psychol    Health    2013;28(9):1082-97,
doi:10.1080/08870446.2013.782405
51.    Armstrong S, Wong CA, Perrin E, et al. Association of Physical Activity With
Income, Race/Ethnicity, and Sex Among Adolescents and Young Adults in the
United States: Findings From the National Health and Nutrition Examination
Survey, 2007-2016. JAMA pediatrics 2018, doi:10.1001/jamapediatrics.2018.1273
[doi]
52.    Piona C, Marigliano M, Mozzillo E, et al. Evaluation of HbA1c and glucose
management indicator discordance in a population of children and adolescents
with type 1 diabetes. Pediatr Diabetes 2022;23(1):84-89, doi:10.1111/pedi.13299
53.    Koerber D, Khan S, Shamsheri T, et al. Accuracy of Heart Rate
Measurement with Wrist-Worn Wearable Devices in Various Skin Tones: a
Systematic    Review.    J    Racial    Ethn    Health    Disparities    2022;1-9,
doi:10.1007/s40615-022-01446-9
54.    Comeras-Chueca C, Villalba-Heredia L, Pérez-Llera M, et al. Assessment
of Active Video Games' Energy Expenditure in Children with Overweight and
Obesity and Differences by Gender. International journal of environmental
research and public health 2020;17(18):6714. doi: 10.3390/ijerph17186714,
doi:10.3390/ijerph17186714 [doi]
743

**Figure S1.** Enrollment flow-chart. *Online advertising consisted of Children with Diabetes posting the advertisement in three issues of their weekly newsletter (February 9th–February 23rd, 2022) as well as their Facebook page on February 15th. The Facebook post received 12 likes, 10 shares, and 23 email inquiries as listed. †One participant required clearance for a medical condition from a specialist (arthritis) prior to enrollment.

**Table S1.** Structure of intervention sessions.

| Component | Saturdays Time of Day[1] | Wednesdays Time of Day |
|---|---|---|
| Icebreaker question | 10:00-10:05 | 19:00-19:05 |
| Sensor glucose check and guidance[2] | 10:05-10:20 | 19:05-19:20[3] |
| Exercise | 10:20-11:00 | 19:20-19:40[3] |
| Sensor glucose check and guidance | 11:00-11:05 | 19:40-19:45[3] |
| Discussion Activities<br>• MVPA goals **(week 1**: set goals; **weeks 2-5:** check progress and provide MVPA and T1D management information supporting goals; **week 6:** reflection on relationship to overall active lifestyle)<br>• Role-playing skits (**weeks 2-5:** planning and acting) | 11:05-11:30 | 19:45-20:00 |

[1]USA eastern time zone, which was the local time of all participants except 1 who was USA mountain time zone (2 hours earlier)

[2]Extended to 30 minutes at first session to give general overview of guidelines. This session had reduced exercise by 8 minutes and discussion activities by 5-10 minutes.

[3]Cohort #1 only had these components on Saturdays (i.e., Wednesday session only met 19:00-19:20). Added these components to Wednesdays per expressed interest of cohort #2.

**Table S2.** Individual Goals (n=15)

| Goal Category | Number of participants | Specific Value | Achieved Goal |
|---|---|---|---|
| Steps | 9 | School to home, 3d/week<br>School to home, every day<br>One mile around neighborhood, every day<br>4000 per day<br>7000 per day<br>8500 per day<br>10000 per day<br>12500 per day<br>13000 per every other day | 2 / 9 (22%) |
| Sport | 5 | Find a soccer group<br>Bike 26.2 miles<br>Run 8:00 mile<br>Run 6:35 mile<br>Run 7:30 pace for 5 miles<br>Dead lift 230 lbs<br>Push-up reps 20<br>Bike 30min per day<br>Pole vault reach minimum club competition clearance<br>Gym twice per week<br>Exercise 3 days per week<br>Run timed mile each week | 4 / 12 (33%) |
| Ring Fit | 3 | 20 minutes per day<br>500 reps of 1 specific movement<br>Reach world 9 | 1 / 3 (33%) |
| Diabetes management | 1 | 7.0% estimated A1c from CGM | 0 / 1 (0%) |
| Total | 15 | | 7 / 25 (28%) |

**Table S3.** Structure of topics, roles, and competencies taught during skits for each cohort.

| Topic | Roles[1] | Educational Points and How Integrated |
|---|---|---|
| Team sports and type 1 diabetes (T1D) | -Athlete with newly diagnosed with T1D<br>-Teammate living with T1D for several years<br>-Continuous glucose monitor with alerts<br>-Coach | -Speaking to coaches who are not familiar with T1D management around MVPA<br>-Peer mentoring<br>-Competitive sports and glycemic management |
| Zombie Apocalypse | -Zombie<br>-Zombie-killing engineer<br>-Teen newly diagnosed with T1D and knows nothing about what they need for T1D management during an apocalypse<br>-Teen with longer duration of T1D<br>-Dr. Rescue<br>-News host<br>-Continuous glucose monitor with alerts | -Essential supplies for T1D<br>-Prioritizing aspects of T1D management<br>-Operating continuous glucose monitors<br>-Impact of MVPA on glycemic management<br>-Impact of stress on glycemic management |
| Wilderness Excursion | -Lilo (Hawaiian girl)<br>-Stitch (helpful alien)<br>-Evil singing crab<br>-Battler of evil singing crab<br>-Diabuddy (diabetes buddy)<br>-Angel<br>-Narrator | -Same as Zombie Apocalypse |

[1]Self-chosen by interested participants, remaining ones assigned.
MVPA, moderate-to-vigorous physical activity

**Table S4.** Summary of CGM for study 1. Reported as mean (SD). Normally distributed except where noted.

| Time Frame | Weeks 1-2 | Weeks 3-4 | Weeks 5-6 |
|---|---|---|---|
| Mean glucose (mg/dL) | 200.5 (53.1) | 199.2 (40.8) | 195.6 (41.9) |
| SD of mean glucose (mg/dL) | 73.6 (17.6) | 76.3 (18.6) | 73.8 (19.4) |
| CV (%) | 37.4 (6.9) | 38.7 (8.1) | 38.0 (4.5) |
| % time < 54 mg/dL[1] | 0.7 (1.0) | 0.8 (1.1) | 0.4 (0.6) |
| % time <70 mg/dL[1] | 2.8 (2.7) | 3.2 (4.4) | 2.7 (2.3) |
| % time 70-180 mg/dL | 46.2 (22.2) | 44.0 (18.6) | 48.6 (21.0) |
| % time >180 mg/dL | 51.0 (22.4) | 52.8 (19.4) | 48.5 (20.1) |
| % time > 250 mg/dL | 27.6 (24.0) | 27.5 (18.6) | 25.1 (20.0) |
| % time with sensor data | 74.3 (33.6) | 73.6 (31.8) | 72.8 (33.8) |

Abbreviations: CGM: continuous glucose monitor, SD: standard deviation, CV: coefficient of variation.
[1]Right-skewed. Specifically, more than one-third of participant-weeks had zero or negligible values: <0.2% for % time <54 mg/dL and/or <1.0% for % time <70 mg/dL.

**Table S5.** Summary of continuous glucose monitoring, insulin, and anthropometric data for study 2. Given as mean (SD).

| Time Frame | Weeks 1-2 | Weeks 3-4 |
|---|---|---|
| Mean glucose (mg/dL) | 189.1 (33.1) | 186.0 (28.8) |
| SD of mean glucose (mg/dL) | 74.3 (18.0) | 73.5 (15.9) |
| CV (%) | 39.0 (4.0) | 39.3 (3.4) |
| % time < 54 mg/dL[1] | 0.7 (1.3) | 0.7 (1.5) |
| % time <70 mg/dL[1] | 2.4 (2.1) | 2.1 (2.2) |
| % time 70-180 mg/dL | 51.9 (16.7) | 53.0 (!6.2) |
| % time >180 mg/dL | 45.7 (16.2) | 44.9 (14.5) |
| % time > 250 mg/dL | 22.0 (15.2) | 21.2 (13.9) |
| % time with sensor data | 75.0 (30.5) | 75.0 (36.7) |

Abbreviations: SD: standard deviation, CV: coefficient of variation.
[1]Right-skewed. Specifically, more than one-third of participant-weeks had zero or negligible values: <0.2% for % time <54 mg/dL and/or <1.0% for % time <70 mg/dL.