1

2   Integrating More Walking into Public Transit Commuting: A Proof-of-Concept Study

3   **Yuval Hadas, PhD**
4   Department of Management
5   Bar-Ilan University, Ramat Gan, 5290002, Israel
6   Email: j.rab@outlook.com
7

8   **Dan Emanuel Katz, MS**
9   Department of Management
10   Bar-Ilan University, Ramat Gan, 5290002, Israel
11   Email: rabinowz@biu.ac.il
12

*Jonathan Rabinowitz, PhD
Wellness Research Lab
School of Social Work
Bar-Ilan University, Ramat Gan, 5290002, Israel
Email: Jonathan.rabinowitz@biu.ac.il
ORCiD: 0000-0002-6845-8064

*Corresponding author: Jonathan Rabinowitz, PhD

Acknowledgements: This work was supported by a grant of the Israeli Smart Transportation

Research Center (ISTRC).

Declarations of interest: none

28
29

30   Abstract

Regular physical activity is essential for maintaining and improving overall health, yet many

individuals struggle to incorporate it into their daily lives. Previous studies, primarily based on

simulations, have suggested that mode shifts in public transportation (PT) can increase walking,

enabling PT commuters to meet some physical activity goals while traveling. However, current

trip planning apps used by PT commuters do not prioritize walking as part of the journey. This

proof-of-concept study investigates the potential for integrating more walking into commutes

using PT, without compromising travel time, accessibility, or level of service. It represents the

first phase of the "More Walking" population-based wellness initiative, which aims to enhance

walking opportunities by modifying trip planning apps. We examined the impact of different

walking thresholds on walking distance using a trip-planning algorithm using the home addresses

of 2,149 commuters traveling to the same workplace in a Tel Aviv suburb. Our findings reveal

that increasing walking thresholds either maintained or reduced total travel time, these results

suggest that introducing a "more walking" option in trip planning could be a feasible way of

incorporating physical activity and wellness as part of daily commutes.

Keywords: Trip planning, transportation, walkability, public health, physical activity,

commuting

1. BACKGROUND

Regular physical activity is essential for maintaining and improving overall health, yet many individuals struggle to incorporate it into their daily lives. Walking has important health benefits, a recent meta-analysis that included almost 50,000 persons found that taking more steps per day was associated with a progressively lower risk of all-cause mortality (1). For many people, walking is part of daily routine, it is frequently the last mile solution for commuters using public transportation (PT).

1.1 LITERATURE REVIEW

The health benefits of walking have been studied thoroughly. A short walk of 20 minutes each day can reduce the chance of early death by 25%. Walking helps prevent chronic diseases such as heart disease, type 2 diabetes, risk of stroke and some cancers (1-3). Walking is easy to include as part of everyday activities. More people walking also benefits the community, it enhances both actual and perceived safety of public spaces, reduces vehicle emissions and traffic congestion, stimulates local businesses and economic growth (4). Moreover, walking as a mode of transport reduces financial burden, leading to transport savings. Promoting walking gives people the option to move around without the need for a private vehicle. Accessibility to PT ensures that individuals have diverse and convenient mobility, contributing to a more inclusive and accessible urban environment (5).

Previous studies, based on simulations (6-15), and projections based on sampling of GPS data (16), have reported the potential for increased walking resulting from mode shifts in public transport. Previous studies did not consider the impact of variable walking distances on overall travel time. In the current study, we used the origins and the destination of actual commuters.

We examined the effects of changing walking thresholds in trip planning on time spent walking

as well as on overall travel time. The study is part of our quest to identify and promote simple,

sustainable and impactful healthy lifestyle changes. Our approach is grounded in time-use

epidemiology which explores effects of health-related time-use patterns in populations and

optimizing distribution of time for population health (17).

Many PT users rely on trip planning apps to navigate their routes and manage fare

payments. These apps do not prioritize walking as part of the journey. Popular trip planners,

such as Google Maps, Moovit, Citymapper, Apple Maps, Bus Nearby (Hopon Mobility, Ltd) and

OneBusAway, lack the options to adjust the maximum allowable walking distance or set a

minimum desired walking threshold. Some apps, like efobus (efobus.com), allow adjusting

maximum aerial distance as a very rough proxy for walking distance. Another group of app's

focus on helping walkers find the most inviting route (e.g., walkonomics.com 2010-2018) but do

not include PT. However, including an upwardly adjustable walking distance parameter in the

trip planning application could potentially provide the user with additional trip options, the

potential of quicker and more reliable trips (less transfers), and improved wellbeing (18).

We conducted a proof-of-concept study with the aim of demonstrating that increasing

walking thresholds among PT users is a practical and effective method for enhancing physical

activity without extending travel time. This study marks the first phase of the 'More Walking'

project, a population-based wellness initiative aimed at promoting walking by modifying trip

planning apps to incorporate increased walking options. In the current phase reported in this

paper, we present a proof-of-concept study in which we examined whether by increasing walking

thresholds in a trip planning app commuters could include walking more in their commute and

getting to their destination without adding travel time. Put simply, we examined whether one

can leave home and get to work at the same time and include a walk.

2. METHODS

To test the concept that commuters could incorporate more walking without increasing travel

time, we used Open Trip Planner (OTP) software (19). OTP is a tool that helps plan itineraries

using public transportation, walking, biking, or other options. It uses data from publicly available

maps and public transportation routes and schedules to plan efficient routes. Using OTP, we

planned travel itineraries to work, using the addresses of the 2,149 employees sharing the same

workplace, Bar-Ilan University, situated in a suburb of Tel Aviv. OTP allows adjusting limits on

walking distance. We varied the amount of walking distance from 100 to 2500 meters. The

research protocol was exempted from full review by the Human Subjects Institutional Review

Board at Bar-Ilan University as the use of this data set which contained only a list of addresses

without other identifying information, had been previously approved for use in another study.

The first step was to generate PT itineraries using OTP. The destination in every itinerary

was Bar-Ilan University and the origins were the employees home addresses. By varying

walking distance and number of stops, multiple itineraries for each employee's commute were

generated. Each itinerary included all trip legs, the start and finish stops, modes, ride/walk times,

waiting times and distance. The travel itineraries generated by OTP were saved into a database

for further analysis.

The second step was to eliminate itineraries that were not likely to be used. This was done in

Python using the Hierarchical Non-Dominated Sorting (NHDS) Pareto Front Identification

algorithm **(20)** which identifies optimal solutions based on multiple criteria. The algorithm first

sorts all itineraries for each employee in ascending order by travel time and walking distance.

Then it compares the first itinerary with all others, to make a rapid distinction between different

itineraries to see whether a solution's total travel time and walk time are greater than or equal to

those of any other solution. If both criteria are met, the solution is considered inferior and

removed from the selection pool. This filtering process helped to eliminate inferior trip options

that passengers would be unlikely to choose. The remaining solutions, which ranged from 2 to 6

for each employee, were grouped by different walking thresholds.

Figure 1 shows the process of applying the algorithm and a route generated. The

employees' address (aka origin) and the university address (aka destination) were input to OTP.

Using local maps and transportation data, OTP generated routes with the chosen relevant

information, including the trip start time, travel time, distance, and other details of the itinerary.

The extracted data was then used to generate a new file that stores the results.

Map in Figure 1 depicts a PT itinerary from OTP with key attributes. We used the

following parameters presented in Figure 1 to analyze the impact of walking time and distance in

the PT system on total travel time: (1) Time to reach the first stop from the origin (aka ingress

time), (2) Transit time, (3) Transfer walk time, (4) Waiting time, and (5) Time to reach the

destination from the last stop (aka egress time). The overall walking time includes walking time

from the employees' address to the boarding point, the transfer walking time (between transfer

stations) and from the alighting point to destination. Transit time is the time spent traveling

within the PT network. The number of transfers made during a trip indicates the complexity and

convenience of interchanging between different PT modes or lines. Waiting time measures the

time passengers spend waiting for the arrival of PT while transferring.

Figure1 Process diagram and route generated

Process diagram of methodology

Route generated

1433.0 RESULTS

Figure 2 maps the location of all 2,149 employees. Each employee's address is

represented by a dot, which is colored based on the travel time by car to the campus. The yellow

dot is the location of the campus. Data on potential itineraries travelling to work at Bar-Ilan

University using public transportation were examined for various walking thresholds ranging

from 100 to 2500 meters.

Figure 2 Origins (Bar-Ilan employees) and Destination (Bar-Ilan Campus) used in the case study

Table 2 and Figure 3 present a descriptive summary by walking distance thresholds for

all employees and for the subset of employees living within a 30-minute drive (Table 2 b, Figure

3, orange line). As seen in Table 2 (a & b) and Figures 3 (a, b & c), when more walking is

introduced, there are more possible itineraries available, more origins are covered, meaning more

employees can walk and use PT to get to work and less transfers are required. For example, only

33 employee addresses had routes requiring less than 100 meters walking whereas almost all of

employees (2,135 out of 2,149) had routes requiring up to 2,500 meters of walking. For those

living within 30 minutes by car travel (Table 2(b) and Figure 3 (d) orange line), transit time also

declines as walking distance threshold increases. These decreases support the concept that the

more passengers are willing to walk, the more direct and efficient the trip options become,

reducing the need for transfer connections and waiting. Beyond the threshold of 1,000 meters of

walking, the average number of transfers had only minor fluctuations. An additional advantage

offered by the increased walking distance is that there are more stops that can be used upon

arrival. There is one station at the entrance to the campus, 2 stations within 300 meters, 4

stations within 500 meters and 7 stations within 1000 meters.

167
168

169 **TABLE 2 Descriptive summary by walking threshold (all times in minutes, rounded to nearest**

170 **minute)**

171(a) All commuters

Walking threshold (meters)	Number of Trips	Covered Origins	Average Total Travel Time	Average Walk Time	Average Transit Time	Average Waiting Time	Average Number of Transfers	Average Car Travel Time
100	92	33	48	1	35	12	1.97	29
500	3,925	1,101	47	5	35	8	1.49	27
1,000	10,021	1,980	52	8	37	8	1.36	30
1,500	11,941	2,127	53	9	37	7	1.33	31
2,000	12,168	2,132	53	9	37	7	1.32	31
2,500	12,228	2,135	53	9	37	7	1.31	31

173(b) Commuters with less than 30 minutes by car (all times in minutes)

Walking threshold (meters)	Number of Trips	Covered Origins	Average Total Travel Time	Average Walk Time	Average Transit Time	Average Waiting Time	Average Number of Transfers	Average Car Travel Time
100	51	18	30	1	23	6	1.53	19
500	2,793	728	32	5	23	5	1.10	18
1,000	7,239	1,239	34	8	23	4	0.94	19
1,500	8,589	1,320	34	9	21	4	0.89	19
2,000	8,730	1,321	34	9	21	4	0.88	19
2,500	8,762	1,323	34	9	21	4	0.88	19

177 **Figure 3 Effects of increasing walking thresholds (in meters) for all commuters (blue line) and those**

178 **living within 30 minutes by car (orange line)**

(a) *Number of trip options (relative, %) as a function of walking distance threshold*

(b) *Covered origins (relative, %) as a function of walking distance threshold*

(c) *Number of transfers (relative, %) as a function of walking distance threshold*

(d) *Transit time as a function of walking distance threshold*

4.1 Discussion of results

As part of a population-based wellness intervention research initiative, we conducted a

proof-of-concept study with the aim of examining the possibility that more walking could be

included in commutes on public transportation. We did this by increasing walking thresholds in

trip planning leveraging home address data on 2,149 employees working at the same location.

We found that for most of the potential routes, as walking thresholds were increased, total travel

time did not increase, and commuters were potentially able to benefit on average from a 9-

minute walk going to work (walking threshold of 1500 meters). Considering that a short walk of

20 minutes each day can reduce the chance of early death by 25% (1-3) increasing walking

thresholds as part of a daily commute has the potential to result in considerable health benefits.

To further contextualize these findings we note that the United Nations has set in its

Sustainability Development Guidelines a target of "Affordable and sustainable transport

systems" (Target 11.2) which they define as convenient access to public transport within 500

meters to a stop or a station (https://sdgs.un.org/goals).

Previous research had used simulated data to demonstrate the concept that more physical

activity could be included in commuting with public transportation (14, 21, 22). We have gone a

step beyond demonstrating this by using real data to support the concept. These findings set the

stage for our next study, which is a population-based intervention encouraging commuters to

increase walking thresholds in their trip planning transportation apps.

4.2 Discussion of methods and limitations
201
Studies of transportation are dependent on infrastructures, services and locations. As

such, each place differs, and the results based on this population may not be applicable to others.

In this proof-of-concept study we were able to control destinations, as all people worked at the

same location. Since each place differs all studies in this area are in some form or another case-

studies.   We believe that the findings from this case study are situationally grounded and

generalizable meeting the duality principle for informative case research (23). Bar-Ilan

University is in the greater Tel Aviv metropolitan area, with no special dedicated buses to the

campus nor a train station. The results from this case are important as they provide additional

evidence based on actual data of origin and a fixed destination that supports what other studies

have found using simulations about the effects of modal switches in transit and walking. We

have gone a step beyond, considering the effects of varying walking thresholds.

While the results are encouraging, we do not know the extent to which commuters will

incorporate more walking; this will be explored in the next phase of the More Walking research

program. Our results are limited as we were not able to take into consideration critical elements

as noted in public policy transport guidelines as critical when attempting to integrate more

walking into public transportation (24). These include elements such as safety, accessibility, how

enjoyable walking routes are, public transport stops and stations, particularly for individuals with

reduced mobility. Specifically, we did not take into account presence of infrastructure for

weather protection, terrain mitigation, quality of street lighting, pedestrian-friendly crossings,

wayfinding signage, obstacle-free pathways, sidewalk life and active frontages, need to carry

cargo, possible difficulty walking long-distances and infrastructure to support individuals with

disabilities. Finally, we did not take into consideration the risk of missing the service or missing

a transfer or the effect of various walking paces. Future work in this area should strive to

address the aforementioned and include measures of walkability (25-27).

5. Conclusions

A key potential benefit of incorporating more walking into public transportation (PT)

commutes is the promotion of physical activity. By including walking segments in their

journeys, commuters are more likely to engage in regular exercise, which can enhance their

health and overall well-being. Integrating walking into daily commutes can play a significant

role in fostering a healthier lifestyle.

The results of this proof-of-concept study suggest that increased walking as part of

commute has the potential to result in shorter overall travel times and a larger variety of journey

options for passengers. By having more choices for their routes, commuters can select paths that

best suit their preferences and schedules. Additionally, this increase in journey options may

enhance the overall reliability of the transportation system. When faced with unexpected delays

or disruptions, passengers have alternative routes to choose from, this may minimize the negative

impact on their daily commutes. Another benefit observed in the analysis is the reduction in the

number of transfers and overall waiting time as walking distance increases. As passengers cover

more distance on foot, the need for multiple transfers decreases. This translates to shorter waiting

times at transfer points, contributing to a more time-efficient and convenient commuting

experience. As a result, passengers can experience quicker and more efficient trips, making PT a

more attractive option for daily commuting, increasing the potential that they will walk more

daily.

In conclusion, this proof-of-concept study highlights the potential benefits of

incorporating increased walking into public transportation (PT) commutes. Beyond the obvious

health advantages, greater integration of walking within PT commutes could enhance reliability,

expand journey options, and even improve overall time efficiency. Additionally, reducing

transfers and waiting times can boost the effectiveness and convenience of the transportation

system. By positioning walking as a key element of PT commutes, this approach shows promise

in optimizing PT services while fostering healthier and more sustainable urban mobility. The

next phase of this research will explore whether promoting increased walking can be effectively

achieved by embedding walking thresholds into PT trip planning applications.

Declarations

Ethics approval and consent to participate: Bar-Ilan University Institutional Review Board

exempted the study from review and consent as this was as an analysis of an existing

administrative data file, previously approved for use in a different study, and did not require

collecting data from human subjects.

Consent for publication: Not applicable.

Availability of data and materials: Due to privacy concerns data can not be made available.

Competing Interests: The authors declared no potential completing of interest with respect to

the research, authorship, and/or publication of this article.

Funding: This work was supported by a grant of the Israeli Smart Transportation Research

Center (ISTRC).

Authors' contributions: The authors confirm contribution to the paper as follows: study

conception and design: YH, DK, JR; data collection: YH; analysis and interpretation of results:

YH, DK; draft manuscript preparation: YH, DK, JR. All authors reviewed the results and

approved the final version of the manuscript.

Acknowledgments: Not applicable.

REFERENCES

1. Paluch AE, Bajpai S, Bassett DR, Carnethon MR, Ekelund U, Evenson KR, et al. Daily steps and all-cause mortality: a meta-analysis of 15 international cohorts. Lancet Public Health. 2022;7(3):e219–e28.

2. Garcia L, Pearce M, Abbas A, Mok A, Strain T, Ali S, et al. Non-occupational physical activity and risk of cardiovascular disease, cancer and mortality outcomes: a dose–response meta-analysis of large prospective studies. British Journal of Sports Medicine. 2023;57(15):979.

3. Powell KE, King AC, Buchner DM, Campbell WW, DiPietro L, Erickson KI, et al. The Scientific Foundation for the Physical Activity Guidelines for Americans, 2nd Edition. J Phys Act Health. 2018:1–11.

4. Zhu X, Lu Z, Yu CY, Lee C, Mann G. Walkable communities: Impacts on residents' physical and social health: Researchers from Texas A&M University studied residents in a newly developed 'walkable community' in Austin, Texas to see how it changed their habits for physical activity and whether it increased social interaction and cohesion in the community. World Health Des. 2013;6(3):68–75.

5. Badawi Y, Maclean F, Mason B. The economic case for investment in walking2018. Available from: https://www.victoriawalks.org.au/Assets/Files/The-Economic-Case-for-Investment-in-Walking-FINAL.pdf.

6. Brondeel R, Kestens Y, Chaix B. An evaluation of transport mode shift policies on transport-related physical activity through simulations based on random forests. International Journal of Behavioral Nutrition and Physical Activity. 2017;14(1):143.

7. Brown V, Barr A, Scheurer J, Magnus A, Zapata-Diomedi B, Bentley R. Better transport accessibility, better health: a health economic impact assessment study for Melbourne, Australia. International Journal of Behavioral Nutrition and Physical Activity. 2019;16(1):89.

8. Kwan SC, Tainio M, Woodcock J, Sutan R, Hashim JH. The carbon savings and health co-benefits from the introduction of mass rapid transit system in Greater Kuala Lumpur, Malaysia. Journal of Transport & Health. 2017;6:187–200.

9. Mahendra A, Rajagopalan L. Evaluating health impacts from a bus rapid transit system implementation in India: case study of indore, Madhya Pradesh. Transportation Research Record. 2015;2531(1):121–8.

10. Morency C, Trépanier M, Demers M. Walking to transit: an unexpected source of physical activity. Transport Policy. 2011;18(6):800–6.

11. Rissel C, Curac N, Greenaway M, Bauman A. Physical activity associated with public transport use—a review and modelling of potential benefits. International journal of environmental research and public health. 2012;9(7):2454–78.

12. Rojas-Rueda D, de Nazelle A, Teixido O, Nieuwenhuijsen MJ. Replacing car trips by increasing bike and public transport in the greater Barcelona metropolitan area: a health impact assessment study. Environment international. 2012;49:100–9.

13. Zapata-Diomedi B, Knibbs LD, Ware RS, Heesch KC, Tainio M, Woodcock J, et al. A shift from motorised travel to active transport: What are the potential health gains for an Australian city? PLoS One. 2017;12(10):e0184799.

14. Del Rosario L, Laffan SW, Pettit CJ. Physical activity associated with public transport commutes and potential feeder mode shifts. Journal of Transport & Health. 2024;35:101751.

15.   Garcia L, Johnson R, Johnson A, Abbas A, Goel R, Tatah L, et al. Health impacts of
changes in travel patterns in Greater Accra Metropolitan Area, Ghana. Environment
international. 2021;155:106680.
16.   Chaix B, Kestens Y, Duncan S, Merrien C, Thierry B, Pannier B, et al. Active
transportation and public transportation use to achieve physical activity recommendations? A
combined GPS, accelerometer, and mobility survey study. International journal of behavioral
nutrition and physical activity. 2014;11:1–11.
17.   International Network of Time-Use Epidemiologists (INTUE) [Available from:
https://www.intue.org/about/our-mission/.
18.   Rabinowitz J. America Walks. 2021. Available from: https://americawalks.org/hacking-
the-map-apps-for-active-transportation-less-walking-vs-more-walking-as-a-means-of-optimized-
movement-and-mobility.
19.   OpenTripPlanner. Multimodal Trip Planning 2020 [Available from:
http://www.opentripplanner.org/.
20.   Bao C, Xu L, Goodman ED, Cao L. A novel non-dominated sorting algorithm for
evolutionary multi-objective optimization. Journal of Computational Science. 2017;23:31–43.
21.   Wasfi RA, Ross NA, El-Geneidy AM. Achieving recommended daily physical activity
levels through commuting by public transportation: Unpacking individual and contextual
influences. Health & place. 2013;23:18–25.
22.   Patterson R, Webb E, Millett C, Laverty AA. Physical activity accrued as part of public
transport use in England. Journal of public health. 2019;41(2):222–30.
23.   Ketokivi M, Choi T. Renaissance of case research as a scientific method. Journal of
Operations Management. 2014;32(5):232–40.
24.   Walk21. Integrating Walking + Public Transport: Policy Brief. 2024.
25.   Shields R, Gomes da Silva EJ, Lima e Lima T, Osorio N. Walkability: a review of trends.
Journal of Urbanism: International Research on Placemaking and Urban Sustainability.
2023;16(1):19–41.
26.   Lee M, Kim S, Kim H, Hwang S. Pedestrian visual satisfaction and dissatisfaction toward
physical components of the walking environment based on types, characteristics, and
combinations. Building and Environment. 2023;244:110776.
27.   Cantley A, Hurley JC, Todd M, McEntee M, Hooker SP, Ohri-Vachaspati P, et al.
Walkability around the worksite and self-reported and accelerometer-measured physical activity
among adults. Health & Place. 2024;85:103143.
350