Title: Phase I study on the pharmacokinetics of intravaginal, self-administered artesunate vaginal

pessaries among women in Kenya.

Short title: Pharmacokinetics of intravaginal, self-administered Artesunate vaginal pessaries

among women in Kenya.

Authors: Chemtai Mungo, MD, MPH[1,2]; Katherine Sorgi[1], Brenda Misiko[3], Cynthia Cheserem[3],

Lisa Rahangdale, MD, MPH[1,2]; George Githongo [3], Cirilus Ogollah[3], Jackton Omoto, MBChB,

MMed[4], Mihaela Plesa, BA, CCRP[5], William Zamboni, Pharm.D., Ph.D.[6]

Affiliations

1. Department of Obstetrics and Gynecology, University of North Carolina Chapel Hill, 321

S Columbia St, Chapel Hill, North Carolina, 27599, USA.

2. Lineberger Comprehensive Cancer Center, University of North Carolina-Chapel Hill, 450

West Dr, Chapel Hill, North Carolina, 27599, USA

3. Kenya Medical Research Institute, Busia Rd, Kisumu, Kenya

4. Department of Obstetrics and Gynecology, Maseno University School of Medicine, P.O,

Kisumu, Kenya

5. Frantz Medical Development Ltd, 7740 Metric Dr, Mentor, OH, 44060

6. Division of Pharmacotherapy and Experimental Therapeutics, UNC Eshelman School of

Pharmacy, University of North Carolina-Chapel Hill, 450 West Dr, Chapel Hill, North

Carolina, 27599, USA

Corresponding Author:

Chemtai Mungo, MD, MPH, chemtai_mungo@med.unc.edu

Abstract (300-word max):

Cervical cancer remains a significant global health issue, especially in low- and middle-income

countries (LMICs), where access to prevention and treatment is limited and women are at a

higher risk of cervical cancer. Artesunate, a widely available drug used to treat malaria, has

shown promise in treating human papillomavirus (HPV)-associated anogenital lesions including

high-grade cervical precancer, in a recent Phase I studies in the United States. Data on the

pharmacokinetics of artesunate following intravaginal use, and its implications on malaria

resistance, are lacking.

Objectives: The primary objective of this study is to investigate the pharmacokinetics of

Artesunate (AS) and its active metabolite, dihydroartemisinin (DHA) following intravaginal use

at the dosing and frequency intended for cervical precancer treatment. A secondary objective is

to assess safety among study participants.

Methods: We are conducting a single-arm, phase I trial with a sample size of 12 female

volunteers. Participants will self-administer artesunate vaginal pessaries in the study clinic daily

for 5 consecutive days. Participants will have their blood drawn prior to receiving the first dose

of artesunate on day one of the study and then will receive 8 blood draws on study day five, prior

to artesunate administration and at 15 minutes, 30 minutes, 1 hour, 2 hours, 4 hours, 6 hours, and

8 hours after pessary administration. Pharmacokinetic parameters of artesunate and DHA will be

calculated by way of quantitative analysis of with determination of maximum concentration

(Cmax), time to Cmax (Tmax), area under the serum concentration versus time curve (AUC),

apparent clearance, and elimination half-life (t1/2).

Keywords: pharmacokinetics, artesunate, dihydroartemisinin, intravaginal use, cervical

precancer, malaria resistance

Introduction:

Cervical cancer is the second most common cancer in women globally, with a high

prevalence in low- and middle-income countries like Kenya. It is caused by human

papillomavirus (HPV) infection but can be prevented through vaccinations or early screening to

find precancerous changes in the cervix – also known as cervical intraepithelial neoplasia. While

these methods can prevent cervical cancer from developing in women, there are access and

resource barriers, especially in LMICs. For instance, HPV vaccinations are new in Kenya and

other LMICs, and very few women under the age of 20 have been vaccinated. There are also cost

barriers and shortages in healthcare providers. In countries like Kenya and other LMICs where

there are few nurses and doctors, many women with cervical precancer are often referred to far

away facilities to access treatment and due to costs and other challenges, many are unable to

access the referral centers with providers who can offer treatment.

These barriers have led to researchers exploring the option of self-administered

treatments for cervical precancer to overcome healthcare provider and healthcare resource

shortages. One treatment being studied is an intravaginal pessary formulation of the drug

Artesunate, which is often used to treat malaria. While a U.S. study has shown its safety and

early efficacy for cervical precancer treatment,[1] it is unknown whether intravaginal use of

artesunate vaginal inserts at the dosing and frequency used, may result in systemic exposure and

hence have implications for malaria resistance. To fill this gap in the literature, this study seeks

to investigate the pharmacokinetics of intravaginal artesunate at the dose and frequency used in

ongoing cervical precancer treatment studies.

Intravaginal Artesunate for Cervical Precancer Treatment

Artesunate (AS), a World Health Organization (WHO) approved malaria drug, is being

explored as a potential topical therapy for cervical precancer as its safety and tolerability has

long been well established.[2-5] Growing evidence suggests its cytotoxic effects against numerous

cancer cell lines both *in vitro* and *in vivo* and have revealed its proposed mechanisms of action

include suppressing cell proliferation by inducing G1 and G2M phrase cell cycle arrest and

modulation of inflammatory pathways characteristic of uncontrolled proliferation and

carcinogenesis.[6-9] Ferroptosis, an iron-dependent cell death type, is thought to be a key

anticancer mechanism for HPV-infected cells.[10] Cancer cells are highly proliferative, requiring a

heavy iron load which acts as a cofactor in synthesizing deoxyriboses before cell division.[11]

Development of both high-grade cervical intraepithelial neoplasia (CIN2/3), the precursor lesion

of cervical cancer, and cervical cancer are associated with the expression of two viral proteins in

the HPV lifecycle, E6 and E7.[12] Epithelial cells that express either or both of these oncoproteins

also overexpress the transferrin receptor, and have been shown to have increased levels of

intracellular iron compared with normal cells.[10] This observation has been exploited to

investigate whether preinvasive cervical cancer (CIN2/3), can be treated with Artesunate, which

contains an endoperoxide bridge that reacts with intracellular ferrous iron to generate free

radicals, capable of inducing direct oxidative damage resulting in cell death.[9] Given the

mechanism of action, artesunate may provide beneficial and therapeutic effects for intraepithelial

HPV disease.

Several studies demonstrate the pro-apoptotic effects of Artesunate by activating the

mitochondria-dependent pathway, including caspase-3/9 activation and cytochrome c release

from the mitochondria.[13] It induces autophagy in uterine corpus endometrial carcinoma cells and

elevates reactive oxygen species in human bladder cancer cells and hepatocellular carcinoma

cells.[14-16] Artesunate has demonstrated anti-angiogenic properties by downregulation VEGF and

angiopoietin-1 in myeloma cells.[17-19] Its reported anti-cancer effects involve the formation of

alkaline radicals through an endoperoxide bridge, reacting with intracellular ferrous iron and

leading to cell death.[20] The overexpression of the transferrin receptor in cervical squamous cell

cancers and their precursors,[21] cervical intraepithelial neoplasia (CIN), prompted a study of

cytotoxic effects of dihydroartemisinin (DHA), the bioactive form of Artesunate, on

papillomavirus-expressing epithelial cells.[22] *In vitro* studies revealed that DHA had minimal

impact on normal cervical epithelial cells but had significantly induced cytotoxicity in HPV-

immortalized cervical cells.[22] Administered as a local treatment in a canine nonclinical model

with a 100% known tumor growth rate with HPV-infection, DHA (2.22 mg dissolved in 100 µl

dimethyl sulfoxide) reported to inhibit papillomavirus-induced tumor formation. In addition,

tumor-negative dogs developed antibodies against the HPV L1 capsid protein.[22]

Safety for Intravaginal Use

In a multi-center dose-escalation phase 1 study in the United States, intravaginal

artesunate inserts (pessaries) were tested for safety, tolerability, and efficacy in women with

CIN2/3.[1] This "first-in-human" study demonstrated that intravaginal Artesunate for CIN2/3

treatment was safe, well tolerated, and resulted in self-limited adverse events that were graded I

and II. Reported AEs among participants who used 3 five-day artesunate cycles included chills

and flu-like symptoms (n=3, grade 1), vaginal (yeast) infection (n=1, grade II), dizziness or

headache (n=2, grade 1), non-infective cystitis (n=1, grade 2), vaginal pain or uterine cramping

(n=9, grade I), vaginal discharge (n=4, grade 1), vaginal pruritis (n=9, grade 1). In summary, 37

drug-related AEs were observed in this Phase I trial, of which 34 (92%) were grade I, and 3 (8%)

were grade 2. Reported grade 2 adverse events included vaginal yeast infection (n = 6), bacterial

vaginosis (n = 2), vaginal inflammation (n = 2), urinary tract infection (n = 2), and noninfective

cystitis (n = 1). Adverse events that were determined to be unrelated to the study medication

included: anxiety, insomnia, suicidal ideation, vaginal twitching, fever, flu-like symptoms in a

patient who developed a cold, body itching, chills, and eczema flare. No subjects withdrew from

the study because of intolerable side effects, and all 28 subjects included in the modified-

intention-to-treat analyses were able to complete their designated dosing regimen. No grade 3 or

4 adverse events were reported, and three subjects reported no noticeable side effects. Currently,

a randomized placebo-controlled phase II study of artesunate vaginal inserts in treatment of

CIN2/3 is enrolling subjects at several US sites, with no serious adverse events reported so far

(NCT04098744). Additionally, a phase II randomized trial of artesunate suppositories for

treating anal high-grade squamous intraepithelial lesions among HIV-negative individuals is

ongoing in the United States (NCT5555862)

Summary of Artesunate Pharmacokinetics

The pharmacokinetics (PK) measures of artesunate (AS) and dihydroartemisinin (DHA)

following intravenous (IV), intramuscular (IM), oral and rectal administration have been well

described.[23] These PK parameters include maximum concentration of the drug in the blood

(Cmax), time to maximum concentration Cmax (Tmax), apparent clearance (CL/F) - the drug

concentration in the body in proportion to the rate of elimination, volume of distribution (V/F),

the area under the serum concentration versus time curve (AUC) which expresses the total

amount of the drug in systemic circulation after administration, and half-life ($t_{1/2}$). IV

administration of AS quickly produces high maximum concentrations (Cmax) of AS, higher than

any other method of administration. One example of this can be seen in a study of adults with

uncomplicated malaria which compared AS and DHA levels following IV and IM

administration, Cmax values for AS when administered intravenously reached over 16,000ng/mL

while only reaching around 884mg/mL when administered intramuscularly.[24] Similarly, the

maximum concentration (Cmax) of AS and DHA peak quickly following IV administration,

followed by a rapid decline. The average half-life of AS following IV administration of less than

fifteen minutes in multiple studies, with an observed clearance range of 2-3 L/kg/hr and a

volume range of 0.1-0.3 L/kg.[23] The hydrolysis of AS into DHA following IV administration is

similarly quick, with maximum DHA levels reached soon after IV AS administration. The Tmax

for DHA after IV administration was consistently less than 25 minutes according to the observed

studies with DHA clearance averaging between 0.5-1.5 L/kg/hr and volume averaging between

0.5-1.0 L/kg.[23] The AUC of AS following IV administration of 120 mg AS in adults ranged

between 876ng*hr/ml in healthy volunteers[25] and 1038 - 1269ng*hr/ml in those with malaria.[26]

The AUC of DHA following IV administration of 120 mg AS in adults ranged from 1850

ng*hr/ml in healthy volunteers[27] to 1845-2377ng*hr/ml in those with malaria.[28]

Compared to IV administration, IM administration produces lower peaks, longer half-life,

and higher volumes of distribution for AS, as well as delayed peaks for DHA. For example, AS

half-life following in IV administration is less than 15 minutes on average, compared to 25.2 to

48.2 minutes with IM administration.[23] Other parameters – in including DHA half-life, volume

of distribution, and clearance rates following IM administration resembled the values recorded

after IV administration in multiple studies, due to the high bioavailability, assessed by exposure

to DHA, associated with IM AS administration (>86%).[23] The AUC of AS following IM

administration of 120 mg AS in adults ranged between 856[29] - 999[25]ng*hr/ml in those with

malaria. The AUC of DHA following IM administration of 120 mg of AS in adults with malaria

was 2474ng*hr/ml in one study.[25]

When AS is administered orally, DHA peak concentrations (Cmax), AUC, and half-life

averages are all notably higher than comparable AS parameters. The average time to DHA Cmax

and half-life following oral AS administration (200 mg/day) are 2 hours, and 0.5-1.5 hours,

respectively, compared to one hour and 20-45 minutes, respectively, for AS.[23] While a similar

pattern is seen following IV and IM administration, namely, elevated levels of DHA half-life and

AUC compared to AS, the difference are notable following oral administration. Also, while oral

administration of AS results in a higher DHA Cmax compared to AS, IV and IM administration

result in a notably higher AS Cmax than DHA Cmax. Morris et al (2011) points out that the

variations observed following oral administration are most likely attributed to AS functioning as

a "pro-drug" for DHA when ingested orally and in response to "first-pass or systemic

metabolism."[23] Essentially, when AS is taken orally, it is converted to DHA at a greater extent

than when it is taken intravenously or intramuscularly. The AUC of AS and DHA in a study of

healthy adult volunteers taking 200mg AS daily for 5 days was 67 ng*hr/ml and 1158 ng*hr/ml

for AS & DHA, respectively, on Day 1, and 60 ng*hr/ml and 1300 ng*hr/ml for AS & DHA,

respectively, on Day 5.[30] Similar Cmax parameters for AS were 67 ng/ml and 58 ng/ml on Day 1

and 5, respectively, and a pooled DHA Cmax of 654 ng/ml, demonstrating the absence of time-

dependent artesunate pharmacokinetics in healthy subjects during 5-day oral administration of

200 mg artesunate.[30] The AUC of AS following oral administration of 200 mg AS in adults

ranged between 60-67ng*hr/ml in healthy volunteers[31] and 310ng*hr/ml in those with malaria.[32]

The AUC of DHA following oral administration of 200 mg of AS in adults ranged from 1158-

1331 ng*hr/ml[31,33] in healthy volunteers to 3027ng*hr/ml in those with malaria.[32] Rectal AS

administration yields pharmacokinetic results similar to those obtained from oral administration,

with the exception of delayed AS Cmax and longer AS half-life. Compared to IV administration,

expectedly, both AS absorption and elimination are prolonged following rectal administration.

Following rectal administration of AS, Tmax average between 0.58-1.43 hours, with a half-life

between 0.9-.95 hours.[23] These averages are based on three different studies: two studies

containing pediatric patients with uncomplicated falciparum malaria (10 - 20mg/Kg dosing)[34,35]

and one study containing healthy Malaysian adults (200 mg rectal suppository, ~4mg/Kg

dosing).[33] Following rectal dosing of a one-time 200 mg AS suppository in healthy adults

(similar to our planned dosing), a Cmax, Tmax, half-life and AUC of 448.5 ng/ml, 1.43 hours,

0.95 hours, and 796 ng*hr/ml of AS were observed, and Cmax, Tmax, half-life and AUC of

385.6 ng/ml, 1.80 hours, 1.21 hours, and 965 ng*hr/ml respectively of DHA were observed in

healthy adults.[33] No data are available on rectal PK in adults with malaria as this route of

administration is not used to treat malaria in adults. The longer half-life of AS following rectal

dosing (average 0.9-0.95 hrs) compared to IM (average 25.2 – 48.2 minutes), or IV (average less

than 5 minutes) may reflect absorption rate-limited elimination of AS.[23] As is expected given

that rectal AS administration avoids by-pass metabolism, the discrepancy in AS and DHA AUC

values (796 ng*hr/ml and 965 ng*hr/ml, respectively), is not as striking with rectal, as compared

with oral administration of AS (119 ng*hr/ml and 1331 ng*hr/ml, respectively).[23] Similar to oral

administration, both DHA Tmax and half-life values were higher than that of AS following rectal

administration.[23]

Effect of malaria infection status on artesunate and DHA pharmacokinetics

Teja-Isavadharm et al., conducted a direct comparison of DHA pharmacokinetics

following oral AS administration in healthy adults and falciparum malaria patients.[36] The

investigators found significantly higher AUC and Cmax of DHA in subjects with malaria as

compared to healthy subjects. Similar results were obtained by Binh et al in a study comparing

the PK in eight patients with falciparum malaria and ten healthy subjects.[37] Due to the small size

of both studies, definitive conclusion regarding differences in PK between healthy and infected

subjects cannot be drawn.[27] However, as DHA clearance is dependent on hepatic blood flow, a

reduction in clearance, and consequently an increase in exposure associated with acute malaria

infection, would be consistent with known DHA's PK properties.[27]

Malaria treatment and artemisinin resistance

Artemisinin-based combination therapies (ACTs) are recommended by the WHO as the

first line treatment for uncomplicated Plasmodium Falciparum.[38] In ACTs, artemisinin quickly,

and drastically, reduces the majority of malaria parasites, with the partner drug clearing the

remaining parasites to prevent recrudescence.[39-40] Artemisinin resistance is defined as delayed

parasite clearance (following treatment with an artesunate monotherapy or ACT) observed as a

parasite clearance half-life greater than five hours or microscopic evidence of parasites on day

three.[41] This represents partial resistance. While artemisinin resistance alone does not necessarily

lead to malaria treatment failure, reduced efficacy of the artemisinin component places greater

demands on the partner drug to clear a larger parasite mass, jeopardizing future efficacy (WHO).

Examples of ACTs include artemether-lumefantrine, artesunate-amodiaquine, artesunate-

mefloquine, among others. Standard ACTs regimen for uncomplicated malaria is an oral 3-day

course. Most studies indicate that current ACTs recommended in national malaria treatment

policies remain effective, with an overall efficacy rate of greater than 95%.[42]

Artemisinin's act exceptionally fast against intra-erythrocytic asexual blood-stage malaria

parasites, affecting up to 10,000- fold reductions in parasite burden every 48 hours.[43] The

primary genetic drivers of artemisinin resistance, both *in vitro* and *in vivo,* are point mutations in

the P. falciparum Kelch13 (*PfK13*) gene during the early ring stage.[44,45] These mutations allow a

subset of early ring-stage parasites to survive cell-cycle arrest brought on by artemisinin

exposure, enabling those parasites to reinitiate transcription and complete their intraerythrocytic

developmental cycle once artemisinin is no longer present at inhibitory concentrations.[46] *In vitro*

resistance is routinely defined as greater than 1% survival of early ring-stage parasites exposed to

700nM dihydroartemisinin (DHA-the primary active metabolite of ART) for 6 hours, followed

by drug-free culture incubation for a further 66 hours.[41] The resistance mechanism appears to

involve a complex interplay of K13 protein abundance, hemoglobin endocytosis, and the parasite

response to stress.[41,47]

In Africa, several studies have identified a number of low-frequency Pfk13 mutations

associated with delayed parasite clearance in four countries: Ghana, Rwanda, Uganda

Tanzania.[48,49] Mutations including M476I, P553L, R561H, P574L, C580Y and A675V, were

observed at low frequencies under 5%.[48] For example, in Tanzania, mutations were found in two

parasites from 764 samples in 2027 (0.3%)[6], and one parasite from 422 samples in 2019

(0.2%).[48] Similarly in Uganda, one parasite was identified from 796 samples in 2018/2019

(0.1%).[50] A 2021 study in Northern Uganda from 2017 to 2019 identified in vivo artemisinin

resistance (parasite clearance half-life >5 hours) in a total of 14 out of 240 patients who received

intravenous artesunate.[50] Of these 14 patients, 13 were infected with P. falciparum parasites with

mutations in the A675V or C469Y allele in the kelch13 gene.[45]

P. falciparum resistance to artemisinin has been documented in five countries in

Southeast Asia; Cambodia, Lao People's Democratic Republic, Myanmar, Thailand and Vietnam

(WHO).[40] With implementation of combination therapy, improvements to health systems and

surveillance systems to monitor first- and second-line treatment, the consequences of the

development of resistance to antimalarial medicines may be less severe today than what was

observed with chloroquine in the 1980s. If parasites develop reduced sensitivity to artemisinin,

ACTS will continue to cure malaria, as long as the partner drug remains effective.[40] To

overcome resistance, potential changes can be made to ACT. Some of these include

modifications such as extending the duration of the ACT course (currently 3 days for oral

treatment), alternating use of different ACT regimens, and addition of another antimalarial drug

to the standard ACTs (triple-ACT).[40] Additionally, adding a malaria vaccine (e.g. RTS, S

vaccine) to mass drug administration campaigns could enhance treatment efficacy and help

prevent further artemisinin resistance development.

Systemic Artesunate absorption and possible implications for developing resistance for

malaria treatment.

Although there is no available pharmacologic data on serum absorption following

intravaginal artesunate administration, this study aims to address this gap. The direct application

of artesunate to the cervical mucosa at the proposed dose of 200mg ($\leq$4 mg/kg based on planned

inclusion criteria of weight$\geq$ 50 Kg) is unlikely to result in systemic absorption. This planned

dosage is 2.5-fold lower than the approved rectal suppository dose (10mg/Kg). In contrast to the

rectal mucosa, which is highly vascular and comprised of single-cell layer of columnar

epithelium making it highly permeable, the cervico-vaginal tissue has a thick, stratified

squamous epithelial cell layer and is significantly less vascular, reducing its systemic

absorption.[51] Similarly, due to the rapid rate of elimination of Artesunate's active metabolite,[52]

no systemic accumulation is expected of Artesunate or its active metabolite with intravaginal

multi-day dosing in the context of cervicovaginal administration.

Methods

Study Objectives

Primary Objective

The primary objective is to determine the area under the serum concentration versus the

time curve (AUC) of DHA following five consecutive days of self-administration of Artesunate

vaginal inserts among healthy women living in Kenya

Secondary Objectives

1. Determine the area under the serum concentration versus time curve (AUC) of Artesunate

(AS) following five consecutive days of self-administration of 200mg Artesunate vaginal

inserts (pessaries) among healthy women in Kenya

2. Determine the maximum concentration of Artesunate (AS) (Cmax) following five

consecutive days of self-administration of 200mg Artesunate vaginal inserts (pessaries)

among healthy women in Kenya

3. Determine the maximum concentration of dihydroartemisinin (DHA) (Cmax) following

five consecutive days of self-administration of 200mg Artesunate vaginal inserts

(pessaries) among healthy women in Kenya

4. Determine the time to maximum concentration (Tmax) of Artesunate (AS) following five

consecutive days of self-administration of 200mg Artesunate vaginal inserts (pessaries)

among healthy women in Kenya

5. Determine the time to maximum concentration (Tmax) of dihydroartemisinin (DHA)

following five consecutive days of self-administration of 200mg Artesunate vaginal

inserts (pessaries) among healthy women in Kenya

6. Determine the half-life ($t_{1/2}$) of Artesunate (AS) following five consecutive days of self-

administration of 200mg Artesunate vaginal inserts (pessaries) among healthy women in

Kenya

7. Determine the half-life ($t_{1/2}$) of dihydroartemisinin (DHA) following five consecutive

days of self-administration of 200mg Artesunate vaginal inserts (pessaries) among

healthy women in Kenya

8. Determine the apparent clearance (CL/F) of Artesunate (AS) following five consecutive

days of self-administration of 200mg Artesunate vaginal inserts (pessaries) among

healthy women in Kenya

9. Determine the apparent clearance (CL/F) of dihydroartemisinin (DHA) following five

consecutive days of self-administration of 200mg Artesunate vaginal inserts (pessaries)

among healthy women in Kenya

10. Determine the volume of distribution (V/F) of Artesunate (AS) following five

consecutive days of self-administration of 200mg Artesunate vaginal inserts (pessaries)

among healthy women in Kenya

11. Determine the volume of distribution (V/F) of dihydroartemisinin (DHA) following five

consecutive days of self-administration of 200mg Artesunate vaginal inserts (pessaries)

among healthy women in Kenya

12. Investigate the safety of 5-day course of self-administered intravaginal artesunate vaginal

inserts (pessary) in women in Kenya

Study design

This is a single-arm, nonrandomized, interventional phase I study among 12 women over

the age of 18 living in Kisumu, Kenya. Participants will self-administer 200 mg artesunate

vaginal inserts for five consecutive days with blood draws performed on day 1 (at baseline) and

on day 5. Day 1 will consist of one blood draw prior to first pessary use and day 5 will consist of

eight blood draws: one prior to final artesunate administration, at time 0, then at 15 minutes, 30

minutes, 1 hour, 2 hours, 4 hours, 6 hours, and 8 hours after final artesunate pessary insertion.

The schedule of events is outlined in Figure 1.

Study setting

This study will be conducted in Kisumu, Kenya at the Lumumba Sub-County Hospital

and at the Victoria Biomedical Research Institute. All study activities will take place at

Lumumba Sub-County Hospital except the day 5 study visit, which will take place at Victoria

Biomedical Research Institute.

Study Drug

In this study, a vaginal insert (pessary) formulation of artesunate will be used. Artesunate

is a derivative of artemisinin which is the active principle of the Chinese medicinal herb

Artemisia annua. The quantitative composition of artesunate inserts utilized in this study is

detailed in Table 1.

352
353 *Table 1: Composition of artesunate inserts.*
354

Ingredient	Function
Artesunate	Active ingredient
PCCA MBK^{TM}* - hydrogenated vegetable oil and PEG-8 distearate.	Base
Butylated Hydroxyl Anisole NF	Preservative

355
The drug is produced by Buderer Drug Co in Perrysburg Ohio and is compounded in

accordance with USP Chapter 795 (Pharmaceutical Compounding – Nonsterile Preparations) and

USP Chapter 797 (Pharmaceutical Compounding – Sterile Preparations). These chapters

establish practice standards and outline the responsibility of the compounder, selection and

appropriate sources of ingredients, quality control, and considerations regarding the stability of

compounded preparations. Additionally, USP Chapter 1075 (Good Compounding Practices) and

USP Chapter 1160 (Pharmaceutical Calculations in Prescription Compounding) provide further

guidance ensuring the quality and accuracy of the compounded Artesunate.

For each compounded lot of vaginal inserts used in this study, potency testing is

conducted to confirm the dosage of Artesunate. Each lot has an expiration date of 6 months after

compounding, although testing on batches beyond this period has demonstrated no significant

loss in strength. This formulation, at various doses, has been used in two Phase 1/2A clinical

trials for cervical and anal HSIL[1,53].

Eligibility and Recruitment

Recruitment

Participants will be recruited from the general population within close proximity to the study location in Kisumu County, which will include recruitment from local health facilities. The study team will conduct local outreach activities and educational talks in the community to harness interest in participating. If an individual is interested in participating, they will be screened for eligibility and, if eligible, will provide their informed consent and will be briefed on other study procedures before any study activities are performed. Inclusion and exclusion criteria for this study is outlined in Table 2. Recruitment started on 29/05/2024.

Table 2: Inclusion and Exclusion Criteria

Inclusion Criteria	Exclusion Criteria
Age 18 years to 65 years	Current pregnancy or breastfeeding status
Negative pregnancy test at screening	History of total hysterectomy
Weight ≥50 Kg at study entry*	Known allergy to Artesunate
Willingness to use contraception (hormonal or barrier) during the 5-day study dosing phase if of childbearing age (less than 50 years of age)	Have a medical comorbidity that in the opinion of the investigator would interfere with study participation
Ability and willingness to provide informed consent	Currently receiving artemisinin-based agents for malaria treatment or completed artemisinin-based treatment within the previous 3 days.**

Plan to reside in the study location during the study period	Male at birth
Agrees for samples without identifiers to be shipped outside of Kenya for testing	Current use of efavirenz antiretroviral therapy
	Positive malaria antigen test at screening

382 *Justification for weight criteria: A minimum body weight of 50Kg will meet the planned
383 artesunate dosing of ≤4 mg/Kg for which excellent safety data is available.
384 **Based on dihydroartesunate (artesunate's active metabolite) half-life of between 0.5-1.5
385 hours[29]
386

Study procedures by visit

Pre-screening visit: Prior to enrollment of a participant, the study staff will pre-screen potential

participants at on-site or off-site locations. During these visits, study staff will explain the aspects

of the study to potential participants and will provide an explanation of eligibility requirements.

Screening/Enrollment visit: To begin the screening process, eligible participants will provide

their written informed consent for the study procedures. Once informed consent is obtained by

the study staff, the screening processes will begin, including a malaria antigen test to screen for

subacute malaria infection. If a participant tests positive for malaria, they will be referred for

treatment and considered screen failures, resulting in their discontinuation from participating in

the study. Given that the study is located in a malaria-endemic region, we have planned for a

high screening-to-enrollment ratio due to the likelihood of high rates of sub-clinical malaria.

Other screening and enrollment activities will involve the collection of basic demographic and

clinical data, study protocol training (including demonstration of intravaginal artesunate

application using a pelvic model), a limited physical and pelvic exam, a urine pregnancy test, and

a review of prohibited medications.

Visit 1: On the same day or up to 14 days after the screening and enrollment visit, the participant

will have their first study visit in the clinic. During this visit, eligibility will be confirmed, the

study protocol will be reviewed, and study staff will provide instructions on self-administering

artesunate. Additionally, the participant will have 2.5ml of blood drawn prior to their first

intravaginal self-administration of artesunate. Once self-administration of artesunate is complete,

the participant will insert a tampon and will be observed for 30 minutes prior to scheduling their

second visit on the following day.

Visits 2-4: Visits 2, 3, and 4 must occur within 24 hours of each other so the study staff will

attempt to schedule the visits at approximately the same time each day. During these visits, the

study staff will review and record any adverse events (AE) using a standardized questionnaire

and the U.S. National Cancer Institute Common Terminology Criteria for Adverse Events

(CTCAE). If an adverse event is scored a grade II or worse, a pelvic exam will be performed.

After review of adverse events, and reiteration of prohibited medications, the participant will be

asked to again self-administer artesunate followed by insertion of a tampon. Once they are

observed for 30 minutes, the following visit will be scheduled.

Visit 5: Visit 5 will occur no more than 24 hours after visit 4, with the study staff aiming to

schedule it at approximately the same time as the prior day's visit. During this visit, the study

staff will review and record adverse events, conducting a pelvic exam as needed based on the

grading of observed AEs. After AEs are reviewed and a reiteration of prohibited medications is

given, the participant will undergo their first 2.5ml blood draw of the visit before the insertion of

artesunate. A peripheral cannula will be inserted at this time for subsequent blood draws.

Following the initial blood draw, the participant will perform their final self-

administration of artesunate, followed by the insertion of a tampon. Once the final self-

administration of artesunate is complete, 2.5ml of blood will be drawn after 15 minutes, 30

minutes, 1 hour, 2 hours, 4 hours, 6 hours, and 8 hours. After the last blood draw, participants

will be scheduled for the final visit, which will take place 4-10 days after visit 5.

Visit 6: Visit 6 will occur 4-10 days after visit 5. During this visit, the study staff will review and

record adverse events since the last visit. Once AEs are recorded, the study staff will initiate

study termination and will provide the participant with any necessary financial reimbursement.

Participant retention plan

Prior to any study procedures, the informed consent document will be reviewed with

participants to ensure a clear understanding of the study purpose, procedures, duration,

compensation, and any other factors that are necessary to understand. To document their

willingness to participate, each participant and a study personnel will sign the informed consent

documents. Once enrolled and consented, the study staff will consistently follow-up with

participants to ensure they attend all scheduled appointments, to report any adverse event

experiences, and to answer any questions that the participant may have about their role in the

study. To compensate for their time, each participant will be reimbursed Kshs 1000

(approximately $10) for each visit to account for potential loss of wages and will be reimbursed

for transport to and from the clinic. For the final visit, which is expected to last up to 8 hours,

participants will be reimbursed an additional Kshs 2,500 ($25) for loss of wages. Participants

may also receive refreshments during the visit depending on how long they are asked to wait. If

there is ever concern that a participant is lost to follow-up, the study staff will make every effort

to regain contact with the participant, whether that be through calls or home visits.

Statistical consideration

Sample size

This study will have a sample size of 12 healthy volunteers.

Sample size justification

Participants will self-administer 200 mg artesunate vaginal inserts for five consecutive

days. Serum PK studies will be performed in 12 patients with blood draws performed on day 1

(single sample at baseline) and day 5 (serial samples). As this is the first PK study evaluating the

serum PK after intravaginal administration of artesunate, there are no directly comparable studies

to guide sample size determination. However, evaluating the PK of artesunate at the same dose

(200mg) in 12 participants is a relatively high number as most studies evaluate the PK of drugs

in n=3 or n=6 participants per dose.[54-56] In addition, other PK studies of oral or intramuscular

artesunate in this population have been performed with approximately 12 patients.[33, 57-59] This,

we expect our sample size of 12 to be sufficient to meet our primary objective. If the PK

variability of artesunate after intravaginal administration is greater than anticipated and cannot be

adequately evaluated in 12 participants, then the protocol can be amended to enroll an additional

n=3-6 participants.

Data analysis

The data analysis plan for this study involves assessing the primary outcome, which is the serum concentration versus time curve (AUC) of dihydroartemisinin following five days of self-administration of 200mg Artesunate pessaries. With the primary endpoint being Mean DHA AUC (ng*hr/ml) with standard deviation on Day 5, statistical analysis will compare this mean DHA AUC to historical studies after intravenous (IV), oral, and rectal administration among adults with similar dosing. Results will be analyzed using a student t-test for one-sample observation and a p-value of 0.05 will be considered statistically significant.

The secondary outcomes include assessing the AUC of Artesunate, maximum concentration of Artesunate and DHA (Cmax), time to reach maximum concentration (Tmax) for Artesunate and DHA, half-life (t1/2) for Artesunate and DHA, apparent clearance of Artesunate and DHA, and the volume of distribution of Artesunate and DHA. Statistical analysis of these variables will include comparing data results to historical studies using relevant test materials. The final outcome to be evaluated is the safety of the 5-day self-administration of Artesunate vaginal inserts. This will be done by monitoring and reporting adverse events and categorizing by their severity, with grade 3 or higher being considered severe. The proportion of participants with a severe AE will be reported along the exact (Clopper-Pearson) one-sided upper 95% confidence bounds. Safety among participants will be monitored and reported starting at the first dose of artesunate.

This study will determine conventional pharmacokinetic parameters for Artesunate and dihydroartemisinin (DHA). Parameters such as AUC, Cmax. Tmax, half-life, apparent clearance, and the volume of distribution will be calculated using non compartmental analysis from the serum concentration-time data. AUC for Artesunate will be determined by linear trapezoidal

summation with extrapolation to infinity, starting from drug administration to the last

observation. All parameters will be calculated using time in hours after the first drug

administration. With respect to DHA, AUC will be calculated to the last drug measurable time

point. The elimination rate constant (beta) will be calculated from the slope of the terminal phase

of the log concentration-time profile, and the elimination half-life (t1/2) calculated from the ratio

of ln 2/beta. Other PK parameters will be calculated using standard model-independent

formulae.[38] The estimates of PK parameters for DHA will assume complete conversion of AS to

DHA as reported previously.[35] Data will be plotted graphically and analyzed using Stata 18

software.[60] Data will be presented as mean with standard deviation (SD).

Quality Control Standards

The sample processing quality control standards are described in Table 3 below.

Table 3: Laboratory Quality Control Standards

Sections	Details
Quality standards	Criteria applied to samples & tests - Analytic methods are validated according to the latest Food and Drug Administration (FDA) bioanalytical method validation guidance[61] - Calibration curve with at least 6 non-zero standards, processed in duplicates per run - Calibration standards at lower and upper limits must be included - Each run includes a double blank, blank sample, and 3 quality control samples (low, medium, high), processed in triplicates
Acceptable limits	Acceptable ranges for tests[62] - For calibration curve use in quantification, at least 6 non-zero standard samples must be within +/- 15% nominal concentration, except lower limit of quantification (LLOQ), which must be within +/- 20% - Quality Control (QC) standards acceptance criteria: +/- 15% of nominal value, with at least 67% passing at low, medium, and high QC levels
Analysis methods	Assay details & limits of quantification for Artesunate and DHA[63] - Liquid chromatography-mass spectrometry (LC-MS/MS)

	method will be used for detection and quantification - Simultaneous measurement of artesunate and DHA from the same sample aliquot - LLOQ: 5 ng/ml for artesunate, 8 ng/ml for DHA
Quality control	Quality control (QC) procedures, acceptable ranges, and batch acceptance criteria - Each batch includes calibration standards and 3 QC samples meeting acceptance criteria outlined above - LLOQ must have a signal-to-noise ratio ≥5 - Low QC: 3xLLOQ - Medium QC: 50% of upper limit of quantification (ULOQ) - High QC: 80% ULOQ

Ethics approval and consent to participate

508 This clinical trial has full ethics review board approval from the University of North Carolina

509 Chapel Hill and the African Medical Research Foundation. Written informed consent will be

510 obtained from all study participants.

Discussion

Cervical cancer is preventable through vaccination against HPV, or screening for cervical

precancerous changes which can be treated. Access to cervical precancer treatment in low- and

middle-income countries is hindered by a shortage of trained healthcare provider and inadequate

health infrastructure. This results in a disproportionately high incidence and mortality rate from

this otherwise preventable disease. Use of self-administered topical therapies for cervical

precancer treatment, if found to be feasible and effective, can be transformative in increasing

access to secondary prevention of cervical cancer for marginalized women globally. Given

recent data demonstrating feasibility of topical Artesunate for treatment of HPV-associated

anogenital lesions, including vulvar and cervical precancer, it is imperative to understand the

pharmacokinetics of intravaginal use to inform studies using this drug in LMICs where malaria is

endemic.

Current Status:

The study opened for accrual in June 2024.

Supporting Information Caption:

S1 Figure. Schedule of Events.

Trial registration:

The trial is registered under U.S Clinical trial registry (clinicaltrials.gov, NCT06263582).

List of abbreviations

LMICs: Low- and middle-income countries:

AS: Artesunate

WHO: World Health Organization

CIN2/3:   High-grade cervical intraepithelial neoplasia

HPV: Human papillomavirus

DHA: Dihydroartemisinin

PK: Pharmacokinetics

IV: Intravenous

IM: Intramuscular

ACTs: Artemisinin-based combination therapies

Declarations:

Consent for publication

Not applicable.

Availability of data and materials

Not applicable.

Competing interests

"The authors declare they have no competing interests."

Funding

This research was supported by the Department of Obstetrics and Gynecology at University of

North Carolina-Chapel Hill, the Women's Reproductive Health Research (WRHR) Career

Development Program under award number 5-K12-HD103085-04 and the University of North

Carolina Center for AIDS Research under award number 5-P30-AI050410. The content is solely

the responsibility of the authors and does not necessarily represent the official views of the

National Institutes of Health. The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

Artesunate vaginal suppositories (pessaries) were provided as in-kind support by Frantz Viral

Therapeutics (Mentor, OH).

Authors' contributions

CM conceived and designed the study, providing subject matter expertise and overseeing all

aspects of protocol development. JO (Co-Principal Investigator) provided guidance on protocol

development and will lead protocol implementation in country. CO (Co-Investigator) contributed

to study design, protocol implementation, and capacity building for providers. KS contributed to

manuscript writing. GG contributes to lab management activities including collection, storage,

and shipment of samples. BM and CC contributed to in-country study coordination activities. LR

contributed to study methodology conceptualization, protocol development, and manuscript

writing. MP contributed to methodology conceptualization, protocol development, manuscript

writing, and implementation. WZ will contribute to formal analysis. All authors, in their

respective roles, contributed to study and manuscript preparation and have collectively approved

the final manuscript.

References

1. Trimble CL, Levinson K, Maldonado L, et al. A first-in-human proof-of-concept trial of
intravaginal artesunate to treat cervical intraepithelial neoplasia 2/3 (CIN2/3). *Gynecol*
Oncol. 2020;157(1):188-194. doi:10.1016/j.ygyno.2019.12.035
2. World Health Organisation. *World Health Organization. Guidelines for Treatment of*
Malaria. 3rd Edition.; 2015.
3. Bar-Zeev N, White N. Evidence behind the WHO guidelines: Hospital Care for Children:
efficacy and safety of artemisinin derivatives in children with malaria. *J Trop Pediatr*.
2006;52(2):78-82. doi:10.1093/tropej/fml005
4. Johann-Liang R, Albrecht R. Safety evaluations of drugs containing artemisinin
derivatives for the treatment of malaria. *Clin Infect Dis*. 2003;36(12):1626-1628.
doi:10.1086/375089
5. Nyaaba N, Andoh NE, Amoh G, et al. Comparative efficacy and safety of the artemisinin
derivatives compared to quinine for treating severe malaria in children and adults: A
systematic update of literature and network meta-analysis. *PLoS One*.
2022;17(7):e0269391. doi:10.1371/journal.pone.0269391
6. Ellis T, Eze E, Raimi-Abraham BT. Malaria and Cancer: a critical review on the
established associations and new perspectives. *Infect Agent Cancer*. 2021;16(1):33.
doi:10.1186/s13027-021-00370-7
7. Ma Z, Woon CYN, Liu CG, et al. Repurposing Artemisinin and its Derivatives as
Anticancer Drugs: A Chance or Challenge? *Front Pharmacol*. 2021;12:828856.
doi:10.3389/fphar.2021.828856
8. Jeong DE, Song HJJ, Lim S, et al. Repurposing the anti-malarial drug artesunate as a

novel therapeutic agent for metastatic renal cell carcinoma due to its attenuation of tumor
growth, metastasis, and angiogenesis. *Oncotarget*. 2015;6(32):33046-33064.
doi:10.18632/oncotarget.5422

9. Efferth T, Sauerbrey A, Olbrich A, et al. Molecular modes of action of artesunate in tumor cell lines. *Mol Pharmacol*. 2003;64(2):382-394. doi:10.1124/mol.64.2.382

10. Siegel EM, Patel N, Lu B, et al. Circulating biomarkers of iron storage and clearance of incident human papillomavirus infection. *Cancer Epidemiol Biomarkers Prev*. 2012;21(5):859-865. doi:10.1158/1055-9965.EPI-12-0073

11. Daniels TR, Bernabeu E, Rodríguez JA, et al. The transferrin receptor and the targeted delivery of therapeutic agents against cancer. *Biochim Biophys Acta*. 2012;1820(3):291-317. doi:10.1016/j.bbagen.2011.07.016

12. Schiffman M, Doorbar J, Wentzensen N, et al. Carcinogenic human papillomavirus infection. *Nat Rev Dis Primers*. 2016;2:16086. doi:10.1038/nrdp.2016.86

13. Wong YK, Xu C, Kalesh KA, et al. Artemisinin as an anticancer drug: Recent advances in target profiling and mechanisms of action. *Med Res Rev*. 2017;37(6):1492-1517. doi:10.1002/med.21446

14. Zhou X, Chen Y, Wang F, et al. Artesunate induces autophagy dependent apoptosis through upregulating ROS and activating AMPK-mTOR-ULK1 axis in human bladder cancer cells. *Chem Biol Interact*. 2020;331:109273. doi:10.1016/j.cbi.2020.109273

15. Zhang J, Zhou L, Xiang JD, Jin CS, Li MQ, He YY. Artesunate-induced ATG5-related autophagy enhances the cytotoxicity of NK92 cells on endometrial cancer cells via interactions between CD155 and CD226/TIGIT. *Int Immunopharmacol*. 2021;97:107705. doi:10.1016/j.intimp.2021.107705

16. Yao X, Zhao CR, Yin H, Wang K, Gao JJ. Synergistic antitumor activity of sorafenib and artesunate in hepatocellular carcinoma cells. *Acta Pharmacol Sin*. 2020;41(12):1609-1620. doi:10.1038/s41401-020-0395-5

17. Gao F, Sun Z, Kong F, Xiao J. Artemisinin-derived hybrids and their anticancer activity. *Eur J Med Chem*. 2020;188:112044. doi:10.1016/j.ejmech.2020.112044

18. Guan X, Guan Y. Artemisinin induces selective and potent anticancer effects in drug resistant breast cancer cells by inducing cellular apoptosis and autophagy and G2/M cell cycle arrest. *J BUON*. 2020;25(3):1330-1336.

19. Zhang B. Artemisinin-derived dimers as potential anticancer agents: Current developments, action mechanisms, and structure-activity relationships. *Arch Pharm (Weinheim)*. 2020;353(2):e1900240. doi:10.1002/ardp.201900240

20. Li D, Zhang J, Zhao X. Mechanisms and Molecular Targets of Artemisinin in Cancer Treatment. *Cancer Invest*. 2021;39(8):675-684. doi:10.1080/07357907.2021.1954190

21. Anfosso L, Efferth T, Albini A, Pfeffer U. Microarray expression profiles of angiogenesis-related genes predict tumor cell response to artemisinins. *Pharmacogenomics J*. 2006;6(4):269-278. doi:10.1038/sj.tpj.6500371

22. Disbrow GL, Baege AC, Kierpiec KA, et al. Dihydroartemisinin is cytotoxic to papillomavirus-expressing epithelial cells in vitro and in vivo. *Cancer Res*. 2005;65(23):10854-10861. doi:10.1158/0008-5472.CAN-05-1216

23. Morris CA, Duparc S, Borghini-Fuhrer I, Jung D, Shin CS, Fleckenstein L. Review of the clinical pharmacokinetics of artesunate and its active metabolite dihydroartemisinin following intravenous, intramuscular, oral or rectal administration. *Malar J*. 2011;10:263. doi:10.1186/1475-2875-10-263

24. Ilett KF, Batty KT, Powell SM, et al. The pharmacokinetic properties of intramuscular artesunate and rectal dihydroartemisinin in uncomplicated falciparum malaria. *Br J Clin Pharmacol*. 2002;53(1):23-30. doi:10.1046/j.0306-5251.2001.01519.x

25. Ilett KF, Batty KT, Powell SM, et al. The pharmacokinetic properties of intramuscular artesunate and rectal dihydroartemisinin in uncomplicated falciparum malaria. *Br J Clin Pharmacol*. 2002;53(1):23-30. doi:10.1046/j.0306-5251.2001.01519.x

26. KF I, KT B, SM P, et al. The pharmacokinetic properties of intramuscular artesunate and rectal dihydroartemisinin in uncomplicated falciparum malaria. *Br J Clin Pharmacol*. 2002;53(1):23-30. doi:10.1046/J.0306-5251.2001.01519.X

27. Morris CA, Duparc S, Borghini-Fuhrer I, Jung D, Shin CS, Fleckenstein L. Review of the clinical pharmacokinetics of artesunate and its active metabolite dihydroartemisinin following intravenous, intramuscular, oral or rectal administration. *Malar J*. 2011;10:263. doi:10.1186/1475-2875-10-263

28. Batty KT, Thu LT, Davis TM, et al. A pharmacokinetic and pharmacodynamic study of intravenous vs oral artesunate in uncomplicated falciparum malaria. *Br J Clin Pharmacol*. 1998;45(2):123-129. doi:10.1046/j.1365-2125.1998.00655.x

29. Hien TT, Davis TME, Chuong L V., et al. Comparative pharmacokinetics of intramuscular artesunate and artemether in patients with severe falciparum malaria. *Antimicrob Agents Chemother*. 2004;48(11):4234-4239. doi:10.1128/AAC.48.11.4234-4239.2004

30. Diem Thuy LT, Ngoc Hung L, Danh PT, Na-Bangchang K. Absence of time-dependent artesunate pharmacokinetics in healthy subjects during 5-day oral administration. *Eur J Clin Pharmacol*. 2008;64(10):993-998. doi:10.1007/s00228-008-0506-6

31. Diem Thuy LT, Ngoc Hung L, Danh PT, Na-Bangchang K. Absence of time-dependent artesunate pharmacokinetics in healthy subjects during 5-day oral administration. *Eur J Clin Pharmacol*. 2008;64(10):993-998. doi:10.1007/S00228-008-0506-6

32. Krudsood S, Looareesuwan S, Tangpukdee N, et al. New fixed-dose artesunate-mefloquine formulation against multidrug-resistant Plasmodium falciparum in adults: a comparative phase IIb safety and pharmacokinetic study with standard-dose nonfixed artesunate plus mefloquine. *Antimicrob Agents Chemother*. 2010;54(9):3730-3737. doi:10.1128/AAC.01187-09

33. Navaratnam V, Mansor SM, Mordi MN, Akbar A, Abdullah MN. Comparative pharmacokinetic study of oral and rectal formulations of artesunic acid in healthy volunteers. *Eur J Clin Pharmacol*. 1998;54(5):411-414. doi:10.1007/S002280050484

34. Sirivichayakul C, Sabchareon A, Pengsaa K, et al. Comparative study of the effectiveness and pharmacokinetics of two rectal artesunate/oral mefloquine combination regimens for the treatment of uncomplicated childhood falciparum malaria. *Ann Trop Paediatr*. 2007;27(1):17-24. doi:10.1179/146532807X170466

35. Halpaap B, Ndjave M, Paris M, Benakis A, Kremsner PG. Plasma levels of artesunate and dihydroartemisinin in children with Plasmodium falciparum malaria in Gabon after administration of 50-milligram artesunate suppositories. *Am J Trop Med Hyg*. 1998;58(3):365-368. doi:10.4269/ajtmh.1998.58.365

36. Teja-isavadharm P, Watt G, Eamsila C, et al. Comparative pharmacokinetics and effect kinetics of orally administered artesunate in healthy volunteers and patients with uncomplicated falciparum malaria. *Am J Trop Med Hyg*. 2001;65(6):717-721. doi:10.4269/AJTMH.2001.65.717

37. Binh TQ, Ilett KF, Batty KT, et al. Oral bioavailability of dihydroartemisinin in
Vietnamese volunteers and in patients with falciparum malaria. *Br J Clin Pharmacol*.
2001;51(6):541-546. doi:10.1046/J.1365-2125.2001.01395.X
38. Guidelines for the Treatment of Malaria - PubMed. Accessed October 22, 2023.
https://pubmed.ncbi.nlm.nih.gov/26020088/
39. Ward KE, Fidock DA, Bridgford JL. Plasmodium falciparum resistance to artemisinin-
based combination therapies. *Curr Opin Microbiol*. 2022;69:102193.
doi:10.1016/j.mib.2022.102193
40. Hanboonkunupakarn B, Tarning J, Pukrittayakamee S, Chotivanich K. Artemisinin
resistance and malaria elimination: Where are we now? *Front Pharmacol*. 2022;13.
doi:10.3389/FPHAR.2022.876282
41. Balikagala B, Fukuda N, Ikeda M, et al. Evidence of Artemisinin-Resistant Malaria in
Africa. *N Engl J Med*. 2021;385(13):1163-1171. doi:10.1056/NEJMoa2101746
42. Mathenge PG, Low SK, Vuong NL, et al. Efficacy and resistance of different
artemisinin-based combination therapies: a systematic review and network meta-analysis.
Parasitol Int. 2020;74. doi:10.1016/J.PARINT.2019.04.016
43. Tilley L, Straimer J, Gnädig NF, Ralph SA, Fidock DA. Artemisinin Action and
Resistance in Plasmodium falciparum. *Trends Parasitol*. 2016;32(9):682-696.
doi:10.1016/J.PT.2016.05.010
44. Siddiqui FA, Liang X, Cui L. Plasmodium falciparum resistance to ACTs: Emergence,
mechanisms, and outlook. *Int J Parasitol Drugs Drug Resist*. 2021;16:102-118.
doi:10.1016/J.IJPDDR.2021.05.007
45. Mok S, Stokes BH, Gnädig NF, et al. Artemisinin-resistant K13 mutations rewire
Plasmodium falciparum's intra-erythrocytic metabolic program to enhance survival. *Nat
Commun*. 2021;12(1). doi:10.1038/S41467-020-20805-W
46. Witkowski B, Amaratunga C, Khim N, et al. Novel phenotypic assays for the detection of
artemisinin-resistant Plasmodium falciparum malaria in Cambodia: in-vitro and ex-vivo
drug-response studies. *Lancet Infect Dis*. 2013;13(12):1043-1049. doi:10.1016/S1473-
3099(13)70252-4
47. Ward KE, Fidock DA, Bridgford JL. Plasmodium falciparum resistance to artemisinin-
based combination therapies. *Curr Opin Microbiol*. 2022;69.
doi:10.1016/J.MIB.2022.102193
48. Ndwiga L, Kimenyi KM, Wamae K, et al. A review of the frequencies of Plasmodium
falciparum Kelch 13 artemisinin resistance mutations in Africa. *Int J Parasitol Drugs
Drug Resist*. 2021;16:155-161. doi:10.1016/J.IJPDDR.2021.06.001
49. Moser KA, Madebe RA, Aydemir O, et al. Describing the current status of Plasmodium
falciparum population structure and drug resistance within mainland Tanzania using
molecular inversion probes. *Mol Ecol*. 2021;30(1):100-113. doi:10.1111/MEC.15706
50. Asua V, Conrad MD, Aydemir O, et al. Changing Prevalence of Potential Mediators of
Aminoquinoline, Antifolate, and Artemisinin Resistance Across Uganda. *J Infect Dis*.
2021;223(6):985-994. doi:10.1093/INFDIS/JIAA687
51. Squier CA, Mantz MJ, Schlievert PM, Davis CC. Porcine vagina ex vivo as a model for
studying permeability and pathogenesis in mucosa. *J Pharm Sci*. 2008;97(1):9-21.
doi:10.1002/jps.21077
52. Meshnick SR. Artemisinin: mechanisms of action, resistance and toxicity. *Int J Parasitol*.
2002;32(13):1655-1660. doi:10.1016/s0020-7519(02)00194-7

53. Fang SH, Plesa M, Carchman EH, Cowell NA, Staudt E, Twaroski KA, Buchwald UK, Trimble CL. A Phase I study of intra-anal artesunate (suppositories) to treat anal high0grade squamous intraepithelial lesions. *PLoS ONE*. 2023;18(12): e0295647. doi: https://doi.org/10.1371/journal.pone.0295647

54. L'homme RF, Dijkema T, Warris A, van der Ven AJ, Gibb DM, Burger DM. Pharmacokinetics of two generic fixed-dose combinations for HIV-infected children (Pedimune Baby & Pedimune Junior) are similar to the branded products in healthy adults. *J Antimicrob Chemother*. 2007 Jan;59(1):92-6. doi: 10.1093/jac/dkl445. Epub 2006 Oct 28. PMID: 17071953.

55. Hosseinipour MC, Corbett AH, Kanyama C, Mshali I, Phakati S, Rezk NL, van der Horst C, Kashuba AD. Pharmacokinetic comparison of generic and trade formulations of lamivudine, stavudine and nevirapine in HIV-infected Malawian adults. AIDS. 2007 Jan 2;21(1):59-64. doi: 10.1097/QAD.0b013e3280117ca0. PMID: 17148968.

56. Stuart GS, Moses A, Corbett A, Phiri G, Kumwenda W, Mkandawire N, Chintedze J, Malunga G, Hosseinipour M, Cohen MS, Stanczyk FZ, Kashuba AD. Combined oral contraceptives and antiretroviral PK/PD in Malawian women: pharmacokinetics and pharmacodynamics of a combined oral contraceptive and a generic combined formulation antiretroviral in Malawi. *J Acquir Immune Defic Syndr*. 2011 Oct 1;58(2):e40-3. doi: 10.1097/QAI.0b013e31822b8bf8. PMID: 21921726; PMCID: PMC3176587.

57. Awad MI, Eltayeb IB, Baraka OZ, Behrens RH, Alkadru AM. Pharmacokinetics of artesunate following oral and rectal administration in healthy Sudanese volunteers. *Trop Doct*. 2004 Jul;34(3):132-5. doi: 10.1177/004947550403400302. PMID: 15267037.

58. Byakika-Kibwika P, Lamorde M, Mayito J, Nabukeera L, Mayanja-Kizza H, Katabira E, Hanpithakpong W, Obua C, Pakker N, Lindegardh N, Tarning J, de Vries PJ, Merry C. Pharmacokinetics and pharmacodynamics of intravenous artesunate during severe malaria treatment in Ugandan adults. Malar J. 2012 Apr 27;11:132. doi: 10.1186/1475-2875-11-132. PMID: 22540954; PMCID: PMC3489518.

59. Ilett KF, Batty KT, Powell SM, Binh TQ, Thu le TA, Phuong HL, Hung NC, Davis TM. The pharmacokinetic properties of intramuscular artesunate and rectal dihydroartemisinin in uncomplicated falciparum malaria. *Br J Clin Pharmacol*. 2002 Jan;53(1):23-30. doi: 10.1046/j.0306-5251.2001.01519.x. PMID: 11849191; PMCID: PMC1874553.

60. StataCorp LLC: 2023. Stata 18. Statistical software. StataCorp LLC.

61. U.S. Department of Health and Human Services Food and Drug Administration, Center for Drug Evaluation and Research and Center for Biologicas Evaluation and Research. *M10 Bioanalytical Method Validation and Study Sample Analysis*. 2022. FDA-2019-D-1469. Accessed October 25, 2024. https://www.fda.gov/regulatory-information/search-fda-guidance-documents/m10-bioanalytical-method-validation-and-study-sample-analysis

62. Ongas M, Standing J, Ogutu B, Waichungo J, Berkley JA, Kipper K. Liquid chromatography–tandem mass spectrometry for the simultaneous quantitation of ceftriaxone, metronidazole and hydroxymetronidazole in plasma from seriously ill, severely malnourished children. *Wellcome Open Re*. 2018; 2(43). https://doi.org/10.12688/wellcomeopenres.11728.2

63. Ongas M, Juma E, Kirimi CG, Oloo F, Kokwaro G, Aman R, Ogutu BR. A selective LC-MS/MS method for simultaneous quantification of Artemether, Lumefantrine and their principle metabolites in human plasma. *ABC Research Alert*. 2018;6(3). doi:

789   https://doi.org/10.18034/ra.v6i3.335.
790
791
792
793