Longitudinal evidence for a mutually reinforcing relationship between white matter hyperintensities and cortical thickness in cognitively unimpaired older adults

Jose Bernal[1,2,3,4], Inga Menze[1,2], Renat Yakupov[1,2], Oliver Peters[5,6], Julian Hellmann-Regen[5,7,8], Silka Dawn Freiesleben[5,6], Josef Priller[4,5,9,10], Eike Jakob Spruth[5,9], Slawek Altenstein[5,9], Anja Schneider[11,12], Klaus Fliessbach[11,12], Jens Wiltfang[13,14,15], Björn H.Schott[13,14,16], Frank Jessen[11,17,18], Ayda Rostamzadeh[17], Wenzel Glanz[2], Enise I. Incesoy[1,2,19], Katharina Buerger[20,21], Daniel Janowitz[21], Michael Ewers[20,21], Robert Perneczky[20,22,23,24], Boris-Stephan Rauchmann[22,25,26], Stefan Teipel[27,28], Ingo Kilimann[27,28], Christoph Laske[29,30], Sebastian Sodenkamp[29,31], Annika Spottke[11,32], Anna Esser[11], Falk Lüsebrink[2], Peter Dechent[33], Stefan Hetzer[34], Klaus Scheffler[35], Stefanie Schreiber[2,36], Emrah Düzel[1,2,T], Gabriel Ziegler[1,2,T]

Author affiliations:

[1]Institute of Cognitive Neurology and Dementia Research, Otto-von-Guericke University Magdeburg, Magdeburg, Germany
[2]German Centre for Neurodegenerative Diseases (DZNE), Magdeburg, Germany
[3]Centre for Clinical Brain Sciences, the University of Edinburgh, UK
[4]UK Dementia Research Institute Centre at the University of Edinburgh, UK
[5]German Centre for Neurodegenerative Diseases (DZNE), Berlin, Germany
[6]Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin-Institute of Psychiatry and Psychotherapy
[7]Charité – Universitätsmedizin Berlin, Department of Psychiatry and Neurosciences, Campus Benjamin Franklin"
[8]German Centre for Mental Health (DZPG), partner site Berlin
[9]Department of Psychiatry and Psychotherapy, Charité, Charitéplatz 1, 10117 Berlin, Germany
[10]School of Medicine, Technical University of Munich; Department of Psychiatry and Psychotherapy, Munich, Germany
[11]German Centre for Neurodegenerative Diseases (DZNE), Bonn, Venusberg-Campus 1, 53127 Bonn, Germany
[12]Department of Old Age Psychiatry and Cognitive Disorders, University Hospital Bonn and University of Bonn, Bonn, Germany
[13]German Centre for Neurodegenerative Diseases (DZNE), Göttingen, Germany
[14]Department of Psychiatry and Psychotherapy, University Medical Centre Göttingen, University of Göttingen, Von-Siebold-Str. 5, 37075 Göttingen
[15]Neurosciences and Signalling Group, Institute of Biomedicine (iBiMED), Department of Medical Sciences, University of Aveiro, Aveiro, Portugal

[16]Leibniz Institute for Neurobiology, Brenneckestr. 6, 39118, Magdeburg, Germany.

[17]Department of Psychiatry, University of Cologne, Medical Faculty, Kerpener Strasse 62, 50924 Cologne,
Germany

[18]Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD), University of
Cologne, Joseph-Stelzmann-Strasse 26, 50931 Köln, Germany

[19]Department for Psychiatry and Psychotherapy, University Clinic Magdeburg, Magdeburg, Germany

[20]German Centre for Neurodegenerative Diseases (DZNE), Feodor-Lynen-Strasse 17, 81377 Munich, Germany

[21]Institute for Stroke and Dementia Research (ISD), University Hospital, LMU Munich, Feodor-Lynen-Strasse 17,
81377 Munich, Germany

[22]Department of Psychiatry and Psychotherapy, University Hospital, LMU Munich, Munich, Germany

[23]Munich Cluster for Systems Neurology (SyNergy) Munich, Munich, Germany

[24]Ageing Epidemiology Research Unit (AGE), School of Public Health, Imperial College London, London, UK

[25]Sheffield Institute for Translational Neuroscience (SITraN), University of Sheffield, Sheffield, UK

[26]Department of Neuroradiology, University Hospital LMU, Munich, Germany

[27]German Centre for Neurodegenerative Diseases (DZNE), Rostock, Germany

[28]Department of Psychosomatic Medicine, Rostock University Medical Center, Gehlsheimer Str. 20, 18147
Rostock

[29]German Centre for Neurodegenerative Diseases (DZNE), Tübingen, Germany

[30]Section for Dementia Research, Hertie Institute for Clinical Brain Research and Department of Psychiatry and
Psychotherapy, University of Tübingen, Tübingen, Germany

[31]Department of Psychiatry and Psychotherapy, University of Tübingen, Tübingen, Germany

[32]Department of Neurology, University of Bonn, Venusberg-Campus 1, 53127 Bonn, Germany

[33]MR-Research in Neurosciences, Department of Cognitive Neurology, Georg-August-University Goettingen,
Germany

[34]Berlin Centre for Advanced Neuroimaging, Charité – Universitätsmedizin Berlin, Berlin, Germany

[35]Department for Biomedical Magnetic Resonance, University of Tübingen, 72076 Tübingen, Germany

[36]Department of Neurology, University Hospital Magdeburg, Magdeburg, Germany

[T]Shared last authorship
65
66
67 Correspondence to: Jose Bernal

68 Full address: German Centre for Neurodegenerative Diseases (DZNE), Magdeburg,

69 Leipziger Str. 44, 39120 Magdeburg, Germany

70 E-Mail: jose.bernalmoyano@dzne.de

Abbreviations

AD: Alzheimer's Disease

BLGCM: Bivariate Latent Growth Curve Model

CERAD: Consortium to Establish a Registry for AD

CFI: Comparative Fit Index

CI: Confidence Interval

CSF: Cerebrospinal Fluid

CSVD: Cerebral Small Vessel Disease

DELCODE: DZNE Longitudinal Cognitive Impairment and Dementia Study

DZNE: German Centre for Neurodegenerative Diseases

FDR: False Discovery Rate

HCP: Human Connectome Project

LGCM: Latent Growth Curve Model

MRI: Magnetic Resonance Imaging

MS: Multiple Sclerosis

RMSEA: Root Mean Square Error of Approximation

SE: Standard Error

SEM: Structural Equation Model

SRMR: Standardised Root Mean Residual

TICV: Total Intracranial Volume

WMH: White Matter Hyperintensities

Abstract

Background

For over three decades, the concomitance of cortical neurodegeneration and white matter hyperintensities (WMH) has sparked discussions about their coupled temporal dynamics. Longitudinal studies supporting this hypothesis remain nonetheless scarce.

Methods

In this study, we applied regional and global bivariate latent growth curve modelling (BLGCM) to longitudinal data from 436 cognitively unimpaired participants (DELCODE cohort; median age 69.70 [IQR 65.44, 74.49] years; 52.98% female) to examine the extent to which WMH and cortical thickness were interrelated over a four-year period.

Results

Our findings were three-fold. First, at baseline, individuals with larger WMH volumes had lower mean cortical thicknesses over the entire brain. Second, individuals who experienced a steeper thinning of their cingulate and temporal cortices over time had larger baseline WMH volumes in the frontal, parietal, and occipital lobes. Third, individuals with thinner cortices at baseline tended to undergo faster WMH progression over four years, particularly in the occipital and parietal lobes.

Conclusions

Our study suggests that cortical thinning and WMH progression could be mutually reinforcing rather than parallel, unrelated processes, which become entangled before cognitive deficits are detectable.

117 **Trial Registration**

118 German Clinical Trials Register (DRKS00007966, 04/05/2015)

120 **Keywords:** White Matter Hyperintensities; Cortical Thickness; Latent Growth Curve
121 Model; Longitudinal Modelling; Structural Magnetic Resonance Imaging

Introduction

Cortical thinning and white matter hyperintensities (WMH) progression are well-known ageing processes that take place throughout middle and late adulthood [1–9]. Both processes appear to be influenced by genetic and lifestyle factors [2,10–15] as well as by the onset and progression of neurodegenerative and cerebrovascular diseases [1,2,9,16–20]. Although overlapping risk factors may offer an initial explanation for their concomitance [3,6,11,21,22], their persistent association after controlling for demographics and traditional cardiovascular risk factors [3,6,10,23–25] has sparked more than three decades of research into coupled temporal dynamics[3,26].

Coupled temporal dynamics between WMH and cortical atrophy are currently discussed from two non-exclusive perspectives: the cerebrovascular and the neurodegenerative hypotheses [17,26]. The cerebrovascular hypothesis posits that ischaemic and hypoxic damages—operationalised as WMH [15,27–29]—may initially result in the depletion of oxygen, nutrients, and trophic support in perilesional regions[16,28]. Subsequently, these damages may also disrupt the function and metabolic demands of compromised white matter tracts and associated cortical regions, leading to cortical atrophy [6,9,17,27,30]. On the other hand, the neurodegenerative hypothesis proposes that cortical neurodegeneration could contribute to WMH formation [17,26,29,31–34], especially in conjunction with tau pathologies [26,29,34]. Excessive tau phosphorylation could promote microtubule destabilisation, thereby causing axonal transport dysfunction, energy depletion, and calcium imbalance—a hallmark of Wallerian degeneration [34]. In the light of the posterior dominance of WMH in Alzheimer's disease (AD) [26,35–38], both

hypotheses would require effects of cortical neurodegeneration and WMH to be
particularly pronounced in parietal and occipital brain regions. Longitudinal evidence
and multivariate modelling substantiating these hypotheses remain nonetheless
scarce, especially in cognitively unimpaired older adults[1].

In this study, we leveraged bivariate latent growth curve modelling (BLGCM) to
examine the bidirectional relationship between lobar WMH and regional cortical
thickness over four years in older individuals without objective cognitive impairment.

Methods

Study participants

We used baseline and annual follow-up data for up to 48 months from participants of
the observational longitudinal multicentre DELCODE (DZNE Longitudinal Cognitive
Impairment and Dementia) Study [39]—a memory-clinic-based observational
multicentre study from the German Centre for Neurodegenerative Diseases (DZNE)
that uses multimodal assessment of preclinical, prodromal, and clinical stages of AD,
with a particular focus on subjective cognitive decline. In the present work, we
focused on cognitively unimpaired participants who underwent at least three MRI
scanning sessions and whose follow-up MRI sessions took place within four months
prior or after their yearly comprehensive examination.

At baseline, participants underwent a thorough evaluation at their local study site,
which included medical history checks, a psychiatric and neurological examination,
neuropsychological testing, blood and cerebrospinal fluid (CSF) collection, and MRI
in accordance with local standards. All DELCODE sites used the Consortium to

Establish a Registry for AD (CERAD-plus) neuropsychological test battery to assess

cognitive function. To be classified as cognitively unimpaired, participants were

defined by performing within at least required to performed better than -1.5 standard

deviations of the age-, sex-, and education-adjusted normal performance on all

subtests of the test battery [39].

Additional inclusion criteria were age $\geq$ 60 years, fluent German language skills,

capacity to provide informed consent, and the the availability of a study partner. The

main exclusion criteria for all groups were conditions clearly interfering with

participation in the study or the study procedures, including significant sensory

impairment. The following medical conditions were considered exclusion criteria:

current or history of major depressive episode and major psychiatric disorders either

at baseline (e.g., psychotic disorder, bipolar disorder, substance abuse),

neurodegenerative diseases other than AD, vascular dementia, history of stroke with

residual clinical symptoms, history of disseminated malignant disease, severe or

unstable medical conditions, and clinically significant vitamin B12 deficiency at

baseline. Prohibited drugs included chronic use of psychoactive compounds with

sedative or anticholinergic effects, use of anti-dementia agents, and investigational

drugs for the treatment of dementia or cognitive impairment one month before study

entry and throughout the duration of the study.

All participants provided their written informed consent in accordance with the

Declaration of Helsinki at baseline. DELCODE has been registered within the

German Clinical Trials Register (DRKS00007966, 04/05/2015). Ethics committees of

the medical faculties of all participating sites (i.e., Berlin (Charité -

Universitätsmedizin Berlin), Bonn, Cologne, Göttingen, Magdeburg, Munich (Ludwig-

Maximilians-University), Rostock, and Tübingen) approved the DELCODE study

protocol before inclusion of the first participants. The ethics committee of the medical

faculty of the University of Bonn led and coordinated the process.

Total cardiovascular risk score

We established a total cardiovascular risk score for each participant by tallying their

dichotomised (y/n) history of smoking, presence of obesity, hyperlipidemia, arterial

hypertension, and diabetes, as reported in their medical records. We corrected the

sum of present risk factors by the amount of available information. For example, if an

individual had a history of arterial hypertension and diabetes but we did not have

data on smoking, obesity, or hyperlipidemia, the final score would be 1.00. The

corrected total cardiovascular risk scores ranged from 0.0 to 1.0, where the lowest

and highest values denoted the absence or presence of all available risk factors,

respectively.

MRI

MRI acquisition took place at nine DZNE sites or associated university medical

centers equipped with 3T Siemens MR scanners. In the present study, we leveraged

the following structural sequences: T1w MPRAGE (full head coverage, 3D

acquisition, GRAPPA factor 2, 1 mm^3 isotropic, 256 × 256 px, 192 sagittal slices,

TR/TE/TI 2500/4.33/1100 ms, FA 7°) and T2w FLAIR (full head coverage, 3D

acquisition, 1 mm^3 isotropic, 256 × 256 px, 192 sagittal slices, TR/TE/TI

5000/394/1800 ms). The DZNE imaging network oversaw operating procedures, as

well as quality assurance and assessment (iNET, Magdeburg) [39].

MRI-based measurements

Cortical thickness

We used the CAT12 longitudinal pipeline[40] (neuro-jena.github.io) to reconstruct cortical thickness surfaces for each subject and for each time point (ageing workflow; default parameters, except for final resolution, which we set to 1 mm^3). We smoothed all surfaces with a 12-mm Gaussian filter and resampled them to the 32k HCP surface template. We also used CAT12 to estimate mean thickness throughout the whole brain cortex and within cortical regions as per the Desikan-Killiany cortical parcellation atlas [41].

WMH segmentation

We manually segmented WMH using the AI-augmented version of the Lesion Segmentation Toolbox (LST-AI) [42–44] and based the segmentation on both T1w MPRAGE and T2w FLAIR imaging data. We then tallied WMH volumes across the frontal, temporal, parietal and occipital lobes using the UCSLobes Atlas [45].

Statistical analyses

We conducted all data analyses in RStudio (v1.3.1073; R v4.0.2) using lavaan (v0.6-16). We created figures using ggplot2 (v3.4.3) and the ENIGMA toolbox [46].

We carried out univariate and bivariate LGCM analyses. (B)LGCMs [47] are a powerful class of structural equation models (SEM) to describe sample average trajectories of one or two constructs over time through the specification of latent intercepts and latent slopes (i.e, initial levels and rates of change). First, we used univariate analyses for contextualisation to examine what covariates were

associated with the baseline measurements, as well as with potential changes over

repeated measures. We then focused on bivariate models to examine

interrelationships between WMH and cortical thickness over time (Figure 1). Intra-

domain and cross-domain relationships can be assessed via the covariance between

these four latent growth parameters [48]. We specifically tested the evidence of four

possible major cross-domain relationships:

1. *intercept-intercept covariance* $\sigma_{\iota_{WMH} \leftrightarrows \iota_{Thick}}$: upon study entry, do individuals

with larger WMH volumes have lower cortical thickness?,

2. *intercept-slope covariance* $\sigma_{\iota_{WMH} \leftrightarrows \Delta_{Thick}}$: do individuals with larger WMH

volumes at study entry experience faster cortical thinning [cerebrovascular

hypothesis]?,

3. *intercept-slope covariance* $\sigma_{\iota_{Thick} \leftrightarrows \Delta_{WMH}}$: do individuals with thinner cortices at

study entry exhibit a faster increase in WMH volumes [neurodegenerative

hypothesis]?, and

4. *slope-slope covariance* $\sigma_{\Delta_{Thick} \leftrightarrows \Delta_{WMH}}$: do individuals exhibiting faster WMH

volume increases also undergo faster cortical thinning over time?

We also studied within-domain *intercept-slope covariances* ($\sigma_{\iota_{WMH} \leftrightarrows \Delta_{WMH}}$ and

$\sigma_{\iota_{Thick} \leftrightarrows \Delta_{Thick}}$) to determine whether baseline levels were associated with ongoing

changes over time.

We conducted global and regional analyses to identify associations at two levels of

granularity. In the global analysis—with no spatial specificity—we focused on the

interrelationship between mean cortical thickness and total WMH volume. In order to

elucidate potential region-specific and cross-domain relationships, we additionally

examined all possible pairs of lobar WMH volumes and regional cortical thicknesses

within Desikan-Killiany atlas regions. Note that our approach is similar to a mass-

univariate analysis scheme, with the difference being that we investigate region-

specific effects through LGCM rather than through GLM. To reduce the

dimensionality and thereby improve the feasibility of our multivariate SEM analysis,

we considered (corresponding) bilateral regions jointly. We used the superscripts

[Global] and [Regional] to indicate the type of analysis utilised to generate the reported

values. We report the completely standardised solutions and provide unstandardised

solutions as Supplementary Material.

Figure 1. BLGCM to probe the coupling of cortical thickness and WMH over repeated measures. Illustration of the longitudinal structural equation modeling (SEM) model used on global regional/lobar level. We employed the conventional notation with squared variables indicating observed and measured variables (manifest variables) and circular ones referring to latent (unobserved) variables. Single-headed solid arrows illustrate a modelled relationship between two variables, with the arrow pointing towards the dependent variable. Single-headed dashed arrows signify a relationship between two variables, where the weight is fixed. Double-headed arrows represent the covariance (hyperparameter) between two variables. Grey triangles represent latent intercept estimates. We further adjusted latent intercepts and slopes for age, sex, years of education, total cardiovascular risk factors, and TICV. We omitted these paths for visualisation purposes.

Adjusting for covariates and confounders

We adjusted latent intercepts and slopes for effects of age, sex, years of education, total cardiovascular risk factor score, and total intracranial volume (TICV) in all models.

Data transformation

We applied a Box-Cox transform to WMH volumes to account for potential skewness and z-scored all variables (pooled across timepoints) prior to model fitting. For the purpose of contextualisation and plotting, we back-transformed the fitted growth curve parameters afterwards.

Model fitting

We employed the maximum likelihood robust estimator to fit the model. We used the full information maximum likelihood estimation to handle missing values. To check for compliance with the assumption of missingness at random, we tested whether missingness in one column (1: missing; 0: not missing) could be predicted from the remaining ones. In all instances, the resulting p-values exceeded 0.05.

Prior to model fitting and solely to ensure model fit, we used Tukey's fences to identify and remove outliers in all data points (threshold of 1.5) [49]. We evaluated

the fit of all global and regional models by analysing their root mean square error of

approximation (RMSEA; values ≤ 0.05 indicate good fit), comparative fit index (CFI;

values exceeding 0.95 indicate good fit), and standardised root mean residual

(SRMR; values < 0.08 suggest good fit) [50]. For the sake of transparency, when

discussing the models, we disclosed their convergence and compliance with the

aforementioned thresholds.

Correction for multiple comparisons

306   We employed the False Discovery Rate (FDR) correction [51] method to account for

307   the issue of multiple comparisons on all region-wise analyses (Figure S1 contains

308   the uncorrected version).

Results

Study participants

311   We included 436 cognitively unimpaired DELCODE participants with imaging data

312   available for at least three visits (1834 MRI sessions; median age 69.70 [IQR 65.44,

313   74.49] years; 52.98% females; median years of education 14 [IQR 13, 17]).

Univariate findings

WMH volumes increased over the course of four years

Model fit

317   All univariate LGCM on total and lobar WMH volumes converged and provided good

318   model fit ($RMSEA \leq 0.05$, $CFI \geq 0.095$, $SRMR \leq 0.05$).

Demographic effects on latent intercept (ι_{WMH})

Baseline total WMH volumes varied significantly among individuals (variance of ι_{WMH}^{Global} = 0.881, SE = 0.033, Z = 25.117, p-value < 0.001).

Total WMH volumes were larger in older individuals ($\beta_{Age \to \iota_{WMH}}^{Global}$ = 0.365, SE = 0.043, Z = 8.462, p-value < 0.001) and in those with higher total cardiovascular risk factor scores ($\beta_{Vascular\ risk \to \iota_{WMH}}^{Global}$ = 0.090, SE = 0.045, Z = 1.978, p-value = 0.048). Moreover, WMH across the temporal lobes tended to be higher in those with fewer years of education (Figure 2; $\beta_{Education \to \iota_{WMH}}^{Regional}$ = -0.099, SE = 0.058, Z = -1.715, p-value = 0.086).

Females had larger total WMH volumes than males ($\beta_{Female \to \iota_{WMH}}^{Global}$ = 0.175, SE = 0.062, Z = 2.830, p-value = 0.005), especially across frontal brain regions ($\beta_{Female \to \iota_{WMH}}^{Regional}$ = 0.184, SE = 0.064, Z = 2.955, p-value = 0.003), despite females in our sample being on average younger ($\sigma_{Age \leftrightarrows Female}^{Global}$ = -0.168, SE = 0.047, Z = -3.598, p-value < 0.001) and having lower total cardiovascular risk scores ($\sigma_{Vascular\ risk \leftrightarrows Female}^{Global}$ = -0.195, SE = 0.045, Z = -4.305, p-value < 0.001) than males. In addition, females had on average fewer years of education ($\sigma_{Education \leftrightarrows Female}^{Global}$ = -0.233, SE = 0.044, Z = -5.346, p-value < 0.001) than males.

Demographic effects on latent slope (Δ_{WMH})

Across participants, total WMH volumes increased over the follow-up period of four years (Figure 2A,B; intercept of Δ_{WMH}^{Global} = 1.066, SE = 0.101, Z = 10.573, p-value < 0.001). The frontal and parietal lobes underwent the most substantial progression in WMH volumes, with an average increase of 0.180 [95%-CI 0.153, 0.207] and 0.175

[95%-CI 0.137, 0.212] ml per year, respectively. Significant WMH progression

occured in occipital and temporal lobes as well (approximated average rates of

change: 0.034 [95%-CI 0.027, 0.042], 0.018 [IQR 0.015, 0.022] ml per year,

respectively).

A strength of the approach is the quantification of WMH progression rate variability

across individuals (Figure 2B; variance of $\Delta_{\text{WMH}}^{\text{Global}}$ = 0.991, SE = 0.012, Z = 80.306,

p-value < 0.001), which we found to be tied to baseline demographics and pre-

existing WMH volumes. WMH progression rates in parietal regions were lower in

individuals with advanced age ($\beta_{\text{Age}\to\Delta_{WMH}}^{\text{Regional}}$ = -0.233, SE = 0.068, Z = -3.431, p-value =

0.001) and with more years of education (Figure 2C; $\beta_{\text{Education}\to\Delta_{WMH}}^{\text{Regional}}$ = -0.166, SE =

0.066, Z = -2.506, p-value = 0.036). Similarly, those with the highest initial regional

WMH volumes tended to exhibit the least progression of WMH in the same region

over time (parietal: $\beta_{\iota WMH\to\Delta_{WMH}}^{\text{Regional}}$ = -0.305, SE = 0.066, Z = -4.654, p-value < 0.001;

occipital: ($\beta_{\iota WMH\to\Delta_{WMH}}^{\text{Regional}}$ = -0.159, SE = 0.078, Z = -2.032, p-value = 0.042; temporal:

$\beta_{\iota WMH\to\Delta_{WMH}}^{\text{Regional}}$ = -0.247, SE = 0.125, Z = -1.982, p-value = 0.048), except in the frontal

lobe ($\beta_{\iota WMH\to\Delta_{WMH}}^{\text{Regional}}$ = -0.102, SE = 0.080, Z = -1.275, p-value = 0.202). Sex differences

in WMH progression rates were evident across the occipital lobe, with females

showing faster occipital WMH progression than males (Figure 2C; $\beta_{\text{Female}\to\Delta_{WMH}}^{\text{Regional}}$ =

0.221, SE = 0.088, Z = 2.510, p-value = 0.012).

360 **Cortical thickness decreased over the course of four years**

361 **Model fit**

362 All univariate LGCM fitted to cortical thickness converged and had good fit indices

363 ($RMSEA \leq 0.05$, $CFI \geq 0.095$, $SRMR \leq 0.05$).

364 **Demographic effects on latent intercept (ι_{Thick})**

365 Cortical thickness values were, on average, lower in individuals with advanced age

366 ($\beta_{\text{Age} \to \iota_{\text{Thick}}}^{\text{Global}}$ = -0.190, SE = 0.024, Z = -8.077, $p\text{-value}$ < 0.001), especially within

367 cortical regions associated with processing sensory information and orchestrating

368 motor functions (peaks - postcentral: $\beta_{\text{Age} \to \iota_{\text{Thick}}}^{\text{Regional}}$ = -0.342, SE = 0.045, Z = -7.666,

369 $p\text{-value}$ < 0.001; superiortemporal: $\beta_{\text{Age} \to \iota_{\text{Thick}}}^{\text{Regional}}$ = -0.352, SE = 0.044, Z = -7.937,

370 $p\text{-value}$ < 0.001; superiorfrontal: $\beta_{\text{Age} \to \iota_{\text{Thick}}}^{\text{Regional}}$ = -0.309, SE = 0.044, Z = -7.035, $p\text{-value}$ <

371 0.001). We did not observe sufficient evidence relating sex or cardiovascular risk

372 factors to initial cortical measurements in the cognitively unimpaired cohort

373 investigated here.

374 **Demographic effects on latent slope (Δ_{Thick})**

375 The thickness of cerebral cortex generally decreased over the course of four years

376 (Figure 2D,E; intercept of $\Delta_{\text{Thick}}^{\text{Global}}$ = -0.269, SE = 0.082, Z = -3.263, $p\text{-value}$ = 0.001).

377 Global thinning rates varied substantially among individuals (Figure 2E; variance

378 $\Delta_{\text{Thick}}^{\text{Global}}$ = 0.958, SE = 0.035, Z = 27.413, $p\text{-value}$ < 0.001). Thinning was particularly

379 pronounced across the cingulate and temporal cortex (Figure 2D, F; intercept of

380 $\Delta_{\text{Thick}}^{\text{Global}}$ peaked in the caudal anterior cingulate cortex = -0.875, SE = 0.107, Z = -

381 8.196, $p\text{-value}$ < 0.001). Caudal anterior and posterior cingulate cortices, for

instance, underwent the most cortical thinning over the course of four years, with an

average decrease in thickness of -0.014 [IQR -0.016, -0.013] and -0.011 [95%-CI -

0.012, -0.009] mm/year, respectively. Annual cortical thinning rates tended to

decelerate with advance age ($\beta_{\text{Age}\rightarrow\Delta_{\text{Thick}}}^{\text{Global}}$ = -0.147, SE = 0.089, Z = -1.652, $p\text{-value}$ =

0.098). Neither global or regional analyses hinted at an association between sex,

years of education, cardiovascular risk score, and changes in cortical thickness in

our cohort.

Figure 2. Changes in WMH volumes and cortical thickness over four years. We obtained latent
intercepts and slopes for each individual through the application of univariate LGCM to WMH volumes
and cortical thickness (separate models for each neuroimaging feature). We used them to compute
latent growth curve parameters and predict individual trajectories, corrected for age, sex, years of
education, total cardiovascular risk scores, and TICV. Prior to plotting and to enhance interpretability,
we back-transformed all predicted measurements. (A) Total WMH volume trajectories, as predicted by
the model. Blue lines represent the predicted trajectories and the pink one the average one. (B) Back-
transformed individual factor scores of latent slopes for WMH ($\Delta_{\text{WMH}}^{\text{Regional}}$), summarised in the density
plots, indicate that WMH volumes generally increased over time. We adjusted density plots such that
the modes attain the highest value, irrespective of the actual frequency. The rate of change varied
substantially across individuals in both cases. WMH progression, from fastest to slowest, occurred in
the frontal, parietal, occipital, and temporal lobes (approximated average rates of change: 0.180
[95%-CI 0.153, 0.207], 0.175 [95%-CI 0.137, 0.212], 0.034 [95%-CI 0.027, 0.042], 0.018 [IQR 0.015,
0.022] ml / year, respectively). Interestingly, while progression was predominant, a few subjects
showed a clear and consistent decrease in global WMH volumes over the course of the study,
especially across the occipital lobes. Visual assessment revealed that enlargement of lateral
ventricles contributed to such volumetric loss. (C) Annual change in parietal and occipital WMH in

relation to education and sex, respectively. (D) Regions exhibiting substantial cortical thinning over
the course of the study. Both the cingulate and lateral temporal cortex underwent the most thinning
(intercept of $\Delta_{\text{Thick}}^{\text{Global}}$ peaked in the caudal anterior cingulate cortex = -0.875, SE = 0.107, Z = -8.196,
p-value < 0.001). (E) Back-transformed individual factor scores of cortical thicknesses ($\Delta_{\text{Thick}}^{\text{Regional}}$)
across regions experiencing the most pronounced thinning over time. Caudal anterior and posterior
cingulate cortices underwent the most decline over the course of four years, with an average
decrease in thickness of -0.014 [IQR -0.016, -0.013] and -0.011 [95%-CI -0.012, -0.009] mm/year,
respectively. The variability in change rates indicated significant inter-individual differences in regional
cortical thinning. (F) Annual change in thickness across the isthmus cingulate and parahippocampal
cortex in relation to age.

Bivariate findings

Model fit

All BLGCMs also converged and had a satisfactory model fit ($RMSEA$ ≤ 0.05, CFI ≥ 0.095, $SRMR$ ≤ 0.05).

Individuals with larger WMH volumes have lower cortical thickness ($\sigma_{\iota_{WMH} \leftrightarrows \iota_{Thick}}$)

At baseline, individuals with larger WMH volumes showed lower mean cortical thickness ($\sigma_{\iota_{WMH} \leftrightarrows \iota_{Thick}}^{\text{Global}}$ = -0.160, SE = 0.049, Z = -3.285, p-value = 0.001). Subsequent analysis of regional measurements revealed that this association was more pronounced within the same lobe (Figure 3A): frontal WMH related more strongly to the frontal cortex (peak at parstriangularis: $\sigma_{\iota_{WMH} \leftrightarrows \iota_{Thick}}^{\text{Regional}}$ = -0.177, SE = 0.050, Z = -3.564, p-value < 0.001), occipital WMH to the occipital cortex (peak at cuneal cortex: $\sigma_{\iota_{WMH} \leftrightarrows \iota_{Thick}}^{\text{Regional}}$ = -0.162, SE = 0.055, Z = -2.957, p-value = 0.003), and parietal WMH to the parietal cortex (peak at postcentral cortex: $\sigma_{\iota_{WMH} \leftrightarrows \iota_{Thick}}^{\text{Regional}}$ = -0.129, SE = 0.057, Z = -2.274, p-value = 0.023). However, we also observed associations across distal regions (Figure 3A), for instance, baseline frontal WMH volumes and

cortical thickness in temporal brain regions, occipital WMH volumes and frontal

cortical thickness.

Individuals with larger WMH volumes show faster cortical thinning

$(\sigma_{\iota_{WMH}\leftrightarrows\Delta_{Thick}})$

Although the association between WMH volumes at baseline and cortical thinning

rates was not observed at a global scale ($\sigma_{\iota_{WMH}\leftrightarrows\Delta_{Thick}}^{\text{Global}}$ = -0.121, SE = 0.076, Z = -

1.595, p-value = 0.111), it was at a regional scale (Figure 3B). Individuals who

exhibited a steeper decrease in thickness of the cingulate cortices showed larger

WMH volumes across the frontal (Figure 4A.1; peak at caudal anterior:

$\sigma_{\iota_{WMH}\leftrightarrows\Delta_{Thick}}^{\text{Regional}}$ = -0.244, SE = 0.079, Z = -3.091, p-value = 0.002), parietal (peak at

caudal anterior: $\sigma_{\iota_{WMH}\leftrightarrows\Delta_{Thick}}^{\text{Regional}}$ = -0.260, SE = 0.079, Z = -3.302, p-value = 0.001), and

occipital lobes (peak at caudal anterior: $\sigma_{\iota_{WMH}\leftrightarrows\Delta_{Thick}}^{\text{Regional}}$ = -0.260, SE = 0.079, Z = -

3.302, p-value = 0.001). Individuals with larger occipital WMH at baseline also

showed a steeper decline in thickness in adjacent cortical regions, such as the

inferior parietal cortex (Figure 3B; $\sigma_{\iota_{WMH}\leftrightarrows\Delta_{Thick}}^{\text{Regional}}$ = -0.247, SE = 0.094, Z = -2.611,

p-value = 0.009), but also in more remote cortical areas like the superior temporal

cortex ($\sigma_{\iota_{WMH}\leftrightarrows\Delta_{Thick}}^{\text{Regional}}$ = -0.316, SE = 0.084, Z = -3.749, p-value < 0.001) and the

insular cortex ($\sigma_{\iota_{WMH}\leftrightarrows\Delta_{Thick}}^{\text{Regional}}$ = -0.312, SE = 0.091, Z = -3.425, p-value = 0.001).

Individuals with thinner cortices exhibit faster increases in WMH volumes ($\sigma_{\iota_{Thick} \leftrightarrows \Delta_{WMH}}$)

Individuals with thinner cortices at baseline tended to undergo faster WMH progression over the span of four years ($\sigma_{\iota_{Thick} \leftrightarrows \Delta_{WMH}}^{\text{Whole-brain}}$ = -0.123, SE = 0.063, Z = -1.964, p-value = 0.049). Upon closer examination of these associations, we found evidence for regional specificities (Figure 3C). Individuals with thicker cingulate cortices at baseline, particularly at the isthmus, experienced a slower progression in parietal WMH volumes compared to those with thinner cortices (Figure 3C; $\sigma_{\iota_{Thick} \leftrightarrows \Delta_{WMH}}^{\text{Regional}}$ = -0.205, SE = 0.064, Z = -3.185, p-value = 0.001). Furthermore, participants with initially thicker precentral, insular, and rostral anterior cingulate cortices showed a slower progression in occipital WMH volumes (Figure 3C, 4A.2; precentral: $\sigma_{\iota_{Thick} \leftrightarrows \Delta_{WMH}}^{\text{Regional}}$ = -0.206, SE = 0.074, Z = -2.791, p-value = 0.005; insular: $\sigma_{\iota_{Thick} \leftrightarrows \Delta_{WMH}}^{\text{Regional}}$ = -0.218, SE = 0.079, Z = -2.743, p-value = 0.006; rostral anterior: $\sigma_{\iota_{Thick} \leftrightarrows \Delta_{WMH}}^{\text{Regional}}$ = -0.183, SE = 0.068, Z = -2.679, p-value = 0.007).

Individuals exhibiting faster WMH volume increases also undergo faster cortical thinning over time ($\sigma_{\Delta_{Thick} \leftrightarrows \Delta_{WMH}}$)

While we found no clear indication for global slope-slope associations ($\sigma_{\Delta_{Thick} \leftrightarrows \Delta_{WMH}}^{\text{Global}}$ = -0.149, SE = 0.145, Z = -1.026, p-value = 0.305), we did observe that individuals with faster progression of WMH in the frontal lobe tended to undergo faster global cortical thinning over time ($\sigma_{\Delta_{Thick} \leftrightarrows \Delta_{WMH}}^{\text{Global}}$ = -0.269, SE = 0.145, Z = -1.861, p-value = 0.063). Further exploration revealed that this relationship was evident in the caudal middle frontal, paracentral, posterior cingulate, and superior

frontal cortex (Figure 4B, S1; peaks – caudal middle frontal: = -0.611,

= 0.243, = -2.516, - = 0.012; paracentral: : = -0.288, =

0.122, = -2.361, - = 0.018; posterior cingulate: : = -0.296, =

0.125, = -2.359, - = 0.018; superior frontal: = -0.336, =

0.152, = -2.216, - = 0.027). Slope-slope associations were generally

characterised by sparsity, with only a few regions showing statistical significance.

Consequently, they collectively did not remain significant after FDR correction and

should thus be interpreted with caution (FDR-corrected version: Figure 3D;

uncorrected version: Figure S1).

Association between growth parameters of WMH and cortical thickness*

(A) Cortical thickness and WMH intercepts ($\sigma_{\iota_{Thick} \leftrightarrows \iota_{WMH}}$)

Frontal   Parietal   Occipital   Temporal

(B) WMH intercept and cortical thickness slope ($\sigma_{\iota_{WMH} \leftrightarrows \Delta_{Thick}}$)

Frontal   Parietal   Occipital   Temporal

(C) Cortical thickness intercept and WMH slope ($\sigma_{\iota_{Thick} \leftrightarrows \Delta_{WMH}}$)

Frontal   Parietal   Occipital   Temporal

(D) Cortical thickness and WMH slopes ($\sigma_{\Delta_{Thick} \leftrightarrows \Delta_{WMH}}$)

Frontal   Parietal   Occipital   Temporal

* Latent growth parameters were adjusted for age, sex, years of education, total cardiovascular risk factor, and total intracranial volume

Figure 3. Spatiotemporal coupling between cortical thickness and WMH. We employed
longitudinal BLGCMs to characterise the spatiotemporal interrelation between lobar WMH and
regional cortical thickness over the span of four years—one model for each pair. We adjusted latent
intercepts and slopes for age, sex, years of education, total cardiovascular risk scores, and TICV. We
applied FDR correction to account for multiple comparisons. In regions highlighted in red, we found a
statistically significant covariance between latent growth curve parameters after FDR correction
( ). (A; *intercept – intercept covariance*). Individuals with larger baseline WMH volumes had
lower mean cortical thicknesses over the entire brain. (B; *WMH intercept – thickness slope*

covariance) Individuals who underwent a steeper decrease in the thickness of their cingulate cortices
had larger WMH volumes across the frontal, parietal, and occipital lobes. (C; *thickness intercept –*
WMH slope covariance). Individuals with thicker cingulate cortices at baseline, particularly at the level
of the isthmus, experienced a slower progression in parietal WMH volumes compared to those with
thinner cortices. Also, those with initially thicker precentral, insular, and rostral anterior cingulate
cortices had a slower progression in occipital WMH volumes. (D; s*lope – slope covariance*) Slope-
slope covariances did not survive FDR correction. However, a rapid increase in frontal, parietal, and
temporal WMH volumes was, in general, associated with accelerated cortical thinning across multiple
brain regions (Figure S1). Associations with temporal WMH volumes did not survive FDR correction
and, thus, we excluded them from the plot. The uncorrected version of the figure is available in
Supplementary Material.

Figure 4. Peak intercept-slope and slope-slope asociations. (A) Average trajectories of cortical
thickness and WMH trajectories over the course of four years, stratified by latent WMH and cortical
thickness intercepts, respectively. We plotted here the four strongest intercept-slope associations,
according to Figure 3B and C. We categorised individuals based on whether their latent intercepts
were below the 33rd or above the 66th percentile, respectively. We then estimated the average
trajectories for each of these groups. We back-transformed all predicted measurements to plotting for
interpretability purposes. Solid and dashed lines depict the predicted average trajectories for these
groups. Shadowed areas represent confidence intervals. (A.1) On average, the reduction in anterior
cingulate cortex thickness among individuals with high frontal WMH volumes would be twice as large

as among those with low WMH volumes (0.069 mm vs 0.029 mm in four years). Likewise, the decrease in thickness across the superior temporal cortices among individuals with high frontal WMH volumes would be approximately 1.5 times greater than that observed in those with low WMH volumes (0.021 mm vs 0.013 mm over four years). (A.2) On average, parietal WMH volume increases among individuals with thick isthmus cingulate cortices would be half as large as that observed in those with thin ones (0.146 ml vs 0.294 ml over four years). Similarly, occipital WMH volume increases across the among individuals with thin anterior cingulate cortex would be approximately four times greater than those observed in those with thick ones (0.108 ml vs 0.026 ml in four years). (B) Individuals with faster frontal WMH progression tended to have faster cortical thinning over time, especially in the caudal middle frontal, paracentral, posterior cingulate, and superior frontal cortex.

Discussion

We studied the interrelationships between WMH and cortical thickness over a four-year period in 436 older adults without objective cognitive impairment (1834 MRI sessions in total) using a longitudinal modelling approach. We made both methodological and clinical contributions to the ongoing efforts to understand the relationship between cerebrovascular dysfunction and neurodegeneration. First, our study demonstrates the potential of integrating surface-based morphometry and BLGCM to investigate interrelationships between neuroimaging markers over time. Second, our findings support the notion that cortical thinning and WMH progression might be mutually reinforcing processes, entangled over a four-year period in a complex and region-specific manner. Our results suggest that this coupling takes place even among individuals with a low vascular risk, given DELCODE's inclusion and exclusion criteria.

WMH progression

WMH generally progressed over the course of four years, reiterating that ageing is associated with WMH increase and constitutes a major risk factor for white matter pathology [2,14,15,28,52]. Progression rates were nonetheless highly subject-specific and dependent on an individual's pre-existing white matter pathology. Parietal, occipital, and temporal WMH progression rates over four years were lower in subjects who had larger WMH volumes at the time of the baseline visit. This suggests that WMH progression may follow a non-linear trajectory, with changes occurring swiftly in its early stages, but gradually reaching a plateau. While non-linear trajectories would be conceivable at late stages of cerebral small vessel

disease (CSVD) spectrum, our finding suggest that this phenomenon also applies to

individuals with low-to-average WMH volumes (compared to CSVD cohorts, see

[52,53]), was somewhat unexpected, albeit not a completely novel finding. The

apparent non-Markovian nature of WMH progression has recently been discussed in

the context of the RUN DMC study [54]. In the RUN DMC study, the total WMH

burden around the age of 64 years predicted their progression over the following 11

years, with severe WMH cases progressing the most, and mild cases progressing

the least [54]. Mild cases rarely progressing to severe stages [54]. Here, we found

that such non-linear trajectories were observable at a study level. However,

significant individual differences in WMH volume changes suggest that there are

numerous other factors that were not accounted for in our study that might contribute

to subject-specific progression of WMH in ageing. For example, heterogeneity of

WMH volumes and progression rates could be reflective of the brain's ability to

respond to and heal from white matter injuries. By extension, heterogeneity of WMH

volumes and progression rates could be reflective of past and current socioeconomic

status and cardiovascular risk factors, as well as the adoption of an unhealthy

lifestyle [2,54,55]. This might explain why lower years of education and greater

cardiovascular risk scores were associated with higher baseline WMH volumes in

our sample.

Interestingly, even though, in our study sample, males were generally older than

females and had higher cardiovascular risk factor scores than females, females

showed significantly greater WMH volumes at baseline compared to males even

after accounting for TICV. Total WMH progression rates over the course of four

years between sexes were nonetheless comparable, except in the occipital lobe,

where females exhibited faster progression. For these two scenarios to be

compatible, WMH would clearly need to evolve faster in females than in males in

other cerebral lobes before the age of 70 years (i.e., the median age in this study).

Menopause may constitute a potential explanation for this sex-specific susceptibility

to WMH. A relatively recent work in the Rhineland study, a large population-based

German cohort, found that while pre-menopausal women and men of similar age did

not differ in WMH volumes, post-menopausal women did have significantly larger

WMH volumes compared to men of similar age [56]. This finding suggests that

indeed menopause and accompanying hormonal and physiological changes might

be behind this sex-difference[56]. Another explanation could be that elderly women

in this ageing cohort had, on average, lower educational attainment, which could

also contribute to their vulnerability to CSVD. The likely multifactorial nature of this

finding requires careful consideration during modelling and reporting as well as

dedicated analysis shedding light on the mechanisms potentially mediating such a

vulnerability.

Albeit less commonly, a small number of participants exhibited clear and consistent

WMH volume regression throughout the study period, as reported in previous

literature [14,57]. The most pronounced case of WMH volume regression was

observed in a female participant in her 60s, with a total cardiovascular risk score of

0.0, and 15 years of education (higher education). Regression in this participant was

most noticeable in occipital brain regions and could be attributed to a loss of

periventricular tissue caused by a substantial enlargement of the occipital horns of

the lateral ventricles over time (Figure S2). While frequently discussed in the context

of a radiological or technical issue [57], our finding adds a new dimension to the

current explanations for WMH regression, wherein genuine changes in one

neuroimaging marker can directly influence another. This finding strongly highlights

the need for multimodal longitudinal strategies to gain a more comprehensive

understanding of the synergistic role of cerebrovascular and neurodegenerative

processes.

Cortical thinning

The thickness of the cerebral cortex decreased over the course of four years,

corroborating that ageing also drives cortical thinning [7,58]. The rate at which

thinning occurred was nonetheless subject- and region-specific. Cortical regions

associated with sensory and sensorimotor functions appeared to be the earliest

affected and most decreased by ageing. A decline in cortical thickness across these

regions might contribute to decreased visual acuity, hearing sensitivity, and motor

abilities [58–61]—all of which are expected during normal ageing, but were not

examined as per the objectives of the current work. We also found substantial

evidence of thinning across the cingulate cortex. This apparent ageing-related

vulnerability is consistent with previous research indicating that both the caudal

anterior and posterior cingulate cortex shrink during normal ageing [62]. The

behavioural consequences of the rate of thinning in terms of decline in cognitive

control and integrating behavioural, affective, and cognitive processes [63], remain to

be elucidated.

Cortical thinning shows considerable heterogeneity across subjects. Somewhat

surprisingly such inter-subject variability could not be fully explained by age, sex,

years of education or cardiovascular risk factors, suggesting that other factors, e.g.,

genetics and lifestyle factors beyond cardiovascular risk factors [10–13], might

influence cortical thinning during late life, possibly to a larger extent than

demographics and established cardiovascular risk factors. Given that the rate of

thinning might affect cognitive performance and activities of daily living, future

research should determine the contribution of brain resilience and (modifiable)

lifestyle factors to abnormal cortical thinning, as such findings could advance the

development of novel interventions [64].

Co-occurrence beyond common risk factors

Even after adjusting for shared risk factors, we found evidence for a negative

correlation between the initial thickness of the cerebral cortex and the initial volume

of WMH, in line with previous work [3,6,10,23–25]. While other factors may

contribute to this relationship, which we did not include in our analysis (e.g., genetics

and lifestyle), this observation, found in a relatively healthy sample, suggests shared

underlying pathological mechanisms.

WMH and cortical thinning

The initial volume of WMH in the brain partly explained the rate of thinning observed

across multiple cortical regions over four years. This observation is consistent with

the cerebrovascular hypothesis [1,65–67] and supports the notion that WMH are the

visible tip of the iceberg [1], a sign of widespread rather than focal cerebrovascular

and metabolic impairment [68,69].The apparent region-specific nature of the

coupling between lobar WMH and regional cortical thickness raises the possibility

that white matter fibres could be involved in the downstream effects of WMH.

Potential secondary effects of WMH along the superior and inferior longitudinal

fasciculus may, for instance, explain why those with occipital WMH would experience

rapid thinning simultaneously across the temporal cortex and the cingulate cortex,

including its rostral anterior regions. Mounting data indeed suggests that abnormal

tissue characteristics can be found in intra- and perilesional white matter regions, but

also in white matter fibres traversing WMH [1,27,69,70]. Also, cross-sectional

investigations conducted in CSVD cohorts have demonstrated that cortical regions

connected to incident lacunes, subcortical lacunar infarcts, and WMH through white

matter fibers exhibit significantly reduced thickness than those that are not [30,65–

67]. Despite the overall compelling evidence for a contribution of WMH to cortical

thinning, additional research leveraging imaging techniques like white matter

tractography as well as animal models is needed to elucidate the role of white matter

fibres in the long-term and remote effects of WMH in the brain.

Cortical thickness and WMH progression

The rate of WMH volumes increase over four years was partly explained by the

thickness of the cerebral cortex at study enrolment, with slower WMH progression

occuring in participants with higher initial cortical thicknesses. This may indicate

potentially higher brain maintanance as a mechanism of healthy ageing [71]. This

relationship was particularly evident when examining WMH progression across

occipital regions in relation to insular and precentral cortex thickness at baseline.

The simultaneous association of insular and precentral cortex thickness with WMH

development may be multifaceted. Neuronal loss in both cortical regions may be

linked to lifestyle adaptations stemming from ageing that contribute to a decline in

sensorimotor functions—a primary risk factor for cardiovascular disease [72].

Considering the involvement of the insular cortex in the regulation of autonomic

functions, a decline in this region could also result in blood pressure dysregulation

[73,74], a condition which has been extensively shown to be associated with

increased progression of WMH, and with more severe manifestations of CSVD

[28,54,75].

The association between baseline cortical thickness and posterior WMH progression

has a fundamental ramification: it supports the spatial heterogeneity of WMH, with

neurodegeneration relating more to the progression of WMH in parietal and occipital

regions than in frontal ones. Since cortical neurodegeneration accelerates with the

pathophysiology of Alzheimer's disease (AD), this would explain why posterior WMH

appear in subjects with minimal vascular pathology across the AD spectrum and why

WMH in deep and periventricular posterior regions appear characteristics of AD

[26,36,38,76]. It is also possible that an early (preclinical) increase in biomarkers

indicative for AD may cause changes in the insular cortex, which then affects the

cardiovascular system [73,74,77] and ultimately speeds up the progression of WMH

in the brain—a possible explanation for Figure 3C. While promising, further research

in other cohorts—especially with available amyloid- or tau- positron emission

tomography [78]—are needed to determine how age- and AD-driven cortical

neurodegeneration influences posterior WMH [76].

WMH progression and cortical thinning

WMH progression and cortical thinning rates were associated with one another,

suggesting a rather consistent and predictable relationship between the two

processes, wherein changes in one marker are accompanied by corresponding

changes in the other and *vice versa*. In our group of cognitively unimpaired

participants, this slope-slope association was particularly evident across frontal brain

regions. This pattern seems to be less confined and more widespread with advanced

stages of AD, as highlighted in a recent work with autosomal dominant AD and late-

onset AD [33]. Further application of our methodology to cohorts at various stages of

AD could, for example, provide further information on the mechanisms underlying the

simultaneous progression of both processes.

Strengths and contextualisation

Longitudinal studies with cognitively unimpaired elderly participants exploring cross-domain associations between WMH and cortical thickness are scarce [1,4,79]. Whenever this kind of research has been done, the evidence supporting any kind of coupling has generally been lacking. In septuagenarian community-dwelling participants, Dickie et al. [4] could not find enough evidence supporting the relationship between total WMH volumes and cortical thickness of cortical grey matter structures neighbouring the Sylvian fissures over a three-year period. In a cohort of cognitively unimpaired participants, Hotz et al. [79] investigated cross-domain associations between total WMH volume and thinning of the entorhinal cortex over a duration of seven years using BLGCM. The authors found no evidence for cross-domain coupling and this absence of association was evident both at the study's baseline and throughout its duration. Our findings are also compatible with the observation that patients with multiple sclerosis (MS), a neuroinflammatory condition primarily affecting white matter fiber tracts through demyelination, exhibit focal thinning of cortical areas [80,81].

Evidence supporting cross-domain has been, nonetheless, growing in participants symptomatic or more severe presentations of cerebrovascular [65–67,78,82] and neurodegenerative pathologies [33,78]. One potential explanation thus far for contradictory results could be the stage of dysfunction at which each participant is situated, i.e., coupling only becomes evident at advanced, symptomatic stages of cerebrovascular and neurodegenerative disease. On the other hand, as emphasised by our study, there are regional nuances to these cross-domain relationships that analyses with a lower level of granularity might fail to capture. Had we solely relied on global analyses for our study, we would have essentially been limited to

720 observing nothing beyond the evident intercept-intercept association. This

721 underscores the significance of employing multimodal and regional approaches to

722 gain a more comprehensive understanding of the local and distant effects of one

723 process on the other.

Limitations

Our research has three main limitations. First, even though our BLGCM aligns with

the data, causality remains elusive due to model equivariance. Latent change score

models might be promising for further study of specific interactions over discrete time

intervals [83]. The mass-univariate application of the BLGCM could be streamlined

by using extended measurement models [84]. We can state, however, that our data

supports a specific and partial spatiotemporal coupling between cortical

neurodegeneration and cerebrovascular dysfunction. The specific circumstances that

might lead to such coupling often remain undetermined and likely require the

inclusion of more extensive biological parameters including complementary imaging

modalities, such as diffusion tensor imaging [27,78,80]. If a Wallerian-like

degeneration is responsible for the observed coupling—as also discussed in the

literature [3,5,9,17,26,34,85]—there should be evidence within the white matter

fibres themselves that mediate the interrelationships between cortical thickness and

WMH. Second, we included a relatively healthy sample and a short time span (48

months, i.e., 4 years), which may have prevented a few cross-domain associations

to become evident. The dynamics over longer time periods, as well as in other

cohorts thus remain elusive, but will be a matter of future investigation. Third, we

have, thus far, not assessed potential cognitive sequelae of WMH progression,

cortical thinning, or their coupling in this study. Because these two processes appear

to be coupled prior to any observable objective cognitive deficiencies, it could be that

cognitive consequences are not detectable at this asymptomatic stage or that

cognitive reserve is still able to compensate for the ongoing pathology or, as a recent

study suggests, that cortical measurements predict well chronological age but not

memory performance [86]. A trivariate latent change score model with WMH, cortical

thickness, and cognitive performance could be used in the future to address this

limitation.

Conclusion

Our work provides longitudinal evidence that cortical thinning and WMH progression

could be mutually reinforcing as opposed to parallel, disassociated processes. The

coupling between these two neuroradiological features appears to be entangled prior

to the onset of any detectable cognitive deficits. Our findings support the ongoing

discussion on perilesional and remote impacts of WMH, but, at the same time,

provide evidence for the effects of cortical neurodegeneration on white matter

integrity. Comprehensive, multimodal approaches, such as the one applied in this

study, have the potential to facilitate the detection of downstream damage

associated with the synergistic interaction among ageing, CSVD, and

neurodegeneration in the brain.

Declarations

Ethics approval and consent to participate

The study protocol was approved by the ethical committees of the medical faculties

of all participating sites: the ethical committees of Berlin (Charite, University

Medicine), Bonn, Cologne, Goettingen, Magdeburg, Munich (Ludwig-Maximilians-

University), Rostock, and Tuebingen. The process was led and coordinated by the

ethical committee of the medical faculty of the University of Bonn. All commitees

gave ethical approval for this work. All participants provided their written informed

consent before inclusion in the study. DELCODE is retrospectively registered at the

German Clinical Trials Register (DRKS00007966, 04/05/2015). The DELCODE

study was conducted in accordance with the Declaration of Helsinki.

Consent for publication

774 Not applicable.

Availability of data and materials

776 The datasets used and analysed during the current study are available from the

777 corresponding author on reasonable request.

Competing interests

779 The authors report no competing interests.

Funding

This research was supported by the German Center for Neurodegenerative

Diseases (Deutsches Zentrum für Neurodegenerative Erkrankungen, DZNE;

reference number BN012) and funded by the German Research Foundation

(Deutsche Forschungsgemeinschaft, DFG; Project IDs 425899996 and

362321501/RTG 2413 SynAGE; CRC 1436, projects A05, B02, B04 and C01). The

funding bodies played no role in the design of the study or collection, analysis, or
interpretation of data or in writing the manuscript.

Author's contributions

Conceptualisation: JB, IM, GZ. Methodology: JB, IM, GZ. Software: JB, IM, GZ.
Formal analysis: JB. Data Acquisition: OP, JHR, SDF, JP, EJS, SA, ASc, KF, JW,
BHS, FJ, AR, WG, EII, KB, DJ, ME, RP, BSR, ST, IK, CL, SSo, ASp, AE, FL, PD,
SH, KS, ED. Image processing: JB, RY. Visualisation: JB. Image analysis and
modelling: JB, GZ. Investigation: JB, GZ. Supervision: GZ, ED. Project
administration: GZ. Funding acquisition: ED. Resources: GZ, ED. Writing original
draft preparation: JB, GZ. Writing – review and editing: All authors. All authors read
and approved the final manuscript.

Acknowledgements

We would like to express our gratitude to all DELCODE study participants. We also
thank the Max-Delbrück-centrum für Molekulare Medizin in der Helmholtz-
Gemeinschaft (MDC), Freie Universität Berlin Center for Cognitive Neuroscience
Berlin (CCNB), Bernstein Center für Computional Neuroscience Berlin,
Universitätsmedizin Göttingen Core Facility MR-Research in Neurosciences, Institut
für Klinische Radiologie Klinikum der Universität München, and Universitätsklinikum
Tübingen MR-Forschungszentrum.

References

References

1. Ter Telgte A, Van Leijsen EMC, Wiegertjes K, Klijn CJM, Tuladhar AM, De Leeuw

FE. Cerebral small vessel disease: From a focal to a global perspective. Nat Rev

Neurol. 2018;14:387–98.

2. Duering M, Biessels GJ, Brodtmann A, Chen C, Cordonnier C, de Leeuw F-E, et

al. Neuroimaging standards for research into small vessel disease-advances since

2013. Lancet Neurol. 2023;4422:2–4.

3. Appelman APA, Exalto LG, Van Der Graaf Y, Biessels GJ, Mali WPTM, Geerlings

MI. White matter lesions and brain atrophy: More than shared risk factors? A

systematic review. Cerebrovascular Diseases. 2009;28:227–42.

4. Dickie DA, Karama S, Ritchie SJ, Cox SR, Sakka E, Royle NA, et al. Progression

of White Matter Disease and Cortical Thinning Are Not Related in Older Community-

Dwelling Subjects. Stroke. 2016;47:410–6.

5. Fiford CM, Manning EN, Bartlett JW, Cash DM, Malone IB, Ridgway GR, et al.

White matter hyperintensities are associated with disproportionate progressive

hippocampal atrophy. Hippocampus. 2017;27:249–62.

6. Lambert C, Benjamin P, Zeestraten E, Lawrence AJ, Barrick TR, Markus HS.

Longitudinal patterns of leukoaraiosis and brain atrophy in symptomatic small vessel

disease. Brain. 2016;139:1136–51.

7. Bethlehem RAI, Seidlitz J, White SR, Vogel JW, Anderson KM, Adamson C, et al.

Brain charts for the human lifespan. Nature. 2022;604:525–33.

8. Narvacan K, Treit S, Camicioli R, Martin W, Beaulieu C. Evolution of deep gray

matter volume across the human lifespan. Hum Brain Mapp. 2017;38:3771–90.

9. Jouvent E, Viswanathan A, Chabriat H. Views and Reviews Cerebral Atrophy in

Cerebrovascular Disorders. 2009;213–8.

10. Enzinger C, Fazekas F, Matthews PM, Ropele S, Schmidt H, Smith S, et al. Risk

factors for progression of brain atrophy in aging: Six-year follow-up of normal

subjects. Neurology. 2005;64:1704–11.

11. Carmelli D, Swan GE, Reed T, Wolf PA, Miller BL, DeCarli C. Midlife

cardiovascular risk factors and brain morphology in identical older male twins.

Neurology. 1999;52:1119–24.

12. Ong M, Foo H, Chander RJ, Wen MC, Au WL, Sitoh YY, et al. Influence of

diabetes mellitus on longitudinal atrophy and cognition in Parkinson's disease. J

Neurol   Sci   [Internet].   2017;377:122–6.   Available   from:

http://dx.doi.org/10.1016/j.jns.2017.04.010

13. Xu J, Li Y, Lin H, Sinha R, Potenza MN. Body mass index correlates negatively

with white matter integrity in the fornix and corpus callosum: A diffusion tensor

imaging study. Hum Brain Mapp. 2013;34:1044–52.

14. Jochems ACC, Arteaga C, Chappell F, Ritakari T, Hooley M, Doubal F, et al.

Longitudinal Changes of White Matter Hyperintensities in Sporadic Small Vessel

Disease: A Systematic Review and Meta-analysis. Neurology. 2022;99:E2454–63.

15. Wardlaw JM, Valdés Hernández MC, Muñoz-Maniega S. What are white matter

hyperintensities made of? Relevance to vascular cognitive impairment. J Am Heart

Assoc. 2015;4:001140.

16. Behl C. Apoptosis and Alzheimer ' s disease Review. Journal of Neural

Transmissio. 2000;1325–44.

17. Nasrabady SE, Rizvi B, Goldman JE, Brickman AM. White matter changes in

Alzheimer's disease: a focus on myelin and oligodendrocytes. Acta Neuropathol

Commun. 2018;6:22.

18. Obulesu M, Lakshmi MJ. Apoptosis in Alzheimer's Disease: An Understanding of

the Physiology, Pathology and Therapeutic Avenues. Neurochem Res.

2014;39:2301–12.

19. Jouvent E, Mangin JF, Duchesnay E, Porcher R, Düring M, Mewald Y, et al.

Longitudinal changes of cortical morphology in CADASIL. Neurobiol Aging [Internet].

2012;33:1002.e29-1002.e36. Available from:

http://dx.doi.org/10.1016/j.neurobiolaging.2011.09.013

20. Brown WR, Moody DM, Thore CR, Challa VR. Apoptosis in leukoaraiosis.

American Journal of Neuroradiology. 2000;21:79–82.

21. Wen W, Sachdev PS, Chen X, Anstey K. Gray matter reduction is correlated with

white matter hyperintensity volume: A voxel-based morphometric study in a large

epidemiological sample. Neuroimage. 2006;29:1031–9.

22. Kim SE, Kim HJ, Jang H, Weiner MW, DeCarli C, Na DL, et al. Interaction

between Alzheimer's Disease and Cerebral Small Vessel Disease: A Review

Focused on Neuroimaging Markers. Int J Mol Sci. MDPI; 2022.

23. Dadar M, Manera AL, Ducharme S, Collins DL. White matter hyperintensities are

associated with grey matter atrophy and cognitive decline in Alzheimer's disease and

frontotemporal dementia. Neurobiol Aging [Internet]. 2022;111:54–63. Available

from: https://doi.org/10.1016/j.neurobiolaging.2021.11.007

24. Rizvi B, Lao PJ, Chesebro AG, Dworkin JD, Amarante E, Beato JM, et al.

Association of Regional White Matter Hyperintensities with Longitudinal Alzheimer-

Like Pattern of Neurodegeneration in Older Adults. JAMA Netw Open. 2021;1–13.

25. Rizvi B, Sathishkumar M, Kim S, Márquez F, Granger SJ, Larson MS, et al.

Posterior white matter hyperintensities are associated with reduced medial temporal

lobe subregional integrity and long-term memory in older adults. Neuroimage Clin.

2023;37.

26. Garnier-crussard A, Krolak-salmon P, Garnier-crussard A, Cotton F, Krolak-

salmon P. White matter hyperintensities in Alzheimer ' s disease▢: Beyond vascular

contribution. Alzheimers Dement. 2023;

27. Dalby RB, Eskildsen SF, Videbech P, Frandsen J, Mouridsen K, Sørensen L, et

al. Oxygenation differs among white matter hyperintensities, intersected fiber tracts

and unaffected white matter. Brain Commun. 2019;1.

28. Ungvari Z, Toth P, Tarantini S, Prodan CI, Sorond F, Merkely B, et al.

Hypertension-induced cognitive impairment: from pathophysiology to public health.

Nat Rev Nephrol. 2021;17:639–54.

29. van Veluw SJ, Arfanakis K, Schneider JA. Neuropathology of Vascular Brain

Health: Insights from Ex Vivo Magnetic Resonance Imaging-Histopathology Studies

in Cerebral Small Vessel Disease. Stroke. 2022;53:404–15.

30. Mayer C, Frey BM, Schlemm E, Petersen M, Engelke K, Hanning U, et al.

Linking cortical atrophy to white matter hyperintensities of presumed vascular origin.

Journal of Cerebral Blood Flow and Metabolism. 2021;41:1682–91.

31. McAleese KE, Firbank M, Dey M, Colloby SJ, Walker L, Johnson M, et al.

Cortical tau load is associated with white matter hyperintensities. Acta Neuropathol

Commun. 2015;3:60.

32. McAleese KE, Walker L, Graham S, Moya ELJ, Johnson M, Erskine D, et al.

Parietal white matter lesions in Alzheimer's disease are associated with cortical

neurodegenerative pathology, but not with small vessel disease. Acta Neuropathol.

2017;134:459–73.

33. Shirzadi Z, Schultz SA, Yau W-YW, Joseph-Mathurin N, Fitzpatrick CD, Levin R,

et al. Etiology of White Matter Hyperintensities in Autosomal Dominant and Sporadic

Alzheimer Disease. JAMA Neurol [Internet]. 2023; Available from:

https://jamanetwork.com/journals/jamaneurology/fullarticle/2810315

34. Salvadores N, Gerónimo-Olvera C, Court FA. Axonal Degeneration in AD: The

Contribution of Aβ and Tau. Front Aging Neurosci. Frontiers Media S.A.; 2020.

35. Bernal J, Schreiber S, Menze I, Ostendorf A, Pfister M, Geisendörfer J, et al.

Arterial hypertension and β-amyloid accumulation have spatially overlapping effects

on posterior white matter hyperintensity volume: a cross-sectional study. Alzheimers

Res Ther. 2023;15.

36. Alber J, Alladi S, Bae HJ, Barton DA, Beckett LA, Bell JM, et al. White matter

hyperintensities in vascular contributions to cognitive impairment and dementia

(VCID): Knowledge gaps and opportunities. Alzheimer's and Dementia: Translational

Research and Clinical Interventions. 2019;5:107–17.

37. Garnier-Crussard A, Bougacha S, Wirth M, Dautricourt S, Sherif S, Landeau B,

et al. White matter hyperintensity topography in Alzheimer's disease and links to

cognition. Alzheimer's and Dementia. 2022;18:422–33.

38. Pålhaugen L, Sudre CH, Tecelao S, Nakling A, Almdahl IS, Kalheim LF, et al.

Brain amyloid and vascular risk are related to distinct white matter hyperintensity

patterns. Journal of Cerebral Blood Flow and Metabolism. 2021;41:1162–74.

39. Jessen F, Spottke A, Boecker H, Brosseron F, Buerger K, Catak C, et al. Design

and first baseline data of the DZNE multicenter observational study on predementia

Alzheimer's disease (DELCODE). Alzheimers Res Ther. 2018;10:1–10.

40. Gaser C, Dahnke R, Kurth K, Luders E, Alzheimer's Disease Neuroimaging

Initiative. A computational Anatomy Toolbox for the Analysis of Structural MRI Data.

bioRxiv. 2022;1–37.

41. Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, et al. An

automated labeling system for subdividing the human cerebral cortex on MRI scans

into gyral based regions of interest. Neuroimage. 2006;31:968–80.

42. Yushkevich PA, Pluta J, Wang H, Wisse LEM, Das S, Wolk D. IC-P-174: Fast

Automatic Segmentation of Hippocampal Subfields and Medial Temporal Lobe

Subregions In 3 Tesla and 7 Tesla T2-Weighted MRI. Alzheimer's & Dementia.

2016;12.

43. Isensee F, Schell M, Pflueger I, Brugnara G, Bonekamp D, Neuberger U, et al.

Automated brain extraction of multisequence MRI using artificial neural networks.

Hum Brain Mapp. 2019;40:4952–64.

44. Wiltgen T, McGinnis J, Schlaeger S, Kofler F, Voon CC, Berthele A, et al. LST-

AI: A deep learning ensemble for accurate MS lesion segmentation. Neuroimage

Clin. 2024;42.

45. Joshi AA, Choi S, Liu Y, Chong M, Sonkar G, Gonzalez-Martinez J, et al. A

hybrid high-resolution anatomical MRI atlas with sub-parcellation of cortical gyri

using resting fMRI. J Neurosci Methods. 2022;374.

46. Larivière S, Paquola C, Park B yong, Royer J, Wang Y, Benkarim O, et al. The

ENIGMA Toolbox: multiscale neural contextualization of multisite neuroimaging

datasets. Nat Methods. Nature Research; 2021. p. 698–700.

47. McArdle JJ, Nesselroade JohnR. Using multivariate data to structure

developmental change." Life-span developmental psychology: Methodological

contributions. In: Cohen SH, Reese HW, editors. Life-Span Developmental

Psychology: Methodological Contributions. 1st ed. New York: Psychology Press;

1994. p. 223–67.

48. Muniz-Terrera G, Robitaille A, Kelly A, Johansson B, Hofer S, Piccinin A. Latent

growth models matched to research questions to answer questions about dynamics

of change in multiple processes. J Clin Epidemiol. 2017;82:158–66.

49. Tukey JW. Exploratory data analysis. Addison-Wesley Pub. Co.; 1977.

50. Hu LT, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis:

Conventional criteria versus new alternatives. Structural Equation Modeling.

1999;6:1–55.

51. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and

powerful approach to multiple testing. Journal of the royal statistical society Series B

(Methodological). 1995;289–300.

52. Van Leijsen EMC, Van Uden IWM, Ghafoorian M, Bergkamp MI, Lohner V,

Kooijmans ECM, et al. Nonlinear temporal dynamics of cerebral small vessel

disease. Neurology. 2017;89:1569–77.

53. Keuss SE, Coath W, Nicholas JM, Poole T, Barnes J, Cash DM, et al.

Associations of β-Amyloid and Vascular Burden with Rates of Neurodegeneration in

Cognitively Normal Members of the 1946 British Birth Cohort. Neurology.

2022;99:E129–41.

54. Cai M, Jacob MA, Van Loenen MR, Bergkamp M, Marques J, Norris DG, et al.

Determinants and Temporal Dynamics of Cerebral Small Vessel Disease: 14-Year

Follow-Up. Stroke. 2022;53:2789–98.

55. Wardlaw JM, Smith C, Dichgans M. Small vessel disease: mechanisms and

clinical implications. Lancet Neurol. 2019;18:684–96.

56. Lohner V, Pehlivan G, Sanroma G, Miloschewski A, Schirmer MD, Stöcker T, et

al. Relation Between Sex, Menopause, and White Matter Hyperintensities: The

Rhineland Study. Neurology. 2022;99:E935–43.

57. Brown RB, Tozer DJ, Egle M, Tuladhar AM, de Leeuw FE, Markus HS. How

often does white matter hyperintensity volume regress in cerebral small vessel

disease? International Journal of Stroke. 2023;00.

58. Salat DH, Buckner RL, Snyder AZ, Greve DN, Desikan RSR, Busa E, et al.

Thinning of the Cerebral Cortex in Aging. Cerebral Cortex [Internet]. 2004;14:721–

30. Available from: https://doi.org/10.1093/cercor/bhh032

59. Cardin V. Effects of aging and adult-onset hearing loss on cortical auditory

regions. Front Neurosci. Frontiers Media S.A.; 2016.

60. Li KZH, Lindenberger U. Relations between aging sensory/sensorimotor and

cognitive functions. Neurosci Biobehav Rev [Internet]. 2002;26:777–83. Available

from: www.elsevier.com/locate/neubiorev

61. Schneider BA, Pichora-Fuller MK. Implications of perceptual deterioration for

cognitive aging research. The handbook of aging and cognition, 2nd ed. Mahwah,

NJ, US: Lawrence Erlbaum Associates Publishers; 2000. p. 155–219.

62. Mann SL, Hazlett EA, Byne W, Hof PR, Buchsbaum MS, Cohen BH, et al.

Anterior and posterior cingulate cortex volume in healthy adults: Effects of aging and

gender differences. Brain Res. 2011;1401:18–29.

63. Shackman AJ, Salomons T V., Slagter HA, Fox AS, Winter JJ, Davidson RJ. The

integration of negative affect, pain and cognitive control in the cingulate cortex. Nat

Rev Neurosci. 2011. p. 154–67.

64. Schwarck S, Voelkle MC, Becke A, Busse N, Glanz W, Ziegler G. Interplay of

physical and cognitive performance using hierarchical continuous-time dynamic

modelling and a 2 dual-task training regime in Alzheimer's patients. Available from:

https://doi.org/10.1101/2022.12.14.22283428

65. Duering M, Righart R, Wollenweber FA, Zietemann V, Gesierich B, Dichgans M.

Acute infarcts cause focal thinning in remote cortex via degeneration of connecting

fiber tracts. Neurology. 2015;84:1685–92.

66. Duering M, Righart R, Csanadi E, Jouvent E, Herve D, Chabriat H, et al. Incident

subcortical infarcts induce focal thinning in connected cortical regions. Neurology.

2012;79:2025–8.

67. Li H, Jacob MA, Cai M, Duering M, Chamberland M, Norris DG, et al. Regional

cortical thinning, demyelination, and iron loss in cerebral small vessel disease. Brain.

2023;

68. Jiaerken Y, Luo X, Yu X, Huang P, Xu X, Zhang M. Microstructural and metabolic

changes in the longitudinal progression of white matter hyperintensities. Journal of

Cerebral Blood Flow and Metabolism. 2019;39:1613–22.

69. Reginold W, Sam K, Poublanc J, Fisher J, Crawley A, Mikulis DJ. Impact of white

matter hyperintensities on surrounding white matter tracts. Neuroradiology.

2018;60:933–44.

70. Wardlaw JM, Makin SJ, Valdés Hernández MC, Armitage PA, Heye AK,

Chappell FM, et al. Blood-brain barrier failure as a core mechanism in cerebral small

vessel disease and dementia: evidence from a cohort study. Alzheimer's and

Dementia. 2017;13:634–43.

71. Cabeza R, Albert M, Belleville S, Craik FIM, Duarte A, Grady CL, et al.

Maintenance, reserve and compensation: the cognitive neuroscience of healthy

ageing. Nat Rev Neurosci. Nature Publishing Group; 2018. p. 701–10.

72. Li J, Siegrist J. Physical activity and risk of cardiovascular disease-a meta-

analysis of prospective cohort studies. Int J Environ Res Public Health. MDPI; 2012.

p. 391–407.

73. Benarroch EE. Insular cortex: Functional complexity and clinical correlations.

Neurology. 2019;93:932–8.

74. Gogolla N. The insular cortex. Current Biology. Cell Press; 2017. p. R580–6.

75. Duering M, Biessels GJ, Brodtmann A, Chen C, Cordonnier C, de Leeuw F-E, et

al. Neuroimaging standards for research into small vessel disease—advances since

2013. Lancet Neurol [Internet]. 2023;22:602–18. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S147444222300131X

76. Desmarais P, Gao AF, Lanctôt K, Rogaeva E, Ramirez J, Herrmann N, et al.

White matter hyperintensities in autopsy-confirmed frontotemporal lobar

degeneration and Alzheimer's disease. Alzheimers Res Ther. 2021;13:1–16.

77. Kitamura J, Nagai M, Ueno H, Ohshita T, Kikumoto M, Toko M, et al. The Insular

Cortex, Alzheimer Disease Pathology, and Their Effects on Blood Pressure

Variability [Internet]. 2020. Available from: www.alzheimerjournal.com

78. Zhang J, Chen H, Wang J, Huang Q, Xu X, Wang W, et al. Linking white matter

hyperintensities to regional cortical thinning, amyloid deposition, and synaptic density

loss in Alzheimer's disease. Alzheimer's & Dementia [Internet]. 2024; Available from:

https://alz-journals.onlinelibrary.wiley.com/doi/10.1002/alz.13845

79. Hotz I, Deschwanden PF, Mérillat S, Jäncke L. Associations between white

matter hyperintensities, lacunes, entorhinal cortex thickness, declarative memory

and leisure activity in cognitively healthy older adults: A 7-year study. Neuroimage.

2023;284.

80. Bussas M, Grahl S, Pongratz V, Berthele A, Gasperi C, Andlauer T, et al. Gray

matter atrophy in relapsing-remitting multiple sclerosis is associated with white

matter lesions in connecting fibers. Multiple Sclerosis Journal. 2022;28:900–9.

81. Sailer M, Fischl B, Salat D, Tempelmann C, Schönfeld MA, Busa E, et al. Focal

thinning of the cerebral cortex in multiple sclerosis. Brain. 2003;126:1734–44.

82. Kim SJ, Lee DK, Jang YK, Jang H, Kim SE, Cho SH, et al. The effects of

longitudinal white matter hyperintensity change on cognitive decline and cortical

thinning over three years. J Clin Med. 2020;9:1–13.

83. Kievit RA, Brandmaier AM, Ziegler G, van Harmelen AL, de Mooij SMM,

Moutoussis M, et al. Developmental cognitive neuroscience using latent change

score models: A tutorial and applications. Dev Cogn Neurosci [Internet]. 2018;33:99–

117. Available from: https://doi.org/10.1016/j.dcn.2017.11.007

84. Wang L, Lyu X, Zhang Z, Li L. High-dimensional Response Growth Curve

Modeling for Longitudinal Neuroimaging Analysis. ArXiv [Internet]. 2023;1–30.

Available from: http://arxiv.org/abs/2305.15751

85. Reginold W, Itorralba J, Luedke AC, Fernandez-Ruiz J, Reginold J, Islam O, et

al. Tractography at 3T MRI of corpus callosum tracts crossing white matter

hyperintensities. American Journal of Neuroradiology. 2016;37:1617–22.

86. Soch J, Richter A, Kizilirmak JM, Schütze H, Feldhoff H, Fischer L, et al.

Structural and Functional MRI Data Differentially Predict Chronological Age and

Behavioral Memory Performance. eNeuro. 2022;9.

1068