



25 **Abstract**

26 *Background*

27 Spinal cord infarction (SCI) is a rare and challenging condition to diagnose. The literature lacks  
28 large studies examining the characteristics and outcomes of SCI. Treatments, such as  
29 thrombolysis, which are considered standard of care in cerebral stroke, have not been previously  
30 studied in SCI. The primary aim was to determine outcomes of hospitalizations with SCI in the  
31 United States (US) by assessing inpatient mortality and all-cause 30-day and 90-day  
32 readmissions. After adjusting for age and comorbidity burden, we analyzed if these outcomes  
33 differed based on thrombolysis, and the most common concomitant conditions, such as  
34 aneurysms, dissections and sepsis.

35 *Methods*

36 Hospitalizations of patients over 18 years of age with SCI were abstracted from the 2016-2021  
37 Nationwide Readmissions Database (NRD). We estimated logistic and lognormal regression  
38 models to evaluate whether outcomes differed based on thrombolysis, or concomitant aneurysm,  
39 dissection, or sepsis. Multivariable models adjusted for age and comorbidity burden.

40 *Results*

41 An estimated 12,548 hospitalizations involved patients diagnosed with SCI; of those, 21.43%  
42 were specifically for SCI. Other common reasons for SCI hospitalization included aneurysms or  
43 dissections (22.22%) and sepsis (9.11%). The average LOS was 13.09 days (95% CI: 12.61-  
44 13.59), with an inpatient mortality rate of 14.66% (95% CI: 13.74-15.59). Age and comorbidity  
45 burden significantly impacted inpatient mortality, LOS, and hospital costs. Hospitalizations of  
46 patients with SCI were predominately at large bed size facilities (72.36%, Weighted N: 9,079)  
47 and metropolitan teaching hospitals (86.57%, Weighted N: 10,862). Thrombolysis was associated

48 with lower odds of 90-day readmissions (adjusted OR=0.35, p-value= 0.009) and no significant  
49 difference in mortality. Concomitant sepsis significantly increased mortality, LOS, cost, and  
50 readmission odds.

#### 51 *Conclusions*

52 This study highlights factors that impact the outcomes of SCI on a large scale. Additionally, the  
53 findings uncover the association between thrombolysis and better long-term outcomes in SCI.

54 This paves the way for further investigation into the role of thrombolysis in SCI.

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

## 71 **Introduction**

72 Spinal cord infarction (SCI) accounts only for 0.3 to 1% of all strokes, and 6% of all acute  
73 myelopathies<sup>1,2</sup>. The most frequently affected spinal cord segments are, in descending order,  
74 thoracic, cervical and the conus medullaris<sup>2,3</sup>. Clinical presentation of SCI consists of an acute  
75 myelopathy, accompanied by sphincteric and autonomic dysfunction, and back pain<sup>2,3</sup>. Previous  
76 work has reported a median age of SCI onset between 52 and 60 years<sup>3</sup>. The diagnosis of SCI  
77 can be challenging to confirm, and misdiagnosis is common<sup>2</sup>. Zalewski et al. described  
78 diagnostic criteria that includes clinical, radiological and CSF findings to aid in the diagnosis of  
79 SCI in clinical practice and for research purposes<sup>3</sup>. Etiologically, SCI can be classified into  
80 spontaneous and peri-procedural. Spontaneous SCI is often attributed to atherosclerosis and  
81 vascular risk factors, whereas aortic repair surgeries contribute to the majority of peri-procedural  
82 SCIs<sup>3-5</sup>. The most reported risk factors for SCI include hypertension, peripheral vascular disease  
83 and advanced age<sup>3,4</sup>. Additionally, spontaneous SCI has been reported in association with  
84 degenerative spine disease, owing to fibrocartilaginous embolization<sup>9,10</sup>. Regarding the risk of  
85 peri-procedural SCI with open versus endovascular aortic repairs, studies showed conflicting  
86 results<sup>6,7</sup>. A meta-analysis by Gaudino et al. showed a rate of SCI of 5.7% for open repair and  
87 3.9% for endovascular repair<sup>8</sup>.

88 The management of SCI is limited, lacking large randomized controlled trials. The role of  
89 thrombolysis in SCI is not established, with only a few case reports and series that showed  
90 potential beneficial effects<sup>12,13</sup>. Lumbar cerebrospinal fluid drains have been proven to aid in the  
91 prevention of SCI in association with aortic repair surgeries<sup>14</sup>. Studies have shown that recovery  
92 after SCI can be slow and gradual over months to years<sup>2,11</sup>. Furthermore, the functional recovery  
93 is much worse after peri-procedural SCI compared to spontaneous SCI<sup>2</sup>. This is attributed to the

94 extent of spinal cord segments involved in peri-procedural SCI<sup>2</sup>. The 1-year and 5-year mortality  
95 rates of SCI were shown to be around 7% and 20.9%, respectively<sup>15</sup>. In addition, no significant  
96 difference was found in the survival of patients affected by SCI due to different etiologies<sup>15</sup>.  
97 The literature lacks large studies analyzing characteristics and outcomes of SCI. Given the rarity  
98 of SCI, standard treatments for cerebral stroke, such as thrombolysis, have not been previously  
99 studied. In this study, we aim to assess the characteristics and outcomes of SCI in a large sample  
100 representative of the United States, using the Nationwide Readmissions Database (NRD). The  
101 advantage of using the NRD as the data source for our study is that readmission rates are  
102 considered good indicators for long-term morbidity in cerebrovascular disease and stroke.  
103 Additionally, it allows us to capture as many hospitalizations as possible when it pertains to rare  
104 conditions and exposures, such as SCI and thrombolysis.

105

## 106 **Methods**

### 107 *Data Source and Identification*

108 Hospitalization data were abstracted from 2016 through 2021 NRD. The NRD is an all-payer,  
109 administrative inpatient care database sponsored by the Agency for Healthcare Research and  
110 Quality (AHRQ) and estimates approximately 35 million yearly discharges in United States<sup>16</sup>.  
111 We identified hospitalization of patients with a diagnosis of SCI using *International*  
112 *Classification of Diseases, Tenth Revision Codes* (ICD-10: G95.11). Hospitalizations were  
113 excluded if the patient was younger than 18 years. Creighton University acknowledged this study  
114 as Not Human Subjects Research (InfoEd Record Number: 2004608); the NRD is HIPAA-  
115 compliant and publicly available. This study followed the STROBE reporting guidelines<sup>17</sup>.

116 *Study Outcomes*

117 The primary aim was to assess the characteristics of hospitalizations of patients with SCI in a  
118 nationally representative sample. First, we described reasons for SCI hospitalization and the  
119 descriptive characteristics of the SCI hospitalization cohort. Then, we assessed whether inpatient  
120 mortality and length of stay (LOS) were associated with age and comorbidity burden. Lastly, we  
121 evaluated whether hospitalization outcomes varied 1) by those who received thrombolysis, 2)  
122 those who experienced an aneurysm or dissection, and 3) those that carried a sepsis diagnosis.  
123 Hospitalization outcomes included inpatient mortality, LOS, hospital cost, discharge disposition  
124 (routine, transfer to short-term hospital, transfer to other facility), 30-day all-cause readmission,  
125 and 90-day all-cause readmission.

126 Most frequent reasons for hospitalization were identified using the primary ICD-10 diagnosis  
127 code. ICD-10 diagnosis and procedures codes were used to identify SCI hospitalizations with  
128 receipt of thrombolysis (ICD-10-CM; Z92.82, ICD-10-PCS: 3E03017, 3E03317). ICD-10  
129 diagnosis codes were used to identify SCI hospitalizations of patients with an  
130 aneurysm/dissection (ICD-10-CM: I71, I72) or sepsis (ICD-10-CM: A40, A41). Hospital cost  
131 was inflation adjusted to July-2021 US dollars<sup>18</sup>. For readmission analysis, if a patient died, the  
132 hospitalization was excluded to allow for complete post-discharge follow-up. Additionally, index  
133 hospitalizations where the patient was discharged in December were excluded for 30-day  
134 readmission analysis and index hospitalizations where the patient was discharge in October,  
135 November, or December were excluded for 90-day readmission analysis to allow complete post-  
136 discharge follow-up since the NRD does not track patients year-to-year.

137 *Descriptives*

138 We extracted descriptives for each hospitalization; descriptives were age, biological sex (male,  
139 female) , primary payer (Medicaid, Medicare, private, other, unknown), income quartile (I, II, III,  
140 IV, unknown), whether the hospital admission was elective, whether the hospital admission  
141 occurred on a weekend, facility bed size (small, medium large), facility location-teaching status  
142 (metropolitan non-teaching, metropolitan teaching, non-metropolitan), comorbidity burden, and  
143 National Institutes of Health Stroke Scale (NIHSS) score. Comorbidity burden was quantified by  
144 the Charlson Comorbidity Index (CCI); a score developed for healthcare research to assess  
145 patient risk<sup>19</sup>. We identified the following Charlson comorbidities: myocardial infarction,  
146 congestive heart failure, peripheral artery disease, stroke, dementia, chronic obstructive  
147 pulmonary disease, rheumatic disease, peptic ulcer, liver disease (mild, severe), diabetes  
148 (uncomplicated, complicated), renal disease (mild, severe), malignancy, metastatic tumor, human  
149 immunodeficiency virus, and acquired immunodeficiency syndrome. We also identified whether  
150 the patient underwent an aortic repair during the hospitalization using ICD-10 procedures codes  
151 (Open Approach: 02QX0ZZ, 02QW0ZZ, 04Q00ZZ; Percutaneous Approach: 02QX3ZZ,  
152 02QW3ZZ, 04Q03ZZ).

### 153 *Statistical Analysis*

154 Continuous descriptive were presented as median and interquartile range and categorical  
155 descriptives were presented as weighted frequency and percent. Logistic regression models were  
156 estimated to evaluate whether inpatient mortality varied by age and CCI. Log-normal models  
157 were estimated to evaluate whether LOS varied by age and CCI.

158 Next, to assess whether inpatient mortality, discharge disposition, and 30-day and 90-day all-  
159 cause readmissions differed by 1) the receipt of thrombolysis, 2) whether the patient experienced  
160 a dissection or aneurysm, and 3) whether the patient suffered from sepsis, we estimated

161 respective logistic regression models. Likewise, we estimated log-normal models to evaluate  
162 whether LOS and hospital cost differed by 1) thrombolysis, 2) dissection or aneurysm, and 3)  
163 sepsis. Multivariable regression models were estimated that controlled for age and comorbidity  
164 burden.

165 The functional form of continuous covariates was evaluated via restricted cubic splines with  
166 knots at the 10<sup>th</sup>, 50<sup>th</sup>, and 90<sup>th</sup> percentiles<sup>20</sup>. The NRD sampling design was accounted for in all  
167 analysis; two-tailed  $p < 0.05$  indicated statistical significances.

168

## 169 **Results**

170 In the United States from 2016 through 2021, there were an estimated 12,548 (95% CI: 12,085-  
171 13,010) hospitalizations where the patient carried a SCI diagnosis. Of those, 21.43% (Weighted  
172 N: 2,689, 95% CI: 2,526-2,853) were hospitalizations specifically for SCI. Other frequent  
173 reasons for hospitalization included hospitalizations for an aneurysm or dissection (22.22%,  
174 Weighted N: 2,537, 95% CI: 2,320-2,754) and sepsis (9.11%, Weighted N: 1,143, 95% CI:  
175 1,131-1,327). **Table 1** presents hospitalization descriptives. Notably, there were no  
176 hospitalizations of patients with SCI where the patient underwent a percutaneous aortic repair.  
177 Hospitalizations of patients with SCI were predominately at large bed size facilities (72.36%,  
178 Weighted N: 9,079) and metropolitan teaching hospitals (86.57%, Weighted N: 10,862). Patients  
179 were primarily insured by Medicare (52.43%, Weighted N= 6,578). Common comorbidities were  
180 peripheral artery disease (37.58%, Weighted N=4,715), chronic obstructive pulmonary disease  
181 (23.00%, Weighted N: 2,886), congestive heart failure (22.22%, Weighted N = 2,788), and  
182 acquired immunodeficiency syndrome (21.46%, Weighted N = 2,693).



183 *Outcomes Overall*

184 Index hospitalizations of patients with a SCI diagnosis had an inpatient mortality rate of 14.66%  
185 (95% CI: 13.74-15.59). The average LOS for hospitalizations of patients with a SCI diagnosis  
186 was 13.09 days (95% CI: 12.61-13.59) and the average hospital cost was \$53,531 (95% CI:  
187 51,615-55,518). Most patients were transferred to a skilled nursing facility, intermediate care  
188 facility, or another type of facility (48.50%, 95% CI: 47.00-49.99). Relatedly, 16.86% (95% CI:  
189 15.78-17.94) of hospitalizations of patients with a SCI diagnosis resulted in a routine discharge  
190 and 3.75% (95% CI: 3.26-4.25) hospitalizations resulted in a transfer to a short-term hospital.  
191 For all-cause readmissions, 21.08% (95% CI: 19.86-22.30) and 34.01% (95% CI: 32.36-35.65)  
192 of hospitalizations of patients with a SCI diagnosis were readmitted within 30 days and 90 days,  
193 respectively.

194 *Outcomes by Age and Comorbidity Burden*

195 **Figure 1** presents the effect of age and CCI on inpatient mortality and LOS. Age was statistically  
196 associated with inpatient mortality and LOS ( $p < 0.001$ ,  $p < 0.001$ , respectively); as age  
197 increased the rate of inpatient mortality increased (**Figure 1A**) and LOS decreased (**Figure 1B**).  
198 CCI was statistically associated with inpatient mortality and LOS ( $p < 0.001$ ,  $p < 0.001$ ,  
199 respectively); as CCI increased the rate of inpatient mortality and LOS increased (**Figure 1C-D**).

200 *Outcomes by Thrombolysis*

201 Notably, 1.17% (Weighted N: 147, 95% CI: 111-183) of hospitalizations with a SCI diagnosis  
202 received thrombolysis. Hospitalization outcomes by whether patients with SCI received  
203 thrombolysis are presented in **Table 2**. Inpatient mortality, length of stay, transfer to a skilled  
204 nursing facility, intermediate care facility, or another type of facility, and all-cause 30-day

205 readmission were statistically similar between hospitalization with thrombolysis and  
206 hospitalization without thrombolysis. After adjusting for age and CCI, the adjusted odds of 90-  
207 day all-cause readmissions were 65% lower for index hospitalizations with thrombolysis  
208 compared to no thrombolysis (aOR 95% CI: 0.16-0.77,  $p = 0.009$ , **Table 2**).

### 209 *Outcomes by Dissections and Aneurysms*

210 From 2016-2021, 29.12% of hospitalizations of patients carrying a diagnosis for SCI also carried  
211 a diagnosis of a dissection or aneurysm (Weighted N: 3,654, 95% CI: 3,393-3,915). **Table 3**  
212 presents hospitalization outcomes by whether patients with SCI had an aneurysm or dissection.  
213 After adjusting for age and CCI, the adjusted odds of inpatient mortality were 52% greater for  
214 SCI hospitalizations of patients with an aneurysm or dissection compared to those without an  
215 aneurysm or dissection (aOR 95% CI: 1.30-1.78,  $p < 0.001$ , **Table 3**). Likewise, adjusted LOS  
216 was 19% greater and adjusted cost was 2.18x greater for SCI hospitalizations of patients with an  
217 aneurysm or dissection compared to those without an aneurysm or dissection (LOS- aOR 95%  
218 CI: 1.09-2.29,  $p < 0.001$ ; Cost- aOR 95% CI: 2.06-2.31,  $p < 0.001$ ; **Table 3**). After adjustment,  
219 discharge disposition and readmissions were similar between SCI hospitalizations with and  
220 without an aneurysm or dissection (**Table 3**).

### 221 *Outcomes by Sepsis*

222 Of the estimated 12,548 SCI hospitalizations, 22.23% also had a sepsis diagnosis (Weighted N:  
223 2,789, 95% CI: 2,618-2,959). Hospitalization outcomes by whether patients with SCI also  
224 suffered from sepsis are presented in **Table 4**. After adjusting for age and CCI, SCI  
225 hospitalizations with sepsis had 3.15x greater adjusted odds of inpatient mortality compared to  
226 hospitalizations without sepsis (aOR 95% CI: 2.64-3.75,  $p < 0.001$ ; **Table 4**). SCI

227 hospitalizations with sepsis had 44% longer adjusted LOS and 54% greater adjusted hospital cost  
228 compared to those without sepsis (LOS- aOR 95% CI: 1.34-1.55,  $p < 0.001$ ; Cost- aOR 95% CI:  
229 1.45-1.65,  $p < 0.001$ ; **Table 4**). Compared to hospitalization without sepsis, hospitalizations with  
230 sepsis had 59% and 33% lower adjusted odds of routine discharge and transfer to a short-term  
231 hospital, respectively (Routine- aOR 95% CI: 0.33-0.52,  $p < 0.001$ ; Short-term Hospital  
232 Transfer- aOR 95% CI: 0.48-0.93,  $p = 0.015$ ; **Table 4**). SCI hospitalizations of patients with  
233 sepsis had 42% and 48% greater adjusted odds of 30-day and 90-day all-cause readmission  
234 compared to those without a sepsis diagnosis, respectively (30-day- aOR 95% CI: 1.19-1.70,  $p <$   
235  $0.001$ ; 90-day- aOR 95% CI: 1.23-1.77,  $p < 0.001$ ; **Table 4**).

236

## 237 **Discussion**

238 Given the rarity of SCI, previous studies have been limited by the number of patients<sup>2,3</sup>. Our  
239 study analyzed 12,548 SCI hospitalizations from 2016 to 2021 using the NRD. Most SCI  
240 hospitalizations were in large and metropolitan teaching hospitals. Diagnosing SCI is not an easy  
241 feat, and most cases require higher levels of care and specialized care<sup>2</sup>. In our study, there were  
242 an estimated 147 SCI hospitalizations where the patient received thrombolysis. Notably,  
243 hospitalizations where the patient received thrombolysis had significantly lower odds of 90-day  
244 all-cause readmissions compared to hospitalizations where the patient did not receive  
245 thrombolysis. This difference remained even after adjusting for age and CCI. However, there was  
246 no significant difference in the rate of inpatient mortality or 30-day all-cause readmissions  
247 between hospitalizations with thrombolysis and hospitalizations without thrombolysis. Studies  
248 have previously shown that recovery after spinal cord infarction can take several months<sup>2,11</sup>. It is,  
249 therefore, plausible that thrombolysis might help in the long-term recovery after SCI. A study

250 using a national registry in Denmark showed that thrombolysis in cerebral stroke had lower  
251 readmission rates for pneumonia, but no overall effect on readmission rates<sup>21</sup>. On the other hand,  
252 Vahidy et al. analyzed the NRD for cerebral strokes and found significantly lower odds of 30-day  
253 readmission with recanalization therapy which included thrombolysis and thrombectomy<sup>29</sup>. Also,  
254 previous work on cerebral stroke showed the long-term benefits of thrombolysis in terms of  
255 functional outcome and independence several months after treatment<sup>22,23</sup>. However, thrombolysis  
256 in cerebral stroke has been shown to be associated with higher in-hospital mortality<sup>24</sup>. This is  
257 also compatible with our findings of 19.45% inpatient mortality of hospitalizations with  
258 thrombolysis compared to 14.60% for hospitalizations without thrombolysis. However, the  
259 difference in mortality was not statistically significant. Our study shows an association between  
260 thrombolysis and better long-term outcomes in SCI. However, large trials are needed to establish  
261 the benefit and risk of thrombolysis in SCI with detailed functional outcomes.

262 Hospitalizations specifically for SCI were the primary reason patients with a SCI diagnosis were  
263 hospitalized; interestingly the other most common reasons a patient with a SCI diagnosis was  
264 hospitalized were aneurysms and dissections followed by sepsis. Hospitalizations primarily for  
265 an aneurysm or dissection had significantly higher inpatient mortality, longer hospital stays, and  
266 higher overall costs compared to hospitalizations where the patient with SCI did not experience  
267 an aneurysm or dissection. These are expected findings given that aortic pathology is one of the  
268 most common conditions leading to SCI<sup>25</sup>. Aortic aneurysms and dissections are also likely to  
269 require surgical intervention, leading to worse outcomes overall<sup>26,27</sup>. These findings carry  
270 prognostic value in patients who suffer from SCI, regardless of age or comorbidity burden.

271 Among all the hospitalizations with SCI, there were no SCI hospitalizations where the patient  
272 had a percutaneous aortic repair. There were 68 SCI hospitalizations where the patient had an

273 open surgical aortic repair. This finding is consistent, yet more remarkable, when compared to  
274 previous studies showing lower rates of peri-procedural SCI after percutaneous aortic repairs  
275 compared to open aortic repairs<sup>8</sup>.

276 We found similar outcomes in hospitalizations that were primarily for sepsis; sepsis was  
277 associated with longer hospital stays and greater costs. The odds of inpatient mortality were  
278 greater for SCI hospitalizations with concomitant sepsis compared to those without sepsis. Sepsis  
279 disproportionately affects older patients with medical comorbidities<sup>30</sup>. Previous studies have  
280 shown that sepsis accounted for 28% of inpatient mortality in spinal cord injuries<sup>28</sup>. The  
281 association between sepsis and SCI is bidirectional. Septic shock can potentially lead to  
282 hypoperfusion and spinal cord ischemia, and septic emboli are known to cause cerebral stroke<sup>31</sup>.

283 Alternatively, it is plausible that SCI is commonly complicated by sepsis. These findings may  
284 highlight the important role sepsis plays in determining inpatient mortality in SCI. Furthermore,  
285 all-cause readmission rates at 30- and 90-days were significantly higher in SCI hospitalizations  
286 with sepsis. Studies that examined admissions solely for sepsis found high rates of readmissions  
287 with most of them occurring within 90 days<sup>32</sup>.

288 We found that as age increased, inpatient mortality increased, contributing to shorter length of  
289 stays and lower costs found in the older age groups of SCI hospitalizations. The median age in  
290 our study was 63 (53-73), which is slightly higher compared to previous reports<sup>3</sup>. Comorbidity  
291 burden played an important role in inpatient mortality and LOS for SCI hospitalization;  
292 comorbidity burden was significantly associated with inpatient mortality and LOS. This is  
293 consistent with reports on cerebral infarction. Han et al. reported a significantly poorer prognosis  
294 in ischemic strokes with a CCI of 3 or more<sup>33</sup>. Our study showed a median CCI of 2 with 50% of  
295 hospitalizations being patients with a CCI between 3 and 7 thus indicating the high comorbidity

296 burden in patients with SCI. In accordance with published reports, we found that peripheral  
297 vascular disease was the most common comorbidity in our SCI hospitalizations<sup>3,4</sup>. Interestingly,  
298 AIDS was one of the common comorbidities in SCI hospitalizations (21.46%). Anecdotal  
299 evidence exists to suggest an association between opportunistic infections, such as Varicella  
300 Zoster Virus, and vasculitis leading to SCI<sup>34</sup>. This finding warrants further investigation into the  
301 association between AIDS and SCI.

302 The NRD is an administrative database reliant on billing codes, non-reimbursable codes may be  
303 underreported. However, the NRD is still a useful database to examine uncommon conditions, as  
304 it captures a large number of hospitalizations on a national and representative scale. The NRD  
305 does not provide information on long-term functional outcomes. Our analysis on thrombolysis is  
306 limited by the low number of hospitalizations, due to possible under-reporting of thrombolysis.

307

## 308 **Conclusions**

309 This study highlights the factors that impact the outcomes of SCI on a large scale. We found that  
310 age, comorbidity burden, and concomitant aneurysms, dissections and sepsis all play major roles  
311 in the outcome of SCI. These findings can assist in evaluating prognosis after SCI. Additionally,  
312 the findings uncover the association between thrombolysis and better long-term outcomes in  
313 SCI. This paves the way for further investigation into the role of thrombolysis in SCI. Large  
314 randomized controlled trials are needed to establish the benefit and risk of thrombolysis in SCI.

315

316

317

318

319 **Acknowledgements and Sources of Funding**

320 None.

321

322 **Disclosures**

323 The authors of this manuscript have nothing to disclose.

324

325 **Conflicts of Interest, Consent and IRB:**

326 We have no conflicts of interest to disclose. Creighton University acknowledged this study as

327 Not Human Subjects Research (InfoEd Record Number: 2004608); the NRD is HIPAA-

328 compliant and publicly available. Therefore, informed consent was deemed unnecessary for this

329 retrospective analysis of de-identified data.

330

331 **Author Contributions:**

332 Ali Al-Salahat: Conception/Design, methodology, literature review, writing manuscript/critical

333 revision

334 Danielle Dilsaver: Design, Statistical Analysis, Writing Manuscript/Critical revision

335 Ali Bin Abdul Jabbar, Conception/Design, Critical revision

336 Abhishek Singh: Conception/ Design, Critical revision

337

338

339

340

341

## 342 **References**

- 343 1. Romi F, Naess H. Spinal Cord Infarction in Clinical Neurology: A Review of  
344 Characteristics and Long-Term Prognosis in Comparison to Cerebral Infarction. *Eur*  
345 *Neurol.* 2016;76(3-4):95-98. doi: 10.1159/000446700.
- 346 2. Stenimahitis V, Fletcher-Sandersjö A, El-Hajj VG, et al. Long-term Outcomes After  
347 Periprocedural and Spontaneous Spinal Cord Infarctions: A Population-Based Cohort  
348 Study. *Neurology.* 2023 Jul 11;101(2):e114-e124. doi:  
349 10.1212/WNL.0000000000207377.
- 350 3. Zalewski NL, Rabinstein AA, Krecke KN, et al. Characteristics of Spontaneous Spinal  
351 Cord Infarction and Proposed Diagnostic Criteria. *JAMA Neurol.* 2019 Jan 1;76(1):56-  
352 63. doi: 10.1001/jamaneurol.2018.2734.
- 353 4. Cheng MY, Lyu RK, Chang YJ, et al. Spinal cord infarction in Chinese patients. Clinical  
354 features, risk factors, imaging and prognosis. *Cerebrovasc Dis.* 2008;26(5):502-8. doi:  
355 10.1159/000155988.
- 356 5. Aucoin VJ, Motyl CM, Novak Z, et al; U.S. Aortic Research Consortium. Predictors and  
357 outcomes of spinal cord injury following complex branched/fenestrated endovascular  
358 aortic repair in the US Aortic Research Consortium. *J Vasc Surg.* 2023 Jun;77(6):1578-  
359 1587. doi: 10.1016/j.jvs.2023.01.205.
- 360 6. MacGillivray TE, Gleason TG, Patel HJ, et al. The Society of Thoracic  
361 Surgeons/American Association for Thoracic Surgery Clinical Practice Guidelines on the  
362 Management of Type B Aortic Dissection. *Ann Thorac Surg.* 2022 Apr;113(4):1073-  
363 1092. doi: 10.1016/j.athoracsur.2021.11.002. Epub 2022 Jan 25. Erratum in: *Ann Thorac*  
364 *Surg.* 2023 Nov;116(5):1120.



- 365 7. Aucoin VJ, Motyl CM, Novak Z, et al; U.S. Aortic Research Consortium. Predictors and  
366 outcomes of spinal cord injury following complex branched/fenestrated endovascular  
367 aortic repair in the US Aortic Research Consortium. *J Vasc Surg.* 2023 Jun;77(6):1578-  
368 1587. doi: 10.1016/j.jvs.2023.01.205.
- 369 8. Gaudino M, Khan FM, Rahouma M, et al. Spinal cord injury after open and endovascular  
370 repair of descending thoracic and thoracoabdominal aortic aneurysms: A meta-analysis. *J*  
371 *Thorac Cardiovasc Surg.* 2022 Feb;163(2):552-564. doi: 10.1016/j.jtcvs.2020.04.126.
- 372 9. Ciceri EF, Opancina V, Pellegrino C, et al. Fibrocartilaginous embolism: a rare cause  
373 leading to spinal cord infarction? *Neurol Sci.* 2023 Jan;44(1):263-271. doi:  
374 10.1007/s10072-022-06398-w.
- 375 10. Thöne J, Hohaus A, Bickel A, et al. Severe spinal cord ischemia subsequent to  
376 fibrocartilaginous embolism. *J Neurol Sci.* 2007 Dec 15;263(1-2):211-3. doi:  
377 10.1016/j.jns.2007.06.009.
- 378 11. Robertson CE, Brown RD Jr, Wijdicks EF, et al. Recovery after spinal cord infarcts: long-  
379 term outcome in 115 patients. *Neurology.* 2012 Jan 10;78(2):114-21. doi:  
380 10.1212/WNL.0b013e31823efc93. Epub 2011 Dec 28.
- 381 12. Focke JK, Seitz RJ. Reversal of Acute Spinal Cord Ischemia by Intravenous  
382 Thrombolysis. *Neurol Clin Pract.* 2021 Dec;11(6):e975-e976. doi:  
383 10.1212/CPJ.0000000000001097.
- 384 13. Haynes J, Shapiro M, Raz E, et al. Intra-arterial thrombolytic therapy for acute anterior  
385 spinal artery stroke. *J Clin Neurosci.* 2021 Feb;84:102-105. doi:  
386 10.1016/j.jocn.2020.11.035.

- 387 14. Isselbacher EM, Preventza O, Hamilton Black J 3rd, et al; Peer Review Committee  
388 Members. 2022 ACC/AHA Guideline for the Diagnosis and Management of Aortic  
389 Disease: A Report of the American Heart Association/American College of Cardiology  
390 Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 Dec 13;146(24):e334-  
391 e482. doi: 10.1161/CIR.0000000000001106.
- 392 15. New PW, McFarlane CL. Survival following spinal cord infarction. *Spinal Cord*. 2013  
393 Jun;51(6):453-6. doi: 10.1038/sc.2013.14.
- 394 16. NRD Overview. Healthcare Cost and Utilization Project (HCUP). Agency for Healthcare  
395 Research and Quality, Rockville, MD. [www.hcup-us.ahrq.gov/nrdoverview.jsp](http://www.hcup-us.ahrq.gov/nrdoverview.jsp). Accessed  
396 April 15, 2024.
- 397 17. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP;  
398 STROBE Initiative. The Strengthening the Reporting of Observational Studies in  
399 Epidemiology (STROBE) statement: guidelines for reporting observational studies.  
400 *Lancet*. 2007 Oct 20;370(9596):1453-7. doi: 10.1016/S0140-6736(07)61602-X. PMID:  
401 18064739.
- 402 18. United States Bureau of Labor Statistics. CPI inflation calculator. Available at:  
403 [https://www.bls.gov/data/inflation\\_calculator.htm](https://www.bls.gov/data/inflation_calculator.htm). Accessed April 15, 2024.
- 404 19. Glasheen, W. P., Cordier, T., et al. (2019). Charlson Comorbidity Index: *ICD-9* Update  
405 and *ICD-10* Translation. *American health & drug benefits*, 12(4), 188–197.
- 406 20. Harrell FE. Regression modeling strategies: With applications to linear models, logistic  
407 and ordinal regression, and survival analysis. 2nd ed. Springer; 2015.
- 408 21. Terkelsen T, Schmitz ML, Simonsen CZ, et al. Thrombolysis in acute ischemic stroke is  
409 associated with lower long-term hospital bed day use: A nationwide propensity score-

- 410 matched follow-up study. *Int J Stroke*. 2016 Oct;11(8):910-916. doi:  
411 10.1177/1747493016654491.
- 412 22. IST-3 collaborative group. Effect of thrombolysis with alteplase within 6 h of acute  
413 ischaemic stroke on long-term outcomes (the third International Stroke Trial [IST-3]): 18-  
414 month follow-up of a randomised controlled trial. *Lancet Neurol*. 2013 Aug;12(8):768-  
415 76. doi: 10.1016/S1474-4422(13)70130-3.
- 416 23. Fekete K, Márton S, Tóth J, et al. Predictors of long-term outcome after intravenous or  
417 intra-arterial recombinant tissue plasminogen activator treatment in the eastern Hungarian  
418 thrombolysis database. *J Stroke Cerebrovasc Dis*. 2015 Jan;24(1):117-24. doi:  
419 10.1016/j.jstrokecerebrovasdis.2014.07.054.
- 420 24. Dubinsky R, Lai SM. Mortality of stroke patients treated with thrombolysis: analysis of  
421 nationwide inpatient sample. *Neurology*. 2006 Jun 13;66(11):1742-4. doi:  
422 10.1212/01.wnl.0000218306.35681.38.
- 423 25. Cheng MY, Lyu RK, Chang YJ, et al. Concomitant spinal cord and vertebral body  
424 infarction is highly associated with aortic pathology: a clinical and magnetic resonance  
425 imaging study. *J Neurol*. 2009 Sep;256(9):1418-26. doi: 10.1007/s00415-009-5126-2.
- 426 26. McClure RS, Brogly SB, Lajkosz K, et al. Epidemiology and management of thoracic  
427 aortic dissections and thoracic aortic aneurysms in Ontario, Canada: A population-based  
428 study. *J Thorac Cardiovasc Surg*. 2018 Jun;155(6):2254-2264.e4. doi:  
429 10.1016/j.jtcvs.2017.11.105.
- 430 27. Gouveia E Melo R, Mourão M, Caldeira D, et al. A systematic review and meta-analysis  
431 of the incidence of acute aortic dissections in population-based studies. *J Vasc Surg*. 2022  
432 Feb;75(2):709-720. doi: 10.1016/j.jvs.2021.08.080.

- 433 28. Chhabra HS, Sharawat R, Vishwakarma G. In-hospital mortality in people with complete  
434 acute traumatic spinal cord injury at a tertiary care center in India-a retrospective  
435 analysis. *Spinal Cord*. 2022 Mar;60(3):210-215. doi: 10.1038/s41393-hfdk021-00657-x.
- 436 29. Vahidy FS, Donnelly JP, McCullough LD, et al. Nationwide Estimates of 30-Day  
437 Readmission in Patients With Ischemic Stroke. *Stroke*. 2017 May;48(5):1386-1388. doi:  
438 10.1161/STROKEAHA.116.016085.
- 439 30. Torvik MA, Nymo SH, Nymo SH, et al. Patient characteristics in sepsis-related deaths:  
440 prevalence of advanced frailty, comorbidity, and age in a Norwegian hospital trust.  
441 *Infection*. 2023 Aug;51(4):1103-1115. doi: 10.1007/s15010-023-02013-y.
- 442 31. Jillella DV, Wisco DR. Infectious causes of stroke. *Curr Opin Infect Dis*. 2019  
443 Jun;32(3):285-292. doi: 10.1097/QCO.0000000000000547.
- 444 32. Dahlberg J, Linder A, Mellhammar L. Use of healthcare before and after sepsis in  
445 Sweden: a case-control study. *BMJ Open*. 2023 Feb 17;13(2):e065967. doi:  
446 10.1136/bmjopen-2022-065967.
- 447 33. Han E, Kim TH, Koo H, et al. Heterogeneity in costs and prognosis for acute ischemic  
448 stroke treatment by comorbidities. *J Neurol*. 2019 Jun;266(6):1429-1438. doi:  
449 10.1007/s00415-019-09278-0.
- 450 34. Kenyon LC, Dulaney E, Montone KT, et al. Varicella-zoster ventriculo-encephalitis and  
451 spinal cord infarction in a patient with AIDS. *Acta Neuropathol*. 1996 Aug;92(2):202-5.  
452 doi: 10.1007/s004010050509.

453

454

455

456 **Figure Legends**

457 **Figure 1.** The effect of age (top row) and comorbidity burden (bottom row) on inpatient  
458 mortality (left column) and length of stay (right column) for hospitalizations of patients with a  
459 spinal cord infarction.

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

**Table 1.** Descriptive characteristics of hospitalizations of patients with a spinal cord infarction

|  |                          |
|--|--------------------------|
| Age, median [IQR]                            | 63 [53-73]               |
| Biological Sex, Weighted N (%)               |                          |
| Male   | 7,377 (58.79)            |
| Female                                       | 5,171 (41.21)            |
| Insurance, Weighted N (%)                    |                          |
| Medicare                                     | 6,578 (52.43)            |
| Medicaid                                     | 1,882 (15)               |
| Private                                      | 3,247 (25.88)            |
| Other  | 821 (6.54)               |
| Unknown                                      | 19 (0.15)                |
| Income Quartile, Weighted N (%)              |                          |
| I  | 3,631 (28.93)            |
| II   | 3,371 (26.87)            |
| III  | 3,134 (24.97)            |
| IV   | 2,268 (18.08)            |
| Unknown                                      | 144 (1.15)               |
| Admission, Weighted N (%)                    |                          |
| Weekend                                      | 2,778 (22.14)            |
| Elective                                     | 2,322 (18.51)            |
| Bed Size, Weighted N (%)                     |                          |
| Small  | 919 (7.33)               |
| Medium                                       | 2,549 (20.32)            |
| Large  | 9,079 (72.36)            |
| Location-Teaching Status, Weighted N (%)     |                          |
| Metropolitan non-teaching                    | 1,296 (10.33)            |
| Metropolitan teaching                        | 10,862 (86.57)           |
| Non-metropolitan                             | 390 (3.11)               |
| Charlson Comorbidity Index, median (IQR)     | 2 (3-7)                  |
| NIHSS score provided, Weighted N (%)         | 464 (3.70)               |
| NIHSS 0-9                                    | 308 (66.37) <sup>a</sup> |
| NIHSS 10-19                                  | 111 (23.85) <sup>a</sup> |
| NIHSS 20-29                                  | 33 (7.06) <sup>a</sup>   |
| NIHSS 30-39                                  | *                        |
| NIHSS 40+                                    | *                        |
| Surgical Aorta Repair during Hospitalization |                          |
| Open   | 68 (0.54)                |
| Percutaneous                                 | 0 (0)                    |
| Percutaneous endoscopic                      | *                        |

<sup>a</sup> Categorical characteristics are presented as weighted count and percent. Continuous characteristics are presented as median and interquartile range.

<sup>b</sup> NIHSS = National Institutes of Health Stroke Scale.

<sup>c</sup> An ^ indicates the percent is out of hospitalizations with a provided National Institutes of Health Stroke Scale score (Weighted N = 464).

<sup>d</sup> An \* indicates the results could not be presented per the NRD Data Use Agreement as there were 10 or less hospitalizations)

484

**Table 2.** Hospitalizations outcomes by whether patients with spinal cord infarctions received thrombolysis

|                                | Unadjusted          |                      |                  |       | Adjusted         |       |
|--------------------------------|---------------------|----------------------|------------------|-------|------------------|-------|
|                                | Thrombolysis:<br>No | Thrombolysis:<br>Yes | Ratio (95% CI)   | p     | Ratio (95% CI)   | p     |
| Index hospitalization          |                     |                      |                  |       |                  |       |
| Inpatient mortality, %         | 14.60               | 19.45                | 1.41 (0.71-2.83) | 0.330 | 1.46 (0.77-2.77) | 0.249 |
| Length of stay, days           | 13.12               | 10.83                | 0.83 (0.61-1.11) | 0.207 | 0.86 (0.65-1.14) | 0.294 |
| Cost, \$                       | 53,551              | 51,869               | 0.97 (0.78-1.18) | 0.747 | 1.01 (0.84-1.20) | 0.941 |
| Discharge disposition, %       |                     |                      |                  |       |                  |       |
| Routine                        | *                   | *                    | *                | *     | *                | *     |
| Transfer (Short-term hospital) | *                   | *                    | *                | *     | *                | *     |
| Transfer (Other)               | 48.41               | 56.24                | 1.36 (0.83-2.26) | 0.217 | 1.32 (0.79-2.21) | 0.284 |
| All-cause Readmission          |                     |                      |                  |       |                  |       |
| 30-day, %                      | 22.44               | 12.93                | 0.55 (0.27-1.15) | 0.114 | 0.58 (0.28-1.20) | 0.142 |
| 90-day, %                      | 34.21               | 15.11                | 0.34 (0.16-0.75) | 0.008 | 0.35 (0.16-0.77) | 0.009 |

<sup>a</sup> An \* indicates the results could be presented per the NRD Data Use Agreement as there were 10 or less hospitalizations.

<sup>b</sup> Transfer (Other) includes skilled nursing facility, intermediate care facility, and other types of facilities

485

486

487

**Table 3.** Hospitalization outcomes by whether patients with a spinal cord infarction suffered from an aneurysm or dissection

|                                | Unadjusted |        |                  |         | Adjusted         |         |
|--------------------------------|------------|--------|------------------|---------|------------------|---------|
|                                | No         | Yes    | Ratio (95% CI)   | p       | Ratio (95% CI)   | p       |
| Index hospitalization          |            |        |                  |         |                  |         |
| Inpatient mortality, %         | 12.48      | 19.98  | 1.75 (1.51-2.04) | < 0.001 | 1.52 (1.30-1.78) | < 0.001 |
| Length of stay, days           | 12.35      | 15.08  | 1.22 (1.12-1.33) | < 0.001 | 1.19 (1.09-1.29) | < 0.001 |
| Cost, \$                       | 42,558     | 93,643 | 2.20 (2.07-2.33) | < 0.001 | 2.18 (2.06-2.31) | < 0.001 |
| Discharge disposition, %       |            |        |                  |         |                  |         |
| Routine                        | 18.57      | 12.69  | 0.64 (0.54-0.75) | < 0.001 | 0.88 (0.74-1.05) | 0.154   |
| Transfer (Short-term hospital) | 3.70       | 3.90   | 1.06 (0.80-1.40) | 0.704   | 1.14 (0.85-1.52) | 0.379   |
| Transfer (Other)               | 48.45      | 48.61  | 1.01 (0.90-1.13) | 0.914   | 0.89 (0.79-1.00) | 0.057   |
| All-cause Readmission          |            |        |                  |         |                  |         |
| 30-day, %                      | 19.95      | 24.13  | 1.28 (1.10-1.48) | 0.002   | 1.16 (0.99-1.36) | 0.067   |
| 90-day, %                      | 32.88      | 37.04  | 1.20 (1.03-1.40) | 0.017   | 1.06 (0.90-1.24) | 0.472   |

<sup>a</sup> An \* indicates the results could be presented per the NRD Data Use Agreement as there were 10 or less hospitalizations.

<sup>b</sup> Transfer (Other) includes skilled nursing facility, intermediate care facility, and other types of facilities

488

489

490




491

**Table 4.** Hospitalization outcomes by whether patients with a spinal cord infarction suffered from sepsis

|                                | Unadjusted |              |                  |         | Adjusted         |         |
|--------------------------------|------------|--------------|------------------|---------|------------------|---------|
|                                | Sepsis: No | Sepsis : Yes | Ratio (95% CI)   | p       | Ratio (95% CI)   | p       |
| Index hospitalization          |            |              |                  |         |                  |         |
| Inpatient mortality, %         | 22.22      | 25.00        | 1.17 (0.65-2.08) | 0.599   | 3.15 (2.64-3.75) | < 0.001 |
| Length of stay, days           | 11.59      | 20.06        | 1.73 (1.61-1.86) | < 0.001 | 1.44 (1.34-1.55) | < 0.001 |
| Cost, \$                       | 46,721     | 86,215       | 1.85 (1.73-1.97) | < 0.001 | 1.54 (1.45-1.65) | < 0.001 |
| Discharge disposition, %       |            |              |                  |         |                  |         |
| Routine                        | 19.33      | 8.22         | 0.37 (0.30-0.47) | < 0.001 | 0.41 (0.33-0.52) | < 0.001 |
| Transfer (Short-term hospital) | 3.99       | 2.92         | 0.72 (0.53-0.99) | 0.044   | 0.67 (0.48-0.93) | 0.015   |
| Transfer (Other)               | 48.50      | 48.51        | 1.00 (0.88-1.14) | 0.993   | 1.00 (0.87-1.14) | 0.975   |
| All-cause Readmission          |            |              |                  |         |                  |         |
| 30-day, %                      | 19.42      | 28.43        | 1.65 (1.39-1.96) | < 0.001 | 1.42 (1.19-1.70) | < 0.001 |
| 90-day, %                      | 31.74      | 44.21        | 1.70 (1.43-2.02) | < 0.001 | 1.48 (1.23-1.77) | < 0.001 |

An \* indicates the results could be presented per the NRD Data Use Agreement as there were 10 or less hospitalizations.

Transfer (Other) includes skilled nursing facility, intermediate care facility, and other types of facilities



**Figure 1.** The effect of age (top row) and comorbidity burden (bottom row) on inpatient mortality (left column) and length of stay (right column) for hospitalizations of patients with a spinal cord infarction.