Performance of LFSPRO *TP53* germline carrier risk predictions compared to standard genetic counseling
practice on prospectively collected probands

Jessica L. Corredor[1*], MS, CGC, Ruonan Li, PhD[2*], Elissa B. Dodd-Eaton[2], MPH, Jacynda Casey[1], MS,
CGC, Ashley H. Woodson[1], MS, CGC, Nam H. Nguyen[2,3], PhD, Gang Peng, PhD[4], Angelica M.
Gutierrez[5], MS, Banu K. Arun[1,5$], MD, Wenyi Wang[2$], PhD

1.   Department of Clinical Cancer Genetics, The University of Texas MD Anderson Cancer Center,
Houston, TX, USA

2.   Department of Bioinformatics and Computational Biology, The University of Texas MD Anderson
Cancer Center, Houston, TX, USA

3.   Department of Statistics, Rice University, Houston, TX, USA

4.   Department of Medical and Molecular Genetics, Indiana University School of Medicine,
Indianapolis, IN, USA

5.   Department of Breast Medical Oncology, The University of Texas MD Anderson Cancer Center,
Houston, TX, USA

*First Authors contributed equally.

$ Last authors contributed equally.

Correspondence to: Wenyi Wang at wwang7@mdanderson.org, Banu Arun at barun@mdanderson.org

Abstract

Genetic counseling and testing for germline mutations are essential for identifying individuals at increased

risk for cancer. Pathogenic variants in *TP53* are diagnostic of Li-Fraumeni syndrome (LFS), a highly

penetrant disorder with diverse, early-onset tumors. Current clinical guidelines, such as Chompret and

Classic criteria, provide frameworks for identifying individuals at risk for likely pathogenic/pathogenic

TP53 variants; however, genetic counselors often encounter patients with features concerning for LFS

that do not clearly meet established criteria, creating challenges for risk assessment and testing

decisions. We evaluated whether LFSPRO, a Mendelian, family-history-based model that estimates the

individual's probability of harboring a deleterious *TP53* variant, improves carrier identification relative to

guideline criteria.

In a prospectively collected cohort of 182 probands who underwent clinical genetic counseling and

germline *TP53* testing, LFSPRO showed superior discrimination compared with Chompret criteria, with

higher sensitivity (81% vs. 33%) and specificity (88% vs. 65%) and improved predictive values (PPV 0.53

vs. 0.14; NPV 0.96 vs. 0.85). Receiver operating characteristic analysis confirmed strong discriminatory

performance (AUC=0.88). Calibration analysis using observed-to-expected ratios indicated good

agreement between predicted and observed carrier frequencies (Observed/Expected=1.07).

These findings demonstrate that LFSPRO outperforms traditional guideline-based criteria for identifying

TP53 mutation carriers in real-world clinical settings. By providing quantitative, well-calibrated carrier

probabilities rather than binary classifications, LFSPRO can enhance genetic counseling and support

testing decisions, particularly for individuals who do not clearly meet existing criteria.

Introduction

Li-Fraumeni syndrome (LFS [MIM: 151623]) is a rare hereditary cancer predisposition syndrome caused by germline likely pathogenic/pathogenic variants (LP/P variants) in the *TP53* (MIM: 191170), a key tumor suppressor gene that safeguards the genome from mutations[1–3]. Individuals with LFS face an increased risk of developing various cancers throughout their lifetime. The most common cancers seen in LFS families are sarcomas, breast cancer, leukemia, brain tumors, and adrenal cortical carcinoma[4,5]. The lifetime cancer risk is approximately 93% for women and 73% for men, with a 50% chance that cancers occur prior to the age of 40[6,7]. Cancer survivors in this population are at higher risk of developing additional primary cancers and treatment-related secondary cancers[7], underscoring the importance of genetic counseling in guiding early detection, cancer risk assessment, and lifelong monitoring to better manage their health outcomes.

However, effective clinical management of LFS is challenging due to its rare nature, the overlap with other hereditary cancer syndromes, and the variable penetrance and phenotypic expression across families[8]. Although early cancer detection, as demonstrated by the 2011 comprehensive Toronto screening protocol, can dramatically improve cancer survival rates in children and adults[9], predicting who will carry a *TP53* LP/P variant remains difficult, making it challenging to start screening at the appropriate time. Currently, risk prediction relies on clinical classification schemes such as Li-Fraumeni classic criteria[10], Birch criteria[5], Eeles Criteria[11], and Chompret criteria[6,12,13]. These schemes include proband's cancer history, age of cancer onset, and limited family cancer history in first- and second-degree relatives to guide recommendations for genetic testing for LFS. However, these criteria do not provide the detailed risk assessment often needed in genetic counseling. In cancer-focused genetic counseling sessions, genetic counselors (GCs) use risk prediction models to estimate a patient's lifetime cancer risk, determine eligibility for surveillance and chemoprevention, and assess the likelihood of having other hereditary cancer predisposition syndromes, such as hereditary breast and ovarian cancer syndrome (MIM: 604370) and Lynch syndrome (MIM: 120435)[14]. However, such risk models did not previously exist for LFS. Given the heterogeneity in cancer presentation of LFS and the significant impact of early identification on

personalized surveillance and treatment strategies, a comprehensive risk model is essential to help GCs

and other healthcare providers communicate more effectively with patients.

To address this gap, Peng *et al*[15] developed LFSPRO, a comprehensive statistical framework designed

to predict both the likelihood of having a *TP53* LP/P variant and the risk of developing LFS-related

cancers. Several key steps were undertaken to move LFSPRO from model development to clinical

applicability (**Fig. 1a**). The development process began by identifying the clinical problem, followed by

initial model development using research datasets[15]. These research datasets differ from clinically

ascertained data in that they often include more detailed personal and family histories and regular follow-

ups, while clinical data tends to have limited information and reflect real-world conditions. To ensure real-

world applicability, the models were trained on clinical data and validated using both internal and external

datasets. LFSPRO was then further validated using retrospective research-based (RB) multicenter

cohorts, which provided penetrance estimates for breast cancer, sarcoma, and other cancers in LFS

patients[16]. Following this, the model was further validated with a single institution's ascertained data,

which was a retrospective clinical-based (RCB) data and collected from genetic counseling sessions,

reflecting real-world clinical settings with potentially less comprehensive personal and family history

information compared to research datasets[17]. This clinical validation is critical for ensuring the model's

applicability in genetic counseling and clinical decision-making in real-world diverse clinical settings.

Subsequently, an R Shiny app (LFSPROShiny[18]) was developed to make the model accessible to GCs

and healthcare providers in a user-friendly format.

While previous studies have demonstrated LFSPRO's strong performance in research and retrospective

clinical datasets, its real-world clinical utility as a complementary tool to existing LFS risk prediction

criteria remains to be fully evaluated. In this prospective study, we assessed the accuracy of LFSPRO in

predicting *TP53* germline variant carrier status compared to established criteria in a real-time clinical

genetic counseling setting. To further evaluate its practical value as a decision-support tool, four

experienced GCs were invited to use LFSPRO following standard genetic counseling sessions with

patients suspected of having LFS. Counselors reviewed the model's output alongside their own clinical

assessments and completed a structured survey providing their detailed feedback, including the model's

concordance with clinical judgment and its perceived usefulness in patient communication and decision-

making.

Methods

Study Design and Study Population

This prospective study was designed to assess LFSPRO's clinical utility in a real-world genetic counseling setting at the University of Texas MD Anderson Cancer Center (MDACC). Building upon previous validations that demonstrated LFSPRO's strong performance in research-based and retrospective clinical datasets, this study aims to evaluate its performance in predicting *TP53* LP/P germline variant carrier status compared to established criteria in a prospective clinical cohort, and to assess GCs' experience with implementing LFSPRO as part of their standard genetic counseling practice.

From December 2021 to March 2025, probands who underwent genetic counseling at MDACC and were identified as having clinical features concerning for LFS were included. Patients were eligible if they either: (1) met Classic and/or Chompret criteria for LFS testing, or (2) were deemed concerning for LFS by GCs based on their personal/family history, despite not meeting existing criteria. The latter category included individuals with early-onset LFS-spectrum tumors but small or limited family history, somatic testing suggestive of germline *TP53* variants, multiple primary cancers, and/or extensive family history of childhood and/or rare cancers.

Data Collection

After receiving standard genetic counseling with an MDACC GC, patients identified as concerning for LFS were flagged in Progeny, which stores the internal family history database. The study team then completed a comprehensive chart review of all identified patients and assessed them using LFSPRO. Data collected included LFSPRO-predicted *TP53* LP/P variant carrier risk, whether the patient met Chompret and/or Classic criteria, decision to test, and genetic test results when available. Additional data documented for each participant included demographic information (age, sex assigned at birth), personal cancer history (type of cancer, age at diagnosis, multiple primaries), and family cancer history in first- and second-degree relatives.

Four MDACC GCs who most frequently see patients concerning for LFS were invited to use LFSPRO on their "concerning for LFS patients" after completing their standard genetic counseling session. After each

instance of running LFSPRO, these counselors completed a survey regarding their experience and

assessment of LFSPRO's clinical utility for that specific patient case. A detailed illustration of the study

can be seen in **Fig. 1b**. A complete survey can be seen in **Supplemental Materials.**

Current Criteria for LFS Risk Assessment

LFS diagnosis is primarily guided by established clinical criteria designed to identify individuals with a

high likelihood of carrying pathogenic *TP53* mutations. The most widely used criteria in genetic

counseling include the Classic LFS criteria and the Chompret criteria (see **Table 1**). The Classic criteria,

originally defined by Li and Fraumeni in 1988, have high specificity (91.0-98.1%) but low sensitivity (25.0-

40.0%), making them the most stringent approach[19]. Chompret criteria were developed by Chompret et

al and revisited in 2009 and 2015. The revised Chompret criteria offer improved sensitivity (83%) while

maintaining reasonable specificity (47%)[13]. These criteria guide GCs in evaluating a proband's personal

cancer history, age of cancer onset, and limited cancer patterns in first- and second-degree relatives to

determine eligibility for genetic testing. While both frameworks have been valuable in identifying LFS

candidates, none of the probands in our study cohort met the Classic LFS criteria, highlighting potential

limitations in real-world clinical settings. Consequently, our analysis focused exclusively on comparing

LFSPRO performance against the Chompret criteria, which has become the more clinically relevant

standard in contemporary practice.

LFSPRO Statistical Framework

LFSPRO was developed as a Mendelian risk prediction model to estimate the probability that an

individual carries a deleterious *TP53* mutation based on detailed family cancer history[15]. The model

adopts a Bayesian formulation in which the carrier probability is obtained by updating the population

prevalence with the likelihood of the observed pedigree and cancer diagnoses:

144
$$P_r(G_0|H) = \frac{P_r(G_0) \times P_r(H|G_0)}{\sum_g P_r(g) \times P_r(H|g)}$$

Here, G_0 denotes the genotype of the counselee, H represents the family cancer history, and $g \in \{0,1\}$

corresponds to the possible genotype states (wildtype or mutation carrier, including both heterozygous

and homozygous mutation genotypes). The prior probability $P_r(G_0)$ is set to the population prevalence of

TP53 mutations (0.0006)[15], following the Hardy-Weinberg equilibrium for homozygotes. The likelihood

term $P_r(H|G_0)$ represents the probability of observing the family cancer history given the counselee's

genotype, updated from integrating over all possible genotype configurations of relatives while

incorporating Mendelian inheritance and age-specific cancer penetrance. Because the number of

genotype configurations grows exponentially with pedigree size, LFSPRO employs the Elston-Stewart or

peeling algorithm[20], which recursively factors the pedigree into anterior (ancestral) and posterior

(descendant) components conditional on a pivot individual, and marginalizes over unobserved genotypes

as individuals are peeled off. This recursion enables efficient likelihood evaluation in extended pedigrees

without enumerating all possible genotype states. Importantly, LFSPRO also models de novo *TP53*

mutations, which arise spontaneously in the germline even when both parents are noncarriers. Such

events account for an estimated 7-20%[21] of LFS cases and are therefore clinically significant. In the

model, they are represented by a transmission probability of 2α, where α is the per-allele mutation rate.

By incorporating both inherited and de novo events, LFSPRO yields more accurate carrier probability

estimates across diverse pedigrees.

Building on the original formulation, the model has been extended to support prospective cancer risk

prediction. The Cancer-Specific (CS) model[22] employs a competing risk approach with cause-specific

hazard functions to estimate the probability of different cancer types as the first primary tumor, and the

Multiple Primary Cancer (MPC) model[23] uses recurrent event models based on non-homogeneous

Poisson processes to capture the elevated risk of subsequent malignancies. By providing individualized,

quantitative risk estimates rather than binary classifications, LFSPRO and its extensions enable more

refined genetic counseling compared with traditional clinical criteria. Validation across multiple

independent cohorts has demonstrated superior sensitivity and specificity for identifying *TP53* mutation

carriers[16,17,24], underscoring its clinical utility in LFS family risk assessment and management.

Clinical Implementation

These statistical models are integrated into LFSPROShiny[18], an interactive R/Shiny application that

provides genetic counselors and healthcare providers with a user-friendly interface for inputting patient

information and generating individualized risk assessments.

For this study, GCs utilized LFSPROShiny to facilitate clinical implementation. The interface allows GCs

to efficiently input patient information from MDACC electronic health records to compute the carrier

probabilities and generate individualized risk assessments. While GCs can select between the CS and

MPC models for specific cases, the MPC model was set as the default based on its better performance

compared to the CS model in prior validation studies[15,17]. Furthermore, we used a binary classification

threshold of 0.2 for LFSPRO risk scores, as established in prior work[15], where probands scoring $\geq$0.2

were classified as screening positive for potential *TP53* LP/P variant carrier status. This threshold

achieved good trade-offs between sensitivity and specificity on multiple LFS datasets[15].

Survey Design and Administration

To evaluate GCs' experience using LFSPRO in clinical practice, we developed a structured REDCap

(Research Electronic Data Capture)[20,21] survey instrument addressing four key domains: (1) concordance

between LFSPRO predictions and standard clinical assessment, (2) additive value of LFSPRO beyond

conventional genetic counseling, (3) comfort level with communicating LFSPRO results to patients, and

(4) practical applications of LFSPRO-derived risk predictions in patient counseling. GCs taking these

surveys could select to complete a survey for 1) a patient undergoing *TP53* germline testing, 2) a patient

with a *TP53* germline variant that wanted to utilize LFSPRO's cancer risk prediction, or 3) a combination

of *TP53* germline testing, risk assessment, and cancer risk prediction. Each survey contained both

structured quantitative measures and open-ended response fields for GCs to provide further feedback

on their use of LFSPRO.

Statistical Analysis

The performance of LFSPRO and Chompret criteria was evaluated using multiple complementary

approaches. Diagnostic accuracy was assessed through sensitivity, specificity, positive predictive value

(PPV), and negative predictive value (NPV) with 95% confidence intervals. Sensitivity measured the

proportion of true *TP53* mutation carriers correctly identified by each model, while specificity measured

the proportion of individuals without *TP53* mutations correctly identified as negative. PPV assessed how

often a positive prediction was correct, and NPV assessed how often a negative prediction was correct,

providing measures of clinical reliability for each prediction outcome. Discriminatory ability was quantified

using receiver operating characteristic (ROC) curve analysis, with area under the curve (AUC) values

and 95% confidence intervals calculated for both LFSPRO and Chompret criteria. An AUC of 0.5

represents no discriminatory ability (equivalent to random chance), while 1.0 represents perfect

discrimination.

Statistical significance of associations was assessed using two-sides Fisher's exact test, which was

selected due to the limited sample size with cell counts below 5 in some contingency table cells. This

non-parametric test was used to evaluate: (1) the association between model-based classifications

(LFSPRO and Chompret criteria) and actual *TP53* mutation status, and (2) the association between

prediction methods and GCs' clinical judgment. The null hypothesis in each case was that no association

exists between the compared variables, with a *p*-value < 0.05 considered statistically significant.

Confusion matrices were visualized as heatmaps to illustrate the distribution of true positives, false

positives, true negatives, and false negatives for each prediction method. For the GC survey data,

descriptive statistics were calculated as percentages for categorical variables.

Results

Cohort Description

From December 2021 to March 2025, a total of 178 probands were included in this concordance study.

The cohort comprised 115 (64.6%) assigned female at birth and 63 (35.4%) assigned male at birth, with

a mean age of 30 years. Among the 148 probands who completed *TP53* genetic testing, 23 (15.5%)

tested positive for a LP/P *TP53* germline variant, including one with a mosaic variant identified in two

separate tissue types, and one with low variant allele frequency (VAF) suggestive of possible mosaicism.

The majority (n=125, 84.5%) tested negative for germline *TP53* variants, though this group included one

individual with a *TP53* variant at 13% VAF that could not be detected in a second tissue source, and

another with a variant of uncertain significance in *TP53* that was likely mosaic. Four individuals had

genetic testing but with results still pending, while 26 probands had no genetic testing results because

they either declined testing or remained undecided. Since the Bayesian Mendelian model used in

LFSPRO does not account for mosaicism, probands with possible mosaic *TP53* variants were excluded

from all downstream analysis. See **Table 2** for a comprehensive summary of proband characteristics

stratified by genetic testing results.

This prospectively collected clinical cohort represents a unique dataset of individuals flagged by GCs as

concerning for LFS in real-world practice. The 15.5% mutation detection rate and inclusion of complex

cases with mosaicism and uncertain variants provides an ideal testing set for evaluating LFSPRO's

clinical utility.

LFSPRO Outperforms Existing Criteria

Among individuals with germline LP/P *TP53* variants (n=21), 15 (71.4%) had at least one primary cancer

and 7 (28.6%) had at least two primary cancers. Of these individuals, only 7 (33%) met Chompret criteria

for genetic testing, while 17 (81.0%) were identified by LFSPRO with a risk score (carrier probability) $\geq$

0.2. In the mutation-negative group (n=123), 112 (91.1%) had at least one primary cancer and 9 (7.3%)

had at least two primary cancers. Within this group, 43 (35.0%) met Chompret criteria while only 15

(12.2%) had an LFSPRO risk score $\geq$ 0.2. Alternatively, 80 (65.0%) individuals did not meet Chompret

criteria, and 108 (87.8%) had LFSPRO scores below the 0.2 threshold. See **Fig. 2a** for detailed confusion

matrices. Also, a two-sided Fisher's exact test analysis confirmed LFSPRO's statistically significant

concordance with genetic testing results ($p<0.001$), whereas Chompret criteria failed to demonstrate

significant association ($p=1.00$). A corresponding Sankey diagram (**Fig. 2b)** further illustrates the

classification flow of probands by LFSPRO and Chompret criteria against genetic testing results.

LFSPRO demonstrated significantly better discriminatory performance over the conventional Chompret

criteria for identifying *TP53* LP/P mutation carriers. LFSPRO achieved higher sensitivity of 0.810 (95%

CI: 0.600-0.923) compared to Chompret's 0.333 (95% CI: 0.172-0.546), and better specificity of 0.878

(95% CI: 0.809-0.925) versus 0.650 (95% CI: 0.563-0.730), as illustrated in **Fig. 2c**. Importantly, LFSPRO

demonstrated substantially higher PPV at 0.531 (95% CI: 0.364-0.691) compared to Chompret's 0.140

(95% CI: 0.070-0.262), indicating greater clinical utility for correctly identifying true mutation carriers.

Similarly, LFSPRO's NPV of 0.964 (95% CI: 0.912-0.986) outperformed Chompret's 0.851 (95% CI:

0.765-0.909), providing greater confidence in ruling out *TP53* mutations when classified as negative.

ROC curve as shown in **Fig. 2d** further demonstrated LFSPRO's discriminatory ability with AUC scores

of 0.88 (95% CI: 0.82-0.95) for MPC model, and 0.84 (95% CI: 0.77-0.91) for CS model. Calibration

analyses using observed-to-expected (O/E) ratios further supported model validity. The MPC model,

which was used as the default for carrier probability estimation based on prior validation studies,

demonstrated good calibration (O/E 1.07, 95% CI: 0.80-1.34), with close agreement between predicted

and observed carrier frequencies. The CS model showed modest underestimation of carrier probabilities

(O/E 1.46, 95% CI: 1.09-1.84).

These findings demonstrate that LFSPRO outperforms Chompret criteria in identifying individuals with

TP53 LP/P variants, showing enhanced ability to predict true positives while reducing false positives, and

strong alignment with genetic testing results. This suggests LFSPRO's potential to assist in clinical

decision-making and assist patients' decisions on genetic testing.

Genetic Counselors Report Positive Experience with LFSPRO

Beyond comparing LFSPRO with established clinical criteria, we evaluated its concordance with GC's

clinical judgment, the actual decision-making standard in practice. Four experienced GCs completed 64

surveys after counseling sessions. None of the probands in these sessions met Classic LFS criteria and

only 20.3% met Chompret criteria. In practice, however, genetic counselors recommended germline

testing for 77% of probands, underscoring that current criteria fail to capture many individuals whom

clinicians consider appropriate for testing. Within these cases, LFSPRO provided additional decision

support in 16%, highlighting its potential to complement clinical judgment when guideline frameworks

alone are insufficient. Furthermore, GCs indicated they were comfortable sharing LFSPRO results with

patients in 75% of cases, most often providing generalized summaries (39.1%) or specific risk estimates

(35.9%), as shown in **Fig. 3a**.

Fisher's exact test demonstrated a significant association between LFSPRO predictions and GCs' clinical

judgment (p=0.01), with differential agreement rates when LFSPRO predicted above the cutoff (44.4%)

versus below the cutoff (86.0%). In contrast, Chompret criteria showed no significant association with

counselor intuition (p=1.0), yielding similar agreement rates regardless of whether criteria were met (**Fig.**

3b). Importantly, GC experience and clinical judgement were not incorporated as features in the LFSPRO

model, making this concordance an independent validation of the model's ability to capture clinically

relevant risk patterns through objective variables alone. Together, these findings suggest that LFSPRO

is usable, aligns more closely with clinical expertise than guideline criteria, and holds promise for

integration into routine practice to support personalized risk assessment in LFS. Additional qualitative

feedback from free-response comments further highlighted counselors' perspectives on LFSPRO's

clinical implementation, with representative quotations presented in **Table 3**.

Discussion

This prospective study provides important and timely clinical validation of LFSPRO, a personalized

cancer risk prediction tool for Li-Fraumeni Syndrome. Compared to the widely used Chompret and

Classic criteria, LFSPRO demonstrated strong predictive performance, with significantly higher

sensitivity, specificity, PPV, and NPV. AUC analysis further confirmed LFSPRO's strong discriminatory

ability, and its predictions were significantly concordant with genetic testing results, in contrast to

Chompret criteria which showed no significant concordance. These findings position LFSPRO as a robust

clinical tool for identifying individuals at risk for carrying LP/P *TP53* variants.

Chompret and Classic criteria have long guided genetic testing for LFS and are endorsed in the National

Comprehensive Cancer Network (NCCN) guidelines. However, these criteria rely on specific and limited

clinical features and can fail to capture individuals with atypical or limited family histories. As a result,

patients who would benefit from testing may be overlooked, particularly when insurance coverage is tied

to satisfying formal guideline criteria. LFSPRO potentially addresses this gap by generating

individualized, probabilistic estimates of carrier mutation status and specific cancer risk based on detailed

family and personal history, offering a more flexible and precise approach that may facilitate access to

testing and lifelong screening protocols.

Beyond its diagnostic accuracy, this study assessed LFSPRO's impact on clinical decision-making

through 64 genetic counseling surveys. In the majority of cases, counselors reported the patient (77%)

did not meet standard testing criteria but were still recommending testing based on clinical judgment, with

LFSPRO providing additional decision support in 16% of these cases. Importantly, LFSPRO predictions

showed significant concordance with GC clinical judgment while current standard criteria did not. These

findings underscore LFSPRO's potential to complement clinical expertise and facilitate more

personalized genetic counseling, particularly in cases that fall outside established testing frameworks.

LFSPRO also offers longitudinal cancer risk estimates at 5-, 10-, and 15-year intervals, enabling GCs

and other providers to deliver more personalized risk counseling. These individualized projections can

help patients better anticipate their likelihood of carrying a *TP53* mutation or developing cancer, and may

assist in making decisions about genetic testing and surveillance. Personalized risk communication has

been shown to improve patient understanding and promote informed decision-making[25]. In related clinical

contexts, more accurate risk estimates have also been associated with reduced anxiety and cancer worry,

particularly in patients with high baseline anxiety[26]. We expect that LFSPRO has similar potential to

improve not only clinical accuracy but also the patient experience in LFS counseling.

Study Limitations

Our study highlights the improved sensitivity and specificity of LFSPRO over existing clinical testing

guidelines; however, several limitations remain. LFSPRO produced both false-positive and false-negative

predictions. In particular, 3 of the 15 false positives involved unaffected family members being test for a

known familial *TP53* variant. LFSPRO assigned a ~50% risk based on Mendelian inheritance, though

these individuals tested negative. These cases may inflate the false-positive rate and underestimate the

model's true performance in diagnostic testing contexts. Additionally, one of the false positives had a

reported familial LP/P variant in other hereditary cancer genes (NF1), likely contributing to a strong family

history misattributed to *TP53* by LFSPRO. Among the four false negatives, one individual carried the

c.638G>A founder mutation, which is associated with reduced cancer risk. Finally, LFSPRO relies on

patient-reported personal and family history collected in clinical settings. Its accuracy may be limited in

cases with incomplete family histories or when a de novo *TP53* mutation occurs in an individual without

prior cancer or sufficient family history. These limitations are shared by current clinical testing guidelines

and highlight opportunities for further refinement of LFSPRO to improve its utility in diverse clinical

scenarios.

Future Directions

This study demonstrates that genetic counselors feel comfortable using LFSPRO in clinical practice and

find it particularly helpful for guiding testing decisions in families not meeting standard criteria. To broaden

its impact, future work should evaluate its usability among non-genetics providers, particularly in

community or resource-limited settings where expertise in LFS may be scarce, and incorporate structured

training to support implementation. The results presented in this work, including LFSPRO's strong

performance compared to existing criteria and alignment with clinical judgement, may also help

encourage patients, their family members, and providers to participate in future clinical trials that

incorporate LFSPRO.

In parallel, methodological advances continue to expand the underlying statistical framework. A new

unified Bayesian semi-parametric model jointly accommodates recurrent and competing risks and

extends applicability to broader cancer survivor populations[27]. Future efforts will focus on validating this

model in large independent cohorts and integrating it into LFSPRO, thereby enhancing its clinical utility

across diverse settings.

Conclusions

In summary, this prospective study demonstrates LFSPRO's real-world clinical accuracy in identifying

TP53 mutation carriers and strong alignment with GCs' clinical judgment. The tool's potential integration

into current standard clinical practice can potentially facilitate GCs' work by providing personalized risk

estimates that enhance decision-making and support effective communication with patients and family,

particularly in complex cases where traditional criteria provide limited guidance.

Data and code availability

The LFSPRO software is publicly available on GitHub at https://github.com/wwylab/LFSPRO. The

LFSPROShiny interactive application used in this study is open-source and available at

https://github.com/wwylab/LFSPRO-ShinyApp, with a live version hosted at

https://namhnguyen.shinyapps.io/lfspro-shinyapp-master/.

De-identified aggregate data supporting the findings of this study are available from the corresponding

authors upon reasonable request and subject to institutional review board approval. Genetic counselor

survey data and analysis script are available from the corresponding authors upon reasonable request,

with participant identifiers removed to protect confidentiality.

Acknowledgements

This study was supported by grants from Cancer Prevention and Research Institute of Texas (RP200383)

and the National Institutes of Health (R01CA239342, P30CA016672).

Author contribution

Conceptualization, B.K.A and W.W.; methodology, software, N.H.N., P.G., and W.W.; data curation,

investigation, J.L.C., J.C., and A.W.; resources, A.G., and B.K.A., formal analysis, J.L.C., R.L., and

E.B.D.; writing-original draft, J.L.C., and E.B.D; writing-review & editing, J.L.C., R.L., B.K.A., and W.W.,

visualization, R.L., supervision, B.K.A., and W.W., funding acquisition, B.K.A., and W.W.

Declaration of interests

The authors declare that there are no potential conflicts of interest.

Web resources

National Comprehensive Cancer Network (NCCN) Guidelines:

https://www.nccn.org/guidelines/

Figure 1. LFSPRO Development and Clinical Validation Study Design.

(a) The systematic development and validation cycle for clinical risk prediction models, progressing from initial planning and model development through clinical tools, validation phases, and maintenance. **(b)** Patients concerning for Li-Fraumeni Syndrome (LFS) were first evaluated by genetic counselors using standard clinical criteria (e.g., Chompret or Classic) and their clinical experience. Probands identified as potentially mutation carriers underwent *TP53* genetic testing. In parallel, genetic counselors (GCs) ran the LFSPRO tool to generate *TP53* carrier probability and cancer risk predictions. GCs completed a post-session survey evaluating LFSPRO's alignment with their clinical judgment and its potential utility. Predicted results from LFSPRO were then compared with actual genetic testing outcomes to assess concordance.

Figure 2. Comparative analysis of LFSPRO and Chompret criteria performance.

(a) Confusion matrices for LFSPRO and Chompret criteria with associated Fisher's exact test p-values.

(b) Sankey diagrams showing the flow relationships between prediction methods and genetic testing

results. **(c)** Performance metric comparison between LFSPRO and Chompret criteria across sensitivity,

specificity, positive predictive value (PPV), and negative predictive value (NPV). **(d)** ROC curves

comparing discriminatory performance of LFSPRO models and Chompret criteria. The red point indicates

the MPC model's performance at the 0.2 threshold.

Figure 3. Statistical Analysis of GC Survey Results.

(a) Genetic counselors' preferred methods for communicating LFSPRO results to patients. **(b)** Contingency table analysis comparing between LFSPRO and Chompret Criteria with genetic counselor clinical judgment using two-sided Fisher's exact test.

412 **Table 1. Description of Li-Fraumeni Syndrome Classic and Chompret Criteria**

Classic LFS criteria
A sarcoma diagnosed before age 45, AND a first-degree relative (i.e., parent, sibling or child) with any cancer before age 45, AND a first-degree relative or second-degree relative (i.e., grandparent, aunt or uncle, niece or nephew, or grandchild) with any cancer before age 45 or a sarcoma at any age
Chompret criteria (2015)
A tumor that belongs to the LFS tumor spectrum (i.e., soft-tissue sarcoma, osteosarcoma, pre-menopausal breast cancer, CNS (central nervous system) tumor, adrenocortical carcinoma) before age 46, AND at least one first-degree or second-degree relative diagnosed with an LFS tumor, except breast cancer if the individual has breast cancer, before age 56 or with multiple tumors OR Multiple tumors, except multiple breast tumors, two of which belong to the LFS spectrum and the first of which occurred before age 46 OR Adrenocortical carcinoma, choroid plexus tumor, or rhabdomyosarcoma of embryonal anaplastic subtype, regardless of family history OR Breast cancer before age 31

415   Table 2. Summary statistics of prospectively collected probands concerning for LFS determined

416   by GCs.

	TP53 germline mutation carrier (n = 21)		TP53 wild type (n = 123)		Unknown/Pending (n =30)	
Gender	Female (n = 15)	Male (n = 6)	Female (n = 78)	Male (n = 45)	Female (n = 19)	Male (n = 11)
Age Average (range)	29 (5-51)	31 (8 - 72)	32 (3 - 75)	24 (1 - 60)	31 (12 - 68)	34 (9 - 64)
Number of primaries Average (range)	1 (0 - 3)	1 (0 - 3)	1 (0 - 3)	1 (0 - 2)	1 (1-1)	2 (1 - 4)
1st Cancer Diagnosis	n=11	n=4	n=74	n=38	n=19	n=11
Breast	4 (36%)	NA	24 (32%)	NA	2 (6%)	NA
Sarcoma	4 (36%)	2 (50%)	25 (34%)	27 (71%)	11 (63%)	6 (55%)
Brain	1 (10%)	NA	14 (19%)	7 (18%)	3 (18%)	2 (18%)
Other	2 (18%)	2 (50%)	11 (15%)	4 (11%)	2 (13%)	3 (27%)
Age at Diagnosis Average (range)	28 (12 - 43)	16 (8 - 24)	28 (2 - 74)	25 (5 - 59)	27 (3 - 65)	27 (8 - 64)
2nd Cancer Diagnosis	n= 4	n = 2	n = 6	n = 3	n = 0	n = 3
Breast	1 (25%)	NA	4 (66%)	NA	NA	NA
Sarcoma	1 (25%)	2 (100%)	NA	2 (67%)	NA	1 (33%)
Brain	NA	NA	1 (17%)	NA	NA	NA
Other	2 (50%)	NA	1 (17%)	1 (33%)	NA	2 (67%)
Age at Diagnosis Average (range)	32 (18 - 50)	48 (38 - 58)	49 (25 - 75)	36 (21-49)	NA	47 (29 - 58)
Meets Chompret	5 (33%)	2 (33%)	34 (44%)	9 (20%)	1 (5%)	2 (18%)
Chompret Concordant with test results	5 (33%)	2 (33%)	44 (56%)	36 (80%)	NA	NA
LFSPRO score ≥ 0.2	11 (73%)	6 (100%)	10 (13%)	5 (11%)	1 (5%)	2 (18%)
LFSPRO score Average (range)	0.43 (0.00 - 1.00)	0.77 (0.49 - 1.00)	0.10 (0.00 - 0.7)	0.10 (0.00 - 0.83)	0.03 (0.00 - 0.21)	0.20 (0.00 - 1.00)
LFSPRO Concordant with test results	11 (73%)	6 (100%)	68 (87%)	40 (89%)	NA	NA

419 **Table 3. Genetic counselor feedback on LFSPRO clinical utility by clinical scenario.**

Clinical Scenario	Genetic Counselors Quotes
Patient anxiety considerations	Patient was highly anxious about cancer history given her and her full siblings have all developed cancer. I think having a risk model would be helpful, but I worry that giving her exact numbers would have made her anxious and/or hyperfocus on those risk estimates. But if this was a less anxious patient, I would probably feel comfortable giving exact risk estimates.
Validating clinical judgement	Patient with uterine rhabdomyosarcoma and no other family history, I felt like the LFSPRO risk was accurate with my gut instinct. This family really wanted specific risk numbers (ex. they wanted specific numbers regarding how often RMS is genetic, they didn't feel like the general 5-10% of cancers are hereditary was specific enough for them). So I think this family would have really liked a personalized risk model where I could have provided the patient with risks tailored to her.
	LFSPRO would have helped me better counsel this family- they came in thinking they likely had LFS, and I provided reassurance that they likely don't, but wasn't able to give them a percentage likelihood- 13% seems like what I would have quoted but LFSPRO's prediction would have validated my gut and made me more comfortable sharing a percentage with them
Supporting testing decisions	Family is undecided on testing- may decide to test in the coming weeks; I think this risk prediction would have been helpful for the family, as they had assumed the risk was MUCH higher
	This family was wanting to know the risk to test positive to make a decision on the risks versus benefits of testing- I shared my prediction of <5%, this would have been helpful to quote more specifically.

References

1. Olivier, M., Hollstein, M. & Hainaut, P. TP53 Mutations in Human Cancers: Origins, Consequences, and Clinical Use. *Cold Spring Harb. Perspect. Biol.* **2**, a001008 (2010).

2. Malkin, D. *et al.* Germ Line p53 Mutations in a Familial Syndrome of Breast Cancer, Sarcomas, and Other Neoplasms. *Science* **250**, 1233–1238 (1990).

3. Srivastava, S., Blattnert, W. & Chang, E. H. Germ-line transmission of a mutated p53 gene in a cancer-prone family with Li-Fraumeni syndrome. **348**, 747–749 (1990).

4. Nichols, K. E., Malkin, D., Garber, J. E., Fraumeni, J. F., Jr. & Li, F. P. Germ-line p53 Mutations Predispose to a Wide Spectrum of Early-onset Cancers. *Cancer Epidemiol. Biomarkers Prev.* **10**, 83–87 (2001).

5. Birch, J. M. *et al.* Relative frequency and morphology of cancers in carriers of germline TP53 mutations. *Oncogene* **20**, 4621–4628 (2001).

6. Chompret, A. *et al.* P53 germline mutations in childhood cancers and cancer risk for carrier individuals. *Br. J. Cancer* **82**, 1932–1937 (2000).

7. Schneider, K., Zelley, K., Nichols, K. E., Schwartz Levine, A. & Garber, J. Li-Fraumeni Syndrome. in *GeneReviews® [Internet]* (eds Adam, M. P. et al.) (University of Washington, Seattle, Seattle (WA), 1993).

8. Frebourg, T., Bajalica Lagercrantz, S., Oliveira, C., Magenheim, R. & Evans, D. G. Guidelines for the Li–Fraumeni and heritable TP53-related cancer syndromes. *Eur. J. Hum. Genet.* **28**, 1379–1386 (2020).

9. Villani, A. *et al.* Biochemical and imaging surveillance in germline TP53 mutation carriers with Li-Fraumeni syndrome: a prospective observational study. *Lancet Oncol.* **12**, 559–567 (2011).

10. Li, F. P. *et al.* A Cancer Family Syndrome in Twenty-four Kindreds. *Cancer Res.* **48**, 101–124 (1988).

11. Eeles, R. A. Germline mutations in the TP53 gene. *Cancer Surv.* **25**, 101–124 (1995).

12. Chompret, A. *et al.* Sensitivity and predictive value of criteria for p53germline mutation screening. *J.*

Med. Genet. **38**, 43–47 (2001).

13. Bougeard, G. *et al.* Revisiting Li-Fraumeni Syndrome From TP53 Mutation Carriers. *J. Clin. Oncol.*

Off. J. Am. Soc. Clin. Oncol. **33**, 2345–2352 (2015).

14. Park, M. S. *et al.* Utilization of breast cancer risk prediction models by cancer genetic counselors in

clinical practice predominantly in the United States. *J. Genet. Couns.* **30**, 1737–1747 (2021).

15. Peng, G. *et al.* Estimating TP53 Mutation Carrier Probability in Families with Li-Fraumeni Syndrome

Using LFSPRO. *Cancer Epidemiol. Biomark. Prev. Publ. Am. Assoc. Cancer Res. Cosponsored Am.*

Soc. Prev. Oncol. **26**, 837–844 (2017).

16. Shin, S. J. *et al.* Penetrance of Different Cancer Types in Families with Li-Fraumeni Syndrome: A

Validation Study Using Multicenter Cohorts. *Cancer Res.* **80**, 354–360 (2020).

17. Nguyen, N. H. *et al.* Validating Risk Prediction Models for Multiple Primaries and Competing Cancer

Outcomes in Families With Li-Fraumeni Syndrome Using Clinically Ascertained Data. *J. Clin. Oncol.*

Off. J. Am. Soc. Clin. Oncol. **42**, 2186–2195 (2024).

18. Nguyen, N. H. *et al.* LFSPROShiny: An Interactive R/Shiny App for Prediction and Visualization of

Cancer Risks in Families With Deleterious Germline TP53 Mutations. *JCO Clin. Cancer Inform.* **8**,

e2300167 (2024).

19. Kumamoto, T. *et al.* Medical guidelines for Li–Fraumeni syndrome 2019, version 1.1. *Int. J. Clin.*

Oncol. **26**, 2161–2178 (2021).

20. Elston, R. C. & Stewart, J. A General Model for the Genetic Analysis of Pedigree Data. *Hum. Hered.*

21, 523–542 (1971).

21. Gonzalez, K. D. *et al.* High frequency of de novo mutations in Li–Fraumeni syndrome. *J. Med.*

Genet. **46**, 689–693 (2009).

22. Shin, S. J., Yuan, Y., Strong, L. C., Bojadzieva, J. & Wang, W. Bayesian Semiparametric Estimation

of Cancer-Specific Age-at-Onset Penetrance With Application to Li-Fraumeni Syndrome. *J. Am. Stat.*

Assoc. **114**, 541–552 (2019).

23. Shin, S. J. *et al.* Bayesian estimation of a semiparametric recurrent event model with applications to

the penetrance estimation of multiple primary cancers in Li-Fraumeni syndrome. *Biostatistics* **21**,

467–482 (2020).

24. Shin, S. J. *et al.* Penetrance Estimates Over Time to First and Second Primary Cancer Diagnosis in

Families with Li-Fraumeni Syndrome: A Single Institution Perspective. *Cancer Res.* **80**, 347–353

(2020).

25. Edwards, A. G. K. *et al.* Personalised risk communication for informed decision making about taking

screening tests. *Cochrane Database Syst. Rev.* **2013**, CD001865 (2013).

26. Xie, Z. *et al.* Risk estimation, anxiety, and breast cancer worry in women at risk for breast cancer: A

single-arm trial of personalized risk communication. *Psychooncology.* **28**, 2226–2232 (2019).

27. Nguyen, N. H., Shin, S. J., Dodd-Eaton, E. B., Ning, J. & Wang, W. Personalized Risk Prediction for

Cancer Survivors: A Bayesian Semi-parametric Recurrent Event Model with Competing Outcomes.

Ann. Appl. Stat. (in press).