Understanding depression in autism: The mediating role of mentalization, attachment, social support, and psychological flexibility

Eszter Komoróczy[1], Dániel Sörnyei[1,2], Ágota Vass[1], Bálint Szuromi[3], Gabriella Vizin[4], János M. Réthelyi[1], **Kinga Farkas**[1*]

[1] Semmelweis University, Department of Psychiatry and Psychotherapy, Budapest, Hungary
[2] Semmelweis University, Department of Clinical Psychology
[3] Panoráma Poliklinika, Budapest, Hungary
[4] ELTE Eötvös Loránd University, Institute of Psychology, Budapest, Hungary

* *corresponding author:* Kinga Farkas, farkas.kinga@semmelweis.hu
Eszter Komoróczy: kom.eszter@gmail.com
Dániel Sörnyei: s.daniel1207@gmail.com
Ágota Vass: agota0207@gmail.com
Bálint Szuromi: balint.szuromi@gmail.com
Gabriella Vizin: vizin.gabriella@ppk.elte.hu
János M. Réthelyi: rethelyi.janos@semmelweis.hu

Keywords: Autism Spectrum Disorder, Depression, Mentalization, Psychological Flexibility, Social Support, Attachment

Data availability statement
The data that support the findings of this study are available on request from the corresponding author.

Funding statement
This research was supported by the Hungarian National Research, Development and Innovation Office Grant OTKA PD 146424 (to K.F.); and the Ministry of Innovation and Technology of Hungary from the National Research, Development and Innovation Fund, financed under the TKP2021-EGA-25 funding scheme (to E.K., Á.V., K.F. and J.M.R).

Conflict of interest statement
The authors declare that there is no conflict of interest.

Ethics approval statement
This study was approved by the Semmelweis University Regional and Institutional Committee of Science and Research Ethics, approval number SE RKEB: 159/2021.

Patient consent statement
Informed consent was obtained from all participants for this study. Participants indicated their consent to participate by clicking the provided button after reading the informed consent form.

Permission to reproduce material from other sources
Not applicable.

Clinical trial registration
Not applicable

Understanding depression in autism: The mediating role of
mentalization, attachment, social support, and psychological flexibility

Abstract

Background: Depression is highly prevalent in autistic individuals, yet its underlying
psychological mechanisms remain insufficiently understood, especially in populations who self-
identify as autistic but lack a formal diagnosis.
Methods: Using a moderated mediation approach (structural equation modeling), we investigated
how autistic traits relate to depressive symptoms in three groups: individuals with clinically
diagnosed autism (ASD, N = 136), self-identified autistic individuals without formal diagnosis
(ASD-sd, N = 100), and neurotypical controls (N = 1566). Five psychosocial mediators were tested:
mentalizing, anxious and avoidant attachment, perceived social support, and psychological
flexibility.
Results: Depressive symptoms were significantly higher in both autistic groups compared to
neurotypical controls. The association between autistic traits and depressive symptoms was fully
mediated by a combination of psychological flexibility, attachment style, mentalizing, and
perceived social support. Among these, psychological flexibility emerged as the strongest and most
consistent mediator across all groups. In contrast, mentalizing showed an indirect effect only in the
non-ASD group, while avoidant attachment was a significant mediator solely in the clinically
diagnosed ASD group. Perceived social support showed a modest indirect effect only in the non-
ASD group.
Conclusions: Our findings support a transdiagnostic model in which psychological flexibility plays
a central protective role against depression across varying levels of autistic traits. Meanwhile,
attachment-related and social-cognitive vulnerabilities contribute in different ways depending on
diagnostic status. These results underscore the clinical importance of individualized interventions
targeting such modifiable factors. Mentalization-based treatment, adapted for neurodiverse
populations, may offer a promising framework for addressing both common and subgroup-specific
mechanisms.

Lay abstract

Many autistic people experience depression, but we still know little about why this happens and
what psychological factors are involved, especially for those who see themselves as autistic but
don't have a formal diagnosis. We studied three groups: people with diagnosed autism, people who
self-identify as autistic, and people without autism. We wanted to understand which psychological
traits might explain why autistic traits are linked to depression. We found that a skill called
psychological flexibility (being able to stay present and adapt to difficult emotions) was the most
important protective factor in all groups. Other factors, like how people think about relationships
(attachment) and how well they understand themselves and others (mentalizing), were only
important for some groups. These findings suggest that therapies which improve flexibility and help
with emotional connection may reduce depression in autistic people and those with high autistic
traits.

Keywords: Autism Spectrum Disorder, Depression, Mentalization, Psychological Flexibility, Social
Support, Attachment

Introduction

What is the relevance of studying depression in autism?

Autism spectrum disorder (ASD) is a neurodevelopmental condition characterised by
difficulties in social communication and interaction, restricted or repetitive behaviours or interests,
and sensory processing alterations (American Psychiatric Association, 2013, 2022). The rising
prevalence of ASD (Zeidan et al., 2022) makes it increasingly crucial to understand its
comorbidities, particularly the frequent and severe co-occurrence of depression (Bougeard et al.,
2021). A number of studies have examined the relationship between ASD and depression,
suggesting that individuals with autism have a higher rate of depression than the population average
(e.g. (Hollocks et al., 2019; Lai et al., 2019; Wigham et al., 2017). While the prevalence of
depressive disorders is estimated between 7-11% in the general population (Lim et al., 2018;
Wittchen et al., 2011; Wittchen & Jacobi, 2005), it is approximately 10-15%, ranging from 2.5% to
47.1% in ASD (Hossain et al., 2020), and individuals with ASD are four times more likely to
experience depression during their lifetime compared to neurotypical individuals (Hudson et al.,
2019) according to recent meta-analyses. Despite the high prevalence of depressive symptoms in
ASD patients, the nature of the relationship between the two conditions remains uncertain.

As depression is a multifactorial disorder and ASD is a heterogeneous spectrum condition, it
is difficult to single out one causal factor. It has been recognised that one in two individuals with
autism has alexithymia (Albantakis et al., 2020); moreover loneliness (Schiltz et al., 2021;
Umagami et al., 2022), emotion regulation deficits, and intolerance of uncertainty have also been
found to be associated with symptoms of anxiety and depression in autistic populations (Cai et al.,
2018). Furthermore, depressive symptoms severely impact the quality of life (Thiel et al., 2024).

Although autism spectrum disorder (ASD) is typically diagnosed by professionals using
standardized criteria (e.g., DSM-5, ICD-11), an increasing number of individuals self-identify as
autistic, particularly those who may have been overlooked by traditional diagnostic pathways, such
as women and gender-diverse individuals who often present atypical or camouflaged symptoms
(Hull et al., 2017; Lai & Baron-Cohen, 2015). These self-diagnosed individuals frequently report
experiences consistent with the autistic phenotype and may experience comparable psychological
distress (Banker et al., 2025; McDonald, 2020). Thus, their inclusion may provide a more
comprehensive understanding of the autistic trait continuum and its association with depression.

In this article, we explore the interconnected phenomena of mentalization, attachment, peer
support and psychological flexibility as potential mediators to understand their combined impact
on the relationship between ASD and depressive symptoms. Our inquiry aligns with the
mentalization theory proposed by Fonagy, Luyten, and Bateman, subsequently extended to
illuminate diverse psychopathological phenomena (Fonagy & Bateman, 2016; Fonagy & Luyten,
2009; Luyten et al., 2020). By this exploration, we endeavour to shed light on the intricate web
connecting ASD and depressive symptomatology. In line with this spectrum perspective, our study

includes individuals with both formal and self-reported ASD diagnoses, allowing us to capture a
broader range of autistic experiences. This is particularly relevant given that many self-identified
autistic individuals exhibit elevated autistic traits and similar patterns of psychological vulnerability
(Banker et al., 2025), even in the absence of formal diagnosis.

What is the role of mediating factors in the relationship between autism and depression?

Mentalization

Social communication deficits are core symptoms of ASD. Mentalizing deficits occur early
but remain observable throughout life (Baron-Cohen, 2000; Chung et al., 2014). Research during
recent years in ASD has consistently shown deficits in theory of mind (ToM; one specific aspect of
mentalization) (Yirmiya et al., 1998), emotion perception and processing (Brewer et al., 2017;
Velikonja et al., 2019), empathy, affective mentalizing, and cognitive mentalizing and related
altered neural functioning (Arioli et al., 2021). ASD is, however, a heterogeneous spectrum; it is
important to note that within the condition, variability in mentalizing abilities is not negligible
(Lombardo et al., 2016).

Mentalization deficits are associated with depressive symptoms as well. Identifying and
interpreting mental states of the self and others have been found to be impaired among patients with
depression, and the degree of mentalizing impairment was associated with the severity of depression
and worse clinical outcomes (Fischer-Kern et al., 2013; L. Lee et al., 2005; Lombardo et al., 2016).
Deficits in theory of mind were shown to be a predictor of relapse and impairment in social
functioning in depression (Inoue et al., 2006) and were associated with cognitive deficits as well
(Wang et al., 2008). Fischer-Kern et al. (2022) also showed that, besides mentalizing deficits,
attachment insecurity and unresolved loss are also important predictors of depression. A current
integrative model of depression, the stress-reward-mentalizing model suggests a developmental
cascade. This sequence involves elevated stress levels and persistent heightened arousal, potentially
coupled with genetic predisposition, culminating in impaired reward sensitivity within attachment
relationships and the realm of agency and self-governance. Consequently, this progression impacts
the capacity for mentalizing and social cognitive functions (Luyten & Fonagy, 2018).

Attachment

Emotion recognition, reciprocal social communication, ToM, and other aspects of
mentalization are essential to form secure attachment and maintain a balanced parent-child
relationship (Fonagy et al., 1991; Keenan et al., 2017). Children with ASD show impairments in
social, and emotional capacities from an early age (American Psychiatric Association, 2022; Baron-
Cohen et al., 1985; Cassel et al., 2007; Dawson et al., 2004; Nuske et al., 2013). They exhibit
heightened attachment insecurity (Naber et al., 2007), as well as decreased sensitivity in their
relationships with caregivers (Rutgers et al., 2004; Van IJzendoorn et al., 2007), including more
contact resistance and less contact-seeking behaviours (Rogers et al., 1993), fewer pro-social

responses to caregivers and difficulty being soothed (Grzadzinski et al., 2014), and less frequent
looking, showing, smiling, and mutual play behaviours (Akdemir et al., 2009; Dissanayake &
Crossley, 1997). According to a recent review, only 47% of children with ASD demonstrated secure
attachments (Teague et al., 2017). The notion that individuals with ASD have difficulties to form
attachments, however, is tempered by the observation that they express a preference for the
caregiver over a stranger, and experience distress when unexpectedly separated from them (Capps
et al., 1994; Dissanayake & Crossley, 1997; Koren-Karie et al., 2009).

Employing questionnaire-based measures, instead of observation-based paradigms like the
Strange Situation (Ainsworth et al., 1978), during middle childhood, no differences in attachment
security were found between autistic and neurotypical participants (Bauminger et al., 2010;
Chandler & Dissanayake, 2014; Wu et al., 2015). Upon reaching adulthood, attachment style
remains relatively stable (Fraley, 2002), but the object of attachment shifts from parents to peers,
including friends and romantic partners (Doherty & Feeney, 2004; Hazan & Zeifman, 1994;
Nickerson & Nagle, 2005). Insecure attachment, especially avoidant attachment style, is more
prevalent in ASD groups than in the general population (Taylor et al., 2008), and associated with
autistic traits in non-clinical samples (Gallitto & Leth-Steensen, 2015), and it is associated with
lower relationship satisfaction (Beffel et al., 2021). A recent study found that adult attachment styles
are important predictors of mental health: higher anxious attachment predicted higher depression
and anxiety, interestingly, higher avoidant attachment predicted lower anxiety and stress in autistic
adults (Lee et al., 2022).

Attachment-related distress and avoidance have been associated with psychopathology
(Mikulincer & Shaver, 2012, 2017), including anxiety (Bosmans et al., 2010) and depression
(Beatson & Taryan, 2003; Besser & Priel, 2003; Cantazaro & Wei, 2010; Dagan et al., 2018; Malik
et al., 2015). Individuals reporting low attachment anxiety and avoidance have the ability to
maintain interpersonal relationships through secure attachment and report higher levels of adaptive
emotion regulation, positive self- and other perceptions, and greater psychological well-being
(Mikulincer & Shaver, 2019; Raque-Bogdan et al., 2011). Attachment style has an impact on
emotional, behavioural, and relational patterns (Mikulincer & Shaver, 2017), resilience to stressful
life events (Mikulincer & Shaver, 2012), and the quality of interpersonal relationships (Collins,
1996; Shaver et al., 2000; Torquati & Vazsonyi, 1999).

Social support

Developmental research on mentalization suggests that environmental influences play a
significant role in the development of mentalization. In recent years, there has been a notable shift
in this respect: while earlier theories (Fonagy et al., 1991) emphasised the specific role of dyadic
(parent-child) attachment in facilitating or inhibiting the development of mentalization, recent
views have taken a more comprehensive approach. Accordingly, in addition to the family
environment, peers, and wider sociocultural factors are in close interaction with attachment,
mentalization and social learning, through the capacity to rely on others to provide social

information (epistemic trust) and the ability to benefit from the positive effects of the environment
(salutogenesis) (Fonagy & Allison, 2014; Luyten et al., 2020). As ASD is linked to difficulties in
social communication and interaction, people with ASD are less able to form social relationships
and access and incorporate the benefits of such peer support. The effect of perceived social support
has been measured mainly among the parents of children with ASD (Alon, 2019; He et al., 2022;
Lu et al., 2021). Studies examining social support in adult ASD found that social support was
associated with psychological, social, and environmental well-being (Bishop-Fitzpatrick et al.,
2018; Leader et al., 2021; Tsermentseli, 2022).

Social support is a protective factor against depression, with parental support as the most
influential during childhood and adolescence, whereas spouses, family, and friends become more
important in adulthood and senior age (Gariépy et al., 2016). Perceived social support mediated
loneliness and depression in the elderly (Liu et al., 2016), lower quality support predicted a sixfold
risk of depressive symptoms among college students (Hefner & Eisenberg, 2009), and elevated the
risk of postpartum depression (Akbari et al., 2020), and more importantly showed to be a moderator
between negative life events and depression in adolescence (Miloseva et al., 2017). On the contrary,
higher levels of perceived social support were linked to positive affect, overall life satisfaction, and
favourable outcomes of mental disorders (Siedlecki et al., 2014; J. Wang et al., 2018).

Psychological flexibility and ASD

Psychological resilience and flexibility are also important factors in mental health (Kashdan
& Rottenberg, 2010; Rutten et al., 2013; Yasinski et al., 2020). Psychological inflexibility is crucial
in understanding the relationship between depression and autism due to its effects on coping
strategies, emotional regulation, and adaptive functioning. Individuals with autism often struggle to
adapt to changing circumstances. Inflexibility can appear as rigid thinking patterns, resistance to
change, and difficulty shifting focus from negative thoughts or experiences, which can intensify
their depressive symptoms. Although ASD is characterised by behavioural inflexibility (American
Psychiatric Association, 2022), the role of psychological inflexibility was measured only once
among autistic adults. A recent study reported that psychological flexibility (measured by AAQ-II)
and attachment anxiety mediated the association between childhood experiences of parenting and
mental health in later life in individuals with autism (Lee et al., 2022). It is conceivable that
psychological flexibility can be instrumental in several ways when considering its reciprocal
relationship with attachment, mentalizing, and perceived social support in individuals with autism.
It can also facilitate considering alternative perspectives, managing conflicting emotions, adapting
their responses in interpersonal challenges, and being open to seeking and receiving support.

Hypotheses and aims of the current paper

In this study, we aim to validate the heightened prevalence of depression in individuals with
autistic traits through a comparative analysis of depressive symptom severity. First, we examine
the associations between autistic traits and depressive symptoms across both neurotypical and

autistic populations. Then, we investigate whether this relationship is mediated by key
psychological constructs: mentalization, attachment style, perceived social support, and
psychological flexibility - factors that are known to influence mental well-being in both autism and
the general population. Finally, we explore whether these pathways differ based on diagnostic
status, including both individuals with a formal ASD diagnosis and those who self-identify as
autistic. By including the self-diagnosed group, we aim to determine whether their psychological
profiles align more closely with clinically diagnosed individuals or represent a distinct subgroup
along the autism spectrum.

Methods

Recruitment procedure, ethical concerns

Participants were requested to complete an online survey between September 2021 and April
2022. Initially, we reached out to outpatient units, institutions, and counsellors specialising in aiding
individuals diagnosed with autism spectrum disorder, with the purpose of facilitating questionnaire
accessibility for those with an established ASD diagnosis.
The language of the questionnaires was Hungarian and the wording was designed for maximum
clarity. Experts from these organisations also checked the questionnaire, provided suggestions and
these were incorporated in the final version of the survey. Subsequently, we also recruited non-
ASD participants through media and social media advertisements.
Participants provided informed consent online. The authors assert that all procedures contributing
to this work comply with the ethical standards of the relevant national and institutional committees
on human experimentation and with the Helsinki Declaration. The study was approved by the
Semmelweis University Regional and Institutional Committee of Science and Research Ethics, SE
RKEB: 159/2021.

Participants

Overall 2409 participants were recruited and started the survey. Completing the whole survey
took about 40 minutes, and 1783 participants answered all the items. Participants were allowed to
stop at any point and resume filling out the questionnaires later. We have constructed four selection
criteria to ensure high quality of our data. First, we have excluded those participants who did not
fill out the BDI, or the AQ questionnaires which were intended to measure the depressive and
autistic symptom severity, respectively. Participants between the ages of 18 and 65 were included.
Further, we have confirmed that all of our participants have completed at least 8 years of elementary
education, or have provided realistic data regarding the years they had spent in education: we have
excluded all participants who claimed to have spent more years in education than their age minus 5
years. Finally, we have also rigorously screened our sample based on the time spent filling out the
questionnaires that were of most interest to us. As participants were allowed to take a break and
resume filling out the questionnaires later at their convenience, there were some extremely high

values in the time spent filling out many of the questionnaires. Thus, we decided to assess the time

it took participants to fill out the respective questionnaires by calculating measures that are more

robust to outliers. We have calculated the median filling time and median absolute deviation of

filling time as a measure of spread for each questionnaire. We have excluded those participants who

have spent less than the median minus one and a half median absolute deviation time on the

following questionnaires: AQ, BDI, MZQ, AAS, MSPSS, and AAQ-II.

Questionnaires

Demographic data collection

Demographic information was collected using a structured questionnaire administered at the

beginning of the study. Participants were asked about their age, gender, education level,

socioeconomic and marital status.

Autism Spectrum Quotient (AQ)

Autism Spectrum Quotient is a 50-item self-report questionnaire to measure the degree of autistic

traits in adults with normal intelligence. The AQ assesses five different areas: social skill, attention

switching, attention to detail, communication, and imagination; 10 items cover each area, on a 4-

point Likert scale with values of 'definitely agree', 'slightly agree', 'slightly disagree' or 'definitely

disagree'. At the end of the questionnaire, each participant was given feedback on their score on

the AQ scale and whether they should seek professional advice. The internal consistency of items

in each of the five domains and Cronbach's alpha coefficients were all moderate to high (Baron-

Cohen et al., 2001). In the present study, the overall reliability was good (total α = 0.89, ASD α =

0.86, ASD-sd α = 0.85 and NTP (non-ASD) α = 0.86).

Beck Depression Inventory (BDI) is a widely used 21-item, self-report inventory measuring

symptoms of depression with good internal consistency (α = 0.86) (Beck et al., 1988). In our

study, this scale had good internal consistency (total α = 0.90, ASD α = 0.91, ASD-sd α = 0.88 and

NTP α = 0.90).

Mentalization Questionnaire (MZQ)

The Mentalization Questionnaire is a 15-item self-report measure of mentalization (Hausberg et al.,

2012). The items of the MZQ cover four domains of mentalization: 'Refusing self-reflection',

'Emotional awareness', 'Psychic equivalence mode', and 'Regulation of affect'. The structural

validity of the Hungarian version of the MZQ was assessed among psychotic patients and the study

confirmed the four-factor structure which differs from healthy samples (Fekete et al., 2019). In our

study, however, we used the overall MZQ score (higher scores meant greater difficulty in

mentalisation), which had good reliability in the sample (total α = 0.86, ASD α = 0.82, ASD-sd α

= 0.77 and NTP α = 0.85).

Adult   attachment   scale   (AAS)

Adult attachment style was measured by the Adult Attachment Scale (AAS), which consists of 18

Likert items that can be grouped into two subscales (Collins, 1996). Psychometric properties of the

AAS were evaluated in a Hungarian sample recently (Őri et al., 2021). In our study, we used the

scores reflecting anxious and avoidant attachment styles, which had good internal consistency

(total α *(anxious)* = 0.88 and α *(avoidant)* = 0.86, ASD α *(anxious)* = 0.87 and α *(avoidant)* = 0.83,

ASD-sd α *(anxious)* = 0.88 and α *(avoidant)* = 0.81, and NTP α *(anxious)* = 0.88 and α *(avoidant)*

= 0.86).

Multidimensional Scale of Perceived Social Support (MSPSS) during adolescence.

Social support during adolescence was measured with the Multidimensional Scale of Perceived

Social Support (MSPSS), which has 12 Likert items (Canty-Mitchell & Zimet, 2000; Papp-

Zipernovszky et al., 2017). Participants were asked to focus on their adolescence. In our study,

this scale had excellent internal consistency (total α = 0.95, ASD α = 0.92, ASD-sd α = 0.90 and

NTP α = 0.95).

Acceptance and Action Questionnaire-II (AAQ-II)

AAQ-II is a measure of psychological flexibility, which has 7 Likert items (7-point scale, from

"never true" to "always true"). Higher total scores mean less flexibility. The tool has satisfactory

psychometric properties (α = 0.84) (Bond et al., 2011), and in a Hungarian sample the internal

consistency was good (α = 0.90) (Eisenbeck & Szabó-Bartha, 2018). In our study, this scale had

good internal consistency (total α = 0.93, ASD α = 0.92, ASD-sd α = 0.90 and NTP α = 0.93).

Data Analysis

R, RStudio 2024.09.1 software was used to perform the analyses and data visualization. First,

the Krukskal-Wallis, Wilcoxon rank sum test and Fisher's exact test statistics were used to compare

the ASD and non-ASD samples (Tables 1 and 3). Second, receiver operating characteristic (ROC)

curve analysis was performed to examine the sensitivity and specificity of the AQ to determine a

scoring cutoff value that discriminates between autistic and non-autistic cases (Hosmer, 2013;

Robin et al., 2011). In the third step, we conducted Spearman correlation analyses to explore the

associations between AQ scores and other psychological constructs. We compared the correlation

coefficients between ASD and non-ASD groups by z-test (Eid et al., 2013). Detailed results and

methods are reported in the Supplementary Material (Figure S5).

Moderated mediation

Finally, to investigate the conditional indirect effects of autistic traits on depressive symptoms, we conducted multivariate moderated mediation analyses using structural equation modeling (SEM) with the *lavaan* package (version 0.6-19) in R (Rosseel, 2012). This approach implements regression-based moderated mediation analysis by modeling all mediators simultaneously within a unified SEM framework. Interaction terms between the predictor (AQ total score) and diagnostic groups (ASD, ASD-sd as dummies) were included to estimate group-specific a paths (predictor → mediator) and b paths (mediator → outcome). Attachment style (AAS scores), mentalization (MZQ total score), perceived social support (MSPSS total score), and psychological flexibility (AAQ-II total score) had been set as mediators and depression severity index (BDI score) was the outcome variable. To control for potential confounding, sex (three categories: male, female, other), age (continuous), and years of education (continuous) were included as covariates predicting both the mediators and the outcome. Standardized estimates are reported. Indirect effects were calculated for each group separately, and significance was determined via bias-corrected bootstrap confidence intervals based on 5000 resamples. This method enabled robust estimation of complex, group-differentiated mediation pathways, accounting for shared variance among mediators and allowing direct comparison of conditional effects across groups. Model fit was evaluated using multiple indices (CFI, TLI, RMSEA, SRMR), although the complexity and moderation structure were expected to lower absolute fit indices (Supplementary Material Table S1)

Sensitivity analysis

To explore whether self-diagnosed and clinically diagnosed individuals could be merged analytically, we also tested models combining both groups (Supplementary material, Sensitivity analysis, Figures S6-7, Tables S7-10).

Results

Descriptive statistics

The basic demographic data of the participants are shown in Table 1. The gender distribution in the non-ASD group reflects the generally higher female response rate (female: 66.2% vs. male: 32.1%), 1.7% of respondents identified themselves as other gender. In the ASD group, by contrast, the proportion of women and men showed a nearly even balance (female: 47.8% vs. male: 43.4%; as men are more likely to be diagnosed with ASD, but women tend to be more willing to answer

questionnaires), and there was a significantly higher proportion of those who defined themselves as

other gender (8.8%). For completeness, we also received responses from individuals who believed

they were on the autism spectrum or were perceived as such by friends or relatives, but who did not

have a formal diagnosis at the time of completing the survey. Their AQ scores (See Supplementary

material Figure S1-S4 & ROC analysis) suggest a real possibility that their assumption is correct;

however, it was not possible to formally investigate or diagnose them. Sensitivity analysis yielded

qualitatively similar results, suggesting robustness, but group-specific analyses revealed important

nuanced psychological differences, justifying the use of separate groups in the main model. Basic

demographic data are shown in Table 1, while detailed demographic data for the three study groups

are presented in Supplementary Material Table S1. As there were significant differences in sex,

age, and education level across the groups, these variables were included as covariates in subsequent

analyses.

Table 1. Basic demographics

		ASD *N*=136	ASD-sd *N*=100	non-ASD *N*=1566	Group comparison statistics			
Sex (N (%))					*Fisher's exact test*ᵃ	*p-value*		
	Female	65 (47.8%)	58 (58.0%)	1037 (66.2%)				
	Male	59 (43.4%)	29 (29.0%)	503 (32.1%)	-	<0.001		
	Other	12 (8.8%)	13 (13.0%)	26 (1.7%)				
					Kruskal-Wallis	*p-value*	ε^2	*Post hoc (Wilcoxon)*
Age (years)		33.25 (9.98)	36.30 (9.00)	38.52 (9.79)	34.572	<0.001	0.018	non-ASD > ASD-sd > ASD
Education (years)		16.87 (4.06)	16.88 (3.83)	17.91 (3.64)	23.048	<0.001	0.012	non-ASD > ASD-sd = ASD

Notes. ASD: Autism Spectrum Disorder diagnosed by clinician; ASD-sd: autism self-diagnosed; non-ASD: neither confirmed
nor presumed diagnosis of ASD; N: number of participants. *A Fisher's exact test (with 10,000 Monte Carlo simulations)
revealed a significant difference in gender distribution across groups, p < .001

Clinical characteristics

The ASD and non-ASD groups scored differently on all questionnaires, with the ASD group

scoring significantly higher on AQ, BDI, MZQ, AAS (both insecure attachment subscale), and AAQ-II,

and lower on the MSPSS scale, the latter indicates lower perceived peer and family support during

adolescence. Detailed data and between-group differences are shown in Table 3 and Figure 1.

In Figure 1A AQ scores were also plotted by both diagnostic group and sex, revealing patterns

similar to those reported by Baron-Cohen et al. (2014), showing an attenuation of typical sex differences

in the autism group.

Table 3. Clinical characteristics – questionnaire scores

	ASD N=133			ASD-sd N=100			non-ASD N=1550		
Questionnaire	*Mean*	*Median*	*SD*	*Mean*	*Median*	*SD*	*Mean*	*Median*	*SD*
AQ sum	32.27	34.0	8.16	32.82	34.0	7.22	20.29	20.0	8.21
social	6.79	8.0	2.64	6.80	7.0	2.31	3.82	3.0	2.81
attention switch	7.91	8.0	2.02	8.01	8.5	2.01	5.20	5.0	2.50
communication	6.42	6.0	2.50	6.38	6.0	1.94	3.03	3.0	2.20
attention to detail	6.35	7.0	2.02	6.65	7.0	2.20	5.10	5.0	2.14
imagination	4.81	4.5	2.22	4.98	5.0	2.19	3.13	3.0	1.95
BDI	16.59	15.5	11.53	17.08	17.0	9.91	11.74	10.0	8.86
MZQ	36.60	38.0	10.70	37.01	37.0	9.10	25.78	26.0	11.03
AAS anxious	20.79	21.0	6.55	20.48	21.0	6.82	16.16	16.0	6.46
AAS avoidant	40.75	41.5	10.03	43.66	45.0	8.63	34.95	35.0	9.45
MSPSS	35.10	36.0	11.62	32.95	31.0	9.87	41.72	43.0	11.87
AAQ-II	31.97	32.0	9.88	31.66	32.0	9.23	22.84	22.0	10.05

Notes. ASD: Autism Spectrum Disorder diagnosed by clinician; non-ASD: neither confirmed nor presumed diagnosis of ASD;
N: number of participants. SD: standard deviation. AQ: Autism Spectrum Quotient; BDI: Beck Depression Inventory; AAS
anxious: Adult Attachment Style: Anxious attachment; AAS avoidant: Adult Attachment Style: Avoidant attachment; MZQ:
Mentalization Questionnaire; MSPSS: Multidimensional Scale of Perceived Social Support; AAQ-II: Acceptance and Action
Questionnaire-II

Group differences in trait measures

Statistically significant group differences were found across all questionnaire measures
(Kruskal–Wallis tests, all ps < .001). Post-hoc comparisons revealed that both individuals with
clinically diagnosed ASD and those with self-diagnosed ASD scored significantly higher than non-
ASD participants on all measures of autistic traits, depression, psychological inflexibility,
attachment insecurity, and mentalization difficulties (all adjusted ps < .001). A similar pattern was
observed in the opposite direction for perceived social support, with lower scores among ASD
groups, consistent with its role as a protective factor. Differences between the ASD and ASD-sd
groups were generally non-significant, except for avoidant attachment (AAS-avoid), where the
ASD-sd group reported higher scores (adjusted p = .025, r = .15). For detailed test statistics, effect
sizes, and adjusted p-values, see Supplementary Table S2.

Figure 1. Distribution of clinical characteristics (questionnaire scores) by group

Notes: Blue: participants diagnosed with ASD; Purple: participants with self-diagnosed autism spectrum disorder; Yellow: participants with neither confirmed nor presumed diagnosis of ASD; Boxplot: the box shows the interquartile range, the horizontal line indicates the median, and the whiskers show the maximum and minimum values.
A) The y axis shows the AQ (Autism Spectrum Quotient) scores of participants. On the x axis participants are assigned into three groups (participants diagnosed with ASD and participants not diagnosed with ASD and are neither self-diagnosed) and participants are further grouped based on their sex. B-G) The y axis shows the further scores of participants. On the x axis participants are assigned into three groups. BDI: Beck Depression Inventory; AAS anxious: Adult Attachment Style: Anxious attachment; AAS avoidant: Adult Attachment Style: Avoidant attachment; MZQ: Mentalization Questionnaire; MSPSS: Multidimensional Scale of Perceived Social Support; AAQ-II: Acceptance and Action Questionnaire-II

Sensitivity and specificity of the AQ

ROC analysis indicates that AQ-50 scores effectively distinguish both clinically diagnosed and self-diagnosed autistic individuals from non-ASD participants, with AUCs of 0.842 and 0.865, respectively. While the self-diagnosed group showed slightly higher discrimination, a DeLong test revealed that this difference was not statistically significant ($D = -0.924$, $p = .355$), indicating comparable classification performance across groups. The optimal cutoff score was higher for the

ASD (clinically diagnosed) group (29.5) than for the self-diagnosed group (24.5). Sensitivity was

higher in the diagnosed group (0.844 vs. specificity: 0.699), whereas specificity was higher in the

self-diagnosed group (0.86 vs. sensitivity: 0.699), suggesting that self-identified individuals show

meaningful autistic traits but may present a more heterogeneous or subclinical profile (see details

in the Supplementary material Figures S1-S4). Accordingly, we included both groups separately in

the main *lavaan* SEM and group comparisons to test whether self-diagnosed individuals represent

a meaningful subtype or a distinct point along the autistic trait continuum. To ensure the robustness

of our findings, we also report a sensitivity analysis using a combined ASD group (clinically

diagnosed and self-diagnosed) in the Supplementary Material (Figures S6-7, Tables S7-S10). This

yielded qualitatively similar results, and model fit did not change significantly. Therefore, in line

with our exploratory aims, we retained the three-group structure in the main analyses to examine

potential nuanced psychological differences among the study groups.

Moderated mediation model

A moderated mediation model was estimated using structural equation modeling (SEM) in

lavaan, controlling for age, years of education, and sex. The model fit was poor ($\chi^2(60) = 3441.42$,

$p < .001$, CFI = 0.58, TLI = 0.14, RMSEA = 0.181, SRMR = 0.056), but the individual pathways

were interpreted according to the pre-specified hypotheses. Indirect effects were calculated for each

mediator across the three groups (non-ASD, ASD, ASD-sd). Mediation analyses revealed that

psychological inflexibility (AAQ-II) significantly mediated the association between autistic traits

(AQ) and depressive symptoms (BDI) across all groups. In the non-ASD group the indirect effect

was $\beta = 0.267$, SE = 0.020, 95% CI [0.229, 0.308], $p < .001$, in the ASD group: $\beta = 0.121$, SE =

0.058, 95% CI [0.026, 0.250], $p = .035$ and in ASD-sd group $\beta = 0.144$, SE = 0.069, 95% CI [0.024,

0.290], $p = .037$. Avoidant attachment (AAS-avoid) significantly mediated the AQ → BDI

relationship only in the clinically diagnosed ASD group: the indirect effect was $\beta = 0.115$, SE

= 0.051, 95% CI [0.026, 0.222], $p = .025$. Indirect effects through AAS-avoidance were smaller and

non-significant in the non-ASD (indirect effect: $\beta = 0.020$, SE = 0.020, 95% CI [-0.019, 0.059], p

= .314) and ASD-sd groups (indirect effect: $\beta = 0.111$, SE = 0.070, 95% CI [-0.009, 0.266], $p =$

.113). Mentalization (MZQ) showed a significant indirect effect only in the non-ASD group ($\beta =$

0.115, SE = 0.021, 95% CI [0.074, 0.157], $p < .001$). The mediation paths via perceived social

support (MSPSS) were weak and non-significant in ASD and ASD-sd groups; a marginal effect

was observed in the non-ASD group ($\beta = 0.022$, SE = 0.011, 95% CI [–0.0002, 0.044], $p = .054$).

Anxious attachment (AAS-anx) did not show significant mediation in any group. (Figure 2, Indirect

mediators). These effects remained consistent after controlling for age, sex, and education. Detailed

parameter estimates, confidence intervals are provided in the Supplementary Material (Tables S2–

S3).

Direct Effect

The direct effect of AQ on BDI was nonsignificant when controlling for mediators and
covariates (standardized estimate: β = -0.014, SE = 0.028, 95% CI [-0.068, 0.041], p = .619) (Figure
2, Direct effect). However, group membership significantly predicted depression: individuals in the
ASD group had higher depression scores compared to the non-ASD group (β = –0.314, p = .038).
No significant direct effects were found for the ASD-self-diagnosed group (p = .499) (Figure 2,
Group effect) or for the interaction terms between AQ and group status (all ps > .4), for details see
Supplementary   Material   Table   S2.
These results suggest that the association between autistic traits and depression was primarily
indirect; psychological inflexibility is the primary mediator linking autistic traits to depressive
symptoms across groups, avoidant attachment plays a specific mediating role in individuals with
ASD, while mentalization in non-ASD.

Figure 2. Moderated mediation model

Notes. Blue: participants diagnosed with ASD; purple: participants with self-diagnosed autism spectrum disorder;
yellow: participants with neither confirmed nor presumed diagnosis of ASD. Predictor: AQ = Autism Spectrum Quotient;
outcome: BDI = Beck Depression Inventory; mediators: AAS anxious = Adult Attachment Scale: attachment anxiety;
AAS avoidant = Adult Attachment Scale: attachment avoidance; MZQ = Mentalization Questionnaire; MSPSS =
Multidimensional Scale of Perceived Social Support; AAQ-II = Acceptance and Action Questionnaire-II. Standardised
regression coefficients (β) are shown on the arrows. Continuous lines indicate significant, and dotted lines indicate non-
significant regression effects. Levels of significance: * p < .05; ** p < .01; *** p < .001

Group Comparisons of Indirect Effects

To visualize how the path estimates vary across diagnostic groups, we plotted the a-paths, b-
paths, and resulting indirect effects for each mediator (see Figure 3).

Figure 3. Standardized estimates of a-paths, b-paths, and indirect effects (a × b) across diagnostic groups   for   each   mediator
Notes. Values are based on SEM estimates adjusted for age, sex, and education. Blue: participants diagnosed with ASD; purple: participants with self-diagnosed autism spectrum disorder; yellow: participants with neither confirmed nor presumed diagnosis of ASD. Predictor: AQ (Autism Spectrum Quotient); outcome: BDI (Beck Depression Inventory); mediators: AAS (anx) = Adult Attachment Scale: attachment anxiety; AAS (avoid) = Adult Attachment Scale: attachment avoidance; MZQ = Mentalization Questionnaire; MSPSS = Multidimensional Scale of Perceived Social Support; AAQ-II = Acceptance and Action Questionnaire-II.

Group comparisons of indirect effects using Wald chi-square tests revealed that, in most cases, the size of the mediation effects did not significantly differ between groups (all *p-values* > .05). An exception was found for psychological inflexibility (AAQ-II), where the indirect effect was significantly larger in the non-ASD group compared to the ASD group ($\chi^2(1) = 5.90$, $p = .015$). The avoidant attachment pathway (AAS-avoid) remained stronger in the ASD group; however, after adjusting for covariates, the group difference was no longer statistically significant, though it approached trend level (Wald $\chi^2 = 3.01$, $p = .083$). No significant differences were observed between groups for the other mediators, including mentalization (MZQ), anxious attachment (AAS-anxious), and perceived social support (MSPSS) (all Wald $\chi^2(1)$ tests, $p > .05$). For detailed descriptions of Wald test comparisons see Supplementary Material Table S4.

Discussion

Our investigation employed between-group comparisons and a spectrum approach, utilizing

a moderated mediation method to analyze the relationship between autistic traits and depressive

symptom severity across three groups: individuals with a formal ASD diagnosis, self-diagnosed

autistic individuals (ASD-sd), and neurotypical controls. We examined five potential mediators:

mentalizing, anxious and avoidant attachment style, perceived social support, and psychological

flexibility. The findings indicated that the strength and pattern of mediation varied across groups.

Despite significant differences in all scores between groups, psychological flexibility emerged as

the strongest and most consistent mediator regardless of diagnostic status, while mentalizing and

avoidant attachment showed significant indirect effects in some groups but not others. Notably, the

mediation was complete: once the mediators were included, the direct association between autistic

traits and depressive symptoms was no longer significant, highlighting the central role of

psychosocial and cognitive processes in mediating this relationship.

Depression in autism presents a notable burden, not solely due to its high frequency, but also

because of challenges in its recognition and accurate diagnosis in clinical practice. The onset of

symptoms may be atypical or misattributed to core features of autism itself, such as social

withdrawal or reduced emotional expression (Chandrasekhar & Sikich, 2015). Our findings confirm

that individuals with higher autistic traits, particularly those in the autism group, report more severe

depressive symptoms than neurotypical controls. However, the psychological mechanisms linking

these traits to depression differ in strength and character across groups. This suggests that while

individuals with elevated autistic traits are broadly at increased risk for depression, as also found in

previous studies (Hollocks et al., 2019; Lai et al., 2019; Wigham et al., 2017), the pathways through

which this risk manifests may vary by diagnostic group, underscoring the importance of identifying

group-specific mechanisms to inform more tailored interventions.

One of the most notable findings of our study was that psychological flexibility was the

strongest and most consistent mediator between autistic traits and depressive symptoms across all

groups. This construct, defined as the ability to stay present while adapting behavior in line with

personal values (Kashdan & Rottenberg, 2010), has been validated for use in autistic adults (Aller

et al., 2022) and has been shown to buffer against emotional distress in both clinical and non-clinical

populations. While autism is characterized by behavioral inflexibility (APA, 2022), psychological

flexibility remains under-studied in this population. Previous research in neurotypical samples has

consistently linked lower flexibility to higher levels of depression and maladaptive coping (Bond

et al., 2011). A recent study by Lee et al. (2022) found that psychological flexibility, as measured

by the AAQ-II, alongside attachment anxiety, mediated the association between adverse parenting

experiences and later mental health outcomes in autistic individuals. This suggests that the capacity

to adaptively respond to distressing internal states may be a crucial resilience factor across

diagnostic boundaries. Consistent with this, theoretical models linking attachment and emotion

regulation to psychological flexibility further support this idea (Mikulincer & Shaver, 2019;

Zilberstein, 2014), highlighting overlapping pathways through which early relational experiences can shape later vulnerability. Despite being under-studied in autism, psychological flexibility appears to be a promising transdiagnostic construct with clear relevance for therapeutic interventions.

While psychological flexibility was the most universal pathway, other mediators, especially mentalization and avoidant attachment, showed group-specific effects. Mentalizing is often cited as a key factor in understanding depression in autism, in our study, it only emerged as a significant mediator in the non-ASD group. This result is somewhat unexpected given the extensive literature linking ASD with impairments in mentalizing and reflective functioning (Arioli et al., 2021; Brewer et al., 2017; Chung et al., 2014; Yirmiya et al., 1998). One possible explanation is that more severe or trait-like impairments in the ASD group may limit individuals' ability to reflect on and accurately report their mentalizing capacity, a phenomenon that is paradoxically consistent with the very definition of compromised reflective functioning. In contrast, subclinical or milder deficits in the non-ASD group may remain accessible to introspection and thus detectable via self-report tools. This divergence highlights a limitation of self-report-based measurement of mentalizing (Wendt et al., 2024) and suggests that more nuanced, performance-based or informant-rated approaches might be necessary to capture its true role in this population. While mentalizing deficits in ASD are well-established, and comparable in severity to those observed in schizophrenia (Bliksted et al., 2016; Fernandes et al., 2018; Oliver et al., 2021), future studies should explore the dimensional profile of mentalization (e.g., cognitive vs. affective, self vs. other) using more sensitive tools, particularly in self-diagnosed populations, where mixed patterns may reflect distinct compensatory or coping strategies.

Avoidant attachment was the only variable that significantly distinguished the ASD and ASD-sd groups in terms of mean scores, with self-diagnosed individuals reporting higher levels than those with a formal diagnosis. Interestingly, however, its mediating effect between autistic traits and depressive symptoms was significant only in the clinically diagnosed ASD group. This contrast raises important questions about the mechanisms underlying depressive vulnerability in self-diagnosed individuals. This divergence suggests that while avoidant tendencies may be more frequently endorsed by self-diagnosed individuals, potentially due to prolonged unmet support needs or social exclusion (Camus et al., 2024; Overton et al., 2024), they may carry greater psychological weight and reflect deeper relational disengagement among formally diagnosed individuals. In the ASD group, avoidant attachment may reflect longstanding relational disengagement, potentially rooted in early social-emotional developmental differences. This aligns with previous studies showing a high prevalence of insecure attachment in autism (Rutgers et al., 2004; Van IJzendoorn et al., 2007) and its association with affect regulation difficulties (Dagan et al., 2018; Mikulincer & Shaver, 2019).

In contrast, in the ASD-sd group, higher avoidant scores may not translate as directly into depressive symptoms, possibly due to different meaning-making (attributional) processes or

identity-related struggles that dominate their psychosocial experience (Davidson & and Henderson, 2010; Lewis, 2017). Instead their depressive symptoms could rather stem from social invalidation, lack of recognition, or difficulty accessing identity-affirming communities (Camus et al., 2024; Cooper et al., 2023). These individuals often report a weaker sense of belonging to the autistic community (Camus et al., 2024), which may reduce the protective effect commonly associated with community identification and support (Cooper et al., 2023; Crompton et al., 2024). Thus, while avoidant attachment is elevated in this group, its explanatory power for depression may be limited compared to that of other psychosocial mechanisms such as perceived isolation or unacknowledged identity. The role of attachment insecurity as a contributing, rather than central, mechanism may suggest indirect clinical relevance. Although this is not a primary therapeutic target, the patient's attachment style could influence treatment engagement and emotional availability during therapy (Fischer-Kern et al., 2022; Reiner et al., 2016).

The observed pattern of indirect effects indicates that high AQ scores alone do not explain the emergence of depression, but rather their interaction with these cognitive-affective mechanisms. These findings are consistent with previous research on neurotypical populations with high autistic traits, who also report more depressive symptoms when mentalizing capacity is impaired or when support is perceived as lacking (Beatson & Taryan, 2003; Besser & Priel, 2003; Carnelley et al., 1994; Malik et al., 2015). The overlap between attachment insecurity, poor mentalizing, and limited psychological flexibility may also hinder individuals in expressing their needs or seeking help—factors that are particularly relevant for those without formal diagnosis and who may not access support services (Davidson & and Henderson, 2010; Lewis, 2017).

Finally, perceived social support, while not the strongest mediator overall, remained an important indirect factor in the non-ASD group. Earlier findings showed that perceived lack of support, combined with social anxiety and camouflaging behaviors, contributes to depressive symptoms (Bernardin et al., 2021; Hull et al., 2017; Lai et al., 2019). For self-diagnosed individuals, these effects may be exacerbated by uncertainty around identity and lower community integration, as suggested in recent studies (Camus et al., 2024; Overton et al., 2024). This supports the need to conceptualize self-diagnosed autism not merely as a milder form of clinical ASD, but as a distinct presentation that may entail equal or greater emotional burden, especially in relation to unacknowledged needs and difficulties accessing appropriate support.

Clinical relevance and therapeutic implications

A significant number of individuals with autism spectrum disorder seek therapy for
depression, presenting unique challenges for effective treatment. Managing (both recognising and
treating) depression in this population is particularly difficult due to the complex interplay of social,
cognitive, and emotional factors inherent to ASD. Our findings underscore the need for
personalized, mechanism-informed interventions for depression in individuals with elevated
autistic traits. Rather than targeting autistic traits directly, therapeutic efforts may be more effective
when focused on transdiagnostic cognitive-affective processes, particularly psychological
flexibility, mentalizing capacities, and attachment-related functioning.

Psychological flexibility emerged as the most robust and universal mediator of depressive
symptoms across all groups, highlighting its value as a core resilience factor and key intervention
target. Enhancing flexibility could reduce emotional rigidity and support adaptive functioning,
particularly through structured approaches such as acceptance and commitment therapy, which has
growing empirical support in both neurotypical and autistic populations (Aller et al., 2022; Pahnke
et al., 2023).

Avoidant attachment, while more prevalent in the self-diagnosed group, exerted its mediating
effect only in the clinically diagnosed ASD group, suggesting that avoidant interpersonal strategies
may carry different meanings and consequences across diagnostic contexts. From a therapeutic
perspective, high levels of avoidant attachment can undermine treatment engagement, emotional
attunement, and the therapeutic alliance. Identifying these patterns early and tailoring relational
dynamics may improve adherence and outcomes (Fischer-Kern et al., 2022; Reiner et al., 2016).
Clinical experience drawn from mentalization-based treatment (MBT) protocols for avoidant
personality disorder suggests that patients with pronounced relational withdrawal benefit from
carefully paced interventions that scaffold mentalizing at a comfortable pace (Bateman et al., 2023).
These strategies emphasize trust-building and indirect engagement and may be highly applicable to
autistic individuals around avoidant attachment, supporting earlier identification and personalized
pacing in therapy.

Although mentalizing did not mediate depressive symptoms in the ASD group, this may
reflect the limits of self-awareness in this population, rather than a lack of clinical relevance.
Mentalization-based treatment, with its focus on enhancing reflective functioning in emotionally
charged contexts, may offer a promising yet underutilized avenue, provided that it is adapted to the
cognitive and social profiles of autistic individuals. Recent efforts to tailor MBT to ASD contexts,
while still limited, show preliminary promise (Cassel et al., 2007; Krämer et al., 2021).

Finally, the findings emphasize the importance of context-sensitive care for self-diagnosed
individuals, who often do not fit into existing diagnostic frameworks but nevertheless face clinically
significant distress. Interventions for this group should consider issues of identity, belonging, and
social validation, which may lie at the core of their depressive vulnerability.

In summary, our results support a shift away from diagnosis-based assumptions and toward

process-based treatment models, aligning with contemporary calls for personalized, transdiagnostic

care in both autism and mental health services more broadly.

Limitations

A major strength of our study is the use of self-report questionnaires administered online,

which enabled the collection of data from a relatively large and diverse sample. However, several

limitations must be acknowledged. First, the validity of self-report measures may be influenced by

participants' introspective accuracy, particularly for constructs like mentalizing, where limited self-

reflection may bias responses. Nonetheless, previous research suggests that subjective perceptions

may be more predictive of psychopathology than objective adversity alone (Danese & Widom, 2020).

Second, while the ASD group included individuals with formal clinical diagnoses, the ASD-

sd group was composed of self-identified autistic individuals without diagnostic confirmation.

Although we analyzed them as a distinct group, the heterogeneity within the ASD-sd population

remains a potential limitation, given that motivations for self-identification and levels of self-

awareness may vary widely. Furthermore, it cannot be ruled out that undiagnosed autistic individuals

may be present in the non-ASD group, potentially diluting group contrasts.

Additionally, the online recruitment strategy may have introduced sampling bias, favoring

individuals who are more self-aware, better resourced, or more engaged with autism-related content.

Importantly, we did not assess the effect of comorbid psychiatric diagnoses or medication use, both

of which may influence depressive symptoms and the mechanisms we examined. Although gender

was included as a covariate in our models, the overall sample showed a gender imbalance, limiting

the generalizability of our findings and highlighting the need for future studies with more

representative and stratified samples.

Finally, the cross-sectional design of our study precludes causal interpretations. Longitudinal

studies are needed to assess the temporal dynamics between autistic traits, mediating psychosocial

factors, and depressive symptoms.

Conclusions

This study provides a nuanced understanding of how psychosocial and cognitive factors

mediate the link between autistic traits and depressive symptoms across formally diagnosed, self-

identified, and neurotypical populations. Among the mediators examined, psychological flexibility

emerged as the most robust and consistent protective factor, while avoidant attachment and

mentalizing difficulties showed group-specific relevance. These findings highlight the need for

targeted, mechanism-based interventions that move beyond categorical diagnoses. Enhancing

psychological flexibility and addressing relational patterns, especially avoidant attachment, may be

particularly effective therapeutic entry points. Mentalization-based treatment, when appropriately

adapted, holds promise as a framework that integrates these mechanisms. Ultimately, our results

underscore the value of personalized approaches that recognize the heterogeneity of autistic

experiences and prioritize transdiagnostic resilience-building strategies in the prevention and

treatment of depression.

References

Ainsworth, M. D. S., Blehar, M. C., Waters, E., & Wall, S. (1978). *Patterns of attachment: A psychological*
study of the strange situation (pp. xviii, 391). Lawrence Erlbaum.
Akbari, V., Rahmatinejad, P., Shater, M. M., Vahedian, M., & Khalajinia, Z. (2020). Investigation of the
relationship of perceived social support and spiritual well-being with postpartum depression. *Journal*
of Education and Health Promotion, *9*(1), 174. https://doi.org/10.4103/jehp.jehp_56_20
Akdemir, D., Pehlivantürk, B., Unal, F., & Ozusta, S. (2009). [Comparison of attachment-related social
behaviors in autistic disorder and developmental disability]. *Turk Psikiyatri Dergisi = Turkish*
Journal of Psychiatry, *20*(2), 105–117.
Albantakis, L., Brandi, M.-L., Zillekens, I. C., Henco, L., Weindel, L., Thaler, H., Schliephake, L.,
Timmermans, B., & Schilbach, L. (2020). Alexithymic and autistic traits: Relevance for comorbid
depression and social phobia in adults with and without autism spectrum disorder. *Autism*, *24*(8),
2046–2056. https://doi.org/10.1177/1362361320936024
Aller, T. B., Barrett, T., Levin, M. E., & McClain, M. B. (2022). Measuring psychological flexibility in
autistic adults: Examining the validity and reliability of the AAQ-II, BEAQ, and VQ. *Journal of*
Contextual Behavioral Science, *26*, 125–133. https://doi.org/10.1016/j.jcbs.2022.09.001
Alon, R. (2019). Social support and post-crisis growth among mothers of children with autism spectrum
disorder and mothers of children with down syndrome. *Research in Developmental Disabilities*, *90*,
22–30. https://doi.org/10.1016/j.ridd.2019.04.010
American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders* (Fifth
Edition). American Psychiatric Association. https://doi.org/10.1176/appi.books.9780890425596
American Psychiatric Association. (2022). *Diagnostic and Statistical Manual of Mental Disorders* (5th ed.,
text rev.). https://doi.org/10.1176/appi.books.9780890425787
Arioli, M., Cattaneo, Z., Ricciardi, E., & Canessa, N. (2021). Overlapping and specific neural correlates for
empathizing, affective mentalizing, and cognitive mentalizing: A coordinate-based meta-analytic
study. *Human Brain Mapping*, *42*(14), 4777–4804. https://doi.org/10.1002/hbm.25570
Banker, S. M., Harrington, M., Schafer, M., Na, S., Heflin, M., Barkley, S., Trayvick, J., Peters, A. W.,
Thinakaran, A. A., Schiller, D., Foss-Feig, J. H., & Gu, X. (2025). Phenotypic divergence between
individuals with self-reported autistic traits and clinically ascertained autism. *Nature Mental Health*,
3(3), 286–297. https://doi.org/10.1038/s44220-025-00385-8
Baron-Cohen, S. (2000). Theory of mind and autism: A review. In *International Review of Research in*
Mental Retardation (Vol. 23, pp. 169–184). Academic Press. https://doi.org/10.1016/S0074-
7750(00)80010-5
Baron-Cohen, S., Cassidy, S., Auyeung, B., Allison, C., Achoukhi, M., Robertson, S., Pohl, A., & Lai, M.-C.
(2014). Attenuation of Typical Sex Differences in 800 Adults with Autism vs. 3,900 Controls. *PLOS*
ONE, *9*(7), e102251. https://doi.org/10.1371/journal.pone.0102251
Baron-Cohen, S., Leslie, A. M., & Frith, U. (1985). Does the autistic child have a 'theory of mind'?
Cognition, *21*(1), 37–46. https://doi.org/10.1016/0010-0277(85)90022-8
Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The Autism-Spectrum
Quotient (AQ): Evidence from Asperger Syndrome/High-Functioning Autism, Malesand Females,
Scientists and Mathematicians. *Journal of Autism and Developmental Disorders*, *31*(1), 5–17.
https://doi.org/10.1023/A:1005653411471
Bateman, A., Campbell, C., Debbané, M., Luyten, P., & Fonagy, P. (Eds.). (2023). Avoidant Personality
Disorder. In *Cambridge Guide to Mentalization-Based Treatment (MBT)* (pp. 203–229). Cambridge
University Press. https://doi.org/10.1017/9781108896054.009
Bauminger, N., Solomon, M., & Rogers, S. J. (2010). Predicting Friendship Quality in Autism Spectrum
Disorders and Typical Development. *Journal of Autism and Developmental Disorders*, *40*(6), 751–
761. https://doi.org/10.1007/s10803-009-0928-8
Beatson, J., & Taryan, S. (2003). Predisposition to Depression: The Role of Attachment. *Australian & New*
Zealand Journal of Psychiatry, *37*(2), 219–225. https://doi.org/10.1046/j.1440-1614.2003.01126.x
Beck, A. T., Steer, R. A., & Carbin, M. G. (1988). Psychometric properties of the Beck Depression

Inventory: Twenty-five years of evaluation. *Clinical Psychology Review*, *8*(1), 77–100. https://doi.org/10.1016/0272-7358(88)90050-5

Beffel, J. H., Cary, K. M., Nuttall, A. K., Chopik, W. J., & Maas, M. K. (2021). Associations between the broad autism phenotype, adult attachment, and relationship satisfaction among emerging adults. *Personality and Individual Differences*, *168*, 110409. https://doi.org/10.1016/j.paid.2020.110409

Bernardin, C. J., Lewis, T., Bell, D., & Kanne, S. (2021). Associations between social camouflaging and internalizing symptoms in autistic and non-autistic adolescents. *Autism: The International Journal of Research and Practice*, *25*(6), 1580–1591. https://doi.org/10.1177/1362361321997284

Besser, A., & Priel, B. (2003). A Multisource Approach to Self-Critical Vulnerability to Depression: The Moderating Role of Attachment. *Journal of Personality*, *71*, 515–555. https://doi.org/10.1111/1467-6494.7104002

Bishop-Fitzpatrick, L., Mazefsky, C. A., & Eack, S. M. (2018). The combined impact of social support and perceived stress on quality of life in adults with autism spectrum disorder and without intellectual disability. *Autism*, *22*(6), 703–711. https://doi.org/10.1177/1362361317703090

Bliksted, V., Ubukata, S., & Koelkebeck, K. (2016). Discriminating autism spectrum disorders from schizophrenia by investigation of mental state attribution on an on-line mentalizing task: A review and meta-analysis. *Schizophrenia Research*, *171*(1), 16–26. https://doi.org/10.1016/j.schres.2016.01.037

Bond, F. W., Hayes, S. C., Baer, R. A., Carpenter, K. M., Guenole, N., Orcutt, H. K., Waltz, T., & Zettle, R. D. (2011). Preliminary psychometric properties of the Acceptance and Action Questionnaire-II: A revised measure of psychological inflexibility and experiential avoidance. *Behavior Therapy*, *42*(4), 676–688. https://doi.org/10.1016/j.beth.2011.03.007

Bosmans, G., Braet, C., & Van Vlierberghe, L. (2010). Attachment and symptoms of psychopathology: Early maladaptive schemas as a cognitive link? *Clinical Psychology & Psychotherapy*, *17*(5), 374–385. https://doi.org/10.1002/cpp.667

Bougeard, C., Picarel-Blanchot, F., Schmid, R., Campbell, R., & Buitelaar, J. (2021). Prevalence of Autism Spectrum Disorder and Co-morbidities in Children and Adolescents: A Systematic Literature Review. *Frontiers in Psychiatry*, *12*, 744709. https://doi.org/10.3389/fpsyt.2021.744709

Brewer, N., Young, R. L., & Barnett, E. (2017). Measuring Theory of Mind in Adults with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*, *47*(7), 1927–1941. https://doi.org/10.1007/s10803-017-3080-x

Cai, R. Y., Richdale, A. L., Dissanayake, C., & Uljarević, M. (2018). Brief Report: Inter-Relationship between Emotion Regulation, Intolerance of Uncertainty, Anxiety, and Depression in Youth with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*, *48*(1), 316–325. https://doi.org/10.1007/s10803-017-3318-7

Camus, L., Jones, K., O'Dowd, E., Auyeung, B., Rajendran, G., & Stewart, M. E. (2024). Autistic Traits and Psychosocial Predictors of Depressive Symptoms. *Journal of Autism and Developmental Disorders*. https://doi.org/10.1007/s10803-024-06361-y

Cantazaro, A., & Wei, M. (2010). Adult Attachment, Dependence, Self-Criticism, and Depressive Symptoms: A Test of a Mediational Model. *Journal of Personality*, *78*(4), 1135–1162. https://doi.org/10.1111/j.1467-6494.2010.00645.x

Canty-Mitchell, J., & Zimet, G. D. (2000). Psychometric properties of the Multidimensional Scale of Perceived Social Support in urban adolescents. *American Journal of Community Psychology*, *28*(3), 391–400. https://doi.org/10.1023/A:1005109522457

Capps, L., Sigman, M., & Mundy, P. (1994). Attachment security in children with autism. *Development and Psychopathology*, *6*(2), 249–261. https://doi.org/10.1017/S0954579400004569

Carnelley, K. B., Pietromonaco, P. R., & Jaffe, K. (1994). Depression, working models of others, and relationship functioning. *Journal of Personality and Social Psychology*, *66*, 127–140. https://doi.org/10.1037/0022-3514.66.1.127

Cassel, T. D., Messinger, D. S., Ibanez, L. V., Haltigan, J. D., Acosta, S. I., & Buchman, A. C. (2007). Early Social and Emotional Communication in the Infant Siblings of Children with Autism Spectrum Disorders: An Examination of the Broad Phenotype. *Journal of Autism and Developmental Disorders*, *37*(1), 122–132. https://doi.org/10.1007/s10803-006-0337-1

Chandler, F., & Dissanayake, C. (2014). An investigation of the security of caregiver attachment during middle childhood in children with high-functioning autistic disorder. *Autism*, *18*(5), 485–492. https://doi.org/10.1177/1362361313486205

Chandrasekhar, T., & Sikich, L. (2015). Challenges in the diagnosis and treatment of depression in autism spectrum disorders across the lifespan. *Dialogues in Clinical Neuroscience*, *17*(2), 219. https://doi.org/10.31887/DCNS.2015.17.2/tchandrasekhar

Chung, Y. S., Barch, D., & Strube, M. (2014). A Meta-Analysis of Mentalizing Impairments in Adults With Schizophrenia and Autism Spectrum Disorder. *Schizophrenia Bulletin*, *40*(3), 602–616. https://doi.org/10.1093/schbul/sbt048

Collins, N. L. (1996). Working models of attachment: Implications for explanation, emotion, and behavior.

Journal of Personality and Social Psychology, *71*(4), 810–832. https://doi.org/10.1037/0022-3514.71.4.810

Cooper, K., Russell, A. J., Lei, J., & Smith, L. G. (2023). The impact of a positive autism identity and autistic community solidarity on social anxiety and mental health in autistic young people. *Autism*, *27*(3), 848–857. https://doi.org/10.1177/13623613221118351

Crompton, C. J., Fotheringham, F., Cebula, K., Webber, C., Foley, S., & Fletcher-Watson, S. (2024). Neurodivergent-designed and neurodivergent-led peer support in school: A feasibility and acceptability study of the neurodivergent peer support toolkit (NEST). *Neurodiversity*, *2*, 27546330241275248. https://doi.org/10.1177/27546330241275248

Dagan, O., Facompré, C. R., & Bernard, K. (2018). Adult attachment representations and depressive symptoms: A meta-analysis. *Journal of Affective Disorders*, *236*, 274–290. https://doi.org/10.1016/j.jad.2018.04.091

Davidson, J., & and Henderson, V. L. (2010). 'Coming out' on the spectrum: Autism, identity and disclosure. *Social & Cultural Geography*, *11*(2), 155–170. https://doi.org/10.1080/14649360903525240

Dawson, G., Webb, S. J., Carver, L., Panagiotides, H., & McPartland, J. (2004). Young children with autism show atypical brain responses to fearful versus neutral facial expressions of emotion. *Developmental Science*, *7*(3), 340–359. https://doi.org/10.1111/j.1467-7687.2004.00352.x

Dissanayake, C., & Crossley, S. A. (1997). Autistic Children's Responses to Separation and Reunion with Their Mothers. *Journal of Autism and Developmental Disorders*, *27*(3), 295–312. https://doi.org/10.1023/A:1025802515241

Doherty, N. A., & Feeney, J. A. (2004). The composition of attachment networks throughout the adult years. *Personal Relationships*, *11*, 469–488. https://doi.org/10.1111/j.1475-6811.2004.00093.x

Eid, M., Gollwitzer, M., & Schmitt, M. (2013). *Statistik und Forschungsmethoden: Lehrbuch* (3., korrigierte Auflage, Online-Ausgabe). Beltz. https://content-select.com/index.php?id=bib_view&ean=9783621278348

Eisenbeck, N., & Szabó-Bartha, A. (2018). Validation of the Hungarian version of the Acceptance and Action Questionnaire-II (AAQ-II). *Journal of Contextual Behavioral Science*, *9*, 80–87. https://doi.org/10.1016/j.jcbs.2018.07.007

Fekete, K., Torok, E., Kelemen, O., Makkos, Z., Csigo, K., & Keri, S. (2019). [Dimensions of mentalization in psychotic disorders]. *Neuropsychopharmacologia Hungarica: A Magyar Pszichofarmakologiai Egyesulet Lapja = Official Journal of the Hungarian Association of Psychopharmacology*, *21*(1), 5–11.

Fernandes, J. M., Cajão, R., Lopes, R., Jerónimo, R., & Barahona-Corrêa, J. B. (2018). Social Cognition in Schizophrenia and Autism Spectrum Disorders: A Systematic Review and Meta-Analysis of Direct Comparisons. *Frontiers in Psychiatry*, *9*. https://www.frontiersin.org/articles/10.3389/fpsyt.2018.00504

Fischer-Kern, M., Fonagy, P., Kapusta, N. D., Luyten, P., Boss, S., Naderer, A., Blüml, V., & Leithner, K. (2013). Mentalizing in Female Inpatients With Major Depressive Disorder. *The Journal of Nervous and Mental Disease*, *201*(3), 202. https://doi.org/10.1097/NMD.0b013e3182845c0a

Fischer-Kern, M., Tmej, A., Naderer, A., Zimmermann, J., & Nolte, T. (2022). Failure to resolve loss and compromised mentalizing in female inpatients with major depressive disorder. *Attachment & Human Development*, *24*(4), 503–524. https://doi.org/10.1080/14616734.2021.2015794

Fonagy, P., & Allison, E. (2014). The role of mentalizing and epistemic trust in the therapeutic relationship. *Psychotherapy*, *51*(3), 372–380. https://doi.org/10.1037/a0036505

Fonagy, P., & Bateman, A. W. (2016). Adversity, attachment, and mentalizing. *Comprehensive Psychiatry*, *64*, 59–66. https://doi.org/10.1016/j.comppsych.2015.11.006

Fonagy, P., & Luyten, P. (2009). A developmental, mentalization-based approach to the understanding and treatment of borderline personality disorder. *Development and Psychopathology*, *21*(4), 1355–1381. https://doi.org/10.1017/S0954579409990198

Fonagy, P., Steele, M., Steele, H., Moran, G. S., & Higgitt, A. C. (1991). The capacity for understanding mental states: The reflective self in parent and child and its significance for security of attachment. *Infant Mental Health Journal*, *12*(3), 201–218. https://doi.org/10.1002/1097-0355(199123)12:3<201::AID-IMHJ2280120307>3.0.CO;2-7

Fraley, R. C. (2002). Attachment Stability From Infancy to Adulthood: Meta-Analysis and Dynamic Modeling of Developmental Mechanisms. *Personality and Social Psychology Review*, *6*(2), 123–151. https://doi.org/10.1207/S15327957PSPR0602_03

Gallitto, E., & Leth-Steensen, C. (2015). Autistic traits and adult attachment styles. *Personality and Individual Differences*, *79*, 63–67. https://doi.org/10.1016/j.paid.2015.01.032

Gariépy, G., Honkaniemi, H., & Quesnel-Vallée, A. (2016). Social support and protection from depression: Systematic review of current findings in Western countries. *The British Journal of Psychiatry: The Journal of Mental Science*, *209*(4), 284–293. https://doi.org/10.1192/bjp.bp.115.169094

Grzadzinski, R. L., Luyster, R., Spencer, A. G., & Lord, C. (2014). Attachment in young children with autism spectrum disorders: An examination of separation and reunion behaviors with both mothers and

fathers. *Autism: The International Journal of Research and Practice*, *18*(2), 85–96. https://doi.org/10.1177/1362361312467235

Hausberg, M. C., Schulz, H., Piegler, T., Happach, C. G., Klöpper, M., Brütt, A. L., Sammet, I., & Andreas, S. (2012). Is a self-rated instrument appropriate to assess mentalization in patients with mental disorders? Development and first validation of the Mentalization Questionnaire (MZQ). *Psychotherapy Research*, *22*(6), 699–709. https://doi.org/10.1080/10503307.2012.709325

Hazan, C., & Zeifman, D. (1994). Sex and the psychological tether. In *Attachment processes in adulthood* (pp. 151–178). Jessica Kingsley Publishers.

He, B., Wongpakaran, T., Wongpakaran, N., & Wedding, D. (2022). Marital Satisfaction and Perceived Family Support in Families of Children with Autistic Spectrum Disorder: Dyadic Analysis. *Healthcare*, *10*(7), Article 7. https://doi.org/10.3390/healthcare10071227

Hefner, J., & Eisenberg, D. (2009). Social support and mental health among college students. *American Journal of Orthopsychiatry*, *79*, 491–499. https://doi.org/10.1037/a0016918

Hollocks, M. J., Lerh, J. W., Magiati, I., Meiser-Stedman, R., & Brugha, T. S. (2019). Anxiety and depression in adults with autism spectrum disorder: A systematic review and meta-analysis. *Psychological Medicine*, *49*(4), 559–572. https://doi.org/10.1017/S0033291718002283

Hosmer, D. (2013). Introduction to the Logistic Regression Model. In *Applied Logistic Regression* (pp. 1–33). John Wiley & Sons, Ltd. https://doi.org/10.1002/9781118548387.ch1

Hossain, M. M., Khan, N., Sultana, A., Ma, P., McKyer, E. L. J., Ahmed, H. U., & Purohit, N. (2020). Prevalence of comorbid psychiatric disorders among people with autism spectrum disorder: An umbrella review of systematic reviews and meta-analyses. *Psychiatry Research*, *287*, 112922. https://doi.org/10.1016/j.psychres.2020.112922

Hudson, C. C., Hall, L., & Harkness, K. L. (2019). Prevalence of Depressive Disorders in Individuals with Autism Spectrum Disorder: A Meta-Analysis. *Journal of Abnormal Child Psychology*, *47*(1), 165–175. https://doi.org/10.1007/s10802-018-0402-1

Hull, L., Petrides, K. V., Allison, C., Smith, P., Baron-Cohen, S., Lai, M.-C., & Mandy, W. (2017). "Putting on My Best Normal": Social Camouflaging in Adults with Autism Spectrum Conditions. *Journal of Autism and Developmental Disorders*, *47*(8), 2519–2534. https://doi.org/10.1007/s10803-017-3166-5

Inoue, Y., Yamada, K., & Kanba, S. (2006). Deficit in theory of mind is a risk for relapse of major depression. *Journal of Affective Disorders*, *95*(1), 125–127. https://doi.org/10.1016/j.jad.2006.04.018

Kashdan, T. B., & Rottenberg, J. (2010). Psychological flexibility as a fundamental aspect of health. *Clinical Psychology Review*, *30*(7), 865–878. https://doi.org/10.1016/j.cpr.2010.03.001

Keenan, B. M., Newman, L. K., Gray, K. M., & Rinehart, N. J. (2017). A qualitative study of attachment relationships in ASD during middle childhood. *Attachment & Human Development*, *19*(1), 1–21. https://doi.org/10.1080/14616734.2016.1246580

Koren-Karie, N., Oppenheim, D., Dolev, S., & Yirmiya, N. (2009). Mothers of securely attached children with autism spectrum disorder are more sensitive than mothers of insecurely attached children. *Journal of Child Psychology and Psychiatry*, *50*(5), 643–650. https://doi.org/10.1111/j.1469-7610.2008.02043.x

Krämer, K., Vetter, A., Schultz-Venrath, U., Vogeley, K., & Reul, S. (2021). Mentalization-Based Treatment in Groups for Adults With Autism Spectrum Disorder. *Frontiers in Psychology*, *12*, 708557. https://doi.org/10.3389/fpsyg.2021.708557

Lai, M.-C., & Baron-Cohen, S. (2015). Identifying the lost generation of adults with autism spectrum conditions. *The Lancet Psychiatry*, *2*(11), 1013–1027. https://doi.org/10.1016/S2215-0366(15)00277-1

Lai, M.-C., Kassee, C., Besney, R., Bonato, S., Hull, L., Mandy, W., Szatmari, P., & Ameis, S. H. (2019). Prevalence of co-occurring mental health diagnoses in the autism population: A systematic review and meta-analysis. *The Lancet Psychiatry*, *6*(10), 819–829. https://doi.org/10.1016/S2215-0366(19)30289-5

Leader, G., Barrett, A., Ferrari, C., Casburn, M., Maher, L., Naughton, K., Arndt, S., & Mannion, A. (2021). Quality of life, gastrointestinal symptoms, sleep problems, social support, and social functioning in adults with autism spectrum disorder. *Research in Developmental Disabilities*, *112*, 103915. https://doi.org/10.1016/j.ridd.2021.103915

Lee, J. Y. S., Whittingham, K., & Mitchell, A. E. (2022). Childhood experiences of being parented, adult attachment, psychological inflexibility, social engagement, and mental health of autistic adults. *Research in Developmental Disabilities*, *130*, 104343. https://doi.org/10.1016/j.ridd.2022.104343

Lee, L., Harkness, K. L., Sabbagh, M. A., & Jacobson, J. A. (2005). Mental state decoding abilities in clinical depression. *Journal of Affective Disorders*, *86*(2), 247–258. https://doi.org/10.1016/j.jad.2005.02.007

Lewis, L. F. (2017). A Mixed Methods Study of Barriers to Formal Diagnosis of Autism Spectrum Disorder in Adults. *Journal of Autism and Developmental Disorders*, *47*(8), 2410–2424.

https://doi.org/10.1007/s10803-017-3168-3

Lim, G. Y., Tam, W. W., Lu, Y., Ho, C. S., Zhang, M. W., & Ho, R. C. (2018). Prevalence of Depression in
the Community from 30 Countries between 1994 and 2014. *Scientific Reports*, *8*(1), Article 1.
https://doi.org/10.1038/s41598-018-21243-x

Liu, L., Gou, Z., & Zuo, J. (2016). Social support mediates loneliness and depression in elderly people.
Journal of Health Psychology, *21*(5), 750–758. https://doi.org/10.1177/1359105314536941

Lombardo, M. V., Lai, M.-C., Auyeung, B., Holt, R. J., Allison, C., Smith, P., Chakrabarti, B., Ruigrok, A.
N. V., Suckling, J., Bullmore, E. T., Bailey, A. J., Baron-Cohen, S., Bolton, P. F., Bullmore, E. T.,
Carrington, S., Catani, M., Chakrabarti, B., Craig, M. C., Daly, E. M., … Baron-Cohen, S. (2016).
Unsupervised data-driven stratification of mentalizing heterogeneity in autism. *Scientific Reports*,
6(1), Article 1. https://doi.org/10.1038/srep35333

Lu, M., Chen, J., He, W., Pang, F., & Zou, Y. (2021). Association between perceived social support of
parents and emotional/behavioral problems in children with ASD: A chain mediation model.
Research in Developmental Disabilities, *113*, 103933. https://doi.org/10.1016/j.ridd.2021.103933

Luyten, P., Campbell, C., Allison, E., & Fonagy, P. (2020). The Mentalizing Approach to Psychopathology:
State of the Art and Future Directions. *Annual Review of Clinical Psychology*, *16*, 297–325.
https://doi.org/10.1146/annurev-clinpsy-071919-015355

Luyten, P., & Fonagy, P. (2018). The stress–reward–mentalizing model of depression: An integrative
developmental cascade approach to child and adolescent depressive disorder based on the Research
Domain Criteria (RDoC) approach. *Clinical Psychology Review*, *64*, 87–98.
https://doi.org/10.1016/j.cpr.2017.09.008

Malik, S., Wells, A., & Wittkowski, A. (2015). Emotion regulation as a mediator in the relationship between
attachment and depressive symptomatology: A systematic review. *Journal of Affective Disorders*,
172, 428–444. https://doi.org/10.1016/j.jad.2014.10.007

McDonald, T. A. M. (2020). Autism Identity and the "Lost Generation": Structural Validation of the Autism
Spectrum Identity Scale and Comparison of Diagnosed and Self-Diagnosed Adults on the Autism
Spectrum. *Autism in Adulthood*, *2*(1), 13–23. https://doi.org/10.1089/aut.2019.0069

Mikulincer, M., & Shaver, P. R. (2012). An attachment perspective on psychopathology. *World Psychiatry:
Official Journal of the World Psychiatric Association (WPA)*, *11*(1), 11–15.
https://doi.org/10.1016/j.wpsyc.2012.01.003

Mikulincer, M., & Shaver, P. R. (2017). *Attachment in Adulthood: Second Edition: Structure, Dynamics, and
Change*. Guilford Press. https://www.guilford.com/books/Attachment-in-Adulthood/Mikulincer-
Shaver/9781462533817

Mikulincer, M., & Shaver, P. R. (2019). Attachment orientations and emotion regulation. *Current Opinion in
Psychology*, *25*, 6–10. https://doi.org/10.1016/j.copsyc.2018.02.006

Miloseva, L., Vukosavljevic-Gvozden, T., Richter, K., Milosev, V., & Niklewski, G. (2017). Perceived social
support as a moderator between negative life events and depression in adolescence: Implications for
prediction and targeted prevention. *EPMA Journal*, *8*(3), 237–245. https://doi.org/10.1007/s13167-
017-0095-5

Naber, F. B. A., Swinkels, S. H. N., Buitelaar, J. K., Bakermans-Kranenburg, M. J., van IJzendoorn, M. H.,
Dietz, C., van Daalen, E., & van Engeland, H. (2007). Attachment in toddlers with autism and other
developmental disorders. *Journal of Autism and Developmental Disorders*, *37*(6), 1123–1138.
https://doi.org/10.1007/s10803-006-0255-2

Nickerson, A. B., & Nagle, R. J. (2005). Parent and Peer Attachment in Late Childhood and Early
Adolescence. *The Journal of Early Adolescence*, *25*, 223–249.
https://doi.org/10.1177/0272431604274174

Nuske, H. J., Vivanti, G., & Dissanayake, C. (2013). Are emotion impairments unique to, universal, or
specific in autism spectrum disorder? A comprehensive review. *Cognition & Emotion*, *27*(6), 1042–
1061. https://doi.org/10.1080/02699931.2012.762900

Oliver, L. D., Moxon-Emre, I., Lai, M.-C., Grennan, L., Voineskos, A. N., & Ameis, S. H. (2021). Social
Cognitive Performance in Schizophrenia Spectrum Disorders Compared With Autism Spectrum
Disorder: A Systematic Review, Meta-analysis, and Meta-regression. *JAMA Psychiatry*, *78*(3), 281–
292. https://doi.org/10.1001/jamapsychiatry.2020.3908

Őri D., Kapornai K., Baji I., & Kiss E. (2021). [Psychometric properties of the Hungarian Adult Attachment
Scale]. *Ideggyógyászati Szemle*, *73*(3–4), 87–98. https://doi.org/10.18071/isz.74.0087

Overton, G. L., Marsà-Sambola, F., Martin, R., & Cavenagh, P. (2024). Understanding the Self-identification
of Autism in Adults: A Scoping Review. *Review Journal of Autism and Developmental Disorders*,
11(4), 682–702. https://doi.org/10.1007/s40489-023-00361-x

Pahnke, J., Jansson-Fröjmark, M., Andersson, G., Bjureberg, J., Jokinen, J., Bohman, B., & Lundgren, T.
(2023). Acceptance and commitment therapy for autistic adults: A randomized controlled pilot study
in a psychiatric outpatient setting. *Autism*, *27*(5), 1461–1476.
https://doi.org/10.1177/13623613221140749

Papp-Zipernovszky, O., Kékesi, M. Z., & Jámbori, S. (2017). A Multidimenzionális Észlelt Társas

Támogatás Kérdőív magyar nyelvű validálása. *Mentálhigiéné és Pszichoszomatika*, *18*(3), 230–262.
https://doi.org/10.1556/0406.18.2017.011

Raque-Bogdan, T. L., Ericson, S. K., Jackson, J., Martin, H. M., & Bryan, N. A. (2011). Attachment and
mental and physical health: Self-compassion and mattering as mediators. *Journal of Counseling*
Psychology, *58*, 272–278. https://doi.org/10.1037/a0023041

Reiner, I., Bakermans-Kranenburg, M. J., Van IJzendoorn, M. H., Fremmer-Bombik, E., & Beutel, M.
(2016). Adult attachment representation moderates psychotherapy treatment efficacy in clinically
depressed inpatients. *Journal of Affective Disorders*, *195*, 163–171.
https://doi.org/10.1016/j.jad.2016.02.024

Robin, X., Turck, N., Hainard, A., Tiberti, N., Lisacek, F., Sanchez, J.-C., & Müller, M. (2011). pROC: An
open-source package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics*, *12*(1),
77. https://doi.org/10.1186/1471-2105-12-77

Rogers, S. J., Ozonoff, S., & Maslin-cole, C. (1993). Developmental Aspects of Attachment Behavior in
Young Children with Pervasive Developmental Disorders. *Journal of the American Academy of*
Child & Adolescent Psychiatry, *32*(6), 1274–1282. https://doi.org/10.1097/00004583-199311000-
00023

Rosseel, Y. (2012). lavaan: An R Package for Structural Equation Modeling. *Journal of Statistical Software*,
48, 1–36. https://doi.org/10.18637/jss.v048.i02

Rutgers, A. H., Bakermans-Kranenburg, M. J., van Ijzendoorn, M. H., & van Berckelaer-Onnes, I. A. (2004).
Autism and attachment: A meta-analytic review. *Journal of Child Psychology and Psychiatry*, *45*(6),
1123–1134. https://doi.org/10.1111/j.1469-7610.2004.t01-1-00305.x

Rutten, B. P. F., Hammels, C., Geschwind, N., Menne-Lothmann, C., Pishva, E., Schruers, K., van den Hove,
D., Kenis, G., van Os, J., & Wichers, M. (2013). Resilience in mental health: Linking psychological
and neurobiological perspectives. *Acta Psychiatrica Scandinavica*, *128*(1), 3–20.
https://doi.org/10.1111/acps.12095

Schiltz, H. K., McVey, A. J., Dolan Wozniak, B., Haendel, A. D., Stanley, R., Arias, A., Gordon, N., & Van
Hecke, A. V. (2021). The role of loneliness as a mediator between autism features and mental health
among autistic young adults. *Autism*, *25*(2), 545–555. https://doi.org/10.1177/1362361320967789

Shaver, P. R., Belsky, J., & Brennan, K. A. (2000). The adult attachment interview and self-reports of
romantic attachment: Associations across domains and methods. *Personal Relationships*, *7*, 25–43.
https://doi.org/10.1111/j.1475-6811.2000.tb00002.x

Siedlecki, K. L., Salthouse, T. A., Oishi, S., & Jeswani, S. (2014). The Relationship Between Social Support
and Subjective Well-Being Across Age. *Social Indicators Research*, *117*(2), 561–576.
https://doi.org/10.1007/s11205-013-0361-4

Taylor, E. L., Target, M., & Charman, T. (2008). Attachment in adults with high-functioning autism.
Attachment & Human Development, *10*(2), 143–163. https://doi.org/10.1080/14616730802113687

Teague, S. J., Gray, K. M., Tonge, B. J., & Newman, L. K. (2017). Attachment in children with autism
spectrum disorder: A systematic review. *Research in Autism Spectrum Disorders*, *35*, 35–50.
https://doi.org/10.1016/j.rasd.2016.12.002

Thiel, T., Riedelbauch, S., Gaigg, S., Roessner, V., & Ring, M. (2024). The impact of depressive and anxious
symptoms on quality of life in adults on the autism spectrum. *Autism Research*, *17*(6), 1161–1174.
https://doi.org/10.1002/aur.3144

Torquati, J. C., & Vazsonyi, A. T. (1999). Attachment as an Organizational Construct for Affect, Appraisals,
and Coping of Late Adolescent Females. *Journal of Youth and Adolescence*, *28*(5), 545–562.
https://doi.org/10.1023/A:1021602609489

Tsermentseli, S. (2022). Self-esteem moderates the impact of perceived social support on the life satisfaction
of adults with autism spectrum disorder. *Autism & Developmental Language Impairments*, *7*,
23969415221147430. https://doi.org/10.1177/23969415221147430

Umagami, K., Remington, A., Lloyd-Evans, B., Davies, J., & Crane, L. (2022). Loneliness in autistic adults:
A systematic review. *Autism*, *26*(8), 2117–2135. https://doi.org/10.1177/13623613221077721

Van IJzendoorn, M. H., Rutgers, A. H., Bakermans-Kranenburg, M. J., Swinkels, S. H. N., Van Daalen, E.,
Dietz, C., Naber, F. B. A., Buitelaar, J. K., & Van Engeland, H. (2007). Parental Sensitivity and
Attachment in Children With Autism Spectrum Disorder: Comparison With Children With Mental
Retardation, With Language Delays, and With Typical Development. *Child Development*, *78*(2),
597–608. https://doi.org/10.1111/j.1467-8624.2007.01016.x

Velikonja, T., Fett, A.-K., & Velthorst, E. (2019). Patterns of Nonsocial and Social Cognitive Functioning in
Adults With Autism Spectrum Disorder: A Systematic Review and Meta-analysis. *JAMA*
Psychiatry, *76*(2), 135–151. https://doi.org/10.1001/jamapsychiatry.2018.3645

Wang, J., Mann, F., Lloyd-Evans, B., Ma, R., & Johnson, S. (2018). Associations between loneliness and
perceived social support and outcomes of mental health problems: A systematic review. *BMC*
Psychiatry, *18*(1), 156. https://doi.org/10.1186/s12888-018-1736-5

Wang, Y., Wang, Y., Chen, S., Zhu, C., & Wang, K. (2008). Theory of mind disability in major depression
with or without psychotic symptoms: A componential view. *Psychiatry Research*, *161*(2), 153–161.

https://doi.org/10.1016/j.psychres.2007.07.018

Wendt, L. P., Zimmermann, J., Spitzer, C., & Müller, S. (2024). Mindreading measures misread? A multimethod investigation into the validity of self-report and task-based approaches. *Psychological Assessment*, *36*(5), 365–378. https://doi.org/10.1037/pas0001310

Wigham, S., Barton, S., Parr, J. R., & Rodgers, J. (2017). A Systematic Review of the Rates of Depression in Children and Adults With High-Functioning Autism Spectrum Disorder. *Journal of Mental Health Research in Intellectual Disabilities*, *10*(4), 267–287. https://doi.org/10.1080/19315864.2017.1299267

Wittchen, H.-U., & Jacobi, F. (2005). Size and burden of mental disorders in Europe—A critical review and appraisal of 27 studies. *European Neuropsychopharmacology*, *15*(4), 357–376. https://doi.org/10.1016/j.euroneuro.2005.04.012

Wittchen, H.-U., Jacobi, F., Rehm, J., Gustavsson, A., Svensson, M., Jönsson, B., Olesen, J., Allgulander, C., Alonso, J., Faravelli, C., Fratiglioni, L., Jennum, P., Lieb, R., Maercker, A., van Os, J., Preisig, M., Salvador-Carulla, L., Simon, R., & Steinhausen, H.-C. (2011). The size and burden of mental disorders and other disorders of the brain in Europe 2010. *European Neuropsychopharmacology*, *21*(9), 655–679. https://doi.org/10.1016/j.euroneuro.2011.07.018

Wu, C.-L., An, C.-P., Tseng, L.-P., Chen, H.-C., Chan, Y.-C., Cho, S.-L., & Tsai, M.-L. (2015). Fear of being laughed at with relation to parent attachment in individuals with autism. *Research in Autism Spectrum Disorders*, *10*, 116–123. https://doi.org/10.1016/j.rasd.2014.11.004

Yasinski, C., Hayes, A. M., Ready, C. B., Abel, A., Görg, N., & Kuyken, W. (2020). Processes of change in cognitive behavioral therapy for treatment-resistant depression: Psychological flexibility, rumination, avoidance, and emotional processing. *Psychotherapy Research: Journal of the Society for Psychotherapy Research*, *30*(8), 983–997. https://doi.org/10.1080/10503307.2019.1699972

Yirmiya, N., Erel, O., Shaked, M., & Solomonica-Levi, D. (1998). Meta-analyses comparing theory of mind abilities of individuals with autism, individuals with mental retardation, and normally developing individuals. *Psychological Bulletin*, *124*, 283–307. https://doi.org/10.1037/0033-2909.124.3.283

Zeidan, J., Fombonne, E., Scorah, J., Ibrahim, A., Durkin, M. S., Saxena, S., Yusuf, A., Shih, A., & Elsabbagh, M. (2022). Global prevalence of autism: A systematic review update. *Autism Research*, *15*(5), 778–790. https://doi.org/10.1002/aur.2696

Zilberstein, K. (2014). Neurocognitive considerations in the treatment of attachment and complex trauma in children. *Clinical Child Psychology and Psychiatry*, *19*(3), 336–354. https://doi.org/10.1177/1359104513486998