Immunometabolic Blood Biomarkers of Developmental Trajectories of Depressive

Symptoms: Findings from the ALSPAC Birth Cohort

Running title: Immunometabolic markers of depression trajectories

Authors:

Ruby S. M. Tsang[1,2,3]

Daniel Stow[4]

Alex S. F. Kwong[1,5]

Nicholas A. Donnelly[1,2,6]

Holly Fraser[1,2,3]

Inês Barroso[7]

Peter A. Holmans[8]

Michael J. Owen[8,9]

Megan L. Wood[10]

LINC Consortium*

Marianne B. M. van den Bree[8,9]

Nicholas J. Timpson[1,2]

Golam M. Khandaker[1,2,3,11]

Affiliations:

[1] MRC Integrative Epidemiology Unit, University of Bristol, Bristol, UK

[2] Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

[3] Centre for Academic Mental Health, Population Health Sciences, University of Bristol, Bristol,

UK

[4] Wolfson Institute of Population Health, Queen Mary University of London, London, UK

[5] Division of Psychiatry, University of Edinburgh, Edinburgh, UK

[6] Avon and Wiltshire NHS Mental Health Partnership NHS Trust, Bristol, UK

[7] Exeter Centre of Excellence for Diabetes Research, University of Exeter, UK

[8] Centre for Neuropsychiatric Genetics and Genomics, Division of Psychological Medicine and

Clinical Neurosciences, Cardiff University, Cardiff, UK

[9] Neuroscience and Mental Health Innovation Institute, Division of Psychological Medicine and

Clinical Neurosciences, Cardiff University, Cardiff, UK

[10] School of Psychology, University of Leeds, Leeds, UK

[11] NIHR Bristol Biomedical Research Centre, Bristol, UK

* A list of members and their affiliations appear at the end of the paper.

Corresponding author:

Dr Ruby Tsang

Population Health Sciences, Bristol Medical School

Oakfield House, Oakfield Grove

Bristol BS8 2BN

United Kingdom

ruby.tsang@bristol.ac.uk

Abstract

Depression is associated with immunological and metabolic alterations, but immunometabolic

characteristics of developmental trajectories of depressive symptoms remain unclear. Studies of

longitudinal trends of depressive symptoms in young people could provide insight into

aetiological mechanisms and heterogeneity behind depression, and origins of possible common

cardiometabolic comorbidities for depression. Using depressive symptoms scores measured on

10 occasions between ages 10 and 25 years in the Avon Longitudinal Study of Parents and

Children (n=7302), we identified four distinct trajectories: low-stable (70% of the sample),

adolescent-limited (13%), adulthood-onset (10%) and adolescent-persistent (7%). We examined

associations of these trajectories with: i) anthropometric, cardiometabolic and psychiatric

phenotypes using multivariable regression (n=1709-3410); ii) 67 blood immunological proteins

and 57 metabolomic features using empirical Bayes moderated linear models (n=2059 and

n=2240 respectively); and iii) 28 blood cell counts and biochemical measures using multivariable

regression (n=2256). Relative to the low-stable group, risk of depression and anxiety in

adulthood was higher for all other groups, especially in the adolescent-persistent

($RR_{depression}$=13.11, 95% CI 9.59-17.90; RR_{GAD}=11.77, 95% CI 8.58-16.14) and adulthood-onset

($RR_{depression}$=6.25, 95% CI 4.50-8.68; RR_{GAD}=4.66, 95% CI 3.29-6.60) groups. The three

depression-related trajectories vary in their immunometabolic profile, with evidence of little or no

alterations in the adolescent-limited group. The adulthood-onset group shows widespread

classical immunometabolic changes (e.g., increased immune cell counts and insulin resistance),

while the adolescent-persistent group is characterised by higher BMI both in childhood and

adulthood with few other immunometabolic changes. These findings point to distinct mechanisms

and prevention opportunities for adverse cardiometabolic profile in different groups of young

people with depression.

Introduction

The first two decades of life represent a critical epoch for human neurodevelopment when most serious mental illnesses of adult life first emerge.[1] Half of all lifetime cases of common mental disorders including depression and anxiety start by 14 years and 75% by 24 years.[2] The first onset of clinically recognised depressive episodes typically occurs between the ages of 12 and 15 years[3] and the increase in new onset of depression peaks between the ages of 15 and 18 years.[4] Depressive symptoms in childhood and adolescence, including those below diagnostic thresholds, are associated with an elevated risk of depression and other psychiatric diagnoses subsequently in adulthood.[2, 5-7] These findings highlight the need for studying depressive symptoms during early life.

Characterisation of longitudinal profiles of depressive symptoms during development could help understand the pathogenesis and heterogeneity of later depression, as different individuals may arrive at the same destination via different routes. There is growing evidence to suggest characteristic depression trajectories in childhood and adolescence are differentially associated with risk factors and outcomes. Existing studies have reported associations of a 'high' or 'increasing' depression trajectory with female sex, lower socioeconomic status, stressful life events, conduct issues, substance use, and parental psychopathology.[3, 8, 9] Trajectories with higher symptom burden have been associated with subsequent depression and other psychiatric diagnoses, lower educational attainment, income and poorer psychosocial adjustment.[8-10] However, less is known about underlying biological correlates of depression trajectories, including blood-based biomarker signatures. A better understanding of the biological correlates may help uncover mechanistic insights and identify accessible predictive markers for depression.

Existing literature suggests that depression and specific symptoms or symptom dimensions of depression are associated with immunometabolic dysfunction, but there is limited work on immunometabolic correlates of depression trajectories. Depression is associated with immunometabolic alterations such as chronic low-grade inflammation,[11, 12] neuroendocrine dysregulations,[13] as well as less favourable metabolic and lipid profiles.[12, 14] Overall effect sizes

for some of these associations are inconsistent, which could be partly due to clinical or

phenotypic heterogeneity within cross-sectional studies.[12] For instance, immunometabolic

alterations appear to be more pronounced or common in individuals endorsing atypical energy-

related symptoms of depression (e.g., hyperphagia, weight gain, hypersomnia, or leaden

paralysis) as opposed to melancholic symptoms.[12, 15] At the symptom level, inflammatory

markers are particularly associated with somatic and neurovegetative symptoms of depression

(e.g., fatigue, altered sleep and appetite) as opposed to psychological symptoms (e.g.,

hopelessness, excessive/inappropriate guilt).[16, 17] Some of these findings are supported by

Mendelian randomization analyses reporting a potentially causal link between inflammatory

markers (e.g., C-reactive protein (CRP) or interleukin 6 (IL-6)) and fatigue, anhedonia, sleep

problems, appetite and psychomotor changes.[18, 19]

The accumulation of risk model for chronic diseases posits that cumulative exposures across the

life course result in diverging health trajectories and widening health inequalities as people age.[20]

By characterising depression trajectories, developmental windows when trajectories begin to

diverge can be identified and we can then examine potential factors driving such divergence and

biological dysregulations linked to subsequent disease risk. By studying the biomarker signatures

of depression trajectories, we may also gain further insight into the origins of higher levels of

cardiometabolic multimorbidity in individuals with depression.[21, 22]

The aims of the current study were threefold: (i) to model depressive symptom trajectories from

childhood to early adulthood to classify individuals into more homogeneous subgroups, (ii) to

examine associations between these subgroups and risk of psychiatric and cardiometabolic

outcomes in early adulthood, and (iii) to examine associations of these subgroups with clinical

and blood immunometabolic markers including proteomic, metabolomic and biochemical

measures in early adulthood. By examining the broader biomarker signature across different

domains including the immune proteome, metabolome, and clinical biochemistry, we aim to

provide more comprehensive insights into biological pathways and systems possibly involved in

the development and persistence of depressive symptoms in young people.

Materials and Methods

Description of cohort

This study uses data from the Avon Longitudinal Study of Parents and Children (ALSPAC).

Pregnant women resident in the former county of Avon, United Kingdom (UK) with expected

dates of delivery between 1[st] April 1991 and 31[st] December 1992 were invited to take part in the

ALSPAC study. The initial recruitment enrolled 14541 pregnancies, which resulted in 14062 live

births and 13988 infants still alive at 12 months. Further recruitment of eligible participants took

place when the oldest children were approximately seven years of age; the total sample size for

analyses using any data collected after the age of seven is therefore 15447 pregnancies; of

these, 14901 children were alive at 12 months of age.[23, 24]

The study website contains details of all the data that is available through a fully searchable data

dictionary and variable search tool: http://www.bristol.ac.uk/alspac/researchers/our-data/

Data

Sociodemographic and health variables

Sociodemographic characteristics used to characterise the identified depressive symptom

trajectories include sex, ethnicity, maternal education, maternal occupational social class,

socioeconomic deprivation, and family adversity during pregnancy. Health characteristics

examined include smoking, risky alcohol use, carotid intima-media thickness, carotid-femoral

pulse wave velocity, metabolic syndrome and its components, obesity, and psychiatric outcomes

and medications. Detailed description of these variables as well as those included as covariables

in the biomarker analyses are presented in **Methods S1**.

Depressive symptoms

Self-reported depressive symptoms were assessed using the 13-item Short Mood and Feelings

Questionnaire (SMFQ).[25] We used data collected on 10 occasions between the ages of 10 and

25 years (ages 10, 12, 13, 16, 17, 18, 21, 22, 23, 25), ending with the last questionnaire

administered in 2017-2018, prior to the start of the COVID-19 pandemic (see **Table S1**).

Questions were answered based on the two-weeks prior to completing the questionnaire. Each

SMFQ item is scored as 0 = "not true", 1 = "sometimes true" and 2 = "always true", resulting in a

total SMFQ sum score 0-26 (higher score reflects more symptoms). For individuals who had

missing data on fewer than three questions, score was imputed to the median value for missing

items. For each time-point, those with missing data on more than three questions had their total

score recoded as missing.

Circulating blood biomarkers

For this analysis, blood biomarkers were assayed in blood samples collected at the face-to-face

research clinic undertaken at 24 years. A total of 92 circulating inflammatory proteins were

measured using the Olink Target 96 Inflammation panel (Olink Analysis Service, Uppsala,

Sweden); proteins with ≥50% values below the limit of detection (LOD) were excluded leaving 67

proteins to be included (**Table S2**). Over 220 metabolomic features (148 metabolites and 77

ratios) were quantified using a high-throughput ^{1}H-NMR spectroscopy-based platform

(Nightingale Health, Helsinki, Finland) using a standardised protocol and parameters described

elsewhere.[26-28] Lipoprotein subclasses were excluded from the analysis to minimise redundancy

of information, leaving a subset of 57 metabolomic features (9 cholesterol measures, 12

apolipoproteins and lipids measures, 3 lipoprotein particle sizes, 16 fatty acids and saturation

measures, 3 glycolysis-related metabolites, 8 amino acids, 3 ketone bodies, 2 fluid balance-

related measures and 1 inflammation-related measure) to be included in the analysis (**Table S3**).

All 26 blood count and chemistry measures collected at the same clinic interaction were included

(**Table S4**). Additionally, we computed the aspartate aminotransferase/alanine aminotransferase

(AST/ALT) ratio and the Homeostatic Model Assessment for Insulin Resistance (HOMA-IR).

Further information on data collection and processing of all blood biomarkers are presented in

Methods S1.

Covariables

Prior to statistical analysis, we plotted a directed acyclic graph (DAG) showing theoretical

relationships between depressive symptom trajectories (independent variable), immunometabolic

markers (dependent variables) and important covariables based on the literature (**Figure S1**).

The minimum adjustment set of confounders included in models was sex at birth, maternal

education, maternal occupational social class, and body mass index (BMI) at age 10.

Statistical analysis

Characterisation of depressive symptom trajectories

Latent class trajectory modelling was performed using the *lcmm* R package[29] to identify

subgroups with distinct SMFQ trajectories. This type of modelling seeks to identify homogenous

groups of individuals with similar trajectories within a heterogeneous population by combining a

latent class model and a mixed model. Models are estimated within the maximum likelihood

framework.[29] The *lcmm* package distinguishes time of measurement and occasion, so individuals

with missing data can still be included; we included those with at least three measurements for

better modelling of non-linear trajectories. A multi-step approach adapted from the model

selection framework suggested by Lennon et al.[30] and van der Nest et al.[31] was used, with the

order of steps changed to address potential overextraction of latent classes from model under-

specification as reported in the simulation literature[32, 33]. The steps followed:

1. Scope literature and inspect plots to inform polynomial order and potential number of classes.

We modelled smfq ~ age + age^2 and estimated models up to six latent classes.

2. Estimate growth mixture models (GMM) with random intercepts and class-specific

proportional random-effect variance-covariance matrix with increasing number of classes.

Select the most appropriate number of classes *k* based on model convergence, model fit

(Bayesian information criterion (BIC), Integrated Completed Likelihood (ICL), and relative

entropy), smallest class size ≥5% and visual inspection of the trajectories.

3. Test alternative model structures with *k* classes – GMM with random intercepts and common

random-effect variance-covariance matrix, and group-based trajectory models (GBTM);

compare model fit indices, smallest class size and visually inspect trajectories as above, and

assess model adequacy (average posterior probabilities ≥0.7 and odds of correct

classification ≥5 for all classes).

4. Refine trajectory shape by testing up to second-degree fractional polynomials including (-2, -

1, -0.5, 0, 0.5, 1, 2, 3) where 0 refers to log X and repeated polynomials refer to $(X^i + X^i * \log$

X). Select final model based on model convergence and model fit.

Age (in years) was used as the time variable in all models. No covariables were included in these

latent class mixed models as the aim is to describe the trajectories; covariables were accounted

for in the next step when testing for associations with phenotypes of interest and biomarkers. For

each model, an automatic grid search with 50 sets of random initial values and up to 10 iterations

was run to reduce the odds of the model converging towards a local maximum and then up to

500 iterations were allowed for the final estimation. Using the selected model, posterior

probabilities for class membership were then estimated and individuals were assigned to the

class of highest posterior probability in the entire sample using the *predictClass* function. We first

performed the latent class mixed modelling on the subsample with three or more data points and

then predicted class membership in the entire sample to reduce uncertainty in the modelling

stage and to maximise sample sizes in the subsequent analyses. Additional information on the

modelling is presented in **Supplementary Methods S1,** R scripts are provided in

Supplementary Methods S2. The reporting of this study adheres to the Guidelines for Reporting

on Latent Trajectory Studies (GRoLTS)[34] (**Table S5**).

Associations with clinical and sociodemographic variables

Sociodemographic characteristics are stratified by trajectory class and summarised using mean

(SD), median [interquartile range] or count (%) as appropriate, with differences between

trajectories tested with chi-square or Kruskal-Wallis tests. Associations of trajectory membership

with psychiatric or cardiometabolic outcomes of interest at age 24 or 28 years were tested using

multivariable linear or logistic regressions, using the largest trajectory class as the reference

group and adjusting for sex, maternal education, occupational social class and BMI at age 10.

These variables are described in detailed in **Supplementary Methods S1**.

Associations with immunometabolic biomarkers

For both proteomic and metabolomic data, associations between depressive symptom

trajectories and markers were evaluated using multiple linear models fitted in the *limma* R

package.[35] *Limma* uses an empirical Bayes method to moderate the standard errors of the

estimated log-fold changes by borrowing strength from linear models of the other analytes and

allowing for different variability between analytes and between samples. Planned contrasts of

each of the intermediary trajectories against the trajectory with the most individuals were

conducted. With the blood count and clinical chemistry data, linear regressions were fitted with

blood markers as dependent variables and SMFQ trajectory class as the independent variable.

For each of these markers, the basic model included sex, maternal education and maternal

occupational social class as covariables and the adjusted model further included BMI at age 10

as a covariable. Correction for multiple testing was performed for each set of models using the

Benjamini-Hochberg procedure, using a false discovery rate (FDR) q-value threshold of <0.1.

This threshold was chosen due to the large number of biomarkers tested, a relatively small

sample size and the exploratory nature of this work. R scripts for these analyses are provided in

Supplementary Methods S2.

Sensitivity analyses

To address potential error carried over from the probabilistic latent class assignment into the

association analyses, we performed two sets of sensitivity analyses, the first set by restricting the

sample to individuals who had a modal posterior probability ≥0.7, and the second set by using

the individuals' posterior probabilities for each latent class as separate terms in the models.

Data extraction and initial data cleaning was performed in StataMP version 17.[36] Further data

preparation and statistical analyses were conducted in R versions 4.1.1 and 4.2.1,[37] using

packages *tidyverse* (v2.0.0), *lcmm* (v2.0.2), *LCTMtools* (v0.1.3), *tableone* (version 0.13.2),

marginaleffects (v0.25.0), *knitr* (v1.43), *kableExtra* (v1.3.4), *limma* (v3.54.2), and *broom* (v1.0.5).

Plots were generated using *ggplot2* (v3.4.2), *ggpubr* (v0.6.0), and *ggrepel* (v0.9.3).

Ethical approval

Ethical approval for the ALSPAC study was obtained from the ALSPAC Ethics and Law

Committee and the Local Research Ethics Committees. Consent for biological samples has been

collected in accordance with the Human Tissue Act (2004). Informed consent for the use of data

collected via questionnaires and clinics was obtained from participants following the

recommendations of the ALSPAC Ethics and Law Committee at the time.

Results

Sample

Latent class trajectory modelling was performed on data from 7302 participants who had SMFQ

scores available from at least three time-points between ages 10 and 25 years. Once the best-

fitting model was identified, posterior probabilities and class membership were estimated in the

entire sample, and 9595 individuals were assigned class membership. Of these 9595 individuals,

2256 had sufficient biomarker and complete covariable data to be included in the biomarker

analyses (**Figure S2**).

Depressive symptom trajectories from childhood to early adulthood

Following comparison of model fit and adequacy statistics and visual inspection of trajectory plots

(**Table S6 and Figures S2-S3**), a four-class group-based trajectory model was identified as best

describing the data. As shown in **Figure 1**, the four identified depressive symptom trajectories

from childhood to early adulthood can be described as follows: low-stable – those who

consistently had no or low levels of depressive symptoms (69.6%, n=6680), adolescent-limited –

those who had elevated depressive symptoms in childhood/adolescence that decreased over

time (13.3%, n=1280), adolescent-persistent – those who had elevated depressive symptoms in

childhood/adolescence that remained high into adulthood (7.0%, n=672) and adulthood-onset –

those who started with low levels of depressive symptoms that increased in late

adolescence/early adulthood (10.0%, n=973).

Characteristics of depressive symptom trajectories

Descriptive statistics for characteristics of these individuals, stratified by trajectory, are presented

in **Table 1** below. There were more women in all three depression-related trajectories:

adolescent-limited (66.3%), adolescent-persistent (76.5%), and adulthood-onset trajectories

(64.6%). Additionally, the adolescent-persistent trajectory was associated with lower maternal

education and greater family adversity during pregnancy. Descriptive statistics for the same

characteristics of the subset of individuals who were included in the biomarker analyses are

presented in **Table S7**.

Figure 1. Predicted Marginal Mean Depressive Symptom Trajectories from Childhood to Early Adulthood

in the ALSPAC Cohort

Lines showing predicted marginal mean depressive symptom trajectories with shaded areas representing 95%

confidence intervals.

315 **Table 1. Characteristics of depressive symptom trajectories in the ALSPAC birth cohort.**

	Low-stable (n=6680)	Adolescent-limited (n=1280)	Adolescent-persistent (n=672)	Adulthood-onset (n=963)	*p*
Sex: Female	3064 (45.9%)	849 (66.3%)	514 (76.5%)	622 (64.6%)	<0.001
Ethnicity: Non-white	240 (4.1%)	44 (3.9%)	36 (6.2%)	40 (4.8%)	0.097
Maternal education					0.026
CSE or none	894 (15.1%)	159 (13.9%)	107 (17.9%)	120 (14.1%)	
Vocational	560 (9.5%)	91 (8.0%)	53 (8.9%)	64 (7.5%)	
O-level	2064 (34.9%)	406 (35.6%)	231 (38.6%)	293 (34.5%)	
A-level	1483 (25.1%)	308 (27.0%)	141 (23.6%)	230 (27.1%)	
Degree	911 (15.4%)	178 (15.6%)	66 (11.0%)	142 (16.7%)	
Maternal occupational social class					0.556
I – highest	356 (7.0%)	63 (6.6%)	28 (5.5%)	45 (6.3%)	
II	1714 (33.8%)	335 (35.0%)	161 (31.8%)	250 (34.9%)	
III (non-manual)	2137 (42.2%)	392 (41.0%)	216 (42.7%)	302 (42.1%)	
III (manual)	350 (6.9%)	66 (6.9%)	34 (6.7%)	57 (7.9%)	
IV or V[1] – lowest	509 (10.0%)	100 (10.5%)	67 (13.2%)	63 (8.8%)	
English IMD 2000 quintile					0.723
1 – least deprived	1665 (32.1%)	355 (35.3%)	146 (30.7%)	242 (33.3%)	
2	1044 (20.1%)	188 (18.7%)	100 (21.0%)	143 (19.7%)	
3	956 (18.4%)	174 (17.3%)	84 (17.6%)	140 (19.3%)	
4	790 (15.2%)	153 (15.2%)	77 (16.2%)	115 (15.8%)	
5 – most deprived	739 (14.2%)	137 (13.6%)	69 (14.5%)	86 (11.8%)	
Family Adversity Index	1 [0, 2]	1 [0, 2]	1 [0, 3]	1 [0, 2]	<0.001
BMI at age 10	17.43 [15.97, 19.73]	17.63 [16.07, 19.96]	18.25 [16.24, 21.00]	17.43 [15.92, 19.60]	<0.001
Number of SMFQ measurements	4 [2, 7]	6 [3, 8]	5 [3, 7]	6 [4, 9]	<0.001

[1] Categories have been collapsed due to small cell counts in subsequent analyses.

Notes: Numbers presented as mean (SD), median [IQR], or n (%). Percentages are column percentages and

computed based on the number of individuals with available data on each variable. Group comparisons were

conducted using Chi-square tests for categorical variables and Kruskal-Wallis tests for non-normal continuous

variables.

Abbreviations: BMI – body mass index; CSE – Certificate of Secondary Education; O-level – Ordinary level; A-

level – Advanced level; IMD – Index of Multiple Deprivation; SMFQ – Short Mood and Feelings Questionnaire

Table 2. Associations of depressive symptom trajectories with anthropometric and cardiometabolic

outcomes at 24 years and psychiatric outcomes at 24 and 28 years

		Adolescent-limited	Adolescent-persistent	Adulthood-onset
Outcomes assessed at age 24	n	Adjusted unstandardised regression coefficient (SE)		
cIMT (continuous)	1565	-0.0002 (0.0033)	-0.0099 (0.0050)	-0.0003 (0.0036)
cfPWV (continuous)	1708	-0.0578 (0.0708)	-0.1429 (0.1109)	0.0108 (0.0790)
		Adjusted relative risk (95% CI)		
Smoking	2731	**1.44 (1.23-1.69)**	**1.92 (1.59-2.31)**	**1.46 (1.23-1.73)**
AUDIT-C score ≥ 5	2704	0.99 (0.91-1.07)	**0.84 (0.72-0.97)**	0.91 (0.83-1.00)
BMI ≥ 30kg/m^2	2731	1.26 (1.00-1.59)	**1.53 (1.14-2.06)**	1.22 (0.94-1.58)
Elevated waist circumference	2707	1.11 (0.96-1.28)	1.12 (0.92-1.38)	1.08 (0.92-1.27)
Triglycerides ≥1.7mmol/L	2241	1.32 (0.89-1.96)	1.13 (0.60-2.13)	1.43 (0.96-2.12)
HDL <1.0mmol/L	2241	0.89 (0.67-1.17)	1.11 (0.77-1.60)	1.26 (0.97-1.63)
SBP ≥ 130mmHg	2751	0.88 (0.66-1.19)	0.66 (0.38-1.16)	0.97 (0.72-1.29)
DBP ≥ 85mmHg	2751	1.30 (0.64-2.63)	0.74 (0.22-2.47)	1.30 (0.61-2.78)
Fasting glucose ≥5.6mmol/L	2241	0.90 (0.73-1.11)	0.94 (0.69-1.29)	0.98 (0.80-1.21)
Metabolic syndrome	2759	0.98 (0.64-1.49)	0.80 (0.42-1.53)	1.20 (0.79-1.80)
ICD-10 depressive episode	2729	**3.65 (2.55-5.22)**	**13.11 (9.59-17.90)**	**6.25 (4.50-8.68)**
ICD-10 GAD	2719	**3.29 (2.29-4.73)**	**11.77 (8.58-16.14)**	**4.66 (3.29-6.60)**
Outcomes assessed at age 28				
Prescribed antidepressants in past 5y	2792	**2.54 (2.02-3.20)**	**4.61 (3.68-5.79)**	**3.71 (3.00-4.57)**
Prescribed anxiolytics in past 5y	2828	**2.25 (1.26-4.00)**	**6.04 (3.54-10.31)**	**3.90 (2.33-6.55)**

Notes: Regression models were adjusted for sex, maternal education, maternal occupational social class, and

BMI at age 10. Effect estimates presented are unstandardised regression coefficients for continuous outcomes

and relative risks (95% confidence intervals) for binary outcomes. The reference group for all analyses is the low-

stable trajectory. Text in bold indicates evidence for the association after FDR correction of p-values.

Metabolic syndrome was defined based on the 2009 consensus definition from the International Diabetes

Federation and the American Heart Association/National Heart, Lung, and Blood Institute, i.e. the presence of any

three of the following five risk factors: elevated waist circumference (≥94cm for white men, ≥90cm for non-white

men, ≥80cm for women); elevated triglycerides (≥1.7mmol/L), reduced high-density lipoprotein-cholesterol

(<1.0mmol/L), elevated blood pressure (systolic ≥130mmHg or diastolic ≥85mmHg), and elevated fasting glucose

(≥5.6mmol/L).

Abbreviations: AUDIT-C – Alcohol Use Disorders Identification Test for Consumption; BMI – body mass index;

cfPWV – carotid-femoral pulse wave velocity; cIMT – carotid intima-media thickness; DBP – diastolic blood

pressure; GAD – generalised anxiety disorder; HDL – high-density lipoprotein; ICD-10 – International

Classification of Diseases 10[th] Revision; SBP – systolic blood pressure

Adulthood cardiometabolic and psychiatric outcomes associated with depressive symptom

trajectories

Compared to the low-stable trajectory, after adjusting for sex, maternal education, maternal

occupational class and BMI at age 10, all three depression-related trajectories were associated

with ICD-10 diagnosis of depression, generalised anxiety disorder, being prescribed an

antidepressant or anxiolytic at ages 24 and 28 years. However, the magnitude of association

varied between the trajectories, with the risk for these outcomes being the highest for the

adolescent-persistent trajectory (approximately 13-fold risk), followed by the adulthood-onset and

adolescent-limited trajectories. The adolescent-persistent trajectory was additionally associated

with obesity (**Table 2**). Unadjusted model results are presented in **Table S8**.

Differentially abundant proteins associated with depressive symptom trajectories

Relative to the low-stable trajectory, after adjusting for sex, maternal education, maternal

occupational class and BMI at age 10, one protein (C-C motif chemokine 25 [CCL25]) was

upregulated in the adolescent-limited trajectory; four proteins (fibroblast growth factor 21 [FGF-

21], hepatocyte growth factor [HGF], eukaryotic translation initiation factor [4E-BP1], and eotaxin-

1 [CCL11]) were upregulated in the adolescent-persistent trajectory; and five proteins (FGF-21,

fibroblast growth factor 19 [FGF-19], CUB domain-containing protein 1 [CDCP1], HGF, CCL11)

were upregulated in the adulthood-onset trajectory (**Figure 2**). Full model results are presented

in **Tables S9-10**.

Differentially abundant metabolites associated with depressive symptom trajectories

Relative to the low-stable trajectory, after adjusting for sex, maternal education and occupational

class, and BMI at age 10, creatinine was decreased in the adolescent-persistent trajectory. Three

metabolite ratios (omega-3 to total fatty acids [omega-3/FA], docosahexaenoic acid to total fatty

acids [DHA/FA], polyunsaturated fatty acids to total fatty acids [PUFA/FA] ratios) were decreased

whereas the monosaturated to total fatty acids (MUFA/FA) and apolipoprotein B to apolipoprotein

A1 (ApoB/ApoA1) ratios were increased in the adult-onset trajectory (**Figure 3**). Full model

results are presented in **Tables S11-12**.

Blood count and clinical chemistry markers

374   Relative to the low-stable trajectory, after adjusting for sex, maternal education and occupational

375   class, and BMI at age 10, there was evidence for decreased AST levels in the adolescent-limited

376   trajectory, and increased HOMA-IR, insulin, neutrophil and white blood cell (WBC) counts in the

377   adulthood-onset trajectory (**Figure 5**). Full model results are presented in **Tables S13-14**.

Sensitivity analyses

380   The sensitivity analyses showed patterns of associations that are largely similar to those

381   observed in the primary analyses, with consistent associations of depression-related trajectories

382   with anthropometric, cardiometabolic and psychiatric outcomes, and top-ranking

383   immunometabolic biomarkers with similar effect sizes in the same directions (**Supplementary**

384   **Methods S3**).

Figure 2. Volcano Plots Showing Differential Immune Protein Abundance Levels in Depressive Symptom Trajectories

Panels **a** and **b** show results from basic and adjusted models respectively. The reference group for all analyses is the low-stable trajectory.

Abbreviations: 4E-BP1 – eukaryotic translation initiation factor; CCL11 – eotaxin-1; CCL25 – C-C motif chemokine 25; CDCP1 – CUB domain-containing protein 1; FGF-19 – fibroblast growth factor 19; FGF-21 – fibroblast growth factor 21; HGF – hepatic growth factor

Figure 3. Volcano Plots Showing Differential Metabolite Abundance Levels in Depressive Symptom Trajectories

Panels **a** and **b** show results from basic and adjusted models respectively. The reference group for all analyses is the low-stable trajectory. Orange points indicate upregulation and blue points indicate downregulation.

Abbreviations: ApoB/ApoA1 – apolipoprotein B to apolipoprotein A1 ratio, DHA/FA – docosahexaenoic acid to total fatty acids ratio, MUFA/FA – monounsaturated fatty acids to total fatty acids ratio, Omega-3/FA – omega-3 fatty acids to total fatty acids ratio, PUFA/FA – polyunsaturated fatty acids/total fatty acids ratio

Figure 4. Differential Levels of Full Blood Count and Clinical Biochemistry Biomarkers in Different Depressive Symptom Trajectories

Dot-and-whisker plots showing effect estimates and 95% confidence intervals for each biomarker. The effect estimates and 95% confidence intervals from the models have

been back-transformed into their original scale for ease of interpretation; effect estimates represent the percentage difference in mean levels of each biomarker for respective

trajectory, in relation to the low-stable trajectory (reference group). Panels a and b show results from basic and adjusted models respectively. Asterisks indicate evidence for the

association after FDR correction of p-values.

Abbreviations: WBC – white blood count, RBC – red blood count, Hb – haemoglobin, Hct – haematocrit, MCV – mean cell volume, MCH – mean cell haemoglobin, MCHC –

mean corpuscular haemoglobin concentration, HOMA-IR – Homeostatic Model Assessment for Insulin Resistance, HDL – high-density lipoprotein, LDL – low-density

lipoprotein, VLDL – very low-density lipoprotein, CRP – C-reactive protein, GGT – gamma-glutamyl transpeptide, ALT – alanine aminotransferase, AST – aspartate

aminotransferase, P3NP – procollagen-3 N-terminal peptide

Discussion

Depression is a complex heterogeneous disorder, which poses a challenge for discovering

biomarkers associated with disease onset and/or progression. We have taken a longitudinal

approach to identifying blood-based biomarkers for depression by examining longitudinal

patterns of depressive symptoms in the population during the critical developmental epoch of

childhood, adolescence and early adulthood. Using data from a prospective birth cohort, we

identified four longitudinal population subgroups based on repeated measures of depressive

symptoms over a 15-year period from ages 10 to 25 years. We show that majority of participants

(approximately 70%) have little or no depressive symptoms (low-stable group). We identified

three depression-related groups which comprise a group with higher symptom levels during

childhood and adolescence which later decrease (adolescent-limited group, 13%), a group with

symptoms emerging during puberty that persist throughout adolescence through to adulthood

(adolescent-persistent group, 7%), and a group with symptoms emerging during late

adolescence/early adulthood and increasing thereafter (adulthood-onset, 10%). Earlier work

examining latent depressive symptom trajectories in ALSPAC have reported varying numbers of

latent trajectories, but all studies consistently identified a group with low/no symptoms, a group

with symptoms emerging in early adolescence that persist, and a group with symptoms emerging

in late adolescence/early adulthood.[10, 38, 39] Differences in the number of trajectories identified

may be due to sample inclusion/exclusion criteria, the number of repeated measures included,

polynomials used to model the trajectories, as well as model selection criteria.

We examined health phenotypes and blood biomarkers associated with these subgroups for

greater insight into the developmental course of depression and associated biomarkers. Our

analyses show that compared to the low-stable group, risk of depression and anxiety in

adulthood is higher for all three depression-related groups. However, such risk is particularly

elevated risk for the adolescent-persistent group (11 to 13-fold risk) followed by the adulthood-

onset group (four to six-fold risk). Interestingly, the group where higher levels of symptoms are

mostly limited to adolescence, they still have a three-fold risk of depression in adulthood. This

suggests that while their depressive symptoms do not persist at the same level as observed in

adolescence, their symptoms do not fully return to the pre-morbid level either, and therefore are

still more likely to meet criteria for a clinical diagnosis than those in the low-stable group. The

literature also suggests that partial remission of depression is common and is a robust predictor

for relapse.[40]

Having examined health phenotypic and blood proteomic, immunological and metabolic

biomarker associations for these three groups, we show that the adolescent-limited group is

distinct from the other two depression-related groups as it showed little immunometabolic

alterations. In contrast, both the adolescent-persistent and adulthood-onset groups are

associated with immunometabolic changes, but the exact pattern of associations varies between

the two groups. The adolescent-persistent group was associated with higher BMI in childhood

and adulthood, whereas the adulthood-onset group did not show this, but rather more

widespread alterations in blood-based metabolic parameters including insulin resistance, insulin

levels and changes in fatty acid ratios. Blood proteomic changes were largely similar between

the two groups and involved proteins that mainly act as growth factors, cytokines and

chemokines. While immunometabolic associations persisted given adjustment for childhood BMI

in the adolescent-persistent group, the presence or absence of this BMI adjustment had an

impact on some association estimates in the adolescent-persistent and adulthood-onset groups.

While this may relate to the apparent differences in BMI by trajectories, it is difficult to distinguish

this as an artefact of adjustment or a true impact of BMI.

Epidemiological studies consistently report a bidirectional relationship (both cross-sectionally and

longitudinally) between obesity and depression,[41] whereas Mendelian randomization studies

support a causal role of BMI on major depressive disorder and depressive symptoms but not vice

versa.[42, 43] The comorbidity between obesity and depression is generally associated with poorer

prognosis, with studies reporting associations with a more chronic course of depression in

adulthood[44, 45] as well as poorer treatment response.[46] Our findings add to this evidence by

showing an association between higher childhood BMI and persistent depressive symptoms

between adolescence and early adulthood.

Many of the alterations observed in the adulthood-onset trajectory are already well studied

markers of cardiometabolic disease risk. The ApoB/ApoA1 ratio is associated with cardiovascular

diseases and metabolic syndrome,[47, 48] and can be used to predict longer-term cardiovascular

risk when measured in early life.[49, 50] Higher values of HOMA-IR are associated with an

increased risk of developing type 2 diabetes mellitus (T2DM), systemic arterial hypertension and

non-fatal major adverse cardiovascular events.[51] Lower blood omega-3 fatty acid concentrations

are associated with poorer cardiovascular outcomes[52, 53] and may also contribute to chronic

systemic inflammation, whereas changes in MUFA, PUFA and DHA concentrations in early

adulthood were associated with incident obesity, insulin resistance and elevated blood pressure

10 years later.[54] Since dietary intake and supplementation are the main predictors of blood levels

of omega-3 fatty acids,[55, 56] with other lifestyle-related factors such as BMI, smoking and alcohol

consumption also playing a role,[57] this suggests that the adulthood-onset group may benefit from

primary prevention strategies such as lifestyle modifications to prevent future cardiometabolic

disease. Given that it is already well established that depression predisposes young people to

accelerated atherosclerosis and early cardiovascular disease,[58] further studies investigating the

use of longitudinal course of depressive symptoms for risk stratification and prevention of

cardiometabolic disease are warranted.

The overlapping proteomic signals observed between the adolescent-persistent and adulthood-

onset trajectories potentially suggest shared underlying mechanisms (genetic or environmental)

or shared biological responses to depression, which warrant further study for their roles in the

pathophysiology of depression and cardiometabolic disease. FGF-21 is a novel regulator of

glucose and lipid metabolism that mainly acts through an FGF receptor 1 (FGFR1)/β-klotho

receptor complex and the Ras/Raf MAPK signalling pathway, which have been implicated in the

pathophysiology of depression and therapeutic effects of antidepressants.[59-61] Elevated

circulating FGF-21 concentrations have also been shown to be associated with a range of

cardiometabolic markers and diseases.[62-66] HGF mediates inflammatory responses to tissue

injury and regulates cell growth and morphogenesis through the activation of the

HGF/mesenchymal-epithelial transition factor (c-Met) signalling pathway, which has downstream

effects on the Raf/MAPK and PI3K/Akt pathways. Altered HGF/c-Met signalling has been

suggested to play a role in the pathogenesis of depression in adolescents through disrupting

interneuron development.[67] CCL11 is a chemokine involved in the selective recruitment of

eosinophils into sites of inflammation and has been implicated in various allergic and

inflammatory conditions.[68] It can be transported across the blood-brain barrier[69] and is an age-

related systemic factor associated with reduced synaptic plasticity and impaired hippocampal-

dependent learning and memory in mice.[70] In humans, CCL11 levels increase with age,[70] and

there is emerging evidence to suggest that CCL11 levels are associated with psychiatric

disorders.[71-75] In summary, the overlapping proteomic signals between the adolescent-persistent

and adulthood-onset trajectories highlight potential roles of physiological stress from lifestyle or

environmental factors, disruptions in neurodevelopment and neurogenesis, and cellular

senescence in the underlying vulnerability or biological response to depression, and may be key

biomarkers relevant to illness pathogenesis.

An advantage of our work is that by examining depressive symptoms longitudinally using a latent

class trajectory approach, we can account for population heterogeneity and obtain better

characterisation of subgroups and their changes over time. Existing literature shows that

depression is associated with alterations in various immunometabolic biomarkers, including

increased inflammatory cytokines,[76, 77] WBC,[78] neutrophils,[78] T-lymphocytes and other immune

cell counts,[78] HOMA-IR,[79] insulin,[79] lipids and fatty acids.[80, 81] Using longitudinal data from young

people, we add to this evidence base by showing that classical immunometabolic changes are

particularly associated with an adulthood-onset trajectory, rather than other developmental

subgroups of depressive symptoms, including one with persistent symptoms since adolescence.

While developmental stages like childhood, adolescence and adulthood are in part socially

constructed, our results show that the timings of divergence in depressive symptom trajectories

are approximately aligned with the transitions between developmental stages when some of the

major biological and environmental changes also take place (e.g., puberty, starting secondary

school, and finishing school).

This study has several limitations. Firstly, the approach of treating assigned class membership as

discrete in assessing relationships with other variables has been shown to underestimate the

strength of the relationships.[82] However, as we are mainly interested in subpopulations with

different depressive symptom trajectories, this approach allows for easier interpretation and

translation. Secondly, depression is episodic in nature and the use of polynomials cannot fully

capture the dynamics of depressive symptom severity over time; however, this approach was

chosen over other methods (e.g., splines) for model parsimony.

Furthermore, the sample size in this study is relatively small and may be underpowered to detect

differences after stratifying individuals into four separate trajectories. For this reason, we did not

further stratify our analyses by sex or other potentially relevant variables (e.g., BMI). While we

have adjusted our biomarker analyses for several potential confounders, residual confounding

could still be an issue. For instance, we have not accounted for medication use or chronic

disease, but these are likely to be uncommon in young people. As the biomarkers were

measured at age 24, which is after the onset of depressive symptoms in many individuals, further

research is required to assess the direction and causality of associations we have identified.

In conclusion, we identified distinct developmental trajectories of depression from childhood to

early adulthood, which show differential associations with cardiometabolic and psychiatric

outcomes, and are characterised by distinct immunometabolic profiles. In particular, individuals

with persistent depressive symptoms from childhood through to early adulthood were more likely

to have higher BMI both in childhood and in early adulthood and few other immunometabolic

changes, whereas individuals who develop depressive symptoms towards early adulthood show

classical immunometabolic alterations in immune cell counts, insulin resistance and fatty acid

profiles. These findings point to distinct mechanisms and prevention opportunities for adverse

cardiometabolic profile in different groups of young people with depressive symptoms.

Acknowledgements

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. We would also like to thank members of the LINC study public advisory group for their contribution.

This work was funded by the Tackling Multimorbidity at Scale Strategic Priorities Fund programme (MR/W014416/1) delivered by the UK Medical Research Council (MRC) and the UK National Institute for Health Research (NIHR) in partnership with the UK Economic and Social Research Council and in collaboration with the UK Engineering and Physical Sciences Research Council. RSMT, DS and MLW are supported by this grant. ASFK is supported by a Wellcome Early Career Award (227063/Z/23/Z). NAD was supported by an NIHR Clinical Lectureship in General Adult Psychiatry. MBMvdB acknowledges additional funding support from the MRC (MR/W028395/1, MR/W020297/1, MR/T033045/1 and MR/S037667/1), NIMH (U01MH119758), the Wellcome Trust (226709/Z/22/Z) and Welsh Government (HS 22 04). NJT is a director of the MRC/ESRC/UKRI supported Population Research UK Coordination Hub (ES/Y008340/1), supported by a Wellcome Trust Investigator award (202802/Z/16/Z), is the PI of the Avon Longitudinal Study of Parents and Children (MRC & WT 217065/Z/19/Z), is supported by the University of Bristol NIHR Biomedical Research Centre (BRC-1215-2001), the MRC Integrative Epidemiology Unit (MC_UU_00011/1), CRUK and works within the CRUK Integrative Cancer Epidemiology Programme (C18281/A29019) and with support from CRUK (PRCPJT-May22\100028). GMK acknowledges funding support from the MRC (MC_UU_00032/6), which forms part of the MRC Integrative Epidemiology Unit at the University of Bristol. This grant also supports HF. GMK acknowledges additional funding from the Wellcome Trust (201486/Z/16/Z and 201486/B/16/Z), the MRC (MR/W014416/1; MR/S037675/1; and MR/Z50354X/1), and the NIHR Bristol Biomedical Research Centre (NIHR 203315). The views expressed are those of the authors and not necessarily those of the UK NIHR or the Department of Health and Social Care.

The UK MRC and Wellcome (217065/Z/19/Z) and the University of Bristol provide core support

for ALSPAC. This publication is the work of the authors and RSMT, NJT and GMK will serve as

guarantors for the contents of this paper. A comprehensive list of grants funding is available on

the ALSPAC website (http://www.bristol.ac.uk/alspac/external/documents/grant-

acknowledgements.pdf).

LINC Consortium members:

Marianne B. M. van den Bree, George Kirov, Michael J. Owen, James T. R. Walters, Peter A.

Holmans, Jane Lynch, Ioanna K. Katzourou, Nabila Ali, Lowri O'Donovan (Cardiff University, UK)

David A. van Heel, Sarah Finer, Daniel Stow (Queen Mary University of London, UK)

Golam M. Khandaker, Nicholas J. Timpson, John A. A. Macleod, Julie P. Clayton, Ruby S. M.

Tsang, Jane Sprackman, Shahid Khan (University of Bristol, UK)

Inês Barroso, Rupert A. Payne (University of Exeter, UK)

Mark Mon-Williams, Megan L. Wood (University of Leeds, UK)

Hilary C. Martin (Wellcome Sanger Institute, UK)

Thomas Werge, Andrés Ingason, Morteza Vaez, Lam O. Huang (Institute of Biological Psychiatry,

Denmark)

Conflict of Interest

None

References

1. Solmi M, Radua J, Olivola M, Croce E, Soardo L, Salazar de Pablo G *et al.* Age at onset of

mental disorders worldwide: large-scale meta-analysis of 192 epidemiological studies.

Molecular Psychiatry 2022; **27**(1)**:** 281-295.

2. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime

Prevalence and Age-of-Onset Distributions of DSM-IV Disorders in the National

Comorbidity Survey Replication. *Archives of General Psychiatry* 2005; **62**(6)**:** 593-602.

3. Shore L, Toumbourou JW, Lewis AJ, Kremer P. Review: Longitudinal trajectories of child

and adolescent depressive symptoms and their predictors – a systematic review and meta-

analysis. *Child and Adolescent Mental Health* 2018; **23**(2)**:** 107-120.

4. Hankin BL, Abramson LY, Moffitt TE, Silva PA, McGee R, Angell KE. Development of

depression from preadolescence to young adulthood: emerging gender differences in a 10-

year longitudinal study. *J Abnorm Psychol* 1998; **107**(1)**:** 128-140.

5. Copeland WE, Adair CE, Smetanin P, Stiff D, Briante C, Colman I *et al.* Diagnostic

transitions from childhood to adolescence to early adulthood. *Journal of Child Psychology*

and Psychiatry 2013; **54**(7)**:** 791-799.

6. Fergusson DM, Woodward LJ. Mental Health, Educational, and Social Role Outcomes of

Adolescents With Depression. *Archives of General Psychiatry* 2002; **59**(3)**:** 225-231.

7. Georgiades K, Lewinsohn PM, Monroe SM, Seeley JR. Major Depressive Disorder in

Adolescence: The Role of Subthreshold Symptoms. *Journal of the American Academy of*

Child & Adolescent Psychiatry 2006; **45**(8)**:** 936-944.

8. Yaroslavsky I, Pettit JW, Lewinsohn PM, Seeley JR, Roberts RE. Heterogeneous trajectories of depressive symptoms: Adolescent predictors and adult outcomes. *Journal of Affective Disorders* 2013; **148**(2)**:** 391-399.

9. Musliner KL, Munk-Olsen T, Eaton WW, Zandi PP. Heterogeneity in long-term trajectories of depressive symptoms: Patterns, predictors and outcomes. *Journal of Affective Disorders* 2016; **192:** 199-211.

10. Weavers B, Heron J, Thapar AK, Stephens A, Lennon J, Bevan Jones R *et al.* The antecedents and outcomes of persistent and remitting adolescent depressive symptom trajectories: a longitudinal, population-based English study. *The Lancet Psychiatry* 2021; **8**(12)**:** 1053-1061.

11. Khandaker GM, Dantzer R, Jones PB. Immunopsychiatry: important facts. *Psychological Medicine* 2017; **47**(13)**:** 2229-2237.

12. Milaneschi Y, Lamers F, Berk M, Penninx BWJH. Depression Heterogeneity and Its Biological Underpinnings: Toward Immunometabolic Depression. *Biological Psychiatry* 2020; **88**(5)**:** 369-380.

13. Nikkheslat N, Pariante CM, Zunszain PA. Chapter 3 - Neuroendocrine Abnormalities in Major Depression: An Insight Into Glucocorticoids, Cytokines, and the Kynurenine Pathway. In: Baune BT (ed). *Inflammation and Immunity in Depression.* Academic Press2018, pp 45-60.

655   14.   Bot M, Milaneschi Y, Al-Shehri T, Amin N, Garmaeva S, Onderwater GLJ *et al.*

656   Metabolomics Profile in Depression: A Pooled Analysis of 230 Metabolic Markers in 5283

657   Cases With Depression and 10,145 Controls. *Biological Psychiatry* 2020; **87**(5)**:** 409-418.

659   15.   Lamers F, Milaneschi Y, Vinkers CH, Schoevers RA, Giltay EJ, Penninx BWJH. Depression

660   profilers and immuno-metabolic dysregulation: Longitudinal results from the NESDA study.

661   *Brain, Behavior, and Immunity* 2020; **88:** 174-183.

663   16.   Milaneschi Y, Kappelmann N, Ye Z, Lamers F, Moser S, Jones PB *et al.* Association of

664   inflammation with depression and anxiety: evidence for symptom-specificity and potential

665   causality from UK Biobank and NESDA cohorts. *Molecular Psychiatry* 2021; **26**(12)**:** 7393-

666   7402.

668   17.   Frank P, Jokela M, Batty GD, Cadar D, Steptoe A, Kivimäki M. Association Between

669   Systemic Inflammation and Individual Symptoms of Depression: A Pooled Analysis of 15

670   Population-Based Cohort Studies. *American Journal of Psychiatry* 2021; **178**(12)**:** 1107-

671   1118.

673   18.   Kappelmann N, Arloth J, Georgakis MK, Czamara D, Rost N, Ligthart S *et al.* Dissecting

674   the Association Between Inflammation, Metabolic Dysregulation, and Specific Depressive

675   Symptoms: A Genetic Correlation and 2-Sample Mendelian Randomization Study. *JAMA*

676   *Psychiatry* 2021; **78**(2)**:** 161-170.

678   19.   Pistis G, Milaneschi Y, Vandeleur CL, Lasserre AM, Penninx BWJH, Lamers F *et al.*

679   Obesity and atypical depression symptoms: findings from Mendelian randomization in two

680   European cohorts. *Translational Psychiatry* 2021; **11**(1)**:** 96.

682   20.   Ben-Shlomo Y, Kuh D. A life course approach to chronic disease epidemiology: conceptual

683   models, empirical challenges and interdisciplinary perspectives. *International Journal of*

684   *Epidemiology* 2002; **31**(2)**:** 285-293.

686   21.   Ditmars HL, Logue MW, Toomey R, McKenzie RE, Franz CE, Panizzon MS *et al.*

687   Associations between depression and cardiometabolic health: A 27-year longitudinal study.

688   *Psychological Medicine* 2022; **52**(14)**:** 3007-3017.

690   22.   Harshfield EL, Pennells L, Schwartz JE, Willeit P, Kaptoge S, Bell S *et al.* Association

691   Between Depressive Symptoms and Incident Cardiovascular Diseases. *JAMA* 2020;

692   **324**(23)**:** 2396-2405.

694   23.   Boyd A, Golding J, Macleod J, Lawlor DA, Fraser A, Henderson J *et al.* Cohort Profile: The

695   'Children of the 90s'—the index offspring of the Avon Longitudinal Study of Parents and

696   Children. *International Journal of Epidemiology* 2013; **42**(1)**:** 111-127.

698   24.   Northstone K, Lewcock M, Groom A, Boyd A, Macleod J, Timpson N *et al.* The Avon

699   Longitudinal Study of Parents and Children (ALSPAC): an update on the enrolled sample

700   of index children in 2019 [version 1; peer review: 2 approved]. *Wellcome Open Research*

701   2019; **4**(51).

703   25.   Angold A, Costello EJ, Messer SC, Pickles A. Development of a short questionnaire for use

704   in epidemiological studies of depression in children and adolescents. *International Journal*

705   *of Methods in Psychiatric Research* 1995; **5**(4)**:** 237-249.

707   26.   Soininen P, Kangas AJ, Würtz P, Tukiainen T, Tynkkynen T, Laatikainen R *et al.* High-

708   throughput serum NMR metabonomics for cost-effective holistic studies on systemic

709   metabolism. *Analyst* 2009; **134**(9)**:** 1781-1785.

711   27.   Inouye M, Kettunen J, Soininen P, Silander K, Ripatti S, Kumpula LS *et al.* Metabonomic,

712   transcriptomic, and genomic variation of a population cohort. *Molecular Systems Biology*

713   2010; **6**(1)**:** 441.

715   28.   Soininen P, Kangas AJ, Würtz P, Suna T, Ala-Korpela M. Quantitative Serum Nuclear

716   Magnetic Resonance Metabolomics in Cardiovascular Epidemiology and Genetics.

717   *Circulation: Cardiovascular Genetics* 2015; **8**(1)**:** 192-206.

719   29.   Proust-Lima C, Philipps V, Liquet B. Estimation of Extended Mixed Models Using Latent

720   Classes and Latent Processes: The R Package lcmm. *Journal of Statistical Software* 2017;

721   **78**(2)**:** 1 - 56.

723   30.   Lennon H, Kelly S, Sperrin M, Buchan I, Cross AJ, Leitzmann M *et al.* Framework to

724   construct and interpret latent class trajectory modelling. *BMJ Open* 2018; **8**(7)**:** e020683.

726   31.   van der Nest G, Lima Passos V, Candel MJJM, van Breukelen GJP. An overview of mixture

727   modelling for latent evolutions in longitudinal data: Modelling approaches, fit statistics and

728   software. *Advances in Life Course Research* 2020; **43:** 100323.

730   32.   van der Nest G, Lima Passos V, Candel MJJM, van Breukelen GJP. Model fit criteria curve

731   behaviour in class enumeration – a diagnostic tool for model (mis)specification in

732   longitudinal mixture modelling. *Journal of Statistical Computation and Simulation* 2022;

733   **92**(8)**:** 1640-1672.

735 33. Diallo TMO, Morin AJS, Lu H. Impact of Misspecifications of the Latent Variance–

736   Covariance and Residual Matrices on the Class Enumeration Accuracy of Growth Mixture

737   Models. *Structural Equation Modeling: A Multidisciplinary Journal* 2016; **23**(4)**:** 507-531.

739 34. van de Schoot R, Sijbrandij M, Winter SD, Depaoli S, Vermunt JK. The GRoLTS-Checklist:

740   Guidelines for Reporting on Latent Trajectory Studies. *Structural Equation Modeling: A*

741   *Multidisciplinary Journal* 2017; **24**(3)**:** 451-467.

743 35. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W *et al.* limma powers differential

744   expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research*

745   2015; **43**(7)**:** e47-e47.

747 36. StataCorp. Stata Statistical Software: Release 17. StataCorp LLC: College Station, TX,

748   USA, 2021.

750 37. R Core Team. R: A language and environment for statistical computing. R Foundation for

751   Statistical Computing: Vienna, Austria, 2022. https://www.R-project.org/.

753 38. Khandaker GM, Stochl J, Zammit S, Goodyer I, Lewis G, Jones PB. Childhood

754   inflammatory markers and intelligence as predictors of subsequent persistent depressive

755   symptoms: a longitudinal cohort study. *Psychological Medicine* 2018; **48**(9)**:** 1514-1522.

757 39. López-López JA, Kwong ASF, Washbrook E, Pearson RM, Tilling K, Fazel MS *et al.*

758   Trajectories of depressive symptoms and adult educational and employment outcomes.

759   *BJPsych Open* 2020; **6**(1)**:** e6.

40.   Gülpen J, Breedvelt JJF, van Dis EAM, Geurtsen GJ, Warren FC, van Heeringen C *et al.*

Psychological interventions for preventing relapse in individuals with partial remission of

depression: a systematic review and individual participant data meta-analysis.

Psychological Medicine 2025; **55:** e50.

41.   Milaneschi Y, Simmons WK, van Rossum EFC, Penninx BWJH. Depression and obesity:

evidence of shared biological mechanisms. *Molecular Psychiatry* 2019; **24**(1)**:** 18-33.

42.   Hartwig FP, Bowden J, Loret de Mola C, Tovo-Rodrigues L, Davey Smith G, Horta BL.

Body mass index and psychiatric disorders: a Mendelian randomization study. *Scientific*

Reports 2016; **6**(1)**:** 32730.

43.   van den Broek N, Treur JL, Larsen JK, Verhagen M, Verweij KJH, Vink JM. Causal

associations between body mass index and mental health: a Mendelian randomisation

study. *Journal of Epidemiology and Community Health* 2018; **72**(8)**:** 708.

44.   Vogelzangs N, Beekman AT, Boelhouwer IG, Bandinelli S, Milaneschi Y, Ferrucci L *et al.*

Metabolic depression: a chronic depressive subtype? Findings from the InCHIANTI study

of older persons. *J Clin Psychiatry* 2011; **72**(5)**:** 598-604.

45.   Nigatu YT, Bültmann U, Reijneveld SA. The prospective association between obesity and

major depression in the general population: does single or recurrent episode matter? *BMC*

Public Health 2015; **15**(1)**:** 350.

46.   Grigolon RB, Trevizol AP, Gerchman F, Bambokian AD, Magee T, McIntyre RS *et al.* Is

Obesity A Determinant Of Success With Pharmacological Treatment For Depression? A

Systematic Review, Meta-Analysis And Meta-Regression. *Journal of Affective Disorders*

2021; **287:** 54-68.

47.   McQueen MJ, Hawken S, Wang X, Ounpuu S, Sniderman A, Probstfield J *et al.* Lipids,

lipoproteins, and apolipoproteins as risk markers of myocardial infarction in 52 countries

(the INTERHEART study): a case-control study. *The Lancet* 2008; **372**(9634)**:** 224-233.

48.   Ulloque-Badaracco JR, Al-kassab-Córdova A, Hernandez-Bustamante EA, Alarcon-Braga

EA, Huayta-Cortez M, Carballo-Tello XL *et al.* Association of apolipoproteins and

lipoprotein(a) with metabolic syndrome: a systematic review and meta-analysis. *Lipids in*

Health and Disease 2023; **22**(1)**:** 98.

49.   Juonala M, Viikari JSA, Kähönen M, Solakivi T, Helenius H, Jula A *et al.* Childhood Levels

of Serum Apolipoproteins B and A-I Predict Carotid Intima-Media Thickness and Brachial

Endothelial Function in Adulthood: The Cardiovascular Risk in Young Finns Study. *Journal*

of the American College of Cardiology 2008; **52**(4)**:** 293-299.

50.   Ojanen X, Cheng R, Törmäkangas T, Rappaport N, Wilmanski T, Wu N *et al.* Towards early

risk biomarkers: serum metabolic signature in childhood predicts cardio-metabolic risk in

adulthood. *eBioMedicine* 2021; **72**.

51.   González-González JG, Violante-Cumpa JR, Zambrano-Lucio M, Burciaga-Jimenez E,

Castillo-Morales PL, Garcia-Campa M *et al.* HOMA-IR as a predictor of Health Outcomes

in Patients with Metabolic Risk Factors: A Systematic Review and Meta-analysis. *High*

Blood Pressure & Cardiovascular Prevention 2022; **29**(6)**:** 547-564.

52.   Harris WS, von Schacky C. The Omega-3 Index: a new risk factor for death from coronary

814   heart disease? *Preventive Medicine* 2004; **39**(1)**:** 212-220.

816   53.   Harris WS, Tintle NL, Etherton MR, Vasan RS. Erythrocyte long-chain omega-3 fatty acid

817   levels are inversely associated with mortality and with incident cardiovascular disease: The

818   Framingham Heart Study. *Journal of Clinical Lipidology* 2018; **12**(3)**:** 718-727.e716.

820   54.   Kaikkonen JE, Jula A, Viikari JSA, Juonala M, Hutri-Kähönen N, Kähönen M *et al.*

821   Associations of Serum Fatty Acid Proportions with Obesity, Insulin Resistance, Blood

822   Pressure, and Fatty Liver: The Cardiovascular Risk in Young Finns Study. *The Journal of*

823   *Nutrition* 2021; **151**(4)**:** 970-978.

825   55.   Harris WS, Pottala JV, Lacey SM, Vasan RS, Larson MG, Robins SJ. Clinical correlates

826   and heritability of erythrocyte eicosapentaenoic and docosahexaenoic acid content in the

827   Framingham Heart Study. *Atherosclerosis* 2012; **225**(2)**:** 425-431.

829   56.   Stark KD, Van Elswyk ME, Higgins MR, Weatherford CA, Salem N. Global survey of the

830   omega-3 fatty acids, docosahexaenoic acid and eicosapentaenoic acid in the blood stream

831   of healthy adults. *Progress in Lipid Research* 2016; **63:** 132-152.

833   57.   de Groot RHM, Emmett R, Meyer BJ. Non-dietary factors associated with n-3 long-chain

834   PUFA levels in humans – a systematic literature review. *British Journal of Nutrition* 2019;

835   **121**(7)**:** 793-808.

837   58.   Goldstein BI, Carnethon MR, Matthews KA, McIntyre RS, Miller GE, Raghuveer G *et al.*

838   Major Depressive Disorder and Bipolar Disorder Predispose Youth to Accelerated

839   Atherosclerosis and Early Cardiovascular Disease. *Circulation* 2015; **132**(10)**:** 965-986.

59. Wang JQ, Mao L. The ERK Pathway: Molecular Mechanisms and Treatment of Depression. *Molecular Neurobiology* 2019; **56**(9)**:** 6197-6205.

60. Duman CH, Schlesinger L, Kodama M, Russell DS, Duman RS. A Role for MAP Kinase Signaling in Behavioral Models of Depression and Antidepressant Treatment. *Biological Psychiatry* 2007; **61**(5)**:** 661-670.

61. Duman RS, Voleti B. Signaling pathways underlying the pathophysiology and treatment of depression: novel mechanisms for rapid-acting agents. *Trends in Neurosciences* 2012; **35**(1)**:** 47-56.

62. Zhang X, Yeung DCY, Karpisek M, Stejskal D, Zhou Z-G, Liu F *et al.* Serum FGF21 Levels Are Increased in Obesity and Are Independently Associated With the Metabolic Syndrome in Humans. *Diabetes* 2008; **57**(5)**:** 1246-1253.

63. Kralisch S, Tönjes A, Krause K, Richter J, Lossner U, Kovacs P *et al.* Fibroblast growth factor-21 serum concentrations are associated with metabolic and hepatic markers in humans. *Journal of Endocrinology* 2013; **216**(2)**:** 135-143.

64. Semba RD, Sun K, Egan JM, Crasto C, Carlson OD, Ferrucci L. Relationship of Serum Fibroblast Growth Factor 21 with Abnormal Glucose Metabolism and Insulin Resistance: The Baltimore Longitudinal Study of Aging. *The Journal of Clinical Endocrinology & Metabolism* 2012; **97**(4)**:** 1375-1382.

65. Chen C, Cheung BMY, Tso AWK, Wang Y, Law LSC, Ong KL *et al.* High Plasma Level of Fibroblast Growth Factor 21 Is an Independent Predictor of Type 2 Diabetes: A 5.4-year

population-based prospective study in Chinese subjects. *Diabetes Care* 2011; **34**(9)**:** 2113-

2115.

66.   Li H, Fang Q, Gao F, Fan J, Zhou J, Wang X *et al.* Fibroblast growth factor 21 levels are

increased in nonalcoholic fatty liver disease patients and are correlated with hepatic

triglyceride. *Journal of Hepatology* 2010; **53**(5)**:** 934-940.

67.   Ciuculete DM, Voisin S, Kular L, Welihinda N, Jonsson J, Jagodic M *et al.* Longitudinal

DNA methylation changes at MET may alter HGF/c-MET signalling in adolescents at risk

for depression. *Epigenetics* 2020; **15**(6-7)**:** 646-663.

68.   Teixeira AL, Gama CS, Rocha NP, Teixeira MM. Revisiting the Role of Eotaxin-1/CCL11 in

Psychiatric Disorders. *Frontiers in Psychiatry* 2018; **9**.

69.   Erickson MA, Morofuji Y, Owen JB, Banks WA. Rapid Transport of CCL11 across the

Blood-Brain Barrier: Regional Variation and Importance of Blood Cells. *Journal of*

Pharmacology and Experimental Therapeutics 2014; **349**(3)**:** 497.

70.   Villeda SA, Luo J, Mosher KI, Zou B, Britschgi M, Bieri G *et al.* The ageing systemic milieu

negatively regulates neurogenesis and cognitive function. *Nature* 2011; **477**(7362)**:** 90-94.

71.   Teixeira AL, Reis HJ, Nicolato R, Brito-Melo G, Correa H, Teixeira MM *et al.* Increased

serum levels of CCL11/eotaxin in schizophrenia. *Progress in Neuro-Psychopharmacology*

and Biological Psychiatry 2008; **32**(3)**:** 710-714.

72.   Grassi-Oliveira R, Brieztke E, Teixeira A, Pezzi JC, Zanini M, Lopes RP *et al.* Peripheral

chemokine levels in women with recurrent major depression with suicidal ideation. *Revista*

Brasileira de Psiquiatria 2012; **34**(1)**:** 71-75.

73.   Panizzutti B, Gubert C, Schuh AL, Ferrari P, Bristot G, Fries GR *et al.* Increased serum

levels of eotaxin/CCL11 in late-stage patients with bipolar disorder: An accelerated aging

biomarker? *Journal of Affective Disorders* 2015; **182:** 64-69.

74.   Simon NM, McNamara K, Chow CW, Maser RS, Papakostas GI, Pollack MH *et al.* A

detailed examination of cytokine abnormalities in Major Depressive Disorder. *European*

Neuropsychopharmacology 2008; **18**(3)**:** 230-233.

75.   Leighton SP, Nerurkar L, Krishnadas R, Johnman C, Graham GJ, Cavanagh J.

Chemokines in depression in health and in inflammatory illness: a systematic review and

meta-analysis. *Molecular Psychiatry* 2018; **23**(1)**:** 48-58.

76.   Goldsmith DR, Rapaport MH, Miller BJ. A meta-analysis of blood cytokine network

alterations in psychiatric patients: comparisons between schizophrenia, bipolar disorder

and depression. *Molecular Psychiatry* 2016; **21**(12)**:** 1696-1709.

77.   Osimo EF, Pillinger T, Rodriguez IM, Khandaker GM, Pariante CM, Howes OD.

Inflammatory markers in depression: A meta-analysis of mean differences and variability in

5,166 patients and 5,083 controls. *Brain, Behavior, and Immunity* 2020; **87:** 901-909.

78.   Foley ÉM, Parkinson JT, Mitchell RE, Turner L, Khandaker GM. Peripheral blood cellular

immunophenotype in depression: a systematic review and meta-analysis. *Molecular*

Psychiatry 2023; **28**(3)**:** 1004-1019.

79. Fernandes BS, Salagre E, Enduru N, Grande I, Vieta E, Zhao Z. Insulin resistance in

depression: A large meta-analysis of metabolic parameters and variation. *Neuroscience &*

Biobehavioral Reviews 2022; **139:** 104758.

80. Davyson E, Shen X, Gadd DA, Bernabeu E, Hillary RF, McCartney DL *et al.* Metabolomic

Investigation of Major Depressive Disorder Identifies a Potentially Causal Association With

Polyunsaturated Fatty Acids. *Biological Psychiatry* 2023.

81. Perry BI, Oltean BP, Jones PB, Khandaker GM. Cardiometabolic risk in young adults with

depression and evidence of inflammation: A birth cohort study. *Psychoneuroendocrinology*

2020; **116:** 104682.

82. Bolck A, Croon M, Hagenaars J. Estimating Latent Structure Models with Categorical

Variables: One-Step Versus Three-Step Estimators. *Political Analysis* 2004; **12**(1)**:** 3-27.