Apathy, intentions, explicit attitudes, and approach-avoidance tendencies in physical activity behavior

Ata Farajzadeh[1,†], François Jabouille[1,†], Nickolas Benoit[1], Olivier Bezeau[1], Tristan Bourgie[1], Benjamin Gerro[1], Jacob Ouimet[1], Matthieu P. Boisgontier[1,2,3,*]

[1]School of Rehabilitation Sciences, Faculty of Health Sciences, University of Ottawa, Canada; [2]Perley Health Centre of Excellence in Frailty-Informed Care, Ottawa, Canada; [3]Bruyère Health Research Institute, Ottawa, Canada; [†]AF and FJ contributed equally to this work; [*]Corresponding author: matthieu.boisgontier@uOttawa.ca

ABSTRACT

Background. Greater behavioral apathy has been shown to be associated with lower engagement in physical activity. However, the mechanisms underlying this association remain overlooked and poorly understood. Intentions, explicit attitudes, and approach-avoidance tendencies toward physical activity may play a central role in the relationship, given their strong links to motivation and physical activity.

Methods. An online study was conducted in 365 participants aged 54±18 years. All measures were assessed using questionnaires, except approach-avoidance tendencies, which were derived from reaction times in an approach-avoidance task. Component mediation analyses based on multiple linear regressions were conducted to examine whether the intention to be physically active mediated the relationship between behavioral apathy levels and usual physical activity levels, and whether explicit attitudes and approach-avoidance tendencies mediated the relationship between behavioral apathy and the intention.

Results. Results showed that weaker intentions to be physically active mediated the association between higher behavioral apathy and lower usual weekly levels of moderate-to-vigorous physical activity. In addition, explicit attitudes mediated the effect of behavioral apathy on intentions to be physically active. Results on approach-avoidance tendencies showed a significant three-way interaction between apathy, stimulus (physical activity vs. sedentary behavior), and action direction (approach vs. avoidance) on corrected reaction time (b = 19.6; 95CI = 2.0 to 37.3; p = .029), with higher apathy being associated with a greater tendency to avoid physical activity stimuli and to approach sedentary stimuli. However, we found no evidence suggesting that these tendencies mediated the effect of apathy on intentions or habitual physical activity. Based on our data, a mean item score greater than 34.5% of the scale range (e.g., >3.07 on a 1–7 scale) is indicative of behavioral apathy.

Conclusion. This study provides new insights into the role of intentions, explicit attitudes, and approach-avoidance tendencies toward physical activity in the relationship between behavioral apathy levels and the engagement in physical activity.

Keywords
Exercise, Health, Motivation, Neuroscience, Psychology

1. Introduction

Behavioral apathy is defined as a state of primary motivational impairment characterized by difficulty in elaborating the action plan required for behavior (Marin, 1990; Marin, 1991; Levy & Dubois, 2006). Apathy is observed in a wide range of conditions, including dementia (Leung et al., 2021), Parkinson's disease (den Brok et al., 2015), and stroke (Zhang et al., 2023), and has been associated with frailty (Mega et al., 1996), functional decline (Ayers et al., 2017), poorer quality of life (Groeneweg-Koolhoven et al., 2014), higher mortality (Vilalta-Franch et al., 2013), and increased healthcare costs (Kruse et al., 2023). Additionally, higher levels of apathy have been associated with lower levels of physical activity (Farajzadeh et al., 2024). This association may further compromise the health of individuals with apathy, as insufficient physical activity is known to be associated with cognitive decline (Cheval et al., 2023), cardiovascular disease (Wahid et al., 2016), cancer (Moore et al., 2016), hypertension (Liu et al., 2017), diabetes (Cheval et al., 2021), obesity (Bleich et al., 2018), depression (Boisgontier et al., 2020), and functional dependence (van Allen et al., 2024a; van Allen et al., 2024b). Therefore, understanding the mechanisms underlying the association between behavioral apathy and physical inactivity may contribute to improving the design and specificity of rehabilitation interventions, ultimately improving the condition of individuals with apathy. However, these mechanisms remain largely overlooked and poorly understood. Considering their close relationship with both motivation and physical activity, intentions, explicit attitudes, and approach-avoidance tendencies toward physical activity may play a pivotal role in the relationship between apathy and physical activity engagement.

Intention is thought to capture the motivational factors that influence a behavior and reflects the effort that individuals are willing to invest for the behavior to occur (Ajzen, 1987). This motivational construct is considered the most proximal antecedent of physical activity behavior (Ajzen, 1987; Biddle et al., 2007). Further, the role of intentions in physical activity engagement is highlighted in a meta-analysis showing that less than 5% of individuals engage in physical activity without intending to be physically active (Feil et al., 2023). However, while intention is necessary to engage in physical activity, it is not sufficient to guarantee this engagement (Rhodes & de Bruijn, 2013). This insufficiency is illustrated by meta-analyses showing that almost half of the individuals who intend to be physically active are unable to fulfill this intention (Rhodes & de Bruijn, 2013; Feil et al., 2023).

In recent years, dual-process models have been developed to address the limitations of social cognitive models in explaining physical activity behavior (Rhodes et al., 2019). These new idiosyncratic models assert that physical activity behavior is governed not only by reflective processes (e.g., intentions, explicit attitudes), but also by automatic processes (e.g., automatic attitudes) (Conroy & Berry, 2017; Brand & Ekkekakis, 2018; Cheval & Boisgontier, 2021; Cheval & Boisgontier, 2024). Reflective processes are typically described as slow, deliberative, inefficient (relying on higher brain functions), and conscious (Strack & Deutsch, 2004), and are typically assessed with questionnaires. However, reflective processes do not always exhibit all these characteristics simultaneously, as certain reflective responses can be efficient or partially automatic (Melnikoff & Bargh, 2018). In contrast, automatic processes are faster and initiated unintentionally, rely on learned associations, do not require conscious awareness, and are most often assessed using reaction times to visual stimuli (e.g., images depicting physical activity vs. sedentary behavior).

Attitude is a psychological tendency to evaluate a stimulus with some degree of favor or unfavor (Eagly & Chaiken, 1993, p. 1). This tendency is considered an indirect antecedent of physical

activity (Brand & Ekkekakis, 2018; Cheval & Boisgontier, 2021; Hagger & Chatzisarantis, 2014), influencing this behavior primarily through its effect on intention (Ajzen, 2011). Attitude can manifest as reflective (i.e., explicit attitudes) and automatic processes (i.e., automatic attitudes). Explicit attitudes are attitudes that people can report and for which activation can be consciously controlled (Rydell & McConnell, 2006). Automatic attitudes are introspectively unidentified traces of past experience that mediate favorable or unfavorable evaluation of a social object (Greenwald & Banaji, 1995). In other words, an automatic attitude is thought to result from the positive or negative value that our brain spontaneously assigns to some concept (e.g., person, place, or behavior), without that value being accurately accessible to cognition (Conroy & Berry, 2017). This implicit value of a stimulus results in an automatic positive or negative inclination toward this stimulus, which influences behavior. An illustration of this influence is approach-avoidance tendencies, an automatic preparation of the organism to execute a motor pattern toward or away from a behavior (Friese et al., 2011).

The aim of this study was to examine the relationships between apathy, the intention to be physically active, explicit attitudes toward physical activity, approach-avoidance tendencies toward physical activity and sedentary behaviors, and weekly physical activity engagement. According to a recent meta-analysis (Farajzadeh et al., 2024), we expected behavioral apathy to influence habitual levels of physical activity. Intention is essential for goal-directed behavior (Gollwitzer, 1997) and a key determinant of physical activity (Ajzen, 1987). Individuals with apathy often have difficulties initiating and maintaining goal-directed actions (Marin et al., 1991, Levy & Dubois, 2006). Therefore, we hypothesized that intention to engage in physical activity would mediate the relationship between apathy and habitual weekly physical activity engagement. The heightened sensitivity to effort-related cues in individuals with apathy leads them to perceive actions as more costly (Bonnelle et al., 2015; Bonnelle et al., 2016) and is likely to influence explicit attitudes toward physical activity as well as their antecedents, approach-avoidance tendencies. Therefore, our secondary hypothesis was that the relationship between apathy and intentions to be physically active would be mediated by explicit attitudes and approach-avoidance tendencies.

2. Methods

2.1. Population
Participants were recruited through social media, posters at the Faculty of Health Sciences, University of Ottawa, and emails to non-profit associations. Inclusion criteria were age 20–90 years and access to a personal computer, a laptop, or a tablet with internet. Informed consent was collected in accordance with the Declaration of Helsinki. The study was approved by University of Ottawa's Research Ethics Boards (H-05-21-6791). All participants provided informed consent. Data were collected between July 2022 and December 2023. Participants were not compensated for their participation.

2.2. Power Analysis
An a priori power analysis was conducted in G*power (Faul et al., 2009) to estimate the minimum sample required for $\alpha = 0.05$, power (1-β) = 90%, and a medium effect size $f^2 = 0.2$ (Cohen, 1988). We used this heuristic value because, at the time the study was designed, no meta-analysis had assessed the effect size of apathy on physical activity or the effect of apathy on the intention to be physically active. The power analysis was based on an F test in a multiple linear regression (R^2 increase) that included the two tested predictors (apathy and approach-avoidance tendencies) and

7 control variables and estimated that a minimum sample size of n = 67 was required. The aim of this approach was to ensure that we would have a 90% or greater chance of finding a statistically significant medium effect, if one exists, in each of the paths (i.e., each linear multiple regression model; Yzerbyt et al., 2018) of the mediation we described in our main hypothesis. However, because heuristics are a weak justification for determining an effect of interest, we aimed to collect as much data as possible within our time and resource constraints (Lakens, 2022). Of note, the meta-analysis by Farajzadeh et al. (2024), published after the data collection for the current study, provided two effect size estimates for the relationship between apathy and physical activity engagement: r = -0.13 (f^2 = 0.017, based on r values in the literature) and r = -0.40 (f^2 = 0.191, based on rho values in the literature). Using these effect sizes, analyses based on an F test in a multiple linear regression that included one tested predictor (apathy) estimated that a minimum sample size of n = 58 to 621 would have been required. Another approach would have been to use the powerMediation R package (Qiu, 2022). However, we decided against this approach because the lack of information available in the literature would have forced us to use heuristics to set the values of several arguments required for the power analysis.

2.3. Experimental Protocol

2.3.1. Procedures

Participants performed approach-avoidance tasks online using Inquisit 6 software (Millisecond Software, 2015), and responded to questions related to apathy, habitual level of moderate-to-vigorous physical activity, age, sex (male, female), gender (man, woman, non-binary, transgender man, transgender woman, other), weight, height, depressive symptoms, and health condition. One attention check question was included in the questionnaires: "Please answer "5" to this question that allows us to verify that you actually read the questions."

2.3.2. Self-Reported Variables

Behavioral Apathy

Behavioral apathy was assessed using the action initiation subscale of the Lille Apathy Rating Scale (LARS) (Sockeel et al., 2006; Bonnelle et al., 2015). This subscale focuses on everyday productivity and initiative, and has been used as an index of behavioral apathy (Bonnelle et al., 2016). This subscale includes 11 questions with responses ranging from 1 (completely untrue) to 7 (completely true), resulting in a total score ranging from 11 to 77. We used the total score in our analyses, with higher scores indicating a higher level of behavioral apathy.

Habitual Level of Moderate-to-Vigorous Physical Activity

The habitual level of physical activity was derived from the short form of the International Physical Activity Questionnaire (IPAQ-SF), a self-administered questionnaire that identifies the frequency and duration of moderate and vigorous physical activity during the past 7 days to estimate habitual level of physical activity and sedentary behavior (Craig et al., 2003). The habitual level of moderate-to-vigorous physical activity (MVPA) in minutes per week was used in the analyses.

Explicit Attitudes

Explicit attitudes toward physical activity were calculated as the mean of two items based on two bipolar semantic differential adjectives on a 7-point scale ["unpleasant" (1) – "pleasant" (7); "unenjoyable" (1) – "enjoyable" (7)]. The statement begins with "For me, to participate in regular physical activity is …" (Hoyt et al., 2009; Farajzadeh et al., 2023). These two scores showed a

very strong correlation ($r = .915$; $p < 2.2 \times 10^{-16}$), and their sum was used as an indicator of explicit attitudes in the statistical analyses.

Intention to Be Physically Active
The intention to be physically active was derived from the response to the question "How much do you agree with the following statement: Over the next 7 days, I intend to do at least 150 minutes of moderate-intensity physical activity; or at least 75 minutes of vigorous intensity physical activity; or an equivalent combination of moderate- and vigorous-intensity physical activity" on a 7-point scale ranging from "strongly disagree" (1) to "strongly agree" (7) (Rhodes & Rebar, 2017).

Depressive Symptoms
As it has been shown to be associated with apathy (Starkstein et al., 2001), depression needed to be controlled for in our analyses. Depressive symptoms were assessed using the Depression subscale of the Depression Anxiety Stress Scale (Lovibond & Lovibond, 1995). Participants were asked to read 14 statements and indicate how much the statement applied to them over the past week using a 4-point Likert scale ranging from 0 ("did not apply to me at all") to 3 ("applied to me very much of the time"), resulting in a total score ranging from 0 to 48. We used the total score as a control variable in our analyses, with higher scores indicating greater depressive symptoms.

Chronic Conditions
Since it has been shown to be related with physical activity (Vancampfort et al., 2017), our analyses included the number of chronic conditions, which was derived from a question based on item PH006 of the Survey of Health, Ageing and Retirement in Europe (Börsch-Supan, 2022). "Has a doctor ever told you that you had any of the following conditions?". The possible answers were "A stroke or cerebral vascular disease", "High blood pressure or hypertension", "High blood cholesterol", "Diabetes or high blood sugar", "arthritis, including osteoarthritis, or rheumatism", "rheumatoid arthritis", "chronic lung disease such as chronic bronchitis or emphysema", "asthma", "osteoporosis", "Cancer or malignant tumour, including leukaemia or lymphoma, but excluding minor skin cancers", "stomach or duodental ulcer, peptic ulcer", "Parkinson disease", "hip fracture or femoral fracture", "Alzheimer's disease, dementia, organic brain syndrome, senility or any other serious memory impairment", "other affective or emotional disorders, including anxiety, nervous or psychiatric problems", "chronic kidney disease", "other conditions, not yet mentioned", and "none".

2.3.3. Approach-Avoidance Tendencies
Approach-Avoidance Task
Approach-avoidance tendencies toward physical activity and sedentary stimuli were tested using an approach-avoidance task, which has shown good reliability ($r = 0.83$) (Farajzadeh et al., 2023) and good validity (Zenko & Ekkekakis, 2019). A video illustrating the task is available online (Boisgontier, 2024). Two experimental conditions and two neutral conditions were tested (Cheval et al., 2018; Farajzadeh et al., 2023). In the experimental conditions of this task, a trial starts with a fixation of a cross presented at the center of the screen for a random time ranging from 500 to 750 ms (Figure 1A). Then, an avatar appears either at the top or bottom third of the screen for one second, before a pictogram representing a physical activity behavior or a sedentary behavior appears in the center of the screen (Figure 1A). The participant sitting in front of the computer with one index finger positioned on the "U" and the other index finger on the "N" key is instructed

that pressing the "U" key moves the avatar up and pressing the "N" key moves the avatar down. Accordingly, the movement of the avatar is always congruent with the pressed key: The top key (i.e., U) moves the avatar up, while the bottom key (i.e., N), moves the avatar down. Importantly, however, the approach or avoidance action depends on the initial position of the avatar at the beginning of the trial. If the avatar appears below the stimulus, the top key is associated with an approach movement, while the bottom key is associated with an avoidance movement. Conversely, if the avatar appears above the stimulus, the approach and avoidance movement are reversed – the top key is associated with an avoidance movement and the bottom key is associated with an approach movement.

Figure 1. A. Illustration of a trial of the approach-avoidance task in the condition where the participant is instructed to approach physical activity stimuli (and avoid sedentary stimuli – not shown). B. Timeline and stimuli of the approach-avoidance task. In the experimental and neutral Condition 1, the participant is instructed to move the avatar toward (i.e., approach) a type of stimuli (i.e., physical activity or rectangles) and to move the avatar away from (i.e., avoid) stimuli depicting the other type of stimuli (i.e., sedentary behavior or ellipses, respectively). In Condition 2, the instruction is reversed: The participant is instructed to move away from physical activity (experimental condition) or rectangle stimuli (neutral condition) and move toward sedentary stimuli or ellipse stimuli.

Experimental Conditions
Two experimental conditions were tested (Figure 1B). In one experimental condition, the

participant is instructed to quickly move the avatar toward (i.e., approach) pictograms depicting physical activity and to move the avatar away from (i.e., avoid) pictograms depicting sedentary behavior. In the other experimental condition, the participant does the opposite: move away from physical activity and move toward sedentary stimuli. The order of the experimental conditions was randomized across participants. In a previous study (Cheval et al., 2018), thirty-two participants were asked to rate the extent to which 24 stimuli expressed "movement and an active lifestyle" and "rest and sedentary lifestyle" (1 = not at all, 7 = a lot). For each stimulus, the "rest and sedentary lifestyle" score was subtracted from the "movement and active lifestyle" score. In the current study, the six stimuli with the largest positive and negative differences were chosen as the stimuli depicting physical activity and sedentary behaviors, respectively.

Neutral Conditions
In addition to the two experimental conditions, two neutral conditions were tested. These conditions were used to account for a potential generic approach-avoidance tendency that could vary across participants and ages (Farajzadeh et al., 2023). In these neutral conditions, the stimuli depicting physical activity and sedentary behaviors were replaced by stimuli made of rectangles or ellipses that matched the number and size of information in 3 physical activity stimuli (swimming, hiking, cycling) and 3 sedentary stimuli (couch, hammock, reading). Two conditions were tested. In one condition, participants are asked to quickly move the manikin toward stimuli with circles and away from stimuli with squares. In the other condition, the participant is given opposite instructions. The order of the neutral conditions was randomized.
One neutral condition was tested before the two experimental conditions, and the other neutral condition was tested after them. Each condition included 96 stimuli, 48 of each class (physical activity and sedentary stimuli in the experimental conditions; rectangles and ellipses in the neutral conditions), that were presented randomly. Familiarization with the task was performed during the first 15 trials of the study, which were removed from the analyses. Familiarization with the subsequent conditions was performed during the first 3 trials of each condition, which were removed from the analyses. The physical activity and sedentary stimuli were presented all together on the screen for seven seconds before each experimental condition. Between conditions, the participant could rest for as long as they wanted before pressing the space key to start the next condition. When the participant pressed the incorrect key ("U" when it should be "N" or "N" when it should be "U"), the message "error" appeared on the screen for 800 ms before the next trial. When the reaction time (i.e., the time between the appearance of the stimuli and the key press) was longer than eight seconds, the message "too slow" appeared on the screen for 800 ms before the next trial (Figure 1A).

Reaction Times
The tendency to approach or avoid a type of stimuli (i.e., physical activity, sedentary, or neutral stimuli) was derived from the time required to press the key in reaction to a type of stimulus (i.e., physical activity vs. sedentary vs. neutral). Corrected reaction times were computed by subtracting the mean reaction time to approach or avoid neutral stimuli from the reaction time on each trial to respectively approach or avoid stimuli depicting a type of behavior (physical activity or sedentary behavior). The bias toward a type of stimulus (physical activity or sedentary behavior) was computed by subtracting the mean corrected reaction time to approach this type of stimulus from the corrected reaction time to avoid it. Therefore, positive scores were indicative of an approach bias, i.e., a higher tendency to approach the stimulus. Incorrect responses, responses faster than

150 ms, and responses slower than 3,000 ms were excluded from the analyses to account for outliers and loss of attention (Farajzadeh et al., 2023). Specifically, because simple reaction times are typically longer than 200 ms (Fozard et al., 1994), and because we were assessing choice reaction times, we considered that reaction times faster than 150 ms could not be considered reactions to the stimuli of our task. In young adults, complex choice reaction times with up to 15 choices and different combinations of limb responses are typically faster than 1,000 ms (Boisgontier et al., 2014). However, in older adults, they can be longer than 3,000 ms (Boisgontier et al., 2016). Therefore, as with the lower threshold (150 ms), we chose a conservative upper threshold to avoid excluding a large proportion of valid reaction times in the older participants, as these observations can have a significant impact on the results (Ulrich & Miller, 1994; Miller, 2023). We did not include reaction times longer than 3000 ms because even in the complex task mentioned above (Boisgontier et al., 2016), the proportion of reaction times longer than 3,000 ms was less than 0.5% of the data. Therefore, we considered reaction times longer than 3,000 ms to be too long to be solely representative of a response to the stimuli of our task.

2.4. Statistical Analyses

To examine the mediation effect of intentions on the association between apathy and the habitual level of moderate-to-vigorous physical activity, we used the component approach (Yzerbyt et al., 2018). This component approach to the assessment of mediation was preferred over the index approach (Hayes, 2022), because the latter has shown a higher risk of false positives (Type I errors) (Yzerbyt et al., 2018). The component approach involves three linear multiple regressions models. Model 1 examines whether the independent variable affects the dependent variable. Model 2 examines the effect of the independent variable on the mediator. Model 3 examines both the independent variable and the mediator as simultaneous predictors of the dependent variable. Mediation was claimed if the "total effect" of the dependent variable in Model 1 is larger in absolute value than its "residual effect" in Model 3. The bias to approach physical activity stimuli, the bias to approach sedentary stimuli, depressive symptoms, age, sex, body mass index, and the number of chronic health conditions were included as control variable in all the models used to test the mediation. The first mediation analysis examined the intention to be active as a mediator of the effect of apathy on the habitual level of physical activity. The second mediation analysis examined explicit attitudes as a mediator of the effect of apathy on the intention to be active. The third mediation analysis examined approach-avoidance tendencies as a mediator of the effect of apathy on the intention to be active.

To further examine the effect of apathy on approach-avoidance attitudes related toward physical activity and sedentary stimuli, a linear and a logistic mixed-effect models (Lohse et al., 2023) were built and fit by maximum likelihood in the R software environment (R Core Team, 2023), using the lme4 (Bates et al., 2021) and lmerTest package (Kuznetsova et al., 2022), which approximates p-values using Satterthwaite's degrees of freedom method. Continuous variables were standardized. For the linear mixed-effects model, restricted maximum likelihood (REML) was used as it provides less biased estimates of variance components than full maximum likelihood (Luke, 2017). Fixed effects included a three-way interaction effect of apathy (continuous), stimulus (physical activity vs. sedentary behavior), and action direction (approach vs. avoid) on corrected reaction time. The other fixed effects controlled for the effect of depressive symptoms, the habitual level of moderate-to-vigorous physical activity, age, sex, body mass index, the number of chronic conditions, and the device used to complete the study (computer vs. tablet). Our balanced design was fully crossed: Each participant was tested in the approach and avoid condition

of four types of stimuli (physical activity, sedentary behavior, rectangles, ellipses), with each type including 6 pictograms. Therefore, we intended to include the random effect of participant, action direction, stimulus, and pictogram (Lohse et al., 2023). However, the model converged only when the random effects of stimulus was removed.

To ensure that the results obtained with the corrected reaction times cannot be explained by the speed-accuracy trade-off (Hick, 1952), we conducted a logistic mixed-effects model with the number of errors in the Approach-Avoidance Task as outcome. The structure of this model was similar to the linear mixed-effects models using reaction time as outcome. However, because the logistic mixed-effects models did not converge when the three-way interaction was included, we conducted a model with a two-way interaction between apathy and action direction on errors when reacting to physical activity stimuli and another model with the same interaction but in the conditions with sedentary stimuli. In addition, only the fixed effects of depressive symptoms, age, and explicit attitudes as well as the random effect of participant were included to allow for model convergence.

3. Results

3.1. Descriptive Results

3.1.1. Participants

Three hundred and ninety two participants initiated the study. Some of them were included in a previous study from our group (Farajzadeh et al., 2022). Twenty three were excluded because they stopped the session before completing the study. Four participants were excluded because they answered the check question incorrectly. When participants reported height <50 cm or >250 cm or weight <30 kg or >250 kg), the data was removed and imputed using the sample mean. The final sample of 365 participants was 53.7 ± 17.9 years (mean ± standard deviation), with a mean score of behavioral apathy of 31.3 ± 9.3 out of a maximum of 77 (mean score per item of 2.9 ± 0.9 on a scale ranging from 1 to 7), a habitual level of moderate-to-vigorous physical activity of 398.3 ± 496.0 min per week, mean depressive symptoms of 0.7 ± 0.4, body mass index of 27.6 ± 12.1 kg.m^{-2}, and 1.5 ± 1.5 chronic health conditions. Two males identified themselves as women. All the other male (n = 146) and female participants (n = 217) identified themselves as men and women, respectively.

Table 1 – Descriptive statistics (n = 365, 148 females)

Variable (minimum score-maximum score)	Mean	SD	Range
Age (years)	53.7	17.9	20-89
Body mass index (kg.m^{-2})	27.6	12.1	15.1-48.7
Moderate-to-vigorous physical activity (min.week^{-1})	398.4	496.0	0-3080
Explicit Attitude (2-14)	11.0	3.2	2-14
Intention (1-7)	5.1	2.1	1-7
Behavioral Apathy (11-77)	31.3	9.3	11-60
Depression (0-48)	9.3	6.2	0-37
Chronic Diseases (0-18)	1.5	0	7
Approach physical activity stimuli (ms)	1094	547	163-2996
Avoid physical activity stimuli (ms)	1183	538	172-2998
Approach sedentary stimuli(ms)	1202	590	174-2996
Avoid sedentary stimuli (ms)	1153	535	192-2989
Approach neutral stimuli (ms)	1021	542	156-2999
Avoid neutral stimuli (ms)	1046	523	154-2997

3.1.2. Cut-off score for behavioral apathy

In our study, 126 participants reported having no chronic condition. In this subsample, the mean score per item was 2.93 on a scale of 1 to 7, with a 95% confidence interval (95CI) ranging from 2.79 to 3.07. This result suggested that a mean score per item > 34.5% [(3.07 − 1) / (7 − 1)] of the scale range is indicative of behavioral apathy. Based on this cut-off score, participants with a score ≥ 34 would be classified as adults with apathy (i.e., 37.5% of the participants). The same approach applied to our full sample (n = 365) yielded similar results (95CI upper bound = 2.98; 33.1% of the scale range).

Figure 2. Flow diagram depicting the distribution of the trials

3.1.3. Observations in the approach-avoidance task
A total of 71,378 trials were performed by the participants (Figure 2), of which 12,350 were excluded: 531 trials were aborted at eight seconds with the "too slow" message appearing on the screen, 768 were removed because the participants failed the attention check question, 3,965 reactions were longer than 3,000 ms and 119 were faster than 150 ms, and 6,967 were familiarization trials. Among the 59,028 trials included in the analyses, 55,050 were in the correct direction (14,523 for sedentary stimuli and 14,724 for physical activity stimuli, 25,803 for neutral stimuli). An error occurred in 3,978 trials. A total of 15,539 trials were reactions to sedentary stimuli and 15,662 to physical activity stimuli (including reactions in the correct and incorrect direction)."

3.2. Statistical Results
3.2.1. Intentions Mediate the Effect of Apathy on Physical Activity Engagement
Model 1 showed an association between apathy and the habitual level of physical activity (b = -67.6; 95CI = -119.1 to -16.2; p = .010). Model 2 showed an association between apathy and the intention to be physically active (b = -0.220; 95CI = -0.431 to -0.009; p = .041). Model 3 showed an association between intentions and physical activity (b = 147.3; 95CI = 97.4 to 197.3; p = 1.48 × 10^{-8}). Model 3 also showed that the association between apathy and the level of physical activity was reduced when intentions were added as a predictor (b = -51.9; 95CI = -101.4 to -2.3; p = .040). From model 1 to 3, this association decreased by 23.3%. Taken together, these results demonstrate that the intention to be physically active partially mediates the relationship between apathy and the habitual level of physical activity (Figure 3).

Figure 3. Mediation effect of intention to be physically active on the association between apathy and the habitual level of physical activity. Note: b' is the estimate of the residual effect of apathy when intention is included in the model.

Figure 4. Mediation effect of explicit affective attitudes on the association between apathy and intentions to be physically active. Note: b' is the estimate of the residual effect of apathy when explicit affective attitudes are included in the model. NS = non significant.

3.2.2. Explicit attitudes Mediate the Effect of Apathy on Intentions

Model 1 showed an association between apathy and the intention to be physically active ($b = -0.220$; 95CI = -0.431 to -0.009; $p = .041$). Model 2 showed an association between apathy and explicit attitudes toward physical activity ($b = -0.707$; 95CI = -1.019 to -0.394; $p = 1.15 \times 10^{-5}$). Model 3 showed an association between explicit attitudes and the intention to physically active ($b = 1.376$; 95CI = 1.202 to 1.550; $p < 2 \times 10^{-16}$). Model 3 also showed that the association between apathy and intentions to be physically active was no longer significant when explicit attitudes were added as a predictor ($b = 0.084$; 95CI = -0.083 to 0.252; $p = .324$). Taken together, these results demontrate that explicit attitudes fully mediate the relationship between apathy and intentions to

be physically active (Figure 4).

3.2.3. Apathy and Approach-Avoidance Tendencies

Results of the linear mixed-effects model showed a significant three-way interaction between apathy (continuous), stimulus (physical activity vs. sedentary behavior), and action direction (approach vs. avoid) on corrected reaction time (b = 19.6; 95CI = 2.0 to 37.3; p = .029) (Figure 5). Simple effect analyses showed that lower apathy scores were associated with faster approach than avoidance of physical activity stimuli, but this difference was no longer significant when the apathy score was ≥ 55 out of a maximum mean score of 77. Additionaly, simple effect analyses showed that lower apathy scores were associated with faster avoidance than approach of sedentary stimuli, but this difference was no longer significant when the apathy scores were ≥ 70. In other words, lower apathy were associated with tendencies to approach physical activity stimuli and to avoid sedentary stimuli. Both these tendencies suggested a positive automatic evaluation of stimuli associated with physical activity. However, such tendencies decreased as apathy increased until they lost their statistical significance.

Results of the logistic mixed-effects models showed no evidence of an interaction between the effects of apathy and action direction on errors when reacting to physical activity (b = .135; 95CI = -.011 to .283; p = .07) or sedentary stimuli (b = .128; 95CI = -.025 to .283; p = .103). Additionally, results showed no evidence of an association between approach-avoidance tendencies and the habitual level of moderate-to-vigorous physical activity, ruling out the possibility for a mediation effect.

Figure 5. Results of the three-way interaction between the effect of apathy (continuous), stimulus (physical activity vs. sedentary behavior), and action direction (approach vs. avoid) on corrected reaction time. Note. LARS = Lille Apathy Rating Scale. NS = Non significant effect of action direction.

4. Discussion

4.1. Main findings

As hypothesized, weaker intention to be physically active mediated the association between higher behavioral apathy and lower habitual levels of moderate-to-vigorous physical activity. In addition, explicit attitudes mediated the relationship between behavioral apathy and intentions to be

physically active. While higher apathy was associated with a stronger tendency to avoid physical activity stimuli and to approach sedentary stimuli, we found no evidence suggesting that this tendency mediated the relationship between apathy and intentions or habitual physical activity. Based on our data, a mean item score greater than 34.5% of the scale range (e.g., >3.07 on a 1–7 scale) is indicative of behavioral apathy.

4.2. Comparison with the Literature

4.2.1. Behavioral Apathy Cut-Off Score

Our data in otherwise healthy older adults (n = 126) suggest that a mean score per item > 34.5% of the scale range is indicative of behavioral apathy. Based on this cut-off score, 37.5% of our participants (n = 365) would be classified as having behavioral apathy. Of note, this cut-off score was similar to the one based on our entire sample (33.1% of the scale range), which is consistent with the absence of statistical correlation between behavioral apathy and chronic conditions (r = -0.017, p = .742) and with the conceptualization of apathy as a syndrome rather than a symptom, since apathy levels were similar regardless of health status. In addition, our cut-off score is consistent with the mean item score of 29.7% of the scale range that was used as a cut-off score in another study of otherwise healthy young and middle-aged adults (n = 108, 94.5% aged 20 to 49 years), although they used a different scale, the Apathy Evaluation Scale ranging, which ranges from 1 to 4 (Kant et al., 1998). However, other results in otherwise healthy young adults (n = 139) showed a mean apathy score per item of 1.46 on a scale of 1 to 5, with an upper bound 95CI of 1.58 (Bonnelle et al., 2016). Accordingly, they considered a mean score per item of 14.5% of the scale range to reflect behavioral apathy. Based on this cut-off score, 88.5% of our sample would have been classified as adults with apathy. We initially hypothesized that this higher threshold for apathy in our study compared to the study by Bonnelle et al. (2016) might be due to the broader age distribution (mean age: 27.4 ± 7.8 vs. 53.7 ± 17.9 years) since a recent meta-analysis showed stronger correlation between apathy and physical activity in older adults compared to younger adults (Farajzadeh et al., 2024). However, our results showed no evidence of a positive correlation between behavioral apathy and age in the healthy subset (r = -0.173, p = .053).

4.2.2. Behavioral Apathy and Physical Activity

The association between apathy and physical activity observed in the current study aligns with results of a recent meta-analysis, which reported a negative correlation based on 22 studies (n = 12,541) (Farajzadeh et al., 2024).

4.2.3. Mediating Role of Intentions

For the first time, we show an association between behavioral apathy and intentions to be physically active. This finding suggests that behavioral apathy affects the intention stage of goal-directed behaviors (Levy & Dubois, 2006), such as physical activity. In addition, this finding is conceptually coherent because behavioral apathy, a motivational impairment (Marin, 1990; Marin, 1991; Levy & Dubois, 2006), is likely to reduce the amount of effort individuals are willing to invest for the physical activity behavior to occur. This reduction could be explained by an increased sensitivity to effort (Bonnelle et al., 2016) and/or a reduced sensitivity to rewards associated with physical activity (Pessiglione et al., 2018), both of which could be linked to changes in brain circuitry, even at subclinical levels of apathy (Derosiere et al., 2024).

Our results also show an association between the intention to be physically active and the engagement in physical activity behavior, which is in line with previous theoretical and experimental literature (Ajzen, 1987; Biddle et al., 2007; Maltagliati et al., 2024a). Recent

experimental results based on the theory of effort minimization in physical activity (Cheval & Boisgontier, 2021; Cheval & Boisgontier, 2024) suggest that this association between intentions and physical activity behavior is influenced by individual differences in the general tendencies to approach and avoid physical effort (Cheval et al., 2024; Maltagliati et al., 2024a). Specifically, the positive association between intentions to be physically active and physical activity engagement was stronger when self-reported tendencies to approach physical effort were higher, and weaker when tendencies to avoid physical effort were higher. These findings encourage future studies to examine whether approach-avoidance tendencies toward physical effort influence the relationship between behavioral apathy and physical activity engagement.

The observed mediating effect of intention suggests that interventions aimed at increasing physical activity levels in patients with apathy should first focus on enhancing their intention to be physically active. While such intention does not guarantee engagement in physical activity, it is an essential prerequisite for this engagement (Rhodes & de Bruijn, 2013; Feil et al., 2023).

4.2.4. Mediating Role of Explicit Attitudes

For the first time, we show an association between behavioral apathy levels and explicit affective attitudes toward physical activity, which mediate the relationship between apathy and intentions to be physically active. These attitudes are based on the aggregation of the likelihood of anticipated affective responses (Williams & Evans, 2014; Stevens et al., 2020). Therefore, to strengthen positive explicit attitudes in individuals with apathy, ensuring that exercise therapy is consistently associated with positive experiences may be important. This suggestion is consistent with existing literature highlighting the role of pleasure and displeasure in physical activity engagement (Ekkekakis, 2017; Maltagliati et al., 2024b). Ultimately, this effect of explicit attitudes on intentions may influence behavior as within-subject increases in positive affect have been shown to be associated with physical activity levels at 3-month follow-up (Kwan & Bryan, 2010).

4.2.5. Physical Activity and Approach-Avoidance Tendencies

Literature has consistently reported a tendency to approach physical activity stimuli and to avoid sedentary stimuli at all ages (Farajzadeh et al., 2023). In the current study, we showed that these tendencies are influenced by apathy. Participants with stronger behavioral apathy showed weaker tendencies to approach physical activity and avoid sedentary behaviors. Approach-avoidance tendencies reflect motivational processes that direct organisms toward beneficial stimuli (approach) and away from detrimental stimuli (avoidance), driving adaptive behavior (Elliot et al., 2013). These tendencies are driven by internal states and action plans that translate perceived urges into behavior (Sander et al., 2018). Specifically, factors such as the perceived reward and the amount of effort involved in a behavior can influence whether a person is motivated to approach or avoid this behavior. Apathy, characterized by reduced motivation and difficulty generating action plans for goal-directed behavior, results from a desensitization to reward (Martínez-Horta et al., 2014; Muhammed et al., 2016) or an increased sensitivity to effort (Bonnelle, 2016; Le Heron, 2018; Le Heron, 2019). Therefore, individuals with apathy may perceive physical activity as more effortful and less rewarding than individuals without apathy. Thus, the threshold of the reward-to-effort ratio that triggers behavior may be harder to reach, reducing the likelihood of approaching physical activity. This example illustrates how apathy can disrupt the approach-avoidance dynamic and dampen the internal drive toward physical activity behavior.

4.3. Limitations and Strengths

The present study has potential limitations. First, the online nature of the study made it impossible to limit the influence of potential distractions in the participant's environment and to control whether participants were using their two index fingers to perform the task as instructed and whether they were sitting or standing, which could have influenced the results (Cheval et al., 2018; Maltagliati et al., 2024c). Second, the participants were recruited in Canada. It is, thus, unclear whether conclusions could generalize to populations from non-Western countries or less active populations of older adults. Third, the habitual level of physical activity was assessed using a self-reported questionnaire, which may not accurately reflect the objective level of physical activity. Assessing physical activity and sedentary behaviors using device-based measures would have provided more reliable estimates. However, these limitations are offset by several strengths, including a large sample size, a measure that accounts for a potential generic approach-avoidance bias that could have confounded the results (Farajzadeh et al., 2023), and the use of statistics that limit information loss (i.e., mixed effects models) (Boisgontier & Cheval, 2016; Lohse et al., 2023).

5. Conclusion

This study supports previous work suggesting that apathy is a syndrome rather than a symptom (Marin et al., 1991), affecting not only individuals with a medical condition but also otherwise healthy adults across all ages. In addition, the results highlight the key role of intentions and explicit affective attitudes in mediating the relationship between behavioral apathy and habitual physical activity. These insights suggest that interventions aimed at improving or maintaining physical activity levels should assess apathy regardless of a patient's or client's health status. For individuals identified as having apathy, incorporating motivational constructs—such as the intention to be physically active and positive affective attitudes toward physical activity—into the intervention design should be considered.

6. Declarations

Data and Code Availability
According to good research practices (Boisgontier, 2022), the dataset and the R script are available in Zenodo (Boisgontier, 2024).

Authorship Contribution Statement
Based on the Contributor Roles Taxonomy (CRediT) (Allen et al., 2019), individual author contributions to this work are as follows: Ata Farajzadeh: Writing – Review and Editing. François Jabouille: Writing – Review and Editing. Nickolas Benoit: Investigation. Olivier Bezeau: Investigation. Tristan Bourgie: Investigation. Benjamin Gerro: Investigation. Jacob Ouimet: Investigation. Matthieu P. Boisgontier: Conceptualization (Lead); Methodology; Formal Analysis; Data Curation; Visualization; Writing – Original Draft; Writing – Review and Editing; Supervision (AF, FJ, NB, OB, TB, BG, JO); Project Administration; Funding Acquisition. All authors have read and agreed to the published version of the manuscript.

Funding
Matthieu P. Boisgontier is supported by Natural Sciences and Engineering Research Council of Canada (NSERC) (RGPIN-2021-03153), the Canada Foundation for Innovation (CFI 43661), and Mitacs.

Conflict of Interest

The authors declare that there are no conflicts of interest related to the content of this article.

7. References

Ajzen, I. (1987). Attitudes, traits, and actions: dispositional prediction of behavior in personality and social psychology. *Advances in Experimental Social Psychology,* 20, 1–63. https://doi.org/10.1016/S0065-2601(08)60411-6

Ajzen, I. (2011). The theory of planned behaviour: reactions and reflections. *Psychology & Health*, 26(9), 1113–1127. https://doi.org/10.1080/08870446.2011.613995

Allen, L., O'Connell, A., & Kiermer, V. (2019). How can we ensure visibility and diversity in research contributions? How the Contributor Role Taxonomy (CRediT) is helping the shift from authorship to contributorship. *Learned Publishing*, 32(1), 71–74. https://doi.org/10.1002/leap.1210

Ayers, E., Shapiro, M., Holtzer, R., Barzilai, N., Milman, S., & Verghese, J. (2017). Symptoms of apathy independently predict incident frailty and disability in community-dwelling older adults. *Journal of clinical psychiatry*, 78(5), e529–e536. https://doi.org/10.4088/JCP.15m10113

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R. H. B., Singmann, H., Dai, B., Scheipl, F., Grothendieck, G., Green, P., Fox, J., Bauer, A., & Krivitsky, P. N. (2021). lme4: linear mixed-effects models using 'eigen' and S4 (version 1.1-27.1). [R package] https://cran.r-project.org/web/packages/lme4/lme4.pdf

Biddle, S. J. H., Hagger, M. S., Chatzisarantis, N. L. D., & Lippke, S. (2007). Theoretical frameworks in exercise psychology. In G. Tenenbaum & R. C. Eklund (Eds.), *Handbook of Sport Psychology* (3rd ed., pp. 537–559). Hoboken, NJ: John Wiley & Sons. https://doi.org/10.1002/9781118270011.ch24

Bleich, S. N., Vercammen, K. A., Zatz, L. Y., Frelier, J. M., Ebbeling, C. B., & Peeters, A. (2018). Interventions to prevent global childhood overweight and obesity: a systematic review. *The Lancet Diabetes & Endocrinology*, 6(4), 332–346. https://doi.org/10.1016/S2213-8587(17)30358-3

Boisgontier, M. P. (2022). Research integrity requires to be aware of good and questionable research practices. *European Rehabilitation Journal*, 2(1), 1–3. https://doi.org/10.52057/erj.v2i1.24

Boisgontier, M. P. (2024). Apathy, motivation, and physical activity behavior: Material, data and R code. Zenodo, v.1.3. [dataset, R script, Material]. https://doi.org/10.5281/zenodo.12750404

Boisgontier, M. P., & Cheval, B. (2016). The anova to mixed model transition. *Neuroscience & Biobehavioral Reviews,* 68, 1004–1005. https://doi.org/10.1016/j.neubiorev.2016.05.034

Boisgontier, M. P., Orsholits, D., von Arx, M., Sieber, S., Miller, M. W., Courvoisier, D., Iversen, M. D., Cullati, S., & Cheval, B. (2020). Adverse childhood experiences, depressive symptoms, functional dependence, and physical activity: a moderated mediation model. *Journal of Physical Activity & Health*, 17(8), 790–799. https://doi.org/10.1123/jpah.2019-0133

Boisgontier, M. P., van Ruitenbeek, P., Leunissen, I., Chalavi, S., Sunaert, S., Levin, O., & Swinnen, S. P. (2016). Nucleus accumbens and caudate atrophy predicts longer action selection times in young and old adults. *Human Brain Mapping*, 37(12), 4629–4639. https://doi.org/10.1002/hbm.23333

Boisgontier, M. P., Wittenberg, G. F., Fujiyama, H., Levin, O., & Swinnen, S. P. (2014). Complexity of central processing in simple and choice multilimb reaction-time tasks. *PLoS One*, 9(2), e90457. https://doi.org/10.1371/journal.pone.0090457

Bonnelle, V., Manohar, S., Behrens, T., & Husain, M. (2016). Individual differences in premotor brain systems underlie behavioral apathy. *Cerebral Cortex*, *26*(2), 807–819. https://doi.org/10.1093/cercor/bhv247

Bonnelle, V., Veromann, K. R., Burnett Heyes, S., Lo Sterzo, E., Manohar, S., & Husain, M. (2015). Characterization of reward and effort mechanisms in apathy. *Journal of Physiology*, *109*(1-3), 16–26. https://doi.org/10.1016/j.jphysparis.2014.04.002

Börsch-Supan A (2022) *Survey of Health, Ageing and Retirement in Europe (SHARE) (Wave 8)* (Version 8.0.0). [Data set] https://doi.org/10.6103/SHARE.w8.800

Brand, R., Ekkekakis, P. Affective–Reflective Theory of physical inactivity and exercise. *Ger J Exerc Sport Res, 48*, 48–58 (2018). https://doi.org/10.1007/s12662-017-0477-9

Cheval, B., & Boisgontier, M. P. (2021). The Theory of Effort Minimization in Physical Activity. *Exercise and Sport Sciences Reviews*, *49*(3), 168–178. https://doi.org/10.1249/JES.0000000000000252

Cheval, B., & Boisgontier, M. P. (2024). Promouvoir une activité physique régulière chez les patients : l'importance de la perception de l'effort. *STAPS*. https://doi.org/10.3917/sta.pr1.0091

Cheval, B., Darrous, L., Choi, K. W., Klimentidis, Y. C., Raichlen, D. A., Alexander, G. E., Cullati, S., Kutalik, Z., & Boisgontier, M. P. (2023). Genetic insights into the causal relationship between physical activity and cognitive functioning. *Scientific Reports*, *13*(1), 5310. https://doi.org/10.1038/s41598-023-32150-1

Cheval, B., Maltagliati, S., Sieber, S., Beran, D., Chalabaev, A., Sander, D., Cullati, S., & Boisgontier, M. P. (2021). Why are individuals with diabetes less active? The mediating role of physical, emotional, and cognitive factors. *Annals of Behavioral Medicine*, *55*(9), 904–917. https://doi.org/10.1093/abm/kaaa120

Cheval, B., Radel, R., Neva, J. L., Boyd, L. A., Swinnen, S. P., Sander, D., & Boisgontier, M. P. (2018). Behavioral and neural evidence of the rewarding value of exercise behaviors: a systematic review. *Sports Medicine*, *48*(6), 1389–1404. https://doi.org/10.1007/s40279-018-0898-0

Cheval, B., Tipura, E., Burra, N., Frossard, J., Chanal, J., Orsholits, D., Radel, R., & Boisgontier, M. P. (2018). Avoiding sedentary behaviors requires more cortical resources than avoiding physical activity: an EEG study. *Neuropsychologia*, *119*, 68–80. https://doi.org/10.1016/j.neuropsychologia.2018.07.029

Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Routledge. https://doi.org/10.4324/9780203771587

Conroy, D. E., & Berry, T. R. (2017). Automatic affective evaluations of physical activity. *Exercise and Sport Sciences Reviews*, *45*(4), 230–237. https://doi.org/10.1249/JES.0000000000000120

Craig, C. L., Marshall, A. L., Sjöström, M., Bauman, A. E., Booth, M. L., Ainsworth, B. E., Pratt, M., Ekelund, U., Yngve, A., Sallis, J. F., & Oja, P. (2003). International Physical Activity Questionnaire: 12-country reliability and validity. *Medicine and Science in Sports and Exercise*, *35*(8), 1381–1395. https://doi.org/10.1249/01.MSS.0000078924.61453.FB

den Brok, M. G., van Dalen, J. W., van Gool, W. A., Moll van Charante, E. P., de Bie, R. M., & Richard, E. (2015). Apathy in Parkinson's disease: a systematic review and meta-analysis. *Movement Disorders*, *30*(6), 759–769. https://doi.org/10.1002/mds.26208

Derosiere, G., Vassiliadis, P., Dricot, L., Dessain, Q., Delinte, N., Zénon, A., & Duque, J. (2024). Fronto-motor circuits linked to subclinical apathy. BioRxiv, 10.07.616861. https://doi.org/10.1101/2024.10.07.616861

Eagly, A. H., & Chaiken, S. (1993). *The psychology of attitudes*. Orlando, FL: Harcourt Brace Jovanovich College Publishers.

Elliot, A. J., Eder, A. B., & Harmon-Jones, E. (2013). Approach–avoidance motivation and emotion: convergence and divergence. *Emotion Review*, 5(3), 308–311. https://doi.org/10.1177/1754073913477517

Ekkekakis, P. (2017). People have feelings! Exercise psychology in paradigmatic transition. *Current Opinion in Psychology*, 16, 84–88. https://doi.org/10.1016/j.copsyc.2017.03.018

Farajzadeh, A., Goubran, M., Beehler, A., Cherkaoui, N., Morrison, P., de Chanaleilles, M., Maltagliati, S., Cheval, B., Miller, M. W., Sheehy, L., Bilodeau, M., Orsholits, D., & Boisgontier, M. P. (2023). Automatic approach-avoidance tendency toward physical activity, sedentary, and neutral stimuli as a function of age, explicit affective attitude, and intention to be active. *Peer Community Journal*, 3, e21. https://doi.org/10.24072/pcjournal.246

Farajzadeh, A., Hébert, A., Lahart, I. M., Bilodeau, M., & Boisgontier, M. P. (2024). Apathy and physical activity: a systematic review and meta-analysis. *MedRxiv*. https://doi.org/10.1101/2024.05.01.24306712

Farajzadeh, A., Orsholits, D., & Boisgontier, M. P. (2022). Automatic approach-avoidance tendency toward physical activity, sedentary, and neutral stimuli as a function of age, explicit affective attitude, and intention to be active: dataset, R scripts, and supplemental material (v1.0.1). https://doi.org/10.5281/zenodo.7051414

Faul, F., Erdfelder, E., Buchner, A., & Lang, A. G. (2009). Statistical power analyses using G*Power 3.1: tests for correlation and regression analyses. *Behavior Research Methods*, *41*(4), 1149–1160. https://doi.org/10.3758/BRM.41.4.1149

Feil, K., Fritsch, J., & Rhodes, R. E. (2023). The intention-behaviour gap in physical activity: a systematic review and meta-analysis of the action control framework. *British Journal of Sports Medicine*, *57*(19), 1265–1271. https://doi.org/10.1136/bjsports-2022-106640

Fozard, J. L., Vercryssen, M., Reynolds, S. L., Hancock, P. A., & Quilter, R. E. (1994). Age differences and changes in reaction time: the Baltimore Longitudinal Study of Aging. *Journal of Gerontology*, 49(4), 179–189. https://doi.org/10.1093/geronj/49.4.p179

Friese, M., Hofmann, W., & Wiers, R. W. (2011). On taming horses and strengthening riders: recent developments in research on interventions to improve self-control in health behaviors. Self and Identity, 10(3), 336–351. https://doi.org/10.1080/15298868.2010.536417

Gollwitzer, P. M., & Brandstätter, V. (1997). Implementation intentions and effective goal pursuit. *Journal of Personality and Social Psychology*, 73(1), 186–199. https://doi.org/10.1037/0022-3514.73.1.186

Greenwald, A. G., & Banaji, M. R. (1995). Implicit social cognition: attitudes, self-esteem, and stereotypes. *Psychological Review*, *102*(1), 4–27. https://doi.org/10.1037/0033-295x.102.1.4

Groeneweg-Koolhoven, I., de Waal, M. W., van der Weele, G. M., Gussekloo, J., & van der Mast, R. C. (2014). Quality of life in community-dwelling older persons with apathy. *American Journal of Geriatric Psychiatry*, *22*(2), 186–194. https://doi.org/10.1016/j.jagp.2012.10.024

Hagger, M. S., & Chatzisarantis, N. L. (2014). An integrated behavior change model for physical activity. *Exercise and Sport Sciences Reviews*, *42*(2), 62–69. https://doi.org/10.1249/JES.0000000000000008

Hayes, A. F. (2022). *Introduction to mediation, moderation, and conditional process analysis: a regression-based approach* (3rd ed.). Guilford Press.

Hick, W. E. (1952). On the rate of gain of information. *Quarterly Journal of Experimental Psychology*, 4(1), 11–26. https://doi.org/10.1080/17470215208416600

Hoyt, A. L., Rhodes, R. E., Hausenblas, H. A., & Giacobbi, P. R. (2009). Integrating five-factor model facet-level traits with the theory of planned behavior and exercise. *Psychology of Sport and Exercise*, *10*(5), 565–572. https://doi.org/10.1016/j.psychsport.2009.02.008

Kant, R., Duffy, J. D., & Pivovarnik, A. (1998). Prevalence of apathy following head injury. *Brain Injury*, *12*(1), 87–92. https://doi.org/10.1080/026990598122908

Kent, M. (2006). Oxford dictionary of sports science and medicine. *Oxford University Press*. https://doi.org/10.1093/acref/9780198568506.001.0001

Kruse, C., Maier, F., Spottke, A., Bach, J. P., Bartels, C., Buerger, K., Fellgiebel, A., Fliessbach, K., Frölich, L., Hausner, L., Hellmich, M., Klöppel, S., Klostermann, A., Kornhuber, J., Laske, C., Peters, O., Priller, J., Richter-Schmidinger, T., Schneider, A., Shah-Hosseini, K., …, & Jessen, F. (2023). Apathy in patients with Alzheimer's disease is a cost-driving factor. *Alzheimer's & Dementia*, *19*(7), 2853–2864. https://doi.org/10.1002/alz.12915

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2022). lmerTest: tests in linear mixed effects models. Version 3.1-3. [R package]. https://cran.r-project.org/web/packages/lmerTest/lmerTest.pdf

Kwan, B. M., & Bryan, A. (2010). In-task and post-task affective response to exercise: translating exercise intentions into behaviour. *British Journal of Health Psychology*, 15(Pt 1), 115–131. https://doi.org/10.1348/135910709X433267

Lakens, D. (2022). Sample size justification. *Collabra: Psychology*, 8(1), 33267. https://doi.org/10.1525/collabra.33267

Le Heron, C., Apps, M. A. J., & Husain, M. (2018). The anatomy of apathy: a neurocognitive framework for amotivated behaviour. *Neuropsychologia*, *118*(Pt B), 54–67. https://doi.org/10.1016/j.neuropsychologia.2017.07.003

Le Heron, C., Holroyd, C. B., Salamone, J., & Husain, M. (2019). Brain mechanisms underlying apathy. *Journal of Neurology, Neurosurgery, and Psychiatry*, *90*(3), 302–312. https://doi.org/10.1136/jnnp-2018-318265

Leung, D. K. Y., Chan, W. C., Spector, A., & Wong, G. H. Y. (2021). Prevalence of depression, anxiety, and apathy symptoms across dementia stages: a systematic review and meta-analysis. *International Journal of Geriatric Psychiatry*, *36*(9), 1330–1344. https://doi.org/10.1002/gps.5556

Levy, R., & Dubois, B. (2006). Apathy and the functional anatomy of the prefrontal cortex-basal ganglia circuits. *Cerebral Cortex*, *16*(7), 916–928. https://doi.org/10.1093/cercor/bhj043

Liu, X., Zhang, D., Liu, Y., Sun, X., Han, C., Wang, B., Ren, Y., Zhou, J., Zhao, Y., Shi, Y., Hu, D., & Zhang, M. (2017). Dose-response association between physical activity and incident hypertension: a systematic review and meta-analysis of cohort studies. *Hypertension*, *69*(5), 813–820. https://doi.org/10.1161/HYPERTENSIONAHA.116.08994

Lohse, K., Kozlowski, A., & Strube, M. J. (2023). Model specification in mixed-effects models: a focus on random effects. *Communications in Kinesiology*, *1*(5), 1–28. https://doi.org/10.51224/cik.2023.52

Lovibond, S. H., & Lovibond, P. F. (1995). Depression Anxiety Stress Scales (DASS--21, DASS--42). *APA PsycTests*. [Database record] https://doi.org/10.1037/t01004-000

Luke, S. G. (2017). Evaluating significance in linear mixed-effects models in R. *Behavior Research Methods*, 49, 1494–1502. https://doi.org/10.3758/s13428-016-0809-y

Maltagliati, S., Raichlen, D., Rhodes, R., & Cheval, B. (2024a). Closing the intention-behavior gap in physical activity: the moderating effect of individual differences in the valuation of physical effort. *SportRxiv*. https://doi.org/10.51224/SRXIV.375

Maltagliati, S., Sarrazin, P., Fessler, L., Lebreton, M., & Cheval, B. (2024b). Why people should run after positive affective experiences instead of health benefits. *Journal of Sport and Health Science*, 13(4), 445–450. https://doi.org/10.1016/j.jshs.2022.10.005

Maltagliati, S., ten Broeke, P., Sarrazin, P., Rhodes, R., Bijleveld, E. H., Dunton, G. F., Bonnet, C., Gardner, B., Raichlen, D., & Cheval, B. (2024c). Uncovering the role of effort in the rewarding value of sedentary behavior. *PsyArXiv*. https://doi.org/10.31234/osf.io/g8mxw

Marcora, S. (2009). Perception of effort during exercise is independent of afferent feedback from skeletal muscles, heart, and lungs. *Journal of Applied Physiology*, 106 (6), 2060–2062. https://doi.org/10.1152/japplphysiol.90378.2008

Marin R. S. (1990). Differential diagnosis and classification of apathy. *The American Journal of Psychiatry*, *147*(1), 22–30. https://doi.org/10.1176/ajp.147.1.22

Marin R. S. (1991). Apathy: a neuropsychiatric syndrome. *The Journal of Neuropsychiatry and Clinical Neurosciences*, *3*(3), 243–254. https://doi.org/10.1176/jnp.3.3.243

Martínez-Horta, S., Riba, J., de Bobadilla, R. F., Pagonabarraga, J., Pascual-Sedano, B., Antonijoan, R. M., Romero, S., Mañanas, M. À., García-Sanchez, C., & Kulisevsky, J. (2014). Apathy in Parkinson's disease: neurophysiological evidence of impaired incentive processing. *Journal of Neuroscience*, *34*(17), 5918–5926. https://doi.org/10.1523/JNEUROSCI.0251-14.2014

Mega, M. S., Cummings, J. L., Fiorello, T., & Gornbein, J. (1996). The spectrum of behavioral changes in Alzheimer's disease. *Neurology*, *46*(1), 130–135. https://doi.org/10.1212/wnl.46.1.130

Melnikoff, D. E., & Bargh, J. A. (2018). The mythical number two. *Trends in Cognitive Sciences*, 22(4), 280–293. https://doi.org/10.1016/j.tics.2018.02.001

Miller, J. (2023). Outlier exclusion procedures for reaction time analysis: The cures are generally worse than the disease. *Journal of Experimental Psychology: General*, *152*(11), 3189–3217. https://doi.org/10.1037/xge0001450

Moore, S. C., Lee, I. M., Weiderpass, E., Campbell, P. T., Sampson, J. N., Kitahara, C. M., Keadle, S. K., Arem, H., Berrington de Gonzalez, A., Hartge, P., Adami, H. O., Blair, C. K., Borch, K. B., Boyd, E., Check, D. P., Fournier, A., Freedman, N. D., Gunter, M., Johannson, M., Khaw, K. T., … Patel, A. V. (2016). Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. *JAMA Internal Medicine*, *176*(6), 816–825. https://doi.org/10.1001/jamainternmed.2016.1548

Muhammed, K., Manohar, S., Ben Yehuda, M., Chong, T. T., Tofaris, G., Lennox, G., Bogdanovic, M., Hu, M., & Husain, M. (2016). Reward sensitivity deficits modulated by dopamine are associated with apathy in Parkinson's disease. *Brain*, *139*(Pt 10), 2706–2721. https://doi.org/10.1093/brain/aww188

Pessiglione, M., Vinckier, F., Bouret, S., Daunizeau, J., & Le Bouc, R. (2018). Why not try harder? Computational approach to motivation deficits in neuro-psychiatric diseases. *Brain,* 141(3), 629–650. https://doi.org/10.1093/brain/awx278

Qiu, W. (2022). powerMediation: Power/sample size calculation for mediation analysis (version 0.3.4). [R package]. https://cran.r-project.org/web/packages/powerMediation/powerMediation.pdf

R Core Team (2023) *R:* A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria (Version 4.3.1). [Computer Software] https://www.R-project.org/

Rhodes, R. E., & de Bruijn, G. J. (2013). How big is the physical activity intention-behaviour gap? A meta-analysis using the action control framework. *British Journal of Health Psychology*, *18*(2), 296–309. https://doi.org/10.1111/bjhp.12032

Rhodes, R. E., McEwan, D., & Rebar, A. L. (2019). Theories of physical activity behaviour change: a history and synthesis of approaches. *Psychology of Sport & Exercise*, *42*, 100–109. https://doi.org/10.1016/j.psychsport.2018.11.010

Rhodes, R. E., & Rebar, A. L. (2017). Conceptualizing and defining the intention construct for future physical activity research. *Exercise and Sport Sciences Reviews*, *45*(4), 209–216. https://doi.org/10.1249/JES.0000000000000127

Rydell, R. J., & McConnell, A. R. (2006). Understanding implicit and explicit attitude change: a systems of reasoning analysis. *Journal of Personality and Social Psychology*, *91*(6), 995–1008. https://doi.org/10.1037/0022-3514.91.6.995

Sander, D., Grandjean, D., & Scherer, K. R. (2018). An appraisal-driven componential approach to the emotional brain. *Emotion Review*, 10(3), 219–231. https://doi.org/10.1177/1754073918765653

Sockeel, P., Dujardin, K., Devos, D., Denève, C., Destée, A., & Defebvre, L. (2006). The Lille Apathy Rating Scale (LARS), a new instrument for detecting and quantifying apathy: validation in Parkinson's disease. *Journal of Neurology, Neurosurgery, and Psychiatry*, *77*(5), 579–584. https://doi.org/10.1136/jnnp.2005.075929

Starkstein, S. E., Petracca, G., Chemerinski, E., & Kremer, J. (2001). Syndromic validity of apathy in Alzheimer's disease. *American Journal of Psychiatry*, *158*(6), 872–877. https://doi.org/10.1176/appi.ajp.158.6.872

Stevens, C. J., Baldwin, A. S., Bryan, A. D., Conner, M., Rhodes, R. E., & Williams, D. M. (2020). Affective determinants of physical activity: a conceptual framework and narrative review. *Frontiers in Psychology*, *11*, 568331. https://doi.org/10.3389/fpsyg.2020.568331

Strack, F., & Deutsch, R. (2004). Reflective and impulsive determinants of social behavior. *Personality and Social Psychology Review, 8*(3), 220–247. https://doi.org/10.1207/s15327957pspr0803_1

Ulrich, R., & Miller, J. O. (1994). Effects of truncation on reaction time analysis. *Journal of Experimental Psychology: General*, 123(1), 34–80. https://doi.org/10.1037/0096-3445.123.1.34

van Allen, Z., Dionne, N., & Boisgontier, M. P. (2024a). Prospective classification of functional dependence: insights from machine learning and 39,927 participants in the Canadian Longitudinal Study on Aging. *MedRxiv*. https://doi.org/10.1101/2024.07.15.24310429

van Allen, Z., Orsholits, D., & Boisgontier, M. P. (2024b). Prestroke physical activity matters for functional limitations: a longitudinal case-control study of 12,860 participants. *Physical Therapy*, 104(10), pzae094. https://doi.org/10.1093/ptj/pzae094

Vancampfort, D., Koyanagi, A., Ward, P. B., Rosenbaum, S., Schuch, F. B., Mugisha, J., Richards, J., Firth, J., & Stubbs, B. (2017). Chronic physical conditions, multimorbidity and physical activity across 46 low- and middle-income countries. *International Journal of Behavioral Nutrition and Physical Activity*, *14*(1), 6. https://doi.org/10.1186/s12966-017-0463-5

Vilalta-Franch, J., Calvó-Perxas, L., Garre-Olmo, J., Turró-Garriga, O., & López-Pousa, S. (2013). Apathy syndrome in Alzheimer's disease epidemiology: prevalence, incidence, persistence, and risk and mortality factors. *Journal of Alzheimer's Disease*, *33*(2), 535–543. https://doi.org/10.3233/JAD-2012-120913

Wahid, A., Manek, N., Nichols, M., Kelly, P., Foster, C., Webster, P., Kaur, A., Friedemann Smith, C., Wilkins, E., Rayner, M., Roberts, N., & Scarborough, P. (2016). Quantifying the Association Between Physical Activity and Cardiovascular Disease and Diabetes: A Systematic Review and Meta-Analysis. *Journal of the American Heart Association*, *5*(9), e002495. https://doi.org/10.1161/JAHA.115.002495

Williams, D. M., & Evans, D. R. (2014). Current Emotion Research in Health Behavior Science. *Emotion Review*, 6(3), 277–287. https://doi.org/10.1177/1754073914523052

Yzerbyt, V., Muller, D., Batailler, C., & Judd, C. M. (2018). New recommendations for testing indirect effects in mediational models: the need to report and test component paths. *Journal of Personality and Social Psychology*, *115*(6), 929–943. https://doi.org/10.1037/pspa0000132

Zenko, Z., & Ekkekakis, P. (2019). Internal consistency and validity of measures of automatic exercise associations. *Psychology of Sport and Exercise*, 43, 4–15. https://doi.org/10.1016/j.psychsport.2018.12.005

Zhang, H., Feng, Y., Lv, H., Tang, S., & Peng, Y. (2023). The prevalence of apathy in stroke patients: A systematic review and meta-analysis. *Journal of Psychosomatic Research*, *173*, 111478. https://doi.org/10.1016/j.jpsychores.2023.111478