Analysis of Genome-Wide Cell-Free DNA Fragment Length Distributions in Colorectal Cancer

Alexandre Matov[1, 2, †]

1   Department of Clinical Medicine, Aarhus University Hospital, Århus, Denmark

2   Present address: DataSet Analysis LLC, 155 Jackson St, San Francisco, CA 94111

†   Corresponding author:

email: matov@datasetanalysis.com

Key words: Patient Stratification, Disease Classification, Cell-Free DNA, Fragmentation Patterns, Low-Coverage Whole Genome Sequencing, Kullback-Leibler Divergence, Broadcast Channel Capacity, Degraded Broadcast Channel, Achievable Rate Regions, Directed Markov Chain with Absorbing States

ABSTRACT

Introduction: Each piece of cell-free DNA (cfDNA) has a length determined by the exact metabolic conditions in the cell it belonged to at the time of cell death. The changes in cellular regulation leading to a variety of patterns, which are based on the different number of fragments with lengths up to several hundred base pairs (bp) at each of the almost three billion genomic positions, allow for the detection of disease and also the precise identification of the tissue of their origin.

Methods: A Kullback-Leibler (KL) divergence computation identifies different fragment lengths and areas of the human genome, depending on the stage, for which disease samples, starting from pre-clinical disease stages, diverge from healthy donor samples. We provide examples of genes related to colorectal cancer (CRC), which our algorithm detected to belong to divergent genomic bins. The staging of CRC can be viewed as a Markov chain and that provides a framework for studying disease progression and the types of epigenetic changes occurring longitudinally at each stage, which might aid the correct classification of a new hospital sample.

Results: In a new look to treat such data as grayscale value images, pattern recognition using artificial intelligence could be one approach to classification. In CRC, Stage I disease does not, for the most part, shed any tumor in circulation, making detection difficult for established machine learning (ML) methods. This leads to the deduction that early detection, where we can only rely on changes in the metabolic patterns, can be accomplished when the information is considered in its entirety, for example by applying computer vision methods.

Conclusions: Longitudinal analysis of patient's genetic datasets can detect the early stages of neoplasm better than population-based methods.

INTRODUCTION

Colorectal cancer (CRC) is the second most common cancer-related cause of death worldwide (Siegel et al., 2020). Approximately two-thirds of newly-diagnosed patients present with a curable disease (Petrelli et al., 2017), but despite curatively intended treatment, up to 40% of them experience relapsed disease after an initial treatment (Jorgensen et al., 2015). In addition, about 86% of Stage I disease do not shed tumor in circulation, which leads to a decreased ability for early detection (Liu et al., 2020). A minimally-invasive analysis based on circulating cfDNA has emerged as a promising nucleic acids biomarker (Cristiano et al., 2019), but this method assumes very homogenous characteristics of the healthy population, which makes it hard to gain regulatory approval. Here, we attempt to consider the processes underlying the evolution of CRC as an electronics communication system and propose to visualize the cfDNA samples as images (Matov, 2024b).

Many cancer drugs target the cellular cytoskeleton, yet examining the cytoskeleton of patient cells prior to making treatment decisions is not incorporated in the clinical practice. We propose the utilization of quantitative microscopy methods for the *ex vivo* evaluation of drug regimens. This can be accomplished by culturing patient-derived organoids from biopsy (tissue or liquid) or resection samples, followed by fluorescent labeling of the desired molecular markers. We have provided examples (Matov and Bacconi, 2024) based on techniques, such as fluorescent speckle microscopy (FSM) (Waterman-Storer et al., 1998), in which a very low labeling ratio of tubulin dimers allows to probe the molecular machinery of the mitotic spindle by computational analysis of faint, stochastic features forming on bundles of often times interconnected microtubules (MTs). In FMS images, four to seven fluorophores cluster within the diffraction limit of the imaging system to form a visible, above background noise, speckle. Each of the fluorophores might be incorporated in the lattice of different MTs, which could be a part of two or more MT bundles, facing different directions within the cell, and form a speckle for several seconds. In the

mitotic spindle, MT plus tips are the ends where MTs add dimers and which undergo a specialized mechanism termed 'dynamic instability' (Mitchison and Kirschner, 1984). This stochastic process of reorganization of the MT cytoskeleton during cell division has been targeted in cancer, yet techniques that measure the precise changes in MT dynamics to result from drug treatment are absent in the clinic.

Aneuploidy likely leads to cancer (Boveri, 1902) and the mutations traditionally considered to be the drivers of neoplasms might be the result of dysregulation in MT regulators. All genes that regulate the MT cytoskeleton have been implicated in conferring drug resistance, in particular to tubulin inhibitors – but not exclusively. In addition, the same MT-regulating genes appear as biomarkers of pathogenesis. When the MT network is dysregulated that leads to inability of the mitotic spindle to properly segregate the duplicated chromosomes and aneuploidy. The consequent proliferation of aneuploid cells with mutant MT regulators cause mutations in oncogenes and tumor suppressors, which causes cancer.

RESULTS

COLORECTAL CANCER AS AN IMAGE

In CRC, patients go through a complex diagnostic paradigm. While healthy, the donors could be characterized as having different stages of co-morbidities. Next, precancerous polyps might lead to adenomas of the colon or the rectum, which also have different stages. About 45% of the patients with advanced adenoma, and oftentimes people with low-grade adenoma or no adenoma at all, develop colorectal neoplasm, which has four stages. The extensive length of the colon makes tissue biopsies and surgical interventions uncertain procedures because of the sheer length of the organ. The physiology of the colon is particular and it is conceivable that changes, reported by the changes in the fragmentation patterns, in the local cellular regulation could offer information on the exact location of early disease. In disease, the variety of cell death fragmentation patterns reflects differential nucleosome packaging,

chromatin remodeling and accessibility (van der Pol and Mouliere, 2019). DNA hypomethylation or the loss of a histone in a nucleosome, for instance, lead to a skewed fragment length distribution (Fig. 1A and Fig. 1B). In contrast, in healthy donor samples (Fig. 1C) the distribution is symmetric, centered around 168 bp. One can observe the consistent number of fragments across the genome for the healthy donor sample (Fig. 1C), which is in stark contrast to the copy-number variation seen as bright horizontal streaks in the two Stage IV samples (Fig. 1A and Fig. 1B). The difference image between a healthy donor sample and a Stage I sample highlights the differences (Fig. 1D) present in cellular regulation and cell death in CRC Stage I (Fig. 1E). This holds true for low- and high-grade adenoma too.

DNA FRAGMENTATION

Cell-free DNA (cfDNA) in blood samples provides a minimally-invasive diagnostic approach for early detection of cancer and other degenerative diseases. A recently developed DELFI (DNA evaluation of fragments for early interception) method for the detection of abnormalities in cfDNA through genome-wide analysis of fragmentation patterns is based on low coverage WGS of isolated cfDNA. Mapped sequences are analyzed in windows covering the whole genome and the windows may range in size from thousands to millions of bases. DELFI uses 5 Mb windows for evaluating cfDNA fragmentation patterns, which provides over 20,000 reads for each window at 1–2x genome coverage. The genome-wide pattern from an individual is compared to a reference (healthy) population utilizing a stochastic gradient boosting model to detect cancers. The classification is based on DNA fragment length ratios between short (100 to 150 bp) and long (151 to 220 bp) cfDNA fragments (Cristiano et al., 2019).

Figure 1. DNA fragmentation patterns in whole genome sequencing datasets. For advanced disease, the distributions are very asymmetric and the histograms are negatively skewed. (A) CRC Stage IV. (B) CRC Stage IV (another patient). (C)

Healthy donor sample, no co-morbidities. (D) A difference image between the images shown on (C) and (E). (E) CRC Stage I. The genomic bins (of 5 Mb) are on the Y-axis. Chromosome 1 is at the top of the image. The DNA fragment length is on the X-axis, from left to right. Pixels with brighter intensity correspond to bins with a higher number of fragments. The peak in intensity is for each sample the vertical streak at 168 bp.

Another method for classifying samples is to compute the relative entropy for each fragmentation length by building a probability density function based on the fragments from all genomic positions. Such distributions consist of about 40 million fragments for each whole genome sequencing sample, which when binned in genomic regions of 5 Mb, for each fragment length, allow us compare histograms of about 45,000 values from each region of the 23 chromosomes. A Kullback-Leibler (KL) divergence computation based on an adaptive minimax rate-optimal estimator (Han et al., 2016) of the changes in disease from healthy state(s) to precancerous lesion(s) to malignant tumor(s) can be presented as a heterogeneous directed Markov chain with absorbing states (Bremaud, 1999). Considering this Markov chain as a degraded broadcast channel, considerations regarding the capacity region (Han, 1981) and its estimation (Bergmans, 1973) for a cohort of CRC samples could aid the classification of a new patient sample in the clinic. Our analysis shows that there are two peaks in the KL divergence at 198 bp and 364 bp. The two main divergence peaks have different healthy-to-CRC number of fragment ratios (Fig. 2), with the ratio being higher for DNA fragment length 364 bp, while the overall number of fragments was significantly higher for length 198 bp (2.5 million) (Fig. S1).

CANCER AS COMMUNICATION ERRORS

To build the boundary of the capacity region for each CRC stage, we assumed that the two peaks in the DNA fragmentation length KL divergence histogram provide the (x, y) coordinates for which the boundary intersects with the x-axis (divergence value for the first peak, 0) and the y-axis (0, divergence value for second peak). As different boundaries of the capacity regions (Cover, 1998) are defined by the

probability density function of the cohort of each disease stage, a new sample will be classified according to the boundary it falls within.

Figure 2. DNA fragmentation length KL divergence (in bits) of a healthy individuals' cohort from a CRC patient cohort. The ratio of number of fragments healthy-to-CRC for each length is color coded. The CRC cohorts consist of 27 patients in Stages I-IV. The healthy individuals' cohort consists of 43 samples. The KL divergence (in bits) is on the Y axis. The maximal value is 2.8 bits divergence (ratio 3.6) for DNA fragment length 364 bp. The second highest divergence value is 2.6 bits (ratio 3.0) for DNA fragment length 198 bp. The fragment length (in bp) is on the X axis, from left to right. The two main divergence peaks have different healthy-to-CRC fragment ratios.

A new sample for an already existing patient which has become an outlier for the disease stage it has been assigned, and falls outside the boundary (Bergmans, 1974), will be classified in the next (or more advanced) disease stage. Oppositely, after a surgical resection or drug treatment, when tumors are removed or recede, and a new sample is collected and classified after an intervention, if it falls within an earlier disease stage boundary of the capacity region, it will be re-classified as a lower burden disease or an earlier disease stage, according to the boundary it falls within.

The KL divergences from different cancer cohorts of healthy donor samples exhibit distributions, which are multi-modal, with different peaks being present for different cancers, defining potentially unique signatures. The two highest peaks on Fig. 3, for 364 base bp and 198 bp, are the result of significant differences in the median number of fragments in the genomic bins for these fragment lengths. While for healthy samples we measured that most genomic bins consist of about 20 fragments with length 364 bp (see the peak in the green histogram on Fig. 3), the CRC samples exhibit a very different distribution in which most genomic bins consist of less than 10 fragments with length 364 bp (see the peak in the pink histogram on Fig. 3).

The distinct per-disease peaks in the divergence from healthy samples (Tab. 1, col. 2) and the divergence of between-cancers signatures (Tab. 1, col. 3) are the result of differential epigenetic regulation of the different cancer types and can be used as diagnostic biomarkers for the detection of disease and identification of the tissue of origin. We measured the divergence in fragment lengths also between the different stages of CRC (Stages I − IV), including between cohorts of samples of precancerous polyps. Our approach to the identification of discriminative biomarkers in disease does not require any user-defined input metrics and it is data-driven. Further divergence analysis of the genomic bins only (for the most divergent fragment lengths - the peaks in the histogram on Fig. 2) demonstrated

the ability of the method to pinpoint the areas of the human genome involved in pathogenesis and drug resistance / susceptibility.

Figure 3. Probability density functions of all genomic bins of a CRC patient cohort (in pink color) and a healthy donor cohort (in green color) for DNA fragment length 364 bp. The CRC cohorts consist of 27 patients in Stages I-IV. The healthy cohort consists of 43 donor samples. The frequency, or the number of genomic bins, is on the Y axis. The number of fragments in each bin, or per bin, is on the X axis. Observe that the healthy samples have roughly 2.2-fold more fragments per bin for DNA fragment length 364 bp. These two histograms, and the corresponding differences in disease, for DNA fragment length 198 bp are very similar. Inset: Distribution of the number of fragments per DNA fragment length for the healthy (green) and CRC (red) cohorts; each cohort comprises over 1 billion fragments. The peaks in the distribution for healthy individuals correspond to mono- (see inset cartoon), di- (see inset cartoon), and tri- (not shown) nucleosome protected DNA. In CRC, there is an additional peak for DNA fragment lengths 137 pb – 143 pb, like due to active section due to metastases (see inset cartoon of healthy (green) and shorter CRC (red) fragments).

Cancer cohort	Healthy cohort	Colorectal cancer cohort
Colorectal cancer	364, 205	-
Ovarian cancer	359, 208	248, 175
Pancreatic cancer	203, 359	111, 269
Gastric cancer	122, 347	193, 138
Breast cancer	177, 333	176, 280
Bile duct cancer	200, 351	111, 164
Lung cancer	193, 343	121, 198

Table 1. Fragment length of maximal KL divergence from seven cancers. DNA fragment lengths (in bp) for the two highest peaks (the primary peak is listed first and the secondary is listed second) of the KL divergence from seven cancers (see the list in the left column) of healthy donor samples (see the peak fragment lengths in the middle column) and of CRC (see the peak fragment lengths in the right column). We measured the divergence in fragment lengths also between the different stages of CRC (Stages I – IV), including between cohorts of samples of precancerous polyps. At least 8% of the fragments in each cohort belong to diverging populations of bins.

We analyzed four adenoma cohorts and identified distinct divergence patterns, which were different from the peaks in CRC (Stages I – IV). The maximal KL divergence we measured for low and high grade colon adenoma as well as low and high grade rectal adenoma cohorts were less than 1 bit (Fig. 4). Interestingly, the DNA fragment lengths of maximum divergence did not coincide with those measured in CRC. In high grade colon adenoma tumors, the maximal KL divergence was measured for DNA fragment length 144 bp, while in low grade colon adenoma tumors, for 186 bp. In high grade rectal adenoma tumors, the maximal KL divergence was measured for DNA fragment length 200 bp, while in low grade rectal adenoma tumors, for 173 bp.

Figure 4. DNA fragmentation length KL divergence (in bits) of a healthy individuals' cohort from four adenoma patient cohorts. The healthy individuals' cohort consists of 74 samples from healthy donors with no co-morbidities. The KL divergence (in bits) is on the Y axis. The fragment length (in bp) is on the X axis, from left to right. (A) The high grade colon adenoma cohort (in red) consists of 20 patients. The maximal value is 0.3 bits divergence for DNA fragment length 144 bp. The low grade colon adenoma cohort (in green) consists of 28 patients. The maximal value is 0.4 bits divergence for DNA fragment length 186 bp. (B) The high grade rectal adenoma cohort (in yellow) consists of 11 patients. The maximal value is 0.7 bits divergence for DNA fragment length 200 bp. The low grade rectal adenoma cohort (in blue) consists of 7 patients. The maximal value is 0.5 bits divergence for DNA fragment length 173 bp.

HIERARCHICAL CLUSTERING

For the most divergence DNA fragment length in CRC of 364 bp, we performed KL divergence analysis

of the genomic bins, which identified 12 bins above 1.5 bits divergence (Fig. S2). The most divergent

genomic bin include genes that are associated with allelic losses from the short arm of chromosome 8

(Takanishi et al., 1997). One of the CRC-related genes, *AXIN2*, mutations in which have been associated

with mismatch repair errors (Liu et al., 2000), falls within the second most divergent genomic bin.

Frequent loss of tumor-suppressors, such as Axin2, a Wnt suppressor (Wu et al., 2012), promotes CRC.

Axin2 has been shown to interact with glycogen synthase kinase-3β (Lejeune et al., 2006), which

regulates MTs in migrating cells (Kumar et al., 2009). Interestingly, two other genomic bins that are

very divergent from the healthy cohort in CRC are those containing *RAP2B* (Yi et al., 2019) and *GPX3*

(Zhang et al., 2020). The second gene is involved in the detoxification (reduction by GPX3, produced

mainly in the kidneys (Avissar et al., 1994)) of soluble reactive oxygen species (Dixon and Stockwell,

2014). This new result indicates the presence of a divergence in the metabolic / redox patterns in CRC.

Further, hierarchical clustering based on genomic bins of patient and healthy samples for fragments with length 364 bp resulted in three false positive selections but, importantly for the detection of sub-clinical disease, no false negative selections (Fig. 5). Clustering of three healthy samples with the CRC cohort suggested that depletion of fragments from di-nucleosome-protected DNA in genomic regions associated with disabled antioxidant program (Barrett et al., 2013) in samples from healthy donors might be indicating pathogenesis and early CRC (Fig. 4). This poses the question whether the three healthy donors from the DELFI (Cristiano et al., 2019) study associated in our analysis with CRC have since the time of blood draw been diagnosed with CRC.

DISEASE PROGNOSTICS

It is conceivable that predictions regarding disease progression can be achieved using generative methods. Generative methods that produce novel samples from high-dimensional data distributions are finding widespread use, for example in speech synthesis. Generative adversarial network (GAN) models (Karras et al., 2017) consist of two CNNs, style-based generator and discriminator, which converge upon reaching Nash equilibrium and can be used to generate synthetic samples. Such sampler posterior distributions can be then used to reduce the complexity and improve the numerical convergence of predicting disease progression for any new sample using the Bayes formula, given there is available longitudinal data and transitional states for several patients and healthy donors. This approach may allow attempting to induce changes in reversal (Mullis, 1968) to the physiological deterioration occurring in disease, if detected early in a pre-clinical stage.

The clustering algorithm separates the samples into healthy donors (upper half of Fig. 5) and patient samples (lower half of Fig. 5). Remarkably, there are no patient samples grouped with the healthy donors, which indicates no false positive selections. There are, however, three false negative selections

(labeled with FN in red color on Fig. 5) within the patient cluster (PGDX labels on Fig. 4). This result is promising in the context of the detection of sub-clinical and early disease; it could be further verified whether these three donors have developed CRC since the time of the blood draw.

Figure 4. Hierarchical clustering of CRC patients and healthy individuals. Hierarchical clustering of a cohort of 27 CRC patients, Stages I – IV (lower half of the image, labeled with PGDX followed by a number – some labeled are not displayed) and 43 healthy samples (upper half of the image, no label, with the exception of the three false negative, clustered with the patient samples, labeled with FN in red color) (Y axis). The clustering is accomplished based on considering the 12 (out of

595) top most divergent 5 Mb genomic bins (divergence histogram is shown in Fig. S2) for DNA fragment length 364 bp (X axis). Observe that the area of the genome containing *AXIN2*, a gene implicated in CRC, is among the most divergent.

KL DIVERGENCE OF INDIVIDUAL PROFILES

Computing KL divergence has been proposed as an approach for detecting sudden deterioration of complex diseases by dynamical network biomarkers (Chen et al., 2012), which can be accomplished with the use of one patient sample. The distribution-embedding scheme transforms the data from the observed state-variables with high level of noise to their distribution-variables with low level of noise, and thus significantly reduces signal fluctuations (Liu et al., 2015). Increasing the dimensions of the observed data by moment expansion that changes the system from state-dynamics to probability distribution-dynamics allows the derivation of new data in a high-dimensional space, but with much lower noise levels. Single-sample KL divergence has thus been proposed as an approach to detect the early-warning signal of critical transition during the progression of lung cancer (LC). The algorithm was applied to datasets of lung squamous cell carcinoma, lung adenocarcinoma, and acute lung injury (Zhong et al., 2020). As the method identifies the critical state (or tipping point) at a single sample level and identifies signaling biomarkers, it can be of great potential in personalized pre-disease diagnosis in LC. The genes it selected were involved in LC signaling and cellular processes, such as cytoskeleton organization, chromosome condensation, regulation of cell division, programmed cell death, among others (Zhong et al., 2020), demonstrating a wide clinical applicability of KL divergence calculations.

Because our results indicated a reduced ability to detect early disease in the adenoma stages, we next evaluated the KL divergence of healthy cohorts from individual (colon cancer (CC) as well as healthy) samples. Our measurements indicated high variability in the individual healthy profiles. Figure 6 shows the individual divergence for 74 healthy donors against the healthy cohort. The KL divergence reached a

peak value of over 3 bits for seven healthy donors (Fig. 6). These data indicate a high level of variability

of DNA fragmentation patterns in healthy individuals, even among those without co-morbidities or any

other findings.

Figure 6. DNA fragmentation length KL divergence (in bits) of a healthy individuals' cohort from a healthy individual. The healthy individuals' cohort consists of 74 samples from healthy donors with no co-morbidities. The KL divergence (in bits) is on the Y axis. The maximal value is 4.2 bits divergence. The profile for each healthy individual is shown in a different color. The fragment length (in bp) is on the X axis, from left to right. (A) KL divergences of samples 1 to 18. (B) KL divergence of samples 19 to 36. (C) KL divergence of samples 37 to 55. (D) KL divergence of samples 56 to 74.

We also measured the KL divergence of a healthy cohort from individual advanced CC samples. For CC Stage IV, the individual divergence profiles indicate that samples are divergent, to a different extent, at around four distinct DNA fragment lengths (Fig. 7). Besides the decrease in the apoptotic fragmentation associated with from mono-, di-nucleosome protected DNA, some (but not all) of the individual samples are very divergent in disease for lengths ~137 bp and ~277 bp. We measured a similar divergence peak at 137 bp in many healthy and Stage I – II samples, and this increase in the number of fragments is further significantly increased in Stage III – IV samples. The divergence peak at 277 bp is unique to Stage IV and such an increase in fragments is not measured in any healthy or Stage I – III samples. Overall, our analyses indicate that while CC Stage IV can be reliably detected against a cohort of healthy samples, the classification of Stages I – III is associated with many errors.

Figure 7. DNA fragmentation length KL divergence (in bits) of a healthy individuals' cohort from individual colon cancer samples. The colon cancer cohort consists of 23 patients in Stage IV. The healthy individuals' cohort consists of 74 samples from healthy donors with no co-morbidities. The KL divergence (in bits) is on the Y axis. The maximal value is 5.2 bits divergence. The profile for each colon cancer patient is shown in a different color. The fragment length (in bp) is on the X axis, from left to right. DNA fragment lengths for which there are more fragments in disease are marked with red rectangles. DNA fragment lengths for which there are less fragments in disease are marked with yellow rectangles.

BOUNDARY OF THE CAPACITY REGION

As a strategy of identifying misclassified samples, we calculated the boundaries of the capacity regions for different CRC stages. We first obtained fragment lengths 70 bp and 253 bp as the (x, y) coordinates for which the boundary intersects with the x-axis (divergence value for the first peak, 0) and the y-axis (0, divergence value for second peak). DNA fragment lengths 70 bp and 253 bp are the two local maxima of the difference vector between the integrated scores for the KL divergence of 27 patients in the CRC cohort and 43 individuals in the healthy cohort. There are different capacity regions for the different disease stages (Fig. 8) based on different channel capacity DNA fragment lengths. The boundary for CRC Stage IV exceeds 7 bits and is not shown. Two of the CRC Stage II samples fall outside the boundary region of CRC Stage II and Stage III, which suggests classification as Stage IV (see the two yellow triangles encircled with dashed purple line on Fig. 8). One of the CRC Stage I samples falls outside the boundary region of CRC Stage I and Stage II, which suggests classification as Stage III (see the blue circle encircled with dashed red line on Fig. 8). This strategy can be utilized for disease classification of low and high adenomas based on identifying outliers within the healthy samples (shown as green rhombs on Fig. 8).

FERROPTOSIS AND ANGIOGENESIS

We next sought to investigate the reasons behind the decrease of fragments for particular fragment lengths. A molecular mechanism that causes peaks at these lengths is that during apoptosis the cleavage of nuclear DNA results into fragments with length proportional to nucleosome size and resulting in patterned fragmentation. A possible mechanism leading to the differential fragmentation is the increase of ferroptotic cell death in CRC. In somatic tissues, the apoptotic cleavage of DNA results in fragments of about 195 bp in lengths and multiples thereof (Ivanov et al., 2015). Ferroptosis, however, is characterized by non-patterned DNA fragmentation and not characterized by caspase-dependent

cleavage and, thus, our analysis results demonstrate that there is a generation of less fragments with lengths 198 bp, 364 bp, and 521 bp. Our hypothesis is that the decrease of DNA fragments with lengths 198 bp, 364 bp, and to a lesser extend 521 bp (resulting in the three divergence peaks on Fig. 2) could be the result of a partial switch from apoptosis in normal physiology to ferroptosis in CRC – thus suppressing the three apoptotic peaks of fragments from mono- (147+2x26, histones plus linkers), di- (2x147+3x23, histones plus linkers), and tri- (3x147+3x27, histones plus linkers) nucleosome protected DNA (Fig. 2 and Fig. 3).

Figure 8. Boundaries of the capacity region for different CRC stages. The boundary of the capacity region for CRC Stage III is shown as a red curve. The 5 Stage III samples are shown as red crosses. The boundary region for CRC Stage II is shown as a yellow curve. The 7 Stage II samples are shown as yellow triangles. Two Stage II samples (see purple dashed circles) fall outside the boundaries for Stage II and Stage III, which suggests reclassification as Stage IV samples. The boundary region for CRC Stage I is shown as a blue curve. The 7 Stage I samples are shown as blue circles. One Stage I sample (see red dashed circle) falls outside the boundaries for Stage I and Stage II, which suggests reclassification as a Stage III sample. The 43 healthy samples are shown as green rhombs. Inset shows the difference between KL divergence for individual healthy samples and individual CRC samples for which the peak histogram values are 70 bp and 253 bp.

Ferroptosis has a dual role in cancer. It plays a role in tumor initiation, tumorigenesis, which depends on inflammation-associated immunosuppression triggered by ferroptotic damage (Chen et al., 2021) and later, during treatment, in tumor suppression (Hangauer et al., 2017). Erastin, for instance, was discovered during a synthetic lethality screen with oncogenic RAS cells (Dolma et al., 2003). It lowers cysteine and, thus, the cells stop the synthesis of antioxidants/glutathione, and this activates voltage-dependent anion channels (VDAC) by reversing tubulin's inhibition on VDAC2/3 (Yagoda et al., 2007). VDAC is a mitochondrial protein, which is a novel target for anti-cancer drugs. Our analysis shows that for the genomic bin where *VDAC2* falls has an increase in the number of fragments with length 195 bp in CC (Fig. S3), which offers a quantitative strategy for drug selection (Lemasters, 2017).

Within the fragments with length 205 bp, we also measured a divergence in CRC in the genomic bin in which falls *TCF7L2* (Fig. S4), which has been implicated in promoting migration and invasive behavior of human CRC cells (Wenzel et al., 2020). Interestingly, when we analyze the data for CRC Stage IV only (without Stages I – III), an additional peak appears on the divergence histogram from healthy donor samples at 129 bp (Fig. S5). Within the fragments with length 129 bp in Stage IV, for a few of the patients, there are about three-fold more fragments than the average in healthy samples in the genomic bin that covers the area of the human genome containing *VEGFC*. This gene has been associated with disseminated epithelial tumor cells to regional lymph nodes (Van Trappen and Pepper, 2002). Thus, it can serve as an endothelial marker likely correlating with angiogenesis due to metastasis or minimal residue disease after a curative-intent surgery and can also offer a strategy for the selection of combination therapy using zaltrap or eylea (Muro et al., 2020).

DISTANT METASTASIS ORGANOIDS

We derived rectal cancer organoids from resected tissue of a metastatic rectal tumor protruding from the sternum. When the organoids exceeded 150 µm in diameter, they began forming intestinal glands or crypts (Lieberkühn, 1782). The intestinal epithelium is an integral component of innate immunity. The intestinal epithelial cells shape multitubular invaginations that form crypts, which increase the absorption surface of the tissue. At the base of the crypts, the intestinal stem cell niche enables the constant regeneration of the intestinal lining (e.g., enterocytes, endocrines, or goblet cells). These cells can proliferate, differentiate and move upwards, where they can be replaced after a couple of days. In our organoid culture, the patient cancer cells readily formed organoids larger than ½ mm in diameter with some of the crypts exceeding 100 µm in diameter (Fig. 9). To evaluate the metastatic potential of the organoids, we used a cell migration transwell assay (Chen, 2005) and observed organoid cells protruding overnight through the insert membrane (Matov, 2024a), which indicated a strong metastatic potential of the tumor.

We sought to identify therapeutically exploitable vulnerabilities in CRC cells. Initial data was obtained with the metastatic RC organoid (we established from 70 mg resected lung tissue containing metastatic tumor) with the standard first line metastatic CRC chemotherapy FOLFOX. We plated organoids with diameters 40-80 µm in 24 wells (Fig. 9) and performed titration with concentration of 1 µM , 10 µM, and 100 µM for two weeks. We repeated the same with docetaxel with concentration of 10 nM, 100 nM, and 1 µM, and the experiments showed clear residual disease measured by CellTiter-Glo assay (Fig. 10). This was the first example of "persister" cancer cells in 3D patient-derived organoids cultured in 30 µL Matrigel drops (van de Wetering et al., 2015), i.e., in their natural 3D environment and not as single patient cells cultured on a flat plastic surface in a 2D Petri dish.

Figure 9. Metastatic rectal cancer organoids obtained from SU2C (chest wall) distant metastasis. Organoids with diameter 40-80 µm (with a few larger organoids also present, see inset to the right) were plated in 30 µL Matrigel drops in six wells per condition of a 24-well plate (see inset to the left). Four weeks after plating, many organoids in the wells without treatment had reached a size of over 200 µm in diameter. Crypts are clearly visible in large organoids (see white ovals). We treated organoids for two or more weeks with docetaxel with concentration of 100 nM and 1 µM, and FOLFOX (folinic acid, fluorouracil, and oxaliplatin) with concentration of 1 µM, 10 µM, and 100 µM to eliminate the vast majority of the organoid cells and induce a persister state in the residual cancer cells. Scale bar equals 50 µm. Inset scale bar equals 70 µm. Phase contrast microscopy, 20x magnification.

Treatment with docetaxel of prostate tumors induces a non-mutational mechanism of acquired resistance in a small subset of cells (<5%), which become quiescent, and thus drug-tolerant persister cells, during a reversible process linked to upregulation of mesenchymal markers and downregulation of epithelial markers. During drug holiday, the cells revert to normal metabolism and expand their population, while becoming again susceptible to drug treatment. This cycle repeats multiple times during patient treatment. While the tumor is under drug pressure, however, the few drug-tolerant persister cells pay a metabolic price and become temporarily vulnerable to GPX4 inhibition. Our work has demonstrated (Matov, 2025a) that prostate tumors with mesenchymal gene expression

Figure 10. ATP values after drug treatment. Per condition, three wells of organoids in Matrigel were treated for two weeks with 10 nM docetaxel and 1 µM FOLFOX. The average values of the measured luminescence were 82,220 and 75,399, respectively.

signatures after docetaxel treatment are susceptible to GPX4 inhibition and that the small molecule ML210, which is electively toxic to persister cells, can eliminate residual disease.

We also derived organoids from organ-confined, drug-naïve intraductal prostate cancer (PC) with strong metastatic potential of the tumor. In our culture, the organoids formed visible prostatic ducts and were fast proliferating, surrounded by a niche of fibroblast cells. Organoids were often shaped like two urethral sinuses. Noticeably, a very large number of ductal PC organoids tended to form at the edge of the Matrigel, in direct contact with the culture medium (Fig. 11).

We derived organoids from high grade (Gleason Score 4+3) acinar adenocarcinoma of the prostate and performed lentiviral transduction with MT markers immediately after tissue dissociation, before we seeded single cells in Matrigel. In our culture, the organoids expressed EB1ΔC-2xEGFP a day after radical prostatectomy and we imaged polymerizing MTs in cancer cells, and also in cancer-associated

fibroblast (CAF) cells. The fibroblasts had 2-fold larger area and over 3.8-fold higher density of polymerizing MTs compared to the cancer cells (Fig. 12). The high level of MT polymerization activity in CAFs and their extensive length (over 100 µm) allow for a rapid paracrine signalling in prostate tumors. Our work has shown the extraordinary ability of small organoids to move in Matrigel and form large structures (Matov, 2025b) in samples from micro-metastatic tumors.

Figure 11. Intraductal prostate cancer organoids obtained from organ-confined disease. Organoids derived from 1 mg tissue (Gleason Score 5+4, punch core biopsy) from radical prostatectomy. Ductal prostate cancer is rare and very aggressive. The organoids formed visible ducts (see left panel) and were surrounded by a niche of fibroblasts. Organoids were often shaped like two urethral sinuses (see the right panel and the lower inset). A very large number of ductal cancer organoids tended to form at the edge of the Matrigel in direct contact with the culture medium (see upper inset). Phase contrast microscopy, 20x magnification. Scale bars equal 50 µm.

Figure 12. Spinning disc confocal microscopy image of a patient organoid seeded and transduced with EB1ΔC-2xEGFP a day after radical prostatectomy. Organoids were derived from 1 mg punch biopsy. Day 3. (A) Primary prostate cancer cells with labeled polymerizing MT ends. The inset shows detection of EB1 comets for an area of the cell. Scale bar equals 15 µm. (B) Cancer-associated fibroblast cell with labeled polymerizing MT ends. The insets show detection of EB1 comets for two areas of the cell. Scale bar equals 10 µm.

The overall most common site of soft tissue (visceral) metastasis in PC and the most common site of secondary metastases of PC patients with bone metastases is the liver (Gandaglia et al., 2014). Further, men with metastasis to the liver have the worst median survival time (Shiner et al., 2024). In this context, we derived PC organoids from liver metastasis needle biopsies. These cultures were characterized by very low proliferation rates and formed distinctly translucent organoids. However, the organoids derived from distant metastases displayed a remarkable ability to move and self-organize. By the third day after tissue dissociation and seeding of single cells in Matrigel, the median organoid size was about 40 µm in diameter, which suggests that there were a maximum of a dozen cells in the organoids.

What we noticed on the third day after seeding was that many of the small organoids had formed a large structure of 0.6 mm in diameter (Fig. 13), which was remarkable. It underscores the incredible ability of these tumors to establish signaling communication in a new environment and protrude to enhance their micro-environment (Fig. 14), all of which required about 72 hours. After the formation of a few large structures on Day 3, the remaining organoids had very reduced proliferation rates with only a small fraction (less than a quarter) increasing in size. By Day 71, the median organoid size had reached 80 µm in diameter, which suggests that there were about 60 cells within the organoids. The large structures had disappeared, likely because the CAFs had died. By Day 139, the organoids had receded to a median size of 70 µm in diameter and had further receded to a median size of 55 µm in diameter by the six months mark.

Figure 13. Metastatic prostate cancer organoids obtained from SU2C (liver) distant metastasis. Organoids were derived from ½ mg needle biopsy. Day 5. While most organoids had reached a size of 40 µm in diameter, a few large aggregates of 0.6 mm in diameter consisting of small organoids had formed. See Fig. 14 for a zoom-in image of the area within the red rectangle. Transmitted light microscopy, 4x magnification. Scale bar equals 100 µm. The left inset shows a translucent organoid, typical for heavily drug-treated tumors. Transmitted light microscopy, 20x magnification. Inset scale bar equals 20

µm. The right inset shows an organoid with irregular shape. Phase contrast microscopy, 20x magnification. Inset scale bar equals 50 µm. The figure at the lower-left corner shows the changes in median organoid diameter (in green) and percent of growing organoids (in red) over a period of six months.

Figure 14. Metastatic prostate cancer organoids obtained from SU2C (liver) distant metastasis. Organoids were derived from ½ mg needle biopsy. Day 5. Large aggregates of 0.6 mm in diameter - consisting of small organoids (see the red insets)

containing very few cells - had formed. Scale bar equals 30 µm. Phase contrast microscopy, 20x magnification. The image represents the same area as the red rectangle on Fig. 13.

DISCRETE BROADCAST CHANNEL

Visible in the difference image on Fig. 1D is the appearance of periodic vertical streaks around 90 to 150 bp. Visible are also horizontal streaks, likely resulting from errors in gene copy-number amplification in CRC. These examples demonstrate the patterns appearing in disease in comparison to normal physiology. Such grayscale images, in the thousands, can be utilized to train a generative transformer (Ren et al., 2024), or another large language model. This classification approach will ensure all epigenetic changes occurring in disease have been taken into account.

One possible avenue for classification is to derive the maximum likelihood estimate of the parameters of Markov Chain Monte Carlo sampling (Kungurtsev et al., 2023) for time series prediction and supervised Bayesian learning (Chandra et al., 2019). Further, let the appearance of co-morbidities in healthy donor samples be a stationary Markov chain and CRC Stages I – IV denote its noisy version as a Hidden Markov Process (HMP), when corrupted by a discrete memoryless channel (DMC) (Marton, 1979), with channel capacity equal to the maximum of the KL divergence. The DMC (Gamal and Meulem, 1981) is completely characterized by the channel transition matrix, also known as the confusion matrix (Lee et al., 2015). Consider the HMP given by a binary symmetric channel with corrupted symmetric binary Markov source. One can approximate the entropy rate of a HMP via approximations of the stationary distribution of a related Markov process with high precision-complexity trade-off (Ordentlich and Weissman, 2011). Therefore, KL divergence can serve as a prognostic biomarker (Zhong et al., 2020) based on longitudinal data prior to diagnosis, i.e., samples collected periodically from the same healthy donor earlier in life.

In this context, we measured the KL divergence of three cohorts of healthy individuals from a CC cohort. The first healthy cohort consisted of samples from healthy individuals with no co-morbidities. This cohort was the most divergent from the CC cohort. The second healthy cohort consisted of samples from healthy individuals with identified co-morbidities, such as hypertension or diabetes. This cohort followed the same patterns of divergence from the CC cohort, but the peak values were lower, i.e., the samples were less divergent than the healthy cohort without co-morbidities. The third cohort consisted of samples from healthy individuals with other findings, such disorders of the glycoprotein metabolism. This cohort exhibited a further decrease in divergence from the CC cohort for DNA fragment lengths around 195 bp, which was the most divergent size. However, for other fragment lengths it was more divergent than the co-morbidities cohort (Fig. 15). Overall, these results indicate that the stages of pathogenesis in CRC form a Markov chain in terms of DNA fragmentation patterns.

Figure 15. DNA fragmentation length KL divergence (in bits) of three healthy individuals' cohorts from a colon cancer cohort. The colon cancer cohort consists of 79 patients in Stages I-IV. The KL divergence (in bits) is on the Y axis. The healthy individuals' cohort shown in green consists of 74 samples from healthy donors with no co-morbidities. The maximal value is 0.8 bits divergence for DNA fragment length 195 bp. The healthy individuals' cohort shown in blue consists of 71 samples from healthy donors with co-morbidities. The maximal value is 0.4 bits divergence for DNA fragment length 195 bp. The healthy individuals' cohort shown in red consists of 71 samples from healthy donors with other findings. The maximal value is 0.4 bits divergence for DNA fragment length 137 bp. The fragment length (in bp) is on the X axis, from left to right.

Further, we analyzed the KL divergence of a cohort of healthy individuals from CRC patient cohorts by disease stage. The cohort of CRC Stage I samples was the least divergent, forming two distinct peaks. Next, the cohort of CRC Stage II samples was more divergent than CRC Stage I and the divergence peaks followed the same distribution but had a higher amplitude. Similarly, the cohort of CRC Stage III samples was more divergent than CRC Stage II and the divergence peaks followed the same distribution but with significantly higher amplitudes. Finally, the KL divergence patterns for the cohort of Stage IV samples followed a different distribution. Besides the two main peaks, the CRC Stage IV samples formed additional divergence peaks, in particular around 290 bp, which was likely due to metastases to the liver (Fig. 16). Overall, these results indicate that the stages of disease progression in CRC form a Markov chain in terms of DNA fragmentation patterns.

If we consider the transformations occurring in the DNA fragmentation patterns within a cohort of healthy samples so that all become alike to disease samples, we can view pathogenesis and disease progression of a population as a broadcast channel with memory and present it in terms of Gaussian multi-user parallel broadcast channels with identical code words (Matov, 1999). The achievable rate for the capacity of a degraded broadcast channel (in bits) is a function of the logarithm of the signal-to-noise ratio of the transmission signal and depends on the quality of the transmission medium. Next, the achievable rates for the capacity region of a family of parallel broadcast channels is given by the union of the overall capacity in each channel (Tse, 1999). Hence, any divergence in the fragmentation

patterning within the healthy cohort (baseline dataset or parallel broadcast channels) would create an elevated noise floor and, thus, an overall unsolvable stochastic heterogeneity.

Figure 16. DNA fragmentation length KL divergence (in bits) of a healthy individuals' cohort from CRC patient cohorts by stage. The CRC cohorts consist of 27 patients of which 7 in Stage I (green), 7 in Stage II (blue), 5 in Stage III (yellow), and 8 Stage IV (red). The healthy individuals' cohort consists of 43 samples. The KL divergence (in bits) is on the Y axis. For CRC Stage I, the maximal value is 2.7 bits divergence for DNA fragment length 216 bp. For CRC Stage II, the maximal value is 3.4 bits for fragment length 213 bp. For CRC Stage III, the maximal value is 5.1 bits divergence for DNA fragment length 210 bp. For CRC Stage IV, the maximal value is 2.4 bits divergence for DNA fragment length 195 bp. The fragment length (in bp) is on the X axis, from left to right. The distribution changes for CRC Stage IV and exhibits three additional peaks (at DNA fragment length 280 bp in particular), likely due to metastases to the liver.

Each time a new patient sample is presented for classification, it would not be compared to samples collected from the same person longitudinally, in which most of the fragmentation pattern would be very similar to the previously collected healthy samples. Instead, it would be compared to a variety of fragmentation patterns in a whole cohort of (different people) healthy donors. Such comparisons will

always generate noisy baseline datasets, even in the case of a very large healthy donor population, because a very few outliers will affect the overall quality of the healthy baseline dataset. For this reason, such population approaches, because of the intrinsic variability in human, inherently impedes the ability of all currently utilized methods to detect disease in its early stage.

MICRO-METASTASIS ORGANOIDS

When prostate tumors escape the capsule, they form micro-metastatic lesions at the retroperitoneal lymph nodes. In our organoid culture, such drug-naïve tumors formed slow-proliferating tumors and had driver mutations, such as homozygous mutations in PTEN (exon2, c.G194>C, p.C65S) and BRCA2 (exon14, c.T7397>C, p.V2466A). Further, we detected a heterozygous ATR mutation (exon4, c.T632>C, p.M211T). ATR has emerged as a promising strategy for cancer treatment that exploits synthetic lethal interactions with proteins involved in DNA damage repair, overcomes resistance to other therapies and enhances antitumour immunity (Ngoi et al., 2024). Figure 17 shows organoids after 152 days in culture, at the time when growth kinetics have plateaued. After an initial couple of days of metabolic recovery after tissue dissociation, the tumor cells began forming organoids and about 65% of the organoids were growing – a number that decreased to about 35% within a month and remained at similar values over the six months we kept the organoids in culture. The organoid growth kinetics were remarkably rapid in the first about 10 days and later slowed down, reaching a plateau in about three months. After that, the organoids continued growing in size at a very slow pace. Larger organoids displayed a twined structure, suggesting a proliferative phenotype related to the mutation in *PTEN* (Fig. S6). Many organoids formed columnar shapes of acinar adenocarcinoma, consistent with luminal differentiation (Agarwal et al., 2015).

Figure 17. Metastatic prostate cancer organoids obtained from micro-metastasis at a retroperitoneal lymph node. Large organoids obtained from resected tissue after radical prostatectomy. Drug-naïve tumor. Day 152. Magnification 4x, transmitted light microscopy. Scale bar equals 100 µm. A larger image of the organoid in the lower-right corner is available in phase contrast on Fig. S7. Lower inset: Day 3; scale bar equals 50 µm. Upper inset: Day 106; scale bar equals 100 µm. Magnification 20x, phase contrast microscopy. The figure at the bottom shows the changes in median organoid diameter (in green) and percent of growing organoids (in red) over a period of six months.

We also detected heterozygous SPOP (exon5, c.T398>G, p.F133C) and TP53 (exon3, c>C98G, p.P33R, exon4, c.C215>G, p.P72R) mutations. *SPOP* is the most commonly mutated gene in PC, mutated in around 10% of primary prostate tumors (Boysen et al., 2015).

We also derived organoids from a micro-metastatic lesion at a retroperitoneal lymph node of tumor treated with androgen deprivation therapy and radiotherapy. The organoids were much more proliferative in comparison to the drug-naïve ones. They exhibited very peculiar growth dynamics and changes in morphology. Initially, the organoid shape was circular and a large number of small organoids was formed in the first days after radical prostatectomy, which appeared distinctly translucent. Organoid shape suddenly changed after two months in culture and many organoids had a hyperplastic phenotype characteristic for *PTEN* deletion and ERG overexpression (Karthaus et al., 2014). The organoids appeared to be embedded in a niche of activated fibroblasts.

Smaller organoids seemed to move in the Matrigel and group into large clumps surrounded by fibroblasts. While two weeks after plating of single cells after tissue dissociation about 80% of the organoids were in a phase of growth, the actual rates of proliferation over the first two months were low. During the third month, when the percent of growing organoids had dropped to between 30% and 40%, the median organoid size suddenly increased 3-fold to 4-fold. This increase in size was coupled with a change in morphology (Fig. 18). Remarkably, at the end of the third month in culture, a couple of very large organoids formed – one of which was almost 1 mm long (Matov, 2025b). The median organoid size at this time was about 120 µm and the formation of organoids 6-fold larger or more was the result of paracrine signaling-induced movement of smaller organoids, which clustered together in one of the Matrigel drops. The ability to move at such speeds was characteristic only for organoids derived from drug-treated metastatic (both micro-metastatic and distant metastasis) tumors. To a certain extent, we

observed similar clumping of smaller organoids in high grade (Gleason Score 5+4) primary tumors (Matov, 2024g).

Figure 18. Metastatic prostate cancer organoids obtained from micro-metastasis at a retroperitoneal lymph node. Two large organoids obtained from resected tissue after radical prostatectomy. Prior treatment - androgen deprivation therapy and radiotherapy. Day 60. Magnification 4x, transmitted light microscopy. Scale bar equals 100 µm. Insets magnification 20x,

transmitted light microscopy. Left inset scale bar equals 30 µm. Right inset scale bar equals 40 µm. The figure at the bottom shows the changes in median organoid diameter (in green) and percent of growing organoids (in red) over a period of three months.

DISCUSSION

In the specific case of CRC, the existence of an intrinsic stochastic heterogeneity of DNA fragmentation in healthy samples means that any non-neoplastic gastrointestinal (GI) metabolic divergence in samples from the healthy donor cohort considered as baseline during classification would impair the ability of traditional ML classifiers to reliably detect neoplastic transformation (Wan et al., 2019). Oppositely, an image-based classification, as we propose here, could be in a position and better equipped to successfully detect and delineate both the DNA fragmentation patterns resulting from tumors and those resulting from non-lethal, transient conditions, such as inflammation (Horiuchi et al., 2024). If this holds true, it would impact, besides CRC diagnosis, our ability to correctly diagnose other cancers of the GI tract as well.

We propose the utilization of the described here DNA fragmentation analytics to generate embeddings (Vaswani et al., 2017) for a generative transformer network in which the tokens are the KL divergence metrics of the samples. The embeddings can include DNA fragment length patterns before and after disease diagnosis. We will retrain a transformer network with a new set of tokens - with genetic mutations rather than words and with tumor burden values rather than sentences of human speech, i.e., we will retrain a large language model with disease progression datasets. This will allow training of the network to classify each new sample and further make predictions regarding disease prognosis. Additional tokens will be the texture metrics of patient-derived organoids (Matov, 2024a; Matov, 2024b; Matov et al., 2016) such as shape descriptors and other morphology data (Galletti et al., 2013; Galletti et al., 2014; Matov, 2024g), which could be obtained from circulating tumor cells (Matov and Modiri,

2024; Sung et al., 2013; Sung et al., 2012; Tasaki et al., 2014). The embeddings can further include cytoskeletal dynamics (Matov, 2024c; Matov, 2024d; Matov, 2024e; Matov, 2024f; Matov, 2025a; Matov, 2025b) before and after treatment with different drugs. This will allow training of the network to classify as sensitive or resistant each organoid and further identify subclasses with different mechanisms of resistance. Further, after training a generative transformer network with organoid texture and live-cell dynamics datasets, we will be able to model disease progression as well.

In the long run, the most reliable way to perform early detection will be to personalize the process by aggregating longitudinal baseline datasets for each individual. We have attempted to do that in an effort of detecting LC in urine samples (Matov, 2024a; Matov, 2024h). Analysis of microRNA profiles extracted from the urine of healthy donors longitudinally indicates a relative consistency in their levels. This result suggests that tracking the levels of nucleic acids in body fluids longitudinally may allow for the delineation of organ- and tissue-specific patterns of changes in dedicated panels of disease-associated biomarkers and, thus, anticipate early disease and inform therapy.

MATERIALS AND METHODS

Data Processing

Whole genome sequencing DELFI raw data from (Cristiano et al., 2019) was processed to extract and bin all available DNA fragments using: https://github.com/Hogfeldt/ctDNAtool.

Data Analysis

All analysis programs for fragmentomics analysis and graphical/image representation of the results were developed in R and Python. The KL divergence method used is described and validated in (Han et al., 2016). The computer code is available for download at:

https://github.com/amatov/FragmentomicsSubclinicalDisease.

Image Analysis

All image analysis programs for EB1 comet detection, MT dynamics, and graphical representation of the results were developed in Matlab and C/C++. The computer code is available for download at: https://github.com/amatov/ClusterTrackTubuline.

Sample Processing

Small RNA were extracted from de-identified urine samples (50 ml) by Norgen Biotek Corp.

Retroperitoneal lymph node organoid whole genome sequencing (736 ng DNA) was done at the UCSF Core Facilities.

Microscopy Imaging

We imaged EB1ΔC-2xEGFP-expressing organoids (Koo et al., 2013) by time-lapse spinning disk confocal microscopy using a long working distance 60x magnification water immersion 1.45 NA objective. We acquired images at half a second intervals for a minute to collect datasets without photobleaching (Gierke and Wittmann, 2012). The 3D z-stacks consisted of 35 2D slices acquired at a 0.5 µm vertical spacing.

Organoids were imaged using transmitted light microscopy at 4x magnification and phase contrast microscopy at 20x magnification on a Nikon Eclipse Ti system with camera Photometrics CoolSnap HQ2.

Cell Culture

Organoids with diameter 40-80 µm were plated in 30 µL Matrigel drops in six wells per condition of a 24-well plate per condition and treated for two or more weeks with docetaxel with concentration of 100 nM and 1 µM to eliminate the vast majority of the organoid cells and induce a persister state in the residual cancer cells. This treatment was followed by co-treatment with docetaxel 100 nM and small molecule ML210 5 µM or docetaxel 100 nM combined with small molecule ML210 5 µM and lipophilic antioxidants ferrostatin-1 2 µM, rescue treatment. Organoids with diameter 40-80 µm were plated in 30 µL Matrigel drops in six wells per condition of a 24-well plate per condition and also treated for two or more weeks with FOLFOX (folinic acid, fluorouracil, and oxaliplatin) with concentration of 1 µM, 10 µM, and 100 µM to eliminate the vast majority of the organoid cells and induce a persister state in the residual cancer cells.

Primary and retroperitoneal lymph node metastatic prostate tumor and sternum metastatic rectal tumor tissues were dissociated to single cells using modified protocols from the Witte lab. Organoids were seeded as single cells in three 30 µL Matrigel drops in 6-well plates. Organoid medium was prepared according to modified protocols from the Clevers lab and the Chen lab.

To obtain a single cell suspension, tissues were mechanically disrupted and digested with 5 mg/ml collagenase in advanced DMEM/F12 tissue culture medium for several hours (between 2 and 12 hours, depending on the biopsy / resection performed). If this step yielded too much contamination with non-epithelial cells, for instance during processing of primary prostate tumors, the protocol incorporated additional washes and red blood cell lysis (Goldstein et al., 2011). Single cells were then counted using a hemocytometer to estimate the number of tumor cells in the sample, and seeded in growth factor-reduced Matrigel drops overlaid with PC medium (Gao et al., 2014). With radical prostatectomy

specimens, we had good success with seeding three thousand cells per 30 µl Matrigel drop, but for metastatic samples organoid seeding could reliably be accomplished with significantly less cells, in the hundreds.

Docetaxel was purchased from Sigma-Aldrich. Folinic acid, fluorouracil, oxaliplatin, RSL3, ML210, and ferrostatin-1 were obtained from the McCormick lab.

Ethics Declaration

IRB (IRCM-2019-201, IRB DS-NA-001) of the Institute of Regenerative and Cellular Medicine. Ethical approval was given.

Approval of tissue requests #14-04 and #16-05 to the UCSF Cancer Center Tissue Core and the Genitourinary Oncology Program was given.

Data Availability Statement

The datasets used and/or analyzed during the current study are available from the corresponding author upon reasonable request.

Conflicts of Interest

The author declares no conflicts of interest.

Funding Statement

No funding was received to assist with the preparation of this manuscript.

ACKNOWEDGEMENTS

I thank Claus Andersen for genome data and his feedback on the shortcomings of the current whole genome sequencing analysis methods. I am grateful to the Genitourinary Tissue Utilization committee and the Genitourinary and Prostate SPORE Tissue Cores at the UCSF Cancer Center for the approval of my tissue requests #14-04 and #16-05, the Stand Up To Cancer / Prostate Cancer Foundation (SU2C/PCF) West Coast Dream Team (WCDT), the Institute of Regenerative and Cellular Medicine for issuing the Institutional Review Board protocol approval IRCM-2019-201, IRB DS-NA-001 for the observational study "Longitudinal analysis of next-generation sequencing of nucleic acids for early detection of degenerative diseases such as cardiovascular, neoplastic and diseases related to the nervous system", and James Faber for his feedback regarding the protocol and the process of approval.

SUPPLEMENTARY MATERIALS

Figure S1 – DNA fragmentation length KL divergence (in bits) of a healthy individuals' cohort from a CRC patient cohort with color-coding for the number of fragments per fragment length.

Figure S2 – Genomic bins KL divergence (in bits) of a healthy individuals' cohort from a CRC patient cohort for DNA fragment length 364 bp.

Figure S3 – Genomic bins KL divergence (in bits) of a healthy individuals' cohort from a CC patient cohort for DNA fragment length 195 bp.

Figure S4 – Genomic bins KL divergence (in bits) of a healthy individuals' cohort from a CRC patient cohort for DNA fragment length 205 bp.

Figure S5 – Genomic bins KL divergence (in bits) of a healthy individuals' cohort from a CRC patient cohort for DNA fragment length 129 bp.

Figure S6 – Metastatic prostate cancer organoids obtained from micro-metastasis at a retroperitoneal lymph node. Day 106.

Figure S7 – Metastatic prostate cancer organoids obtained from micro-metastasis at a retroperitoneal lymph node. Day 152.

Figure S1. DNA fragmentation length KL divergence (in bits) of a healthy individuals' cohort from a CRC patient cohort. The ratio of number of fragments for each length is color coded. The CRC cohorts consist of 27 patients in Stages I-IV. The healthy individuals' cohort consists of 43 samples. The KL divergence (in bits) is on the Y axis. The maximal value is 2.8 bits divergence for DNA fragment length 364 bp (150,000 fragments). The second highest divergence value is 2.6 bits for DNA fragment length 198 bp (2.5 million fragments). The fragment length (in bp) is on the X axis, from left to right. The two main divergence peaks have different fragment numbers.

Figure S2. Genomic bins KL divergence (in bits) of a healthy individuals' cohort from a CRC patient cohort for DNA fragment length 364 bp. Above a threshold of 1.5 bits, 12 genomic bins were identified as divergence in CRC.

Figure S3. Genomic bins KL divergence (in bits) of a healthy individuals' cohort from a CC patient cohort for DNA fragment length 195 bp. Above a threshold of 1.2 bits, 12 genomic bins were identified as divergence in CC.

Figure S4. Genomic bins KL divergence (in bits) of a healthy individuals' cohort from a CRC patient cohort for DNA fragment length 205 bp. Above a threshold of 0.56 bits, 12 genomic bins were identified as divergence in CRC.

Figure S5. Genomic bins KL divergence (in bits) of a healthy individuals' cohort from a CRC patient cohort for DNA fragment length 129 bp. Above a threshold of 1.05 bits, 12 genomic bins were identified as divergence in CRC.

Figure S6. Metastatic prostate cancer organoids obtained from micro-metastasis at a retroperitoneal lymph node. Large organoids obtained from resected tissue after radical prostatectomy. Drug-naïve tumor. Day 106. Scale bar equals 100 µm. Magnification 20x, phase contrast microscopy.

Figure S7. Metastatic prostate cancer organoids obtained from micro-metastasis at a retroperitoneal lymph node. Large organoids obtained from resected tissue after radical prostatectomy. Drug-naïve tumor. Day 152. Scale bar equals 100 µm. Magnification 20x, phase contrast microscopy. This is the same organoid as in the lower-right corner on Fig. 17.

REFERENCES

Agarwal, S., P.G. Hynes, H.S. Tillman, R. Lake, W.G. Abou-Kheir, L. Fang, O.M. Casey, A.H. Ameri, P.L. Martin, J.J. Yin, P.J. Iaquinta, W.R. Karthaus, H.C. Clevers, C.L. Sawyers, and K. Kelly. 2015. Identification of Different Classes of Luminal Progenitor Cells within Prostate Tumors. *Cell reports*. 13:2147-2158.

Avissar, N., D.B. Ornt, Y. Yagil, S. Horowitz, R.H. Watkins, E.A. Kerl, K. Takahashi, I.S. Palmer, and H.J. Cohen. 1994. Human kidney proximal tubules are the main source of plasma glutathione peroxidase. *The American journal of physiology*. 266:C367-375.

Barrett, C.W., W. Ning, X. Chen, J.J. Smith, M.K. Washington, K.E. Hill, L.A. Coburn, R.M. Peek, R. Chaturvedi, K.T. Wilson, R.F. Burk, and C.S. Williams. 2013. Tumor suppressor function of the plasma glutathione peroxidase gpx3 in colitis-associated carcinoma. *Cancer Res*. 73:1245-1255.

Bergmans, P. 1973. Random Coding Theorem for Broadcast Channels with Degraded Components. *IEEE Transactions on Information Theory*. 19(2):197 - 207.

Bergmans, P. 1974. A Simple Converse for Broadcast Channel with Additive White Gaussian Noise. *IEEE Transactions on Information Theory*. 20 (2):279 - 280.

Boveri, T. 1902. Über mehrpolige Mitosen als Mittel zur Analyse des Zellkerns. *Verhandlungen der Physikalisch-medizinische Gesellschaft in Würzburg*. 35:67–90.

Boysen, G., C.E. Barbieri, D. Prandi, M. Blattner, S.-S. Chae, A. Dahija, S. Nataraj, D. Huang, C. Marotz, L. Xu, J. Huang, P. Lecca, S. Chhangawala, D. Liu, P. Zhou, A. Sboner, J.S. de Bono, F. Demichelis, Y. Houvras, and M.A. Rubin. 2015. SPOP mutation leads to genomic instability in prostate cancer. *eLife*. 4:e09207.

Bremaud, P. 1999. Gibbs Fields and Monte Carlo Simulation. *Markov Chains*. 31:253–322.

Chandra, R., K. Jain, R.V. Deo, and S. Cripps. 2019. Langevin-gradient parallel tempering for Bayesian neural learning. *Neurocomputing*. 359:315-326.

Chen, H.-C. 2005. Boyden Chamber Assay. *In* Cell Migration: Developmental Methods and Protocols. J.-L. Guan, editor. Humana Press, Totowa, NJ. 15-22.

Chen, L., R. Liu, Z.P. Liu, M. Li, and K. Aihara. 2012. Detecting early-warning signals for sudden deterioration of complex diseases by dynamical network biomarkers. *Sci Rep*. 2:342.

Chen, X., R. Kang, G. Kroemer, and D. Tang. 2021. Broadening horizons: the role of ferroptosis in cancer. *Nat Rev Clin Oncol*. 18:280-296.

Cover, T. 1998. Comments on Broadcast Channels. *IEEE Transactions on Information Theory*. 44 (6):2524-2530.

Cristiano, S., A. Leal, J. Phallen, J. Fiksel, V. Adleff, D.C. Bruhm, S. Jensen, J.E. Medina, C. Hruban, J.R. White, D.N. Palsgrove, N. Niknafs, V. Anagnostou, P. Forde, J. Naidoo, K. Marrone, J. Brahmer, B.D. Woodward, H. Husain, K.L. van Rooijen, M.W. Ørntoft, A.H. Madsen, C.J.H. van de Velde, M. Verheij, A. Cats, C.J.A. Punt, G.R. Vink, N.C.T. van Grieken, M. Koopman, R.J.A. Fijneman, J.S. Johansen, H.J. Nielsen, G.A. Meijer, C.L. Andersen, R.B. Scharpf, and V.E. Velculescu. 2019. Genome-wide cell-free DNA fragmentation in patients with cancer. *Nature*. 570:385-389.

Dixon, S.J., and B.R. Stockwell. 2014. The role of iron and reactive oxygen species in cell death. *Nature chemical biology*. 10:9-17.

Dolma, S., S.L. Lessnick, W.C. Hahn, and B.R. Stockwell. 2003. Identification of genotype-selective antitumor agents using synthetic lethal chemical screening in engineered human tumor cells. *Cancer Cell*. 3:285-296.

Galletti, G., K. Cleveland, A. Matov, J.E. Hayes, R.J. Klein, D.C. Hassane, L.V. Nicacio, and P. Giannakakou. 2013. Clinical and preclinical evaluation of taxane sensitivity in gastric cancer (GC): Relevance of GC histology. *Journal of Clinical Oncology*. 31:37-37.

Galletti, G., K. Cleveland, C. Zhang, A. Gjyrezi, A. Matov, D. Betel, M.A. Shah, and P. Giannakakou. 2014. Elucidating the molecular basis of intrinsic taxane resistance in gastric cancer. *Cancer Research*. 74:897.

Gamal, A., and E. Meulem. 1981. A Proof of Marton's Coding Theorem for the Discrete Memoryless Broadcast Channel. *IEEE Transactions on Information Theory*. 27 (1):120 - 122.

Gandaglia, G., F. Abdollah, J. Schiffmann, V. Trudeau, S.F. Shariat, S.P. Kim, P. Perrotte, F. Montorsi, A. Briganti, Q.D. Trinh, P.I. Karakiewicz, and M. Sun. 2014. Distribution of metastatic sites in patients with prostate cancer: A population-based analysis. *The Prostate*. 74:210-216.

Gao, D., I. Vela, A. Sboner, P.J. Iaquinta, W.R. Karthaus, A. Gopalan, C. Dowling, J.N. Wanjala, E.A. Undvall, V.K. Arora, J. Wongvipat, M. Kossai, S. Ramazanoglu, L.P. Barboza, W. Di, Z. Cao, Q.F. Zhang, I. Sirota, L. Ran, T.Y. MacDonald, H. Beltran, J.M. Mosquera, K.A. Touijer, P.T. Scardino, V.P. Laudone, K.R. Curtis, D.E. Rathkopf, M.J. Morris, D.C. Danila, S.F. Slovin, S.B. Solomon, J.A. Eastham, P. Chi, B. Carver, M.A. Rubin, H.I. Scher, H. Clevers, C.L. Sawyers, and Y. Chen. 2014. Organoid cultures derived from patients with advanced prostate cancer. *Cell*. 159:176-187.

Gierke, S., and T. Wittmann. 2012. EB1-recruited microtubule +TIP complexes coordinate protrusion dynamics during 3D epithelial remodeling. *Current biology : CB*. 22:753-762.

Goldstein, A.S., J.M. Drake, D.L. Burnes, D.S. Finley, H. Zhang, R.E. Reiter, J. Huang, and O.N. Witte. 2011. Purification and direct transformation of epithelial progenitor cells from primary human prostate. *Nat Protoc*. 6:656-667.

Han, T.S. 1981. The Capacity Region for the Deterministic Broadcast Channel with a Common Message. *IEEE Transactions on Information Theory*. 27(1):122 - 125.

Han, Y., J. Jiao, and T. Weissman. 2016. Minimax Rate-optimal Estimation of KL Divergence between Discrete Distributions. *International Symposium on Information Theory and its Applications*. 2016:256-260.

Hangauer, M.J., V.S. Viswanathan, M.J. Ryan, D. Bole, J.K. Eaton, A. Matov, J. Galeas, H.D. Dhruv, M.E. Berens, S.L. Schreiber, F. McCormick, and M.T. McManus. 2017. Drug-tolerant persister cancer cells are vulnerable to GPX4 inhibition. *Nature*. 551:247-250.

Horiuchi, D., H. Tatekawa, T. Oura, S. Oue, S.L. Walston, H. Takita, S. Matsushita, Y. Mitsuyama, T. Shimono, Y. Miki, and D. Ueda. 2024. Comparing the Diagnostic Performance of GPT-4-based ChatGPT, GPT-4V-based ChatGPT, and Radiologists in Challenging Neuroradiology Cases. *Clinical neuroradiology*.

Ivanov, M., A. Baranova, T. Butler, P. Spellman, and V. Mileyko. 2015. Non-random fragmentation patterns in circulating cell-free DNA reflect epigenetic regulation. *BMC genomics*. 16 Suppl 13:S1.

Jorgensen, M.L., J.M. Young, and M.J. Solomon. 2015. Optimal delivery of colorectal cancer follow-up care: improving patient outcomes. *Patient related outcome measures*. 6:127-138.

Karras, T., T. Aila, S. Laine, and J.J.A. Lehtinen. 2017. Progressive Growing of GANs for Improved Quality, Stability, and Variation. abs/1710.10196.

Karthaus, Wouter R., Phillip J. Iaquinta, J. Drost, A. Gracanin, R. van Boxtel, J. Wongvipat, Catherine M. Dowling, D. Gao, H. Begthel, N. Sachs, Robert G.J. Vries, E. Cuppen, Y. Chen, Charles L. Sawyers, and Hans C. Clevers. 2014. Identification of Multipotent Luminal Progenitor Cells in Human Prostate Organoid Cultures. *Cell*. 159:163-175.

Koo, B.K., V. Sasselli, and H. Clevers. 2013. Retroviral gene expression control in primary organoid cultures. *Current protocols in stem cell biology*. 27:Unit 5A.6.

Kumar, P., K.S. Lyle, S. Gierke, A. Matov, G. Danuser, and T. Wittmann. 2009. GSK3β phosphorylation modulates CLASP–microtubule association and lamella microtubule attachment. *The Journal of Cell Biology*. 184:895-908.

Kungurtsev, V., A. Cobb, T. Javidi, and B. Jalaian. 2023. Decentralized Bayesian learning with Metropolis-adjusted Hamiltonian Monte Carlo. *Machine Learning*. 112:2791-2819.

Lee, B., T. Moon, S. Yoon, and T. Weissman. 2015. DUDE-Seq: Fast, flexible, and robust denoising for targeted amplicon sequencing. *PLoS ONE*. 12.

Lejeune, S., F. Guillemot, J.P. Triboulet, S. Cattan, C. Mouton, N. Porchet, S. Manouvrier, and M.P. Buisine. 2006. Low frequency of AXIN2 mutations and high frequency of MUTYH mutations in patients with multiple polyposis. *Human mutation*. 27:1064.

Lemasters, J.J. 2017. Evolution of Voltage-Dependent Anion Channel Function: From Molecular Sieve to Governator to Actuator of Ferroptosis. *Frontiers in oncology*. 7:303.

Lieberkühn, J.N. 1782. De Valvula Coli et Usu Processus Vermicularis. *Dissertationes Quatuor*.

Liu, M.C., G.R. Oxnard, E.A. Klein, C. Swanton, and M.V. Seiden. 2020. Sensitive and specific multi-cancer detection and localization using methylation signatures in cell-free DNA. *Annals of oncology : official journal of the European Society for Medical Oncology*. 31:745-759.

Liu, R., P. Chen, K. Aihara, and L. Chen. 2015. Identifying early-warning signals of critical transitions with strong noise by dynamical network markers. *Sci Rep*. 5:17501.

Liu, W., X. Dong, M. Mai, R.S. Seelan, K. Taniguchi, K.K. Krishnadath, K.C. Halling, J.M. Cunningham, L.A. Boardman, C. Qian, E. Christensen, S.S. Schmidt, P.C. Roche, D.I. Smith, and S.N. Thibodeau. 2000. Mutations in AXIN2 cause colorectal cancer with defective mismatch repair by activating beta-catenin/TCF signalling. *Nature genetics*. 26:146-147.

Marton, A. 1979. Coding Theorem for the Discrete Memoryless Broadcast Channel. *IEEE Transactions on Information Theory*. 25 (3):306 - 311.

Matov, A. 1999. Capacity Region Characterization of Multi-User Parallel Gaussian Broadcast Channels. *Swiss Federal Institute of Technology Lausanne*:1-41.

Matov, A. 2024a. Analysis of Multiple Myeloma Drug Efficacy. *medRxiv*:2024.24311450.

Matov, A. 2024b. Colorectal Cancer DNA Fragmentation Patterns as Image Datasets. *SSRN*:2024.4909309.

Matov, A. 2024c. Computational Analysis of Treatment Resistant Cancer Cells. *medRxiv*:2024.24312813.

Matov, A. 2024d. Microtubule Regulation in Cancer Cells *SSRN*:2024.5072927.

Matov, A. 2024e. Mitosis, Cytoskeleton Regulation, and Drug Resistance in Receptor Triple Negative Breast Cancer. *arXiv*:2407.19112.

Matov, A. 2024f. Modulation of the Cytoskeleton for Cancer Therapy *SSRN*:2024.5061950.

Matov, A. 2024g. Quantitative Video Microscopy in Medicine. *SSRN*:2024.4909311.

Matov, A. 2024h. Urinary Biomarkers for Lung Cancer Detection. *medRxiv*:2024.24311186.

Matov, A. 2025a. Microtubule Dysregulation in Prostate Cancer Metastasis to the Vertebrae *SSRN*:2025.5112427.

Matov, A. 2025b. Quantitative Cell Division and Migration in Medicine. *SSRN*:2025.5130515.

Matov, A., and A. Bacconi. 2024. Measurements of Contractility in Actin Convergence Zone and KIF11-Inhibited Mitotic Spindles. *bioRxiv*:2024.601275.

Matov, A., J. de Rooij, J.C. Gatlin, J. Rohrberg, N. Ahmad, R. Aggarwal, J. Simko, A. Goga, C.J. Ryan, and T. Wittmann. 2016. Investigating microtubule growth dynamics in patient-derived metastatic prostate cancer cells. *Cancer Research*. 76:243.

Matov, A., and S. Modiri. 2024. Quantitative Microscopy in Medicine. *medRxiv*:2024.24311304.

Mitchison, T., and M. Kirschner. 1984. Dynamic Instability of Microtubule Growth. *Nature*. 312:237-242.

Mullis, K.B. 1968. Cosmological Significance of Time Reversal. *Nature*. 218:663.

Muro, K., T. Salinardi, A.R. Singh, and T. Macarulla. 2020. Safety of Aflibercept in Metastatic Colorectal Cancer: A Literature Review and Expert Perspective on Clinical and Real-World Data. *Cancers*. 12.

Ngoi, N.Y.L., P.G. Pilié, D.J. McGrail, M. Zimmermann, K. Schlacher, and T.A. Yap. 2024. Targeting ATR in patients with cancer. *Nature Reviews Clinical Oncology*. 21:278-293.

Ordentlich, E., and T. Weissman. 2011. Bounds on the entropy rate of binary hidden Markov processes. *Entropy of Hidden Markov Processes and Connections to Dynamical Systems*.

Petrelli, F., G. Tomasello, K. Borgonovo, M. Ghidini, L. Turati, P. Dallera, R. Passalacqua, G. Sgroi, and S. Barni. 2017. Prognostic Survival Associated With Left-Sided vs Right-Sided Colon Cancer: A Systematic Review and Meta-analysis. *JAMA oncology*. 3:211-219.

Ren, Z., Y. Su, and X. Liu. 2024. ChatGPT-Powered Hierarchical Comparisons for Image Classification. *NIPS '23: Proceedings of the 37th International Conference on Neural Information Processing Systems*:69706 - 69718.

Shiner, A., R.C. Sperandio, M. Naimi, and U. Emmenegger. 2024. Prostate Cancer Liver Metastasis: An Ominous Metastatic Site in Need of Distinct Management Strategies. *Journal of clinical medicine*. 13.

Siegel, R.L., K.D. Miller, and A. Jemal. 2020. Cancer statistics, 2020. *CA: a cancer journal for clinicians*. 70:7-30.

Sung, M.S., A. Gjyrezi, G.Y. Lee, A. Matov, G. Galletti, M. Loftus, Y. Syed, T. Lannin, A. Hristov, C. Mason, S.T. Tagawa, B.J. Kirby, D.M. Nanus, and P. Giannakakou. 2013. Using CTCs to interrogate mechanisms of taxane resistance in the prospective TAXYNERGY clinical trial in prostate cancer. *Cancer Research*. 73:3492.

Sung, M.S., A. Gjyrezi, Y. Syed, T. Lannin, A. Hristov, A. Matov, M. Loftus, G. Galletti, G.Y. Lee, C. Mason, B.J. Kirby, D.M. Nanus, S.T. Tagawa, and P. Giannakakou. 2012. Molecular determinants of taxane activity using circulating tumor cells (CTCs) from castrate-resistant prostate cancer patients. *Understanding Cancer: Genomes to Devices Symposium, New York Academy of Medicine*.

Takanishi, D.M., Jr., S.Y. Kim, P.R. Kelemen, M.L. Yaremko, A.H. Kim, J.E. Ramesar, S.K. Horrigan, A. Montag, F. Michelassi, and C.A. Westbrook. 1997. Chromosome 8 Losses in Colorectal Carcinoma: Localization and Correlation With Invasive Disease. *Molecular diagnosis : a journal devoted to the understanding of human disease through the clinical application of molecular biology*. 2:3-10.

Tasaki, S., M. Sung, A. Matov, G. Galletti, E. Diamond, N. Bander, K. Zhou, S. Tagawa, D. Nanus, and P. Giannakakou. 2014. Analysis of microtubule perturbations and androgen receptor localization in circulating tumor cells from castration resistant prostate cancer patients as predictive biomarkers of clinical response to docetaxel chemotherapy. *Cancer Research*. 74:923-923.

Tse, D.N. 1999. Optimal Power Allocation Over Parallel Gaussian Broadcast Channel. *Proceedings of IEEE International Symposium on Information Theory*:27.

van de Wetering, M., H.E. Francies, J.M. Francis, G. Bounova, F. Iorio, A. Pronk, W. van Houdt, J. van Gorp, A. Taylor-Weiner, L. Kester, A. McLaren-Douglas, J. Blokker, S. Jaksani, S. Bartfeld, R. Volckman, P. van Sluis, V.S. Li, S. Seepo, C. Sekhar Pedamallu, K. Cibulskis, S.L. Carter, A. McKenna, M.S. Lawrence, L. Lichtenstein, C. Stewart, J. Koster, R. Versteeg, A. van Oudenaarden, J. Saez-Rodriguez, R.G. Vries, G. Getz, L. Wessels, M.R. Stratton, U. McDermott, M. Meyerson, M.J. Garnett, and H. Clevers. 2015. Prospective derivation of a living organoid biobank of colorectal cancer patients. *Cell*. 161:933-945.

van der Pol, Y., and F. Mouliere. 2019. Toward the Early Detection of Cancer by Decoding the Epigenetic and Environmental Fingerprints of Cell-Free DNA. *Cancer cell*. 36:350-368.

Van Trappen, P.O., and M.S. Pepper. 2002. Lymphatic dissemination of tumour cells and the formation of micrometastases. *The Lancet. Oncology*. 3:44-52.

Vaswani, A., N. Shazeer, N. Parmar, J. Uszkoreit, L. Jones, A.N. Gomez, Ł. Kaiser, and I. Polosukhin. 2017. Attention is all you need. *In* Proceedings of the 31st International Conference on Neural Information Processing Systems. Curran Associates Inc., Long Beach, California, USA. 6000–6010.

Wan, N., D. Weinberg, T.Y. Liu, K. Niehaus, E.A. Ariazi, D. Delubac, A. Kannan, B. White, M. Bailey, M. Bertin, N. Boley, D. Bowen, J. Cregg, A.M. Drake, R. Ennis, S. Fransen, E. Gafni, L. Hansen, Y. Liu, G.L. Otte, J. Pecson, B. Rice, G.E. Sanderson, A. Sharma, J. St John, C. Tang, A. Tzou, L. Young, G. Putcha, and I.S. Haque. 2019. Machine learning enables detection of early-stage colorectal cancer by whole-genome sequencing of plasma cell-free DNA. *BMC cancer*. 19:832.

Waterman-Storer, C.M., A. Desai, J.C. Bulinski, and E.D. Salmon. 1998. Fluorescent speckle microscopy, a method to visualize the dynamics of protein assemblies in living cells. *Curr. Biol.* 8:1227-1230.

Wenzel, J., K. Rose, E.B. Haghighi, C. Lamprecht, G. Rauen, V. Freihen, R. Kesselring, M. Boerries, and A. Hecht. 2020. Loss of the nuclear Wnt pathway effector TCF7L2 promotes migration and invasion of human colorectal cancer cells. *Oncogene*. 39:3893-3909.

Wu, Z.Q., T. Brabletz, E. Fearon, A.L. Willis, C.Y. Hu, X.Y. Li, and S.J. Weiss. 2012. Canonical Wnt suppressor, Axin2, promotes colon carcinoma oncogenic activity. *Proceedings of the National Academy of Sciences of the United States of America*. 109:11312-11317.

Yagoda, N., M. von Rechenberg, E. Zaganjor, A.J. Bauer, W.S. Yang, D.J. Fridman, A.J. Wolpaw, I. Smukste, J.M. Peltier, J.J. Boniface, R. Smith, S.L. Lessnick, S. Sahasrabudhe, and B.R. Stockwell. 2007. RAS-RAF-MEK-dependent oxidative cell death involving voltage-dependent anion channels. *Nature*. 447:864-868.

Yi, L., X. Zhong, Z. Chen, Q. Wang, Y. Yan, J. Wang, and X. Deng. 2019. MicroRNA-147b Promotes Proliferation and Invasion of Human Colorectal Cancer by Targeting RAS Oncogene Family (RAP2B). *Pathobiology : journal of immunopathology, molecular and cellular biology*. 86:173-181.

Zhang, M.L., H.T. Wu, W.J. Chen, Y. Xu, Q.Q. Ye, J.X. Shen, and J. Liu. 2020. Involvement of glutathione peroxidases in the occurrence and development of breast cancers. *Journal of translational medicine*. 18:247.

Zhong, J., R. Liu, and P. Chen. 2020. Identifying critical state of complex diseases by single-sample Kullback-Leibler divergence. *BMC genomics*. 21:87.