1

Optimizing telework in an epidemic context: contrasting the infectious and non-communicable diseases perspectives

Competing interests

The authors declare that they have no competing interests.

1. Abstract

Objectives: In a global context of increasing telework, this study explores its health effects, to determine if there is an optimal teleworking frequency during an epidemic. We aim to quantify the relationship between teleworking frequency and both infectious disease (ID) transmission and non-communicable disease (NCD) risk.

Methods: We developed a mathematical model simulating ID transmission and NCD acquisition in a medium-sized company, analysing how different teleworking levels impact workers health. We conducted a rapid literature review to identify potential exposure-response relationships between teleworking and NCD risk and inform this model. We then simulated infection dynamics over a three-month epidemic wave to contrast ID and NCD risks in relation to the extent of telework engagement.

Results: Evidence from the literature showed varying patterns of NCD risk across different teleworking frequencies. Depending on these relationships, we observed that risk may peak at low, intermediate, or high teleworking levels. We demonstrated the existence of a benefit-risk balance between reducing ID transmission and potentially increasing NCD burden.

Conclusions: Our study highlights that the definition of an "optimal" teleworking frequency to maximise health benefits is inconsistent, and depends on the NCD considered. Effective teleworking strategies must consider both ID prevention and the risk of NCDs, and weigh these health impacts accordingly to optimise societal health outcomes.

Keywords: teleworking, infectious diseases, non-communicable diseases, exposure-response relationship, mathematical model

2. Introduction

Teleworking describes work that is fully or partially carried out at a location other than the default place of work, typically at home (1). In the European Union, the proportion of workers who sometimes or usually telework rose from 9% in 2019 before the COVID-19 pandemic, to 12.2% in 2022 (15.8% to 21.3% in France) (2). Recently, several large organisations have switched to a full-remote work organisation, where workers whose job can be performed remotely no longer have an office (3,4). During the same period, the willingness of employees to telework has increased (5,6). Overall, the increasing prevalence of teleworking implies the necessity to understand the impact of this activity on worker health (7).

From the perspective of infectious diseases (IDs), there is a consensus that teleworking in the context of an epidemic reduces the risk of infection (8,9). An important proportion of contacts occur at work, hence reducing these can have a substantial impact on ID incidence (10–14). The value of teleworking against ID transmission was particularly highlighted during the COVID-19 pandemic, during which public health and social measures led to a global transition towards telework across numerous sectors (15). Through these elements, we can establish an exposure-response relationship between teleworking and IDs, whereby an increased teleworking frequency mechanistically leads to a reduction of ID risk.

On the other hand, the shape of this potential exposure-response relationship between teleworking and non-communicable diseases (NCDs) risk remains unclear (16,17). This is notably due to conflicting evidence for the impact of teleworking on NCD risk, within which we can distinguish between physical and mental health. In a well-organised manner, teleworking can enhance the equilibrium between work-related commitments and personal life, diminish road congestion and commuting time, and decrease atmospheric pollutants, all of which indirectly contribute to the improvement of physical and mental well-being (18,19). However,

suboptimal physical settings and workplace design can cause musculoskeletal disorders

(MSDs), ocular strain (20), and an associated psychological burden (21). Other major health

determinants can be affected by teleworking such as obesity, alcohol abuse, physical

inactivity, and tobacco use (22). Combined, these elements suggest that the relationship

between teleworking and NCD risk is not as clear as for IDs. By extension, this implies that

the overall impact of teleworking on health and the potential existence of an optimal

teleworking frequency to maximise health benefits are unclear.

In this study, we aimed to explore the health impacts associated with telework in an epidemic

context, including both ID and NCD. We designed a novel mathematical model simulating the

incidence of an ID amongst employees of a non-specific company. This model accounts for

the impact of teleworking frequency on both the ID transmission and the incidence of NCD.

We parameterised the model by conducting a rapid review to identify exposure-response

relationships that have been quantified between teleworking frequency and physical or

mental health. We then illustrated how different exposure-response functions may lead to

contrasting impacts of teleworking on health.

3. Methods

3.1. Mathematical model

94   To simultaneously capture the relationship between teleworking and both ID and NCD risk,

95   we designed a compartmental deterministic model to represent the population of employees

96   from a non-specific company during an epidemic wave of a SARS-CoV-2-like virus (Figure

97   1a). This model is summarised in the following set of ordinary differential equations driving

98   the dynamics of numbers of individuals in each model compartment:

$$\frac{dS}{dt} = -\lambda \cdot S$$

$$\frac{dE}{dt} = \lambda \cdot S - \sigma \cdot E$$

$$\frac{dI_A}{dt} = p_A \cdot \sigma \cdot E - \gamma_A \cdot I_A$$

$$\frac{dP}{dt} = (1 - p_a) \cdot \sigma \cdot E - \rho \cdot P$$

$$\frac{dI_S}{dt} = \rho \cdot P - \gamma_S \cdot I_S$$

$$\frac{dR}{dt} = \gamma_A \cdot I_A + \gamma_S \cdot I_S$$

We considered that all employees of the company are initially susceptible (S) to the ID and

can become exposed (E), *i.e.* infected but not yet infectious, following a transmission event at

rate λ. Exposed individuals progress through a latent period at rate σ before either becoming

infectious asymptomatic (I_A, with probability p_A), or infectious presymptomatic (P).

Presymptomatic individuals eventually become infectious symptomatic (I_S) at a rate ρ. All

infectious individuals eventually become recovered (R) at a rate which varies depending on

whether they are asymptomatic (γ_A) or symptomatic (γ_S). The total number of employees

N=S+E+I_A+P+I_s+R is assumed constant over the time period of interest.

111   3.1.1.   Rate of infection

Employees can become infected either via contacts with 1) other infectious employees during

working days when they are present in the workplace, 2) contacts with individuals outside the

company during working days when they are teleworking, or 3) contacts with individuals

outside the company during non-working days (Figure 1b). We assumed that five days per

week are working days, and calculated the overall force of infection as a weighted average of

forces of infection in the workplace (λ_W), during telework (λ_T), and on non-working days (λ_C),

as follows:

$$\lambda = \frac{5}{7} \cdot (1 - \alpha) \cdot \lambda_W + \frac{5}{7} \cdot \alpha \cdot \lambda_T + \frac{2}{7} \cdot \lambda_C$$

121   with

$$\lambda_w = \beta \frac{v \cdot I_A + P}{N - I_S}$$

Here, α indicates the average proportion of time spent in telework rather than in the office. If

employees are working from the office, they can be infected at a rate β either by the

proportion of infected asymptomatic (I_A) or presymptomatic (P) employees also present. We

assumed that infected   symptomatic (I_S) individuals are systematically on sick leave, and

hence cannot infect other employees since they are absent from the workplace. We further

assumed that asymptomatic individuals are less infectious than presymptomatic by a factor v

(23).

The term λ_C corresponds to the force of infection which an employee is subjected to outside

the workplace, on a non-working day. We considered that this represents the wider epidemic

in the community, which is independent of the workplace epidemic since it is much larger.

We calibrated λ_C to available incidence data from the second wave of SARS-CoV-2 in France

(01/09/2020 - 01/12/2020) (Figure 1c). On teleworking days, we assumed that employees

would also have contacts with other individuals in the community, although at a lower rate

than on non-working days, hence we defined the force of infection on teleworking days λ_T as

a fraction ε of λ_C.

We calculated the within-company basic reproduction number R_0 using the next generation

matrix method (24) (Supplementary Text 1) from which we derived the per-employee

transmission rate β for presymptomatic individuals in the workplace as

$$\beta = \frac{R_0 \cdot \rho \cdot \gamma_A}{(1 - p_A)\gamma_A + \rho \cdot \upsilon \cdot p_A}$$

Figure 1: Model structure. a) Model diagram. With regards to the infectious disease,

individuals can either be Susceptible (S), Exposed (E), Infected Asymptomatic (I_A),

Presymptomatic (P), Infected Symptomatic (I_S) or Recovered (R). Compartments with the

superscript "C" indicate individuals who will eventually develop a NCD due to teleworking. **b)**

Components of the total force of infection λ. c) Community epidemic curve (bars, left

axis) and corresponding force of infection λ_c (line, right axis).

3.2.2. Rate of non-communicable disease incidence

To represent the incidence of the selected NCD due to telework in this model, we duplicated

the compartments listed above to further stratify individuals according to whether they will

develop a NCD due to teleworking. We considered that individuals in the infected

symptomatic state cannot move to the corresponding compartment, since they do not

telework but are on sick leave. For all other states, the transition towards the NCD status

occurs at an average daily rate defined by the exposure-response function $f(\alpha)$ which

depends on the proportion of telework α. Possible shapes for this function were informed by

the studies identified in our rapid review (22,25,26). To ensure comparability between the

different forms of $f(\alpha)$, we systematically set the baseline value $f(0)$ equal to 8.9 x 10^{-5} per

day. This corresponds to an annual probability of approximately 3.2% (= 1 – exp(-8.9*10^{-}

^{5}*365)) to develop a NCD in absence of teleworking, similar to the Global Burden of Disease

2021 estimates for the cumulative incidence of MSDs and mental disorders (27). Parameter

values for the model are summarised in Table 1.

Table 1: Model parameters.

Name	Symbol	Value	Range explored	Source
Reproduction number	R_0	2.66	2 - 4	(28)
Coefficient of relative infectiousness of asymptomatics	v	0.35	0.1 – 1.27	(23)
Coefficient of relative community force of infection on teleworking days	ε	0.21	0.17 – 0.25 (assumed +/- 20%)	(15)
Progression rate from exposed to infectious (= 1/incubation period)	σ	1/6.57 days^{-1}	1/18.87 - 1/1.80 days^{-1}	(29)
Probability of asymptomatic infection	p_A	0.2	0.17 – 0.25	(23)
Recovery rate from infection for asymptomatics	γ_A	1/5 days^{-1}	N/A	(30)
Progression rate from presymptomatic to symptomatic	ρ	1/1.5 days^{-1}	N/A	(30)

Recovery rate for symptomatics	γ_S	1/5 days^{-1}	N/A	(30)
Baseline incidence rate of non-communicable disease in absence of telework	f(0)	8.9 x 10^{-5} days^{-1}	N/A	(27)
Maximum change in relative risk of non-communicable disease due to telework*	ω	+/- 0.7	+/- 0.05-0.7	This study

170   *This parameter is only used in the sensitivity analysis

173   Model simulations over a 3-month time period were performed using R version 4.2.2. The

174   code for the model and all the analysis presented in this article is included in a GitHub

175   repository (https://github.com/MESuRS-Lab/telework_health).

177   3.2.   Rapid review

178   We conducted a rapid review of research articles investigating quantifiable aspects of

179   telework (e.g. frequency measured in days per month or hours per week) in association with

180   the risk of NCDs, such as mental health disorders and MSDs (31).

182   3.2.1. Search strategy and study selection

Previous reviews on the impact of telework on health have found that the most notable

outcomes are related to behavioural risks, mental health or MSDs (32–34). Therefore, we

restricted our review to the impact of telework on these outcomes.

We searched three databases (Pubmed, Scopus and Google Scholar) to identify original

studies assessing the quantitative association between telework and health. We then

synthesised the evidence regarding the shape and magnitude of the relationship between

telework exposure and the various health outcomes studied.

3.2.2. <u>Inclusion and exclusion criteria</u>

We used the following inclusion and exclusion criteria to select studies:

● Publication type: we included original research articles only.

● Population: we included articles studying workers exposed to some level of telework,

whether they were part of a specific sub-population (such as specific professions) or

not.

● Exposure: we included studies that compared at least two levels of exposure to

telework, in addition to no telework.

● Outcomes: we included studies that reported MSDs, mental health conditions or

behavioural risks as outcomes, and excluded studies reporting only other outcomes.

We did not restrict ourselves to a single specific MSD, mental health condition or

behavioural risk.

We placed no restriction on the date of publication or the language.

3.2.3. <u>Data collection</u>

From the selected studies, we extracted the following information: study years, country,

working population (sector), study design, quantification of telework exposure, the health-

related outcomes examined, and the shape of the observed relationship. Importantly, we

extracted relationships regarding health-related outcomes for any telework level different

from 0 to portray plausible exposure-response functions. To inform the model, we only

considered exposure-response relationships based on significant ($p < 0.05$) values.

4. Results

4.1. Studies on NCD and telework identified to inform the model

From our rapid review, we identified only three studies that met our selection criteria and presented exposure-response relationships between teleworking and NCD risk which could inform the mathematical model (Table 2). Of those studies, one focused on work-related outcomes (25), one on physical health (26), and one on health determinants (including behavioural changes) impacting both mental and physical health (22). Importantly, two of the identified studies were entirely conducted before the COVID-19 pandemic (22,25).

The study that focused on work-related outcomes notably examined depressive symptoms and self-rated health (25). The results of this longitudinal study conducted on United States employees highlighted exposure-response relationships for several outcomes. A U-shaped (US) relationship was found between the number of teleworking days per week and depressive symptoms, with a higher risk among non-teleworkers (0 days per week) and those with a higher frequency of telework (5 days per week). The association between number of teleworking days per week and self-rated health also displayed a US relationship.

For physical health, the study we identified only focused on lower back pain as a MSD-related outcome (26). The findings from this cross-sectional study conducted in Japan revealed an inverted U-shaped (IU) exposure-response relationship between teleworking frequency and lower back pain, with the highest risk in case of intermediate teleworking frequencies (2-3 days/week).

Finally, the last study described exposures related to both mental and physical health such as alcohol abuse or physical inactivity (22). This longitudinal study was conducted in the United States on insurance company employees. Retrieved results (significant values at $p <$

239   0.05) suggested a broadly decreasing relationship between teleworking intensity and risks of

240   alcohol abuse and tobacco use. For the Edington risk score (a summary score accounting for

241   several risk factors), we observed an L-shaped (LS) relationship with increased telework

242   intensity, with an initial rapid decrease in risk at low telework frequencies followed by a more

243   stable relationship at higher frequencies.

 Table 2: Summary of identified studies on the exposure-response relationship

 between teleworking frequency and non-communicable disease (NCD) risk. Here, we

 classified (or retrieved) the shape exposure-response relationships into five types: U-shaped

 (US), inverted U-shaped (IU), L-shaped (LS), broadly decreasing (D) and broadly increasing

 (I). Relationships in italic are tested in our mathematical model in the next section. "Baseline

 level" values are values from the studies for groups not exposed to telework (0 days per

 week).

Article	Study period & country	Exposure to telework	Health outcome	Health outcome measurement & shape of relationship	Baseline level
Chen et al., 2023 (25)	2018-2019, USA	Days per week (3 groups): • 1-4 • 5	**Self-reported (range: 0-10)**		
			Depressive symptoms	*US: 1.71* ; 2.08**	*2.33*
			Self-rated health	US: 6.01* ; 5.96*	5.69
Matsugaki et al., 2021 (26)	2020, Japan	Days per week (3 groups): • 1 • 2-3 • 4+	**Self-reported symptom (% of Yes)**		
			Lower back pain	*IU: 52.8%* ; 53.2%* ; 47.5%**	*49.9%*
Henke et al., 2016 (22)	2010-2011, USA	Hours per month (4 groups) : • 8- • 9-32	**Regression coefficient and percentage at baseline**		
			Alcohol abuse (at least 2 or 3 drinks per day)	D: -0.061 ; -0.503 ; -0.024 ; -1.423*	4.9%
			Physical	US: -0.189 ; -0.249* ; -0.140 ; -	40.4%

	inactivity (<3 days of exercise per week)	0.015	
	Tobacco use (yes/no)	D: -0.100 ; -0.175 ; -0.426* ; -0.263	12.3%
	Edington risk (>5 health risk factors)	*LS: -0.571* ; -0.997* ; -0.321 ; -1.233**	*5.7%*
	Obesity risk	US: -0.150 ; -0.146 ; -0.202 ; -0.113	33.1%
	Depression	D: -0.083 ; -0.056 ; -0.202 ; -0.151	16.4%
	Stress	I: 0.033 ; -0.037 ; 0.040 ; 0.137	29.5%
	Poor nutrition	US: -0.017 ; -0.110 ; 0.009 ; 0.194	86.2%

- 33-72
- 73+

251 *Reported p value <0.05 in the referenced study

4.2. Impact of varying teleworking frequency on health outcomes

Our review shows that the exposure-response relationship between teleworking and NCD

risk may vary depending on the mental or physical health focus, as well as on other factors.

This is bound to impact the level at which teleworking best prevents both ID and NCD. Based

on our literature review and the statistically significant relationships we identified, we

explored three relationships between telework frequency and the risk of NCD in our model: L-

shaped (LS, for Edington risk), U-shaped (US, for depressive symptoms), and inverted U-

shaped (IU, for lower back pain). In the model, the corresponding exposure-response

functions *f(α)* giving the daily rate at which individuals will develop a NCD for a given

frequency of teleworking *α* were parameterized using data from the three studies analysed in

the rapid review (22,25,26) (values in italic in Table 2, see Supplementary Text 2 for details).

Importantly, since we used values from the literature to parameterize these functions, the

strengths of the association and therefore their upper and lower bounds are different.

Implementing telework throughout the epidemic wave substantially reduced the cumulative

incidence of ID amongst employees (yellow curve), with greater reductions at higher

teleworking frequencies and no substantial differences between frequencies greater than 0.8,

i.e. 4/5 days per week (Figure 2a). However, the cumulative incidence of NCD at different

teleworking frequencies varied depending on the assumed exposure-response relationship

function, with the predicted peak incidence of NCD occurring either at low (LS, US) or

intermediate (IU) teleworking frequencies (Figure 2b).

275
Figure 2: Epidemic dynamics and cumulative incidence of non-communicable disease

(NCD) at varying telework frequencies. Model simulations over 3 months using a telework

frequency $\alpha = 0$ (no telework), 0.2 (1 day in 5), 0.4 (2 days), 0.6 (3 days), 0.8 (4 days) and 1

(full telework), for **a)** changes in the number of susceptible (S), exposed (E),

asymptomatically infected (Ia), presymptomatic (P), symptomatically infected (Is) and

recovered (R) employees, and **b)** the relative risk for employees to eventually develop a

NCD, in case of a L-shaped (LS, circles), U-shaped (US, triangles) or inverted U-shaped (IU,

squares) exposure-response relationship between telework and NCD risk.

4.3. Impact of varying teleworking start date on health outcomes

The impact of teleworking on cumulative disease incidence further varied depending on the timing of its implementation from the start of the epidemic wave (Figure 3). Since we performed simulations in the context of an epidemic wave, implementing teleworking too late prevented any significant reduction of the ID cumulative incidence (orange lines). If we set an arbitrary objective to reduce this incidence below 50% of the baseline value obtained without teleworking, at least two days of teleworking (frequency of 0.4) were necessary to reach this target, with teleworking implemented within the first 40 days of the epidemic (solid orange lines).

Regarding the change in NCD risk caused by teleworking, as expected, early implementation of teleworking systematically led to a higher change in cumulative incidence (blue lines). However, the nature of this change depended on the assumed exposure-response function shape, upper and lower bounds, and on the teleworking frequency. For example, in the case of the L-shaped relationship for Edington risk parameterised according to our review (first row), the cumulative incidence of NCD was lowest when teleworking was implemented at a frequency of 1 on the first day (last column), while for the U-shaped relationship for depressive symptoms (second row) the lowest incidence was achieved at a frequency of 0.6 (fourth column). On the other hand, teleworking led to an increase in cumulative NCD incidence (relative incidence > 100%) when considering the inverted U-shaped relationship with lower back pain (last row), particularly at intermediate teleworking frequencies (0.4-0.6, third and fourth columns). In that case, slightly delaying the implementation of teleworking to avoid increasing the risk of NCD while still having an impact on ID could be better, and/or implementing 100% teleworking, which would not increase the risk of NCD while keeping the ID risk at a minimum. For the US relationship, timing is also important because early teleworking implementation for intermediate frequencies reduces both health risks below their respective targets, which is not the case for lower or higher frequencies. Additionally,

since there is no major difference for ID relative risk between frequencies of 0.8 and 1, a

teleworking frequency of 0.8 would be preferable to achieve strong reductions in both NCD

and ID risks.

The contrasting relationship between teleworking and ID versus NCD risks implies that

reducing both incidences simultaneously may not always be feasible, depending on the

characteristics of the NCD. In our illustrative example here, where we aimed to reduce ID

incidence by at least 50% and NCD incidence by at least 10%, we observed only limited

conditions where teleworking could simultaneously reduce both diseases incidences below

these targets (Figure 3, grey shaded areas). The definition of an "optimal" teleworking

frequency to improve health is therefore inconsistent and will vary depending on (*i*) the

observed exposure-response relationship between teleworking and health outcomes, and (*ii*)

the relative importance granted to both ID and NCD risks when defining target thresholds.

Figure 3: Impact of teleworking timing and frequency on the relative cumulative incidences of infectious (ID) and non-communicable diseases (NCD) using observed exposure-response functions. Cumulative incidence corresponds to the number of employees infected over the three months of the simulated epidemic. Here, we represent the cumulative incidences of ID and NCD relative to the cumulative incidences predicted by each model without teleworking (first column). For example, the lowest incidence of NCD (blue line) for the L-shaped function (first row) is when teleworking is implemented with a frequency of 1 (5 days a week) on day 0 (last column), while for the U-shaped exposure-response function (second row) this occurs at a frequency of 0.6 (fourth column). The grey dashed line indicates 100%, i.e. the baseline incidences. Lines are solid when the relative incidences of ID and NCD are respectively lower than 50% and 90% (chosen arbitrarily as examples), and faded otherwise. The shaded grey areas indicate conditions where both relative cumulative

incidences are below the defined targets of 50% and 90% (for ID and NCD incidence,
respectively).

4.4. Exploring different exposure-response relationships between
teleworking and NCDs

While the exposure-response functions used in the previous section were directly informed
by values we identified in the literature, their upper and lower bounds varied, which
introduces a bias when comparing the incidence of different NCDs since the differences are
not only attributable to the shape of the functions. For example, the greatest change in
relative risk of NCD for the IU function (lower back pain) was approximately +6%, while for
the LS function (Edington risk) this reached -70%. We therefore reproduced our analysis
using five theoretical functions with shapes corresponding to those we identified in our review
(see Supplementary Text 2 for details, and Supplementary Figures 2 and 3 for the shapes of
the functions), but each with the same greatest absolute change in relative risk of NCD (+/-
70%). The results obtained underline how the shape of these functions alone affects the
optimal telework frequency and implementation date, independently of upper/lower bounds
(Figure 4). This analysis also highlights scenarios where the relative increase in NCD
incidence can be equivalent to the relative decrease in ID incidence, as can be seen for the
theoretical IU function at a telework frequency of 0.6, and the theoretical linear increasing
function at a frequency of 1 (Figure 4, rows 3 and 4).

360
Figure 4: Impact of teleworking timing and frequency on the relative cumulative

incidences of infectious (ID) and non-communicable diseases (NCD) using theoretical

exposure-response functions. Cumulative incidence corresponds to the number of

employees infected over the three months of the simulated epidemic. Here, we represent the

cumulative incidences of ID and NCD relative to the cumulative incidences predicted by each

model without teleworking (first column). The grey dashed line indicates 100%, i.e. the

baseline incidences. Lines are solid when the relative incidences of ID and NCD are

respectively lower than 50% and 90% (chosen arbitrarily as examples), and faded otherwise.

The shaded grey areas indicate conditions where both relative cumulative incidences are

below the defined targets of 50% and 90% (for ID and NCD incidence, respectively).

4.5.   Sensitivity analysis

We calculated partial correlation coefficients to examine the impact of model parameter

values on cumulative incidence of ID and NCD over the period, assuming a telework

frequency of 0.5 (Supplementary Figure 3). The only differences in correlation coefficients

that we observed between scenarios of exposure-response relationships affected one

parameter, the maximum change in relative risk of NCD due to telework (ω); as expected,

this parameter was strongly positively correlated (coefficient $\approx$ 1) with NCD incidence for IU

and LI curves, and strongly negatively correlated (coefficient $\approx$ -1) with NCD incidence for LD,

LS and US curves.

Regardless of which exposure-response function was used, the relative infectiousness of

asymptomatic individuals (v) was only slightly negatively correlated with ID cumulative

incidences (coefficient $\approx$ 0.2). The coefficient of relative community force of infection on

teleworking days (ε) was only slightly positively correlated with ID incidence (coefficient $\approx$

0.2). The reproduction number (R_0) and the progression rate from exposed to infectious (σ)

were strongly positively correlated with ID incidence (coefficient > 0.7). These two

parameters were negatively correlated with NCD incidence (coefficient $\leq$ -0.5), which is

expected since we assume that infected individuals cannot develop a NCD during the period

when they are infected and symptomatic. Inversely, the proportion of asymptomatic infections

(p_A) was slightly positively correlated with NCD incidence (coefficient $\approx$ 0.2), since it leaves a

greater proportion of infected individuals at risk of developing a NCD in parallel due to

reduced sick leaves.

5. Discussion

Recent years have seen an unprecedented increase in teleworking frequency in many countries worldwide. In this study, we aimed to determine if there existed an optimal teleworking frequency to maximise health benefits, by designing a mathematical model accounting for both ID transmission and NCD incidence in employees of a non-specific company. To parameterise this model, we first reviewed the evidence on the consequences this may have in terms of NCDs for teleworking employees, notably underlining potential impacts on mental health and MSDs. Our rapid review uncovered a wide variety of possible exposure-response relationships between teleworking intensity and NCD risk. By incorporating this data in our model, we showed that optimal frequency and timing of implementation of teleworking during an epidemic wave could vary widely. For instance, for health impacts associated with teleworking through a L-shaped function with a strength of association such as the one we identified for Edington risk (22), rapid and wide implementation of teleworking during the first few days of an epidemic can reduce both ID and NCD incidences. On the other hand, for a U-shaped relationship with parameters such as the one we identified for depression (25), intermediate (3-4 days per week) teleworking frequencies may be more optimal to maximise health benefits, while for inverted U-shaped relationships with parameters as observed for lower back pain (26), it may be necessary to weigh the increased NCD risk attributable to teleworking against the decreased ID risk. Importantly, both the shape and the upper/lower bound of these exposure-response functions must be taken into consideration when contrasting ID and NCD risk.

5.1. Implications

Through our rapid review, we confirmed that telework may impact both mental and physical health, although we identified several important knowledge gaps based on the selected

studies. First, there is a lack of uniformity across the measurements of health outcomes

(clinical diagnoses and declarative statements) explored in relation with teleworking, ranging

from physical to mental health (depression, anxiety, burnout). Second, exposures are poorly

characterised. Exposure to teleworking is not coded in a standardised manner; depending on

the study, it may be expressed in terms of number of days per week, or number of hours per

day, with different categories. Third, study contexts are heterogenous. Studies encompassing

short- and long-term perspectives, within both pandemic and non-pandemic settings, yield

outcomes potentially contingent on the phase of the COVID-19 pandemic (35). Confounding

factors (such as the region, pre-existing comorbidities or professional status) are not well

described or controlled for, and differ across studies. Fourth, very few studies met our

inclusion criteria, and we found no longitudinal study that focused on MSDs, leading to a lack

of quality in evidence especially regarding the temporal relationship between exposure and

MSDs. Our findings are in agreement with two recent systematic reviews that also outlined

the low quality of evidence regarding health impacts of teleworking (36,37).

Overall, the identified studies led to divergent results on the shape of the exposure-response

function, depending on the considered health outcome (U-shaped, inverted U-shaped, L-

shaped, broadly decreasing, and broadly increasing curves). In addition, even for the same

health outcome, reported impacts of teleworking could be heterogeneous. For example,

Henke and colleagues found a broadly decreasing relationship between hours/month of

telecommuting and depression, while Chen and colleagues found a U-shaped relationship.

This conflicting evidence may be due to modifying effects as demonstrated by one study

according to which organisational factors within a company could alter the shape/direction of

the relationship, depending on the COVID-19 wave (35).

Our mathematical model suggests that the shape and range of variation of the exposure-

response relationship function between teleworking and NCD risk may influence the optimal

teleworking frequency. For a given target of incidence reduction, the choice of teleworking

frequency and the timing of its implementation as an intervention during an epidemic wave

varies. The implications in terms of performances and costs are also different as we expect

sick-leaves due to IDs to be numerous but short while sick-leaves due to NCDs would be

fewer but longer. Furthermore, MSDs and psychological disorders are both heterogeneous in

severity, which makes quantitative assessments difficult. Finally, implications may largely

depend on the professional sector, for example, workers in healthcare settings or densely-

populated workplaces are expected to be more exposed to IDs, while desk-based workers

will be more exposed to NCDs.

5.2. Limitations

As shown in our rapid review, the relationship between telework frequency and the risk of

developing common NCDs such as lower back pain or psychological distress in the long-term

depends on many unmeasured individual and environmental characteristics. Instead of

accounting for all these specificities, we illustrated with our model the impact of different

relationships considering one average risk for all individuals. The lack of significant

associations between telework and some NCDs is also a source of uncertainty in the

exposure-response relationships used in the model, as shown in Table 2. Alternative

modelling strategies could be used to integrate individual heterogeneity, but this would

require additional evidence regarding the distribution of exposures and risks based on

individual characteristics (age, gender, job…) that is not yet available in the literature.

The majority of evidence regarding telework arises from the COVID-19 pandemic. In this

context, telework was frequently unplanned and imposed on individuals, which does not

necessarily reflect conditions where teleworking would be planned and adapted at an

individual level (7). In addition, relatively few countries were represented in these studies,

while we would expect the impact of teleworking to vary across regions, between urban and

rural settings and living conditions. Finally, the impact of teleworking on health likely depends

on the socio-professional categories considered and its desirability.

In our illustrative example, we simulated the impact of telework policies in the context of the

second wave of the SARS-CoV-2 pandemic. Thus, the results should be interpreted in the

context of an emerging ID for which no pharmaceutical intervention (e.g., vaccination) is

available, and for which no behavioural change among workers is observed apart from the

telecommuting policy decided by the employer. We expect different results in case these two

assumptions are not met.

Finally, we assumed homogeneous mixing within the company, whereby all employees could

be in contact, without considering more complex work organisations. Similarly, we

considered a simple telework policy according to which a fixed percentage of the total

workforce in the company is teleworking every day, but more refined policies have been

implemented during the pandemic, such as rotating telework (13). In these strategies,

employees are evenly distributed in groups that alternate on a daily or weekly basis. This is

expected to reduce the overall number of contacts per individual and potentially the risk of

transmission, whilst maintaining a reduced average frequency of telework.

5.3.   Perspective and future directions

495
Several options could be considered to mitigate the NCD risk among teleworking individuals.

First, sedentary behaviours, prolonged computer use, and poor ergonomics during telework

have been linked to increased musculoskeletal disorders (MSDs), particularly affecting the

neck, shoulders, and lower back. To mitigate MSD risks, studies recommend ergonomic

workspaces, regular posture changes, and proactive telework preparations by companies,

ensuring adequate support and proper work conditions. Second, telework has mixed effects

on mental health, enhancing positive emotions, job satisfaction, and organizational

commitment while reducing emotional exhaustion, but also contributing to anxiety, stress,

fatigue, and social isolation. Negative effects could be mitigated by measures like technical

support, flexible hours, social communication, and health interventions, with part-time

telework offering benefits for work-life balance and social relationships.

In practice, the optimal frequency and implementation timing of telework will depend on the

relative importance given to both NCDs and ID. The decision on respective weights given to

the ID and NCD incidence should consider the different timelines at which these diseases

occur: typically, short-term for IDs and mid- to long-term for NCDs. IDs represent a

substantial socioeconomic burden (38), which can justify the implementation of teleworking

as an intervention to reduce disease spread. Part of this burden is related to sick leave (39),

which can lead companies to act in order to minimise disease incidence among their

employees. However, when designing a teleworking policy, deciders need to account for

feasibility, legal and ethical criteria. For instance, teleworkers need to have access to

telework equipment, which was not always straightforward during the COVID-19 pandemic

(40).

Future studies should better characterize the distribution of telework among population and

identify its association with NCDs and mental health outcomes based on individual

characteristics (age, gender, job…). It is also important to explore how telework dynamics

can influence the transmission of infectious diseases among workers.

6. Conclusions

In our rapid review, we identified three studies, two being longitudinal. This very low number underlines the need for more data to monitor the health impacts of teleworking. In particular, further studies should characterise the relationship between telework frequency and NCDs. To this end, it is crucial to collect data outside the context of the COVID-19 pandemic, since additional stressors during this period may have modulated the relationship between telework and health. In addition, the mechanisms by which teleworking impacts mental and physical health should be better characterised (e.g. unadapted workstation for MSDs). Lastly, further efforts are needed to identify the individual factors affecting the exposure-response relationships both during and outside of epidemic contexts.

Our innovative approach, which attempts to combine short- and longer-term consequences of teleworking in a unique framework, could serve as a basis to develop tools for employers and policymakers. Such tools could be used to quantify the impact of telework on employee health and identify optimal telework strategies to limit health adverse events and improve employee well-being.

7. References

1. International Labour Organization. COVID-19: Guidance for labour statistics data collection: Defining and measuring remote work, telework, work at home and home-based work [Internet]. 2020 [cited 2023 Aug 24]. Available from: https://www.ilo.org/global/statistics-and-databases/publications/WCMS_747075/lang--en/index.htm

2. Eurostat. Employed persons working from home as a percentage of the total employment, by sex, age and professional status [Internet]. 2023 [cited 2023 Aug 29]. Available from: https://ec.europa.eu/eurostat/databrowser/view/lfsa_ehomp/default/table?lang=en

3. Chocron V. Télétravail «total» chez Boursorama : les salariés commencent à quitter Paris pour la province. Le Monde [Internet]. 2021 Sep 28; Available from: https://www.lemonde.fr/economie/article/2021/09/28/teletravail-total-chez-boursorama-les-salaries-commencent-a-quitter-paris-pour-la-province_6096226_3234.html

4. Hyder B. Return & Remote: How Salesforce Brings Employees Together for Modern Work [Internet]. 2023 [cited 2023 Aug 29]. Available from: https://www.salesforce.com/news/stories/return-remote-how-salesforce-brings-employees-together-for-modern-work/

5. OECD. Teleworking in the COVID-19 pandemic: Trends and prospects [Internet]. OECD; 2021 [cited 2023 Oct 31]. Available from: https://www.oecd.org/coronavirus/policy-responses/teleworking-in-the-covid-19-pandemic-trends-and-prospects-72a416b6/#biblio-d1e1079

6. Weber C, Golding SE, Yarker J, Lewis R, Ratcliffe E, Munir F, et al. Future Teleworking Inclinations Post-COVID-19: Examining the Role of Teleworking Conditions and Perceived Productivity. Front Psychol [Internet]. 2022 [cited 2023 Oct 31];13. Available from: https://www.frontiersin.org/articles/10.3389/fpsyg.2022.863197

7. Bouziri H, Smith DRM, Descatha A, Dab W, Jean K. Working from home in the time of COVID-19: how to best preserve occupational health? Occup Environ Med. 2020 Jul;77(7):509–10.

8. Fisher KA. Telework Before Illness Onset Among Symptomatic Adults Aged ≥18 Years With and Without COVID-19 in 11 Outpatient Health Care Facilities — United States, July 2020. MMWR Morb Mortal Wkly Rep [Internet]. 2020 [cited 2023 Nov 7];69. Available from: https://www.cdc.gov/mmwr/volumes/69/wr/mm6944a4.htm

9. Galmiche S, Charmet T, Schaeffer L, Paireau J, Grant R, Chény O, et al. Exposures associated with SARS-CoV-2 infection in France: A nationwide online case-control study. Lancet Reg Health - Eur. 2021 Jun 7;7:100148.

10. Ahmed F, Zviedrite N, Uzicanin A. Effectiveness of workplace social distancing measures in reducing influenza transmission: a systematic review. BMC Public Health. 2018 Apr 18;18:518.

11. Hill EM, Atkins BD, Keeling MJ, Dyson L, Tildesley MJ. A network modelling approach to assess non-pharmaceutical disease controls in a worker population: An application to SARS-CoV-2. PLoS Comput Biol. 2021 Jun 16;17(6):e1009058.

12. Jarvis CI, Gimma A, van Zandvoort K, Wong KLM, Edmunds WJ. The impact of local and
national restrictions in response to COVID-19 on social contacts in England: a
longitudinal natural experiment. BMC Med. 2021 Feb 19;19:52.

13. Mauras S, Cohen-Addad V, Duboc G, Tour MD la, Frasca P, Mathieu C, et al. Mitigating
COVID-19 outbreaks in workplaces and schools by hybrid telecommuting. PLOS Comput
Biol. 2021 Aug 26;17(8):e1009264.

14. Mousa A, Winskill P, Watson OJ, Ratmann O, Monod M, Ajelli M, et al. Social contact
patterns and implications for infectious disease transmission – a systematic review and
meta-analysis of contact surveys. Rodriguez-Barraquer I, Serwadda DM, editors. eLife.
2021 Nov 25;10:e70294.

15. Bosetti P, Huynh BT, Abdou AY, Sanchez M, Eisenhauer C, Courtejoie N, et al.
Lockdown impact on age-specific contact patterns and behaviours, France, April 2020.
Eurosurveillance. 2021 Dec 2;26(48):2001636.

16. Nowrouzi-Kia B, Haritos AM, Long BZS, Atikian C, Fiorini LA, Gohar B, et al. Remote
work transition amidst COVID-19: Impacts on presenteeism, absenteeism, and worker
well-being—A scoping review. PLOS ONE. 2024 Jul 18;19(7):e0307087.

17. Perelman J, Serranheira F, Castanheira F, Raposo JF, Aguiar P, Neves P, et al.
Teleworking: does it make workers healthier and productive? A cross-sectional study on
a Southern European population. BMC Public Health. 2024 Jul 20;24(1):1946.

18. OECD. Assessing teleworking strategies for local development: a framework proposal
[Internet]. Paris; 2022 [cited 2023 Aug 31]. Available from:
https://www.oecd.org/cfe/leed/Assessing-teleworking-strategies-for-local-development-
PAT.pdf

19. World Health Organization and the International Labour Organization. Healthy and safe
telework: technical brief [Internet]. Geneva; 2021. Available from:
https://www.who.int/publications/i/item/9789240040977

20. Tezuka M, Nagata T, Saeki K, Tsuboi Y, Fukutani N. Association Between Abrupt
Change to Teleworking and Physical Symptoms During the Coronavirus Disease 2019
(COVID-19) Emergency Declaration in Japan. J Occup Environ Med. 2022 Jan
1;64(1):1–5.

21. De Macêdo TAM, Cabral ELDS, Silva Castro WR, De Souza Junior CC, Da Costa Junior
JF, Pedrosa FM, et al. Ergonomics and telework: A systematic review. Work. 2020 Sep
17;66(4):777–88.

22. Henke RM, Benevent R, Schulte P, Rinehart C, Crighton KA, Corcoran M. The Effects of
Telecommuting Intensity on Employee Health. Am J Health Promot. 2016 Nov
1;30(8):604–12.

23. Buitrago-Garcia D, Egli-Gany D, Counotte MJ, Hossmann S, Imeri H, Ipekci AM, et al.
Occurrence and transmission potential of asymptomatic and presymptomatic SARS-
CoV-2 infections: A living systematic review and meta-analysis. PLOS Med. 2020 Sep
22;17(9):e1003346.

24. Diekmann O, Heesterbeek JAP, Metz JAJ. On the definition and the computation of the
basic reproduction ratio R0 in models for infectious diseases in heterogeneous
populations. J Math Biol. 1990 Jun 1;28(4):365–82.

25. Chen Y, Weziak-Bialowolska D, Lee MT, Bialowolski P, Cowden RG, McNeely E, et al.
Working from home and subsequent work outcomes: Pre-pandemic evidence. PloS One.
2023;18(4):e0283788.

26. Matsugaki R, Muramatsu K, Tateishi S, Nagata T, Tsuji M, Hino A, et al. Association
Between Telecommuting Environment and Low Back Pain Among Japanese
Telecommuting Workers: A Cross-Sectional Study. J Occup Environ Med. 2021
Dec;63(12):e944–8.

27. Institute for Health Metrics and Evaluation (IHME). Global Burden of Disease
Collaborative Network (GBD 2021) [Internet]. 2024. Available from:
https://vizhub.healthdata.org/gbd-results/

28. Dhungel B, Rahman MS, Rahman MM, Bhandari AKC, Le PM, Biva NA, et al. Reliability
of Early Estimates of the Basic Reproduction Number of COVID-19: A Systematic
Review and Meta-Analysis. Int J Environ Res Public Health. 2022 Jan;19(18):11613.

29. Wu Y, Kang L, Guo Z, Liu J, Liu M, Liang W. Incubation Period of COVID-19 Caused by
Unique SARS-CoV-2 Strains: A Systematic Review and Meta-analysis. JAMA Netw
Open. 2022 Aug 22;5(8):e2228008.

30. Kissler SM, Fauver JR, Mack C, Olesen SW, Tai C, Shiue KY, et al. Viral dynamics of
acute SARS-CoV-2 infection and applications to diagnostic and public health strategies.
Riley S, editor. PLOS Biol. 2021 Jul 12;19(7):e3001333.

31. Grant MJ, Booth A. A typology of reviews: an analysis of 14 review types and associated
methodologies. Health Inf Libr J. 2009 Jun;26(2):91–108.

32. Beckel JLO, Fisher GG. Telework and Worker Health and Well-Being: A Review and
Recommendations for Research and Practice. Int J Environ Res Public Health. 2022 Mar
24;19(7):3879.

33. Furuya Y, Nakazawa S, Fukai K, Tatemichi M. Health impacts with telework on workers:
A scoping review before the COVID-19 pandemic. Front Public Health. 2022 Oct
31;10:981270.

34. Liu W, Xu Y, Ma D. Work-Related Mental Health Under COVID-19 Restrictions: A Mini
Literature Review. Front Public Health. 2021;9:788370.

35. Wels J, Wielgoszewska B, Moltrecht B, Booth C, Green MJ, Hamilton OK, et al. Home
working and social and mental wellbeing at different stages of the COVID-19 pandemic in
the UK: Evidence from 7 longitudinal population surveys. Thorley J, editor. PLOS Med.
2023 Apr 27;20(4):e1004214.

36. Fadel M, Bodin J, Cros F, Descatha A, Roquelaure Y. Teleworking and Musculoskeletal
Disorders: A Systematic Review. Int J Environ Res Public Health. 2023 Mar
11;20(6):4973.

37. Lunde LK, Fløvik L, Christensen JO, Johannessen HA, Finne LB, Jørgensen IL, et al.
The relationship between telework from home and employee health: a systematic review.
BMC Public Health. 2022 Jan 7;22(1):47.

38. Samsudin EZ, Yasin SM, Ruslan NH, Abdullah NN, Noor AFA, Hair AFA. Socioeconomic
impacts of airborne and droplet-borne infectious diseases on industries: a systematic
review. BMC Infect Dis. 2024 Jan 16;24(1):93.

39. Smith DRM, Jijón S, Oodally A, Shirreff G, Bouziad KA, Ante-Testard PA, et al. Sick
leave due to COVID-19 during the first pandemic wave in France, 2020. Occup Environ
Med. 2023 May 1;80(5):268–72.

40. Belzunegui-Eraso A, Erro-Garcés A. Teleworking in the Context of the Covid-19 Crisis.
Sustainability. 2020 Jan;12(9):3662.

8. List of abbreviations

ID: Infectious disease

IU: Inverted U-shaped

LD: Linear decrease

LI: Linear increase

LS: L-shaped

MSD: Musculoskeletal disorder

NCD: Non-communicable disease

US: U-shaped