Optimizing telework in an epidemic context: contrasting the infectious and non-communicable diseases perspectives

Léo Moutet † [1], Quentin J. Leclerc † [1,2,3,4], Maylis Layan[1,2,3,4], Karim Aït Bouziad[1], William Dab[1], Paul Henriot[1,2], Elise Hodbert[1,5], Narimène Louati[1], Aurélie Maurin[1,2], Frédérique Thonon[1], Sylvie Znaty[1], Mohamed Ben Halima[1], Kévin Jean[1,2,6], Laura Temime[1,2]

† These authors contributed equally

† Correspondence to: leo.moutet@gmail.com and quentin.leclerc@pasteur.fr

[1] Laboratoire Modélisation, Épidémiologie et Surveillance des Risques Sanitaires (MESuRS), Conservatoire National des Arts et Métiers, 75003 Paris, France

[2] Institut Pasteur, Conservatoire National des Arts et Métiers, Unité PACRI, 75015 Paris, France

[3] Institut Pasteur, Université Paris Cité, Epidemiology and Modelling of Bacterial Escape to Antimicrobials (EMEA), 75015 Paris, France

[4] INSERM, Université Paris-Saclay, Université de Versailles St-Quentin-en-Yvelines, Team Échappement aux Anti-infectieux et Pharmacoépidémiologie U1018, CESP, 78000 Versailles, France

[5] Cibles et médicaments des infections et de l'immunité, IICiMed, UR 1155, Nantes Université, Nantes, France)

[6] Eco-Evolutionary Mathematics team, IBENS, École Normale Supérieure, CNRS, INSERM, Université Paris Science & Lettres, Paris, France

1. Abstract

Objectives: In a global context of increasing telework, this study explores its health effects, to determine if there is an optimal teleworking frequency during an epidemic. We aim to quantify the relationship between teleworking frequency and both infectious disease (ID) transmission and non-communicable disease (NCD) risk.

Methods: We developed a mathematical model simulating ID transmission and NCD acquisition in a medium-sized company, analysing how different teleworking levels impact workers health. We conducted a rapid literature review to identify potential exposure-response relationships between teleworking and NCD risk and inform this model. We then simulated infection dynamics over a three-month epidemic wave to contrast ID and NCD risks in relation to the extent of telework engagement.

Results: Evidence from the literature showed varying patterns of NCD risk across different teleworking frequencies. Depending on these relationships, we observed that risk may peak at low, intermediate, or high teleworking levels. We demonstrated the existence of a benefit-risk balance between reducing ID transmission and potentially increasing NCD burden.

Conclusions:. Based on current scientific knowledge, no single optimal teleworking frequency can be identified, as the optimum varies depending on the NCD outcome considered. Our study highlights the need for stronger evidence to estimate robust exposure-response functions linking teleworking frequency and NCDs, and ultimately to inform prevention strategies for both infectious and NCD risks in an epidemic context.

Keywords: teleworking, infectious diseases, non-communicable diseases, exposure-response relationship, mathematical model

2. Introduction

Teleworking describes work that is fully or partially carried out at a location other than the default place of work, typically at home (1). In the European Union, the proportion of workers who sometimes or usually telework rose from 9% in 2019 before the COVID-19 pandemic, to 12.2% in 2022 (15.8% to 21.3% in France) (2). Recently, several large organisations have switched to a full-remote work organisation, where workers whose job can be performed remotely no longer have an office (3,4). During the same period, the willingness of employees to telework has increased (5,6). Overall, the increasing prevalence of teleworking implies the necessity to understand the impact of this activity on worker health (7).

From the perspective of infectious diseases (IDs), there is a consensus that teleworking in the context of an epidemic reduces the risk of infection (8,9). An important proportion of contacts occur at work, hence reducing these can have a substantial impact on ID incidence (10–14). The value of teleworking against ID transmission was particularly highlighted during the COVID-19 pandemic, during which public health and social measures led to a global transition towards telework across numerous sectors (15). Through these elements, we can establish an exposure-response relationship between teleworking and IDs, whereby an increased teleworking frequency mechanistically leads to a reduction of ID risk.

On the other hand, the shape of this potential exposure-response relationship between teleworking and non-communicable diseases (NCDs) risk remains unclear (16,17). This is notably due to conflicting evidence for the impact of teleworking on NCD risk, within which we can distinguish between physical and mental health. In a well-organised manner, teleworking can enhance the equilibrium between work-related commitments and personal life, diminish road congestion and commuting time, and decrease atmospheric pollutants, all of which indirectly contribute to the improvement of physical and mental well-being (18,19). However,

suboptimal physical settings and workplace design can cause musculoskeletal disorders

(MSDs), ocular strain (20), and an associated psychological burden (21). Other major health

determinants can be affected by teleworking such as obesity, alcohol abuse, physical

inactivity, and tobacco use (22). Combined, these elements suggest that the relationship

between teleworking and NCD risk is not as clear as for IDs. By extension, this implies that

the overall impact of teleworking on health and the potential existence of an optimal

teleworking frequency to maximise health benefits are unclear. In this study, we define as

"optimal" a teleworking frequency that would yield the lowest possible number of COVID-19

cases, and the lowest possible number of NCD cases.

In this study, we aimed to explore the health impacts associated with telework in an epidemic

context, including both ID and NCD. We designed a novel mathematical model simulating the

incidence of an ID amongst employees of a non-specific company. This model accounts for

the impact of teleworking frequency on both the ID transmission and the incidence of NCD.

We parameterised the model by conducting a rapid review to identify exposure-response

relationships that have been quantified between teleworking frequency and physical or

mental health. We then illustrated how different exposure-response functions may lead to

contrasting impacts of teleworking on health.

3. Methods

3.1. Mathematical model

To simultaneously capture the relationship between teleworking and both infectious disease

(ID) and non-communicable disease (NCD) risk, we designed a compartmental deterministic

model to represent the population of employees from a non-specific company during an

epidemic wave of a SARS-CoV-2-like virus (Figure 1a). This model is summarised in the

following set of ordinary differential equations driving the dynamics of numbers of individuals

in each model compartment:

$$\frac{dS}{dt} = -\lambda \cdot S$$

$$\frac{dE}{dt} = \lambda \cdot S - \sigma \cdot E$$

$$\frac{dI_A}{dt} = p_A \cdot \sigma \cdot E - \gamma_A \cdot I_A$$

$$\frac{dP}{dt} = (1 - p_a) \cdot \sigma \cdot E - \rho \cdot P$$

$$\frac{dI_S}{dt} = \rho \cdot P - \gamma_S \cdot I_S$$

$$\frac{dR}{dt} = \gamma_A \cdot I_A + \gamma_S \cdot I_S$$

We considered that all employees of the company are initially susceptible (S) to the ID and

can become exposed (E), *i.e.* infected but not yet infectious, following a transmission event at

rate λ. Exposed individuals progress through a latent period at rate σ before either becoming

infectious asymptomatic (I_A, with probability p_A), or infectious presymptomatic (P).

Presymptomatic individuals eventually become infectious symptomatic (I_S) at a rate ρ. All

infectious individuals eventually become recovered (R) at a rate which varies depending on

whether they are asymptomatic (γ_A) or symptomatic (γ_S). The total number of employees

N=S+E+I_A+P+I_s+R is assumed constant over the time period of interest. The cumulative

incidence is calculated over a 3-month period.

3.1.1. Rate of infection

Employees can become infected either via contacts with 1) other infectious employees during

working days when they are present in the workplace, 2) contacts with individuals outside the

company during working days when they are teleworking, or 3) contacts with individuals

outside the company during non-working days (Figure 1b). We assumed that five days per

week are working days, and calculated the overall force of infection as a weighted average of

forces of infection in the workplace (λ_W), during telework (λ_T), and on non-working days (λ_C),

as follows:

$$\lambda = \frac{5}{7} \cdot (1 - \alpha) \cdot \lambda_W + \frac{5}{7} \cdot \alpha \cdot \lambda_T + \frac{2}{7} \cdot \lambda_C$$

128   with

$$\lambda_w = \beta \frac{v \cdot I_A + P}{N - I_S}$$

Here, α indicates the average proportion of time spent in telework rather than in the office. If

employees are working from the office, they can be infected at a rate β either by the

proportion of infected asymptomatic (I_A) or presymptomatic (P) employees also present. We

assumed that infected   symptomatic (I_S) individuals are systematically on sick leave, and

hence cannot infect other employees since they are absent from the workplace. We further

assumed that asymptomatic individuals are less infectious than presymptomatic by a factor v

(23).

The term λ_C corresponds to the force of infection which an employee is subjected to outside

the workplace, on a non-working day. We considered that this represents the wider epidemic

in the community, which is independent of the workplace epidemic since it is much larger.

We calibrated λ_C to available incidence data from the second wave of SARS-CoV-2 in France

(01/09/2020 - 01/12/2020) (Figure 1c). On teleworking days, we assumed that employees

would also have contacts with other individuals in the community, although at a lower rate

than on non-working days, hence we defined the force of infection on teleworking days λ_T as

a fraction ε of λ_C.

We calculated the within-company basic reproduction number R_0 using the next generation

matrix method (24) (Supplementary Text 1 (25)) from which we derived the per-employee

transmission rate β for presymptomatic individuals in the workplace as

$$\beta = \frac{R_0 \cdot \rho \cdot \gamma_A}{(1 - p_A)\gamma_A + \rho \cdot \upsilon \cdot p_A}$$

152
Figure 1: Model structure. a) Model diagram. With regards to the infectious disease,

individuals can either be Susceptible (S), Exposed (E), Infected Asymptomatic (I_A),

Presymptomatic (P), Infected Symptomatic (I_S) or Recovered (R). Compartments with the

superscript "C" indicate individuals who will eventually develop a NCD due to teleworking. **b)**

Components of the total force of infection λ. c) Community epidemic curve (bars, left

axis) and corresponding force of infection λ_c (line, right axis).

3.1.2. Rate of non-communicable disease incidence

To represent the incidence of the selected NCD due to telework in this model, we duplicated

the compartments listed above to further stratify individuals according to whether they will

develop a NCD due to teleworking. We considered that individuals in the infected

symptomatic state cannot move to the corresponding compartment, since they do not

telework but are on sick leave. For all other states, the transition towards the NCD status

occurs at an average daily rate defined by the exposure-response function *f(α)* which

depends on the proportion of telework *α*. Possible shapes for this function were informed by

the studies identified in our rapid review (22,26,27). We derived the exposure-response

curves using linear interpolation between the data points retrieved from the identified studies.

To ensure comparability between the different forms of *f(α)*, we systematically set the

baseline value *f(0)* equal to 8.9 x 10^{-5} per day. This corresponds to an annual probability of

approximately 3.2% (= 1 − exp(-8.9*10^{-5}*365)) to develop a NCD in absence of teleworking,

similar to the Global Burden of Disease (GBD) 2021 estimates for the cumulative incidence

of MSDs and mental disorders (28). Parameter values for the model are summarised in

Table 1.

Table 1: Model parameters.

Name	Symbol	Value	Range explored	Source
Reproduction number	R_0	2.66	2 - 4	(29)
Coefficient of relative infectiousness of asymptomatics	ν	0.35	0.1 − 1.27	(23)
Coefficient of relative community force of infection on teleworking days	ε	0.21	0.17 − 0.25 (assumed +/- 20%)	(15)
Progression rate from exposed to infectious (= 1/incubation period)	σ	1/6.57 days^{-1}	1/18.87 - 1/1.80 days^{-1}	(30)
Probability of asymptomatic infection	p_A	0.2	0.17 − 0.25	(23)
Recovery rate from infection for asymptomatics	$γ_A$	1/5 days^{-1}	N/A	(31)

Progression rate from presymptomatic to symptomatic	ρ	1/1.5 days^{-1}	N/A	(31)
Recovery rate for symptomatics	γ$_S$	1/5 days^{-1}	N/A	(31)
Baseline incidence rate of non-communicable disease in absence of telework	f(0)	8.9 x 10^{-5} days^{-1}	N/A	(28)
Maximum change in relative risk of non-communicable disease due to telework*	ω	+/- 0.7	+/- 0.05-0.7	This study

*This parameter is only used in the sensitivity analysis

Model simulations over a 3-month time period were performed using R version 4.2.2. The

code for the model and all the analysis presented in this article is freely available online (32).

3.1.3. Converting incidences to health burden

In order to compare the health burden of ID and NCD cases using the same metric, we

estimated the number of disability-adjusted life years (DALYs) caused by these diseases,

placing ourselves in a scenario where the ID was COVID-19 and the NCD was lower back

pain. For COVID-19, DALYs were computed by adding up the years of life lost (YLL) from

deceased individuals and the years of healthy life lost due to disability (YLD) from

symptomatic cases, while the DALYs for the NCD were simply the YLDs. Disease-specific

disability weights were obtained from the 2021 GBD (28). In the GBD, these weights are

stratified by the level of severity of the disease. In order to obtain disease-specific average

disability weights which account for the severity spectrum of each disease, we divided

disease-specific YLD-estimates by the number of prevalent cases among individuals aged 20

to 64 years in France, as provided by the GBD. We also derived average disease durations

from the 2021 GBD estimates using the ratio of prevalent to incident cases. For both

diseases, we obtained the YLD term by multiplying the model-predicted number of cases by

the average disease-specific disability weight, and the average disease duration. Finally, we

estimated the YLLs from deaths following SARS-CoV-2 infections by multiplying the model-

predicted number of COVID-19 cases by the age-specific infection fatality rate (33) and the

age-specific life expectancy. Based on the statistics of the French public finance services on

the working population, the mean age was assumed to be 41 years

(https://www.insee.fr/fr/outil-interactif/6794598/EVDA/FRANCE). In 2023, life expectancy at

41 was 45.6 years for women and 40.5 for men, averaging to 43.1 years.

3.2. Rapid review

We conducted a rapid review of research articles investigating quantifiable aspects of

telework (e.g. frequency measured in days per month or hours per week) in association with

the risk of NCDs, such as mental health disorders and MSDs (34).

3.2.1. Search strategy and study selection

Previous reviews on the impact of telework on health have found that the most notable

outcomes are related to behavioural risks, mental health or MSDs (35–37). Therefore, we

restricted our review to the impact of telework on these outcomes.

We searched three databases (Pubmed, Scopus and Google Scholar) to identify original

studies assessing the quantitative association between telework and health. We then

synthesised the evidence regarding the shape and magnitude of the relationship between

telework exposure and the various health outcomes studied.

The search terms used were ("telework" OR "remote work" OR "work from home" OR

"telecommuting") AND ("mental health" OR "musculoskeletal disorder").

3.2.2. <u>Inclusion and exclusion criteria</u>

We used the following inclusion and exclusion criteria to select studies:

● Publication type: we included original research articles only.

● Population: we included articles studying workers exposed to some level of telework,

whether they were part of a specific sub-population (such as specific professions) or

not.

● Exposure: we included studies that compared at least two levels of exposure to

telework, in addition to no telework.

● Outcomes: we included studies that reported MSDs, mental health conditions or

behavioural risks as outcomes, and excluded studies reporting only other outcomes.

We did not restrict ourselves to a single specific MSD, mental health condition or

behavioural risk.

We placed no restriction on the date of publication or the language.

3.2.3. <u>Data collection</u>

From the selected studies, we extracted the following information: study years, country,

working population (sector), study design, quantification of telework exposure, the health-

related outcomes examined, and the shape of the observed relationship. Importantly, we

extracted relationships regarding health-related outcomes for any telework level different

from 0 to portray plausible exposure-response functions. To inform the model, we only

considered exposure-response relationships based on significant ($p < 0.05$) values.

3.3. Sensitivity analysis

3.3.1. Theoretical exposure-response functions

To compare the exposure-response relationship shapes independently of the upper/lower bounds of the functions, we developed a set of theoretical functions which reproduced the shapes we identified in our review, whilst allowing us to flexibly change the upper/lower bounds. These functions all take as input the teleworking frequency α (α ∈ {0,1}) and return the relative risk of non-communicable disease, with the constraint that f(0) = 1 and the upper/lower bound is equal to ω.

For linear increasing or decreasing functions:

$$f(\alpha) = (\omega - 1) \cdot \alpha + 1$$

Where the function is linear increasing if ω > 1 or decreasing if ω < 1.

For inverted-U or U-shaped functions:

$$f(\alpha) = 4(1 - \omega) \cdot (\alpha - 0.5)^2 + \omega$$

Where the function is inverted-U shaped if ω > 1 or U-shaped if ω < 1. The value ω is reached for α=0.5.

For L-shaped functions:

$$f(\alpha) = (1 - \omega) \cdot e^{-10\alpha} + \omega$$

The choice of value 10 controls the rate of decay of the function, to obtain a function similar

to the observed LS function from the literature with $\omega=0.3$ represented in Supplementary

Figure 1.

These theoretical functions are represented in Supplementary Figure 2.

3.3.2. Impact of model parameter values on incidence

We calculated partial correlation coefficients to examine the impact of model parameter

values on cumulative incidence of ID and NCD over the period, assuming a teleworking

frequency of 0.5. A total of 400 parameter sets were drawn, with each parameter

independently sampled from a uniform distribution within the ranges specified in Table 1.

279
280

4. Results

4.1. Studies on NCD and telework identified to inform the model

From our rapid review, we identified only three studies that met our selection criteria and presented exposure-response relationships between teleworking and NCD risk which could inform the mathematical model (Table 2). Of those studies, one focused on work-related outcomes (26), one on physical health (27), and one on health determinants (including behavioural changes) impacting both mental and physical health (22). Importantly, two of the identified studies were entirely conducted before the COVID-19 pandemic (22,26).

The study that focused on work-related outcomes notably examined depressive symptoms and self-rated health (26). The results of this longitudinal study conducted on United States employees highlighted exposure-response relationships for several outcomes. A U-shaped (US) relationship was found between the number of teleworking days per week and depressive symptoms, with a higher risk among non-teleworkers (0 days per week) and those with a higher frequency of telework (5 days per week). The association between number of teleworking days per week and self-rated health also displayed a US relationship.

For physical health, the study we identified only focused on lower back pain as a MSD-related outcome (27). The findings from this cross-sectional study conducted in Japan revealed an inverted U-shaped (IU) exposure-response relationship between teleworking frequency and lower back pain, with the highest risk in case of intermediate teleworking frequencies (2-3 days/week).

Finally, the last study described exposures related to both mental and physical health such as alcohol abuse or physical inactivity (22). This longitudinal study was conducted in the United States on insurance company employees. Retrieved results (significant values at $p <$

0.05) suggested a broadly decreasing relationship between teleworking intensity and risks of

alcohol abuse and tobacco use. For the Edington risk score (a summary score accounting for

several risk factors), we observed an L-shaped (LS) relationship with increased telework

intensity, with an initial rapid decrease in risk at low telework frequencies followed by a more

stable relationship at higher frequencies.

Table 2: Summary of identified studies on the exposure-response relationship
between teleworking frequency and non-communicable disease (NCD) risk. Here, we
classified (or retrieved) the shape exposure-response relationships into five types: U-shaped
(US), inverted U-shaped (IU), L-shaped (LS), broadly decreasing (D) and broadly increasing
(I). Relationships in italic are tested in our mathematical model in the next section. "Baseline
level" values are values from the studies for groups not exposed to telework (0 days per
week).

Article	Study period & country	Exposure to telework	Health outcome	Health outcome measurement & shape of relationship		Baseline level
Chen et al., 2023 (26)	2018-2019, USA	Days per week (3 groups): • 1-4 • 5	**Self-reported (range: 0-10)**			
			Depressive symptoms	*US: 1.71* ; 2.08**		*2.33*
			Self-rated health	US: 6.01* ; 5.96*		5.69
Matsugaki et al., 2021 (27)	2020, Japan	Days per week (3 groups): • 1 • 2-3 • 4+	**Self-reported symptom (% of Yes)**			
			Lower back pain	*IU: 52.8%* ; 53.2%* ; 47.5%**		*49.9%*
Henke et al., 2016 (22)	2010-2011, USA	Hours per month (4 groups) : • 8- • 9-32	**Regression coefficient and percentage at baseline**			
			Alcohol abuse (at least 2 or 3 drinks per day)	D: -0.061 ; -0.503 ; -0.024 ; -1.423*		4.9%
			Physical	US: -0.189 ; -0.249* ; -0.140 ; -		40.4%

			inactivity (<3 days of exercise per week)	0.015	
		• 33-72 • 73+	Tobacco use (yes/no)	D: -0.100 ; -0.175 ; -0.426* ; -0.263	12.3%
			Edington risk (>5 health risk factors)	*LS: -0.571* ; -0.997* ; -0.321 ; -1.233**	*5.7%*
			Obesity risk	US: -0.150 ; -0.146 ; -0.202 ; -0.113	33.1%
			Depression	D: -0.083 ; -0.056 ; -0.202 ; -0.151	16.4%
			Stress	I: 0.033 ; -0.037 ; 0.040 ; 0.137	29.5%
			Poor nutrition	US: -0.017 ; -0.110 ; 0.009 ; 0.194	86.2%

319   *Reported p value <0.05 in the referenced study

4.2. Impact of varying teleworking frequency on health outcomes

Our review shows that the exposure-response relationship between teleworking and NCD
risk may vary depending on the mental or physical health focus, as well as on other factors.
This is bound to impact the level at which teleworking best prevents both infectious disease
(ID) and NCD. Based on our literature review and the statistically significant relationships we
identified, we explored three relationships between telework frequency and the risk of NCD in
our model: L-shaped (LS, for Edington risk), U-shaped (US, for depressive symptoms), and
inverted U-shaped (IU, for lower back pain). In the model, the corresponding exposure-

329 response functions $f(α)$ giving the daily rate at which individuals will develop a NCD for a

330 given frequency of teleworking $α$ were parameterized using data from the three studies

331 analysed in the rapid review (22,26,27) (values in italic in Table 2, see Supplementary Text 2

332 for details). Importantly, since we used values from the literature to parameterize these

333 functions, the strengths of the association and therefore their upper and lower bounds are

334 different.

336 Implementing telework throughout the epidemic wave substantially reduced the cumulative

337 incidence of ID amongst employees (yellow curve), with greater reductions at higher

338 teleworking frequencies and no substantial differences between frequencies greater than 0.8,

339 i.e. 4/5 days per week (Figure 2a). However, the cumulative incidence of NCD at different

340 teleworking frequencies varied depending on the assumed exposure-response relationship

341 function, with the predicted peak incidence of NCD occurring either at low (LS, US) or

342 intermediate (IU) teleworking frequencies (Figure 2b).

344
Figure 2: Epidemic dynamics and cumulative incidence of non-communicable disease

(NCD) at varying telework frequencies. Model simulations over 3 months using a telework

frequency α = 0 (no telework), 0.2 (1 day in 5), 0.4 (2 days), 0.6 (3 days), 0.8 (4 days) and 1

(full telework), for **a)** changes in the number of susceptible (S), exposed (E),

asymptomatically infected (Ia), presymptomatic (P), symptomatically infected (Is) and

recovered (R) employees, and **b)** the cumulative incidence of employees who will eventually

develop a NCD, in case of a L-shaped (LS, circles), U-shaped (US, triangles) or inverted U-

shaped (IU, squares) exposure-response relationship between telework and NCD risk.

4.3. Impact of varying teleworking start date on health outcomes

The impact of teleworking on cumulative disease incidence further varied depending on the timing of its implementation from the start of the epidemic wave (Figure 3). Since we performed simulations in the context of an epidemic wave, implementing teleworking too late prevented any significant reduction of the ID cumulative incidence (orange lines). If we set an arbitrary objective to reduce this incidence below 50% of the baseline value obtained without teleworking, at least two days of teleworking (frequency of 0.4) were necessary to reach this target, with teleworking implemented within the first 40 days of the epidemic (solid orange lines).

Regarding the change in NCD risk caused by teleworking, as expected, early implementation of teleworking systematically led to a higher change in cumulative incidence (blue lines). However, the nature of this change depended on the assumed exposure-response function shape, upper and lower bounds, and on the teleworking frequency. For example, in the case of the L-shaped relationship for Edington risk parameterised according to our review (first row), the cumulative incidence of NCD was lowest when teleworking was implemented at a frequency of 1 on the first day (last column), while for the U-shaped relationship for depressive symptoms (second row) the lowest incidence was achieved at a frequency of 0.6 (fourth column). On the other hand, teleworking led to an increase in cumulative NCD incidence (relative incidence > 100%) when considering the inverted U-shaped relationship with lower back pain (last row), particularly at intermediate teleworking frequencies (0.4-0.6, third and fourth columns). In that case, slightly delaying the implementation of teleworking to avoid increasing the risk of NCD while still having an impact on ID could be better, and/or implementing 100% teleworking, which would not increase the risk of NCD while keeping the ID risk at a minimum. For the US relationship, timing is also important because early teleworking implementation for intermediate frequencies reduces both health risks below their respective targets, which is not the case for lower or higher frequencies. Additionally,

since there is no major difference for ID relative risk between frequencies of 0.8 and 1, a

teleworking frequency of 0.8 would be preferable to achieve strong reductions in both NCD

and ID risks.

The contrasting relationship between teleworking and ID versus NCD risks implies that

reducing both incidences simultaneously may not always be feasible, depending on the

characteristics of the NCD. In our illustrative example here, where we aimed to reduce ID

incidence by at least 50% and NCD incidence by at least 10%, we observed only limited

conditions where teleworking could simultaneously reduce both diseases incidences below

these targets (Figure 3, grey shaded areas). The definition of an "optimal" teleworking

frequency to improve health is therefore inconsistent and will vary depending on (*i*) the

observed exposure-response relationship between teleworking and health outcomes, and (*ii*)

the relative importance granted to both ID and NCD risks when defining target thresholds.

Figure 3: Impact of teleworking timing and frequency on the relative cumulative incidences of infectious (ID) and non-communicable diseases (NCD) using observed exposure-response functions. Cumulative incidence corresponds to the number of employees infected over the three months of the simulated epidemic. Here, we represent the cumulative incidences of ID and NCD relative to the cumulative incidences predicted by each model without teleworking (first column). For example, the lowest incidence of NCD (blue line) for the L-shaped function (first row) is when teleworking is implemented with a frequency of 1 (5 days a week) on day 0 (last column), while for the U-shaped exposure-response function (second row) this occurs at a frequency of 0.6 (fourth column). The grey dashed line indicates 100%, i.e. the baseline incidences. Lines are solid when the relative incidences of ID and NCD are respectively lower than 50% and 90% (chosen arbitrarily as examples), and faded otherwise. The shaded grey areas indicate conditions where both relative cumulative

407   incidences are below the defined targets of 50% and 90% (for ID and NCD incidence,

408   respectively).

4.4. Exploring different exposure-response relationships between teleworking and NCDs

While the exposure-response functions used in the previous section were directly informed

by values we identified in the literature, their upper and lower bounds varied, which

introduces a bias when comparing the incidence of different NCDs since the differences are

not only attributable to the shape of the functions. For example, the greatest change in

relative risk of NCD for the IU function (lower back pain) was approximately +6%, while for

the LS function (Edington risk) this reached -70%. To overcome heterogeneity in exposure

categorisation and data limitations regarding exposure-response curves, we reproduced our

analysis using five theoretical functions with shapes corresponding to those we identified in

our review (see Supplementary Text 2 for details, and Supplementary Figures 2 and 3 for the

shapes of the functions), but each with the same greatest absolute change in relative risk of

NCD (+/- 70%). The results obtained underline how the shape of these functions alone

affects the optimal telework frequency and implementation date, independently of

upper/lower bounds (Figure 4). This analysis also highlights scenarios where the relative

increase in NCD incidence can be equivalent to the relative decrease in ID incidence, as can

be seen for the theoretical IU function at a telework frequency of 0.6, and the theoretical

linear increasing function at a frequency of 1 (Figure 4, rows 3 and 4).

430
Figure 4: Impact of teleworking timing and frequency on the relative cumulative

incidences of infectious (ID) and non-communicable diseases (NCD) using theoretical

exposure-response functions. Cumulative incidence corresponds to the number of

employees infected over the three months of the simulated epidemic. Here, we represent the

cumulative incidences of ID and NCD relative to the cumulative incidences predicted by each

model without teleworking (first column). The grey dashed line indicates 100%, i.e. the

baseline incidences. Lines are solid when the relative incidences of ID and NCD are

respectively lower than 50% and 90% (chosen arbitrarily as examples), and faded otherwise.

The shaded grey areas indicate conditions where both relative cumulative incidences are

below the defined targets of 50% and 90% (for ID and NCD incidence, respectively).

4.5. Estimating the health burden of infectious and non-communicable diseases at different teleworking frequencies

At baseline, without any telework, we estimate that an infectious disease such as COVID-19

could be responsible for approximately 257 DALYs in our worker population (96% of total

health burden, Figure 5). In the same scenario, a non-communicable disease such as lower

back pain could be responsible for approximately 11 DALYs (4% of total burden).

As teleworking frequency increases, the burden of the ID decreases, down to 16 DALYs at a

frequency of 1. However, depending on the exposure-response function, the burden of the

NCD can either increase or decrease, while its value relative to the burden of the ID always

increases. For example, assuming an L-shaped or linear decreasing shaped function, the

NCD burden remains small, but represents a higher share of total health burden (3 DALYs,

15% of total burden at 100% teleworking frequencies). However, with a linear increasing

shaped function, the burden of the NCD can increase up to 18 DALYs, representing 53% of

the total health burden of the ID and NCD combined.

Figure 5: Impact of teleworking frequency on the health burden of infectious (ID) and non-communicable diseases (NCD) using theoretical exposure-response functions. The burden is expressed in disability-adjusted life years (DALYs). The health burden is estimated using COVID-19 as the infectious disease, and lower back pain as the non-communicable disease, in a population of 5 000 workers with an average age of 41 years.

4.6. Sensitivity analysis

The only differences in correlation coefficients that we observed between scenarios of exposure-response relationships affected one parameter, the maximum change in relative risk of NCD due to telework (ω) (Supplementary Figure 3); as expected, this parameter was strongly positively correlated (coefficient ≈ 1) with NCD incidence for IU and LI curves, and strongly negatively correlated (coefficient ≈ -1) with NCD incidence for LD, LS and US curves.

Regardless of which exposure-response function was used, the relative infectiousness of asymptomatic individuals (v) was only slightly negatively correlated with ID cumulative incidences (coefficient ≈ 0.2). The coefficient of relative community force of infection on

teleworking days (ε) was only slightly positively correlated with ID incidence (coefficient ≈

0.2). The reproduction number (R_0) and the progression rate from exposed to infectious (σ)

were strongly positively correlated with ID incidence (coefficient > 0.7). These two

parameters were negatively correlated with NCD incidence (coefficient ≤ -0.5), which is

expected since we assume that infected individuals cannot develop a NCD during the period

when they are infected and symptomatic. Inversely, the proportion of asymptomatic infections

(p_A) was slightly positively correlated with NCD incidence (coefficient ≈ 0.2), since it leaves a

greater proportion of infected individuals at risk of developing a NCD in parallel due to

reduced sick leaves.

5. Discussion

Recent years have seen an unprecedented increase in teleworking frequency in many countries worldwide. In this study, we aimed to determine if there existed an optimal teleworking frequency to maximise health benefits, by designing a mathematical model accounting for both infectious disease (ID) transmission and NCD incidence in employees of a non-specific company. To parameterise this model, we first reviewed the evidence on the consequences this may have in terms of NCDs for teleworking employees, notably underlining potential impacts on mental health and MSDs. Our rapid review uncovered a wide variety of possible exposure-response relationships between teleworking intensity and NCD risk. By incorporating this data in our model, we showed that optimal frequency and timing of implementation of teleworking during an epidemic wave could vary widely. For instance, for health impacts associated with teleworking through a L-shaped function with a strength of association such as the one we identified for Edington risk (22), rapid and wide implementation of teleworking during the first few days of an epidemic can reduce both ID and NCD incidences. On the other hand, for a U-shaped relationship with parameters such as the one we identified for depression (26), intermediate (3-4 days per week) teleworking frequencies may be more optimal to maximise health benefits, while for inverted U-shaped relationships with parameters as observed for lower back pain (27), it may be necessary to weigh the increased NCD risk attributable to teleworking against the decreased ID risk. Importantly, both the shape and the upper/lower bound of these exposure-response functions must be taken into consideration when contrasting ID and NCD risk. We further demonstrate the need for this weighing by showing that telework can increase the relative importance of the health burden of NCDs, highlighting the importance of accounting for both ID and NCD risk when evaluating the impact of telework.

5.1. Implications

Through our rapid review, we confirmed that telework may impact both mental and physical health, although we identified several important knowledge gaps based on the selected studies. First, there is a lack of uniformity across the measurements of health outcomes (clinical diagnoses and declarative statements) explored in relation with teleworking, ranging from physical to mental health (depression, anxiety, burnout). Second, exposures are poorly characterised. Exposure to teleworking is not coded in a standardised manner; depending on the study, it may be expressed in terms of number of days per week, or number of hours per day, with different categories. Third, study contexts are heterogenous. Studies encompassing short- and long-term perspectives, within both pandemic and non-pandemic settings, yield outcomes potentially contingent on the phase of the COVID-19 pandemic (38). Confounding factors (such as the region, pre-existing comorbidities or professional status) are not well described or controlled for, and differ across studies. Fourth, very few studies met our inclusion criteria, and we found no longitudinal study that focused on MSDs, leading to a lack of quality in evidence especially regarding the temporal relationship between exposure and MSDs. Our findings are in agreement with two recent systematic reviews that also outlined the low quality of evidence regarding health impacts of teleworking (39,40). A recent study also shows that the relationship between telework and well-being can be dynamic over the duration of a pandemic and will differ according to the baseline level of telework (41).

Overall, the identified studies led to divergent results on the shape of the exposure-response function, depending on the considered health outcome (U-shaped, inverted U-shaped, L-shaped, broadly decreasing, and broadly increasing curves). In addition, even for the same health outcome, reported impacts of teleworking could be heterogeneous. For example, Henke and colleagues found a broadly decreasing relationship between hours/month of telecommuting and depression, while Chen and colleagues found a U-shaped relationship. This conflicting evidence may be due to modifying effects as demonstrated by one study

according to which organisational factors within a company could alter the shape/direction of

the relationship, depending on the COVID-19 wave (38).

Our mathematical model suggests that the shape and range of variation of the exposure-

response relationship function between teleworking and NCD risk may influence the optimal

teleworking frequency. When using this kind of model in an epidemic context, it is crucial to

initially define the risk levels considered optimal for both ID and NCD. For a given target of

incidence reduction, the choice of teleworking frequency and the timing of its implementation

as an intervention during an epidemic wave varies. The implications in terms of

performances and costs are also different as we expect sick-leaves due to IDs to be

numerous but short while sick-leaves due to NCDs would be fewer but longer. Furthermore,

MSDs and psychological disorders are both heterogeneous in severity, which makes

quantitative assessments difficult. Finally, implications may largely depend on the

professional sector, for example, workers in healthcare settings or densely-populated

workplaces are expected to be more exposed to IDs, while desk-based workers will be more

exposed to NCDs. The risk of ID and NCD, and therefore the definition of the "optimal"

teleworking frequency could therefore vary depending on the job category and the associated

tasks.

5.2. Limitations

As shown in our rapid review, the relationship between telework frequency and the risk of

developing common NCDs such as lower back pain or psychological distress in the long-term

depends on many unmeasured individual and environmental characteristics. Instead of

accounting for all these specificities, we illustrated with our model the impact of different

relationships considering one average risk for all individuals. The lack of significant

associations between telework and some NCDs is also a source of uncertainty in the

exposure-response relationships used in the model, as shown in Table 2. Alternative

modelling strategies could be used to integrate individual heterogeneity, but this would

require additional evidence regarding the distribution of exposures and risks based on

individual characteristics (age, gender, job…) that is not yet available in the literature. In

addition, one of our associations (inverted U-shaped relationship for lower back pain) is

derived from a single cross-sectional study, limiting causal inference and highlighting the

scarcity of evidence on teleworking and NCD incidence. Another key limitation of the current

evidence base linking telework and NCDs is its reliance on arbitrarily defined categories of

telework exposure, as no study to date has reported fully disaggregated or continuous

measures. This may bias the estimation and shape of dose–response relationships, and

affect the conclusions of studies like ours.

The majority of evidence regarding telework arises from the COVID-19 pandemic. In this

context, telework was frequently unplanned and imposed on individuals, which does not

necessarily reflect conditions where teleworking would be planned and adapted at an

individual level (7). In addition, relatively few countries were represented in these studies,

while we would expect the impact of teleworking to vary across regions, between urban and

rural settings and living conditions. Finally, the impact of teleworking on health likely depends

on the socio-professional categories considered and its desirability.

In our illustrative example, we simulated the impact of telework policies in the context of the

second wave of the SARS-CoV-2 pandemic. Thus, the results should be interpreted in the

context of an emerging ID for which no pharmaceutical intervention (e.g., vaccination) is

available, and for which no behavioural change among workers is observed apart from the

telecommuting policy decided by the employer. We expect different results in case these two

assumptions are not met.

Finally, we assumed homogeneous mixing within the company, whereby all employees could

be in contact, without considering more complex work organisations. Similarly, we

considered a simple telework policy according to which a fixed percentage of the total

workforce in the company is teleworking every day, but more refined policies have been

implemented during the pandemic, such as rotating telework (13). In these strategies,

employees are evenly distributed in groups that alternate on a daily or weekly basis. This is

expected to reduce the overall number of contacts per individual and potentially the risk of

transmission, whilst maintaining a reduced average frequency of telework. Building a

discrete-time model to represent alternating teleworking days and on-site days would be a

valuable extension, providing a more realistic description of telework patterns and

transmission dynamics.

5.3. Perspective and future directions

Several options could be considered to mitigate the NCD risk among teleworking individuals.

First, sedentary behaviours, prolonged computer use, and poor ergonomics during telework

have been linked to increased musculoskeletal disorders (MSDs), particularly affecting the

neck, shoulders, and lower back. To mitigate MSD risks, studies recommend ergonomic

workspaces, regular posture changes, and proactive telework preparations by companies,

ensuring adequate support and proper work conditions. Second, telework has mixed effects

on mental health, enhancing positive emotions, job satisfaction, and organizational

commitment while reducing emotional exhaustion, but also contributing to anxiety, stress,

fatigue, and social isolation. Negative effects could be mitigated by measures like technical

support, flexible hours, social communication, and health interventions, with part-time

telework offering benefits for work-life balance and social relationships.

In practice, the optimal frequency and implementation timing of telework will depend on the

relative importance given to both NCDs and ID, as we show by estimating the health burden

of both diseases under varying teleworking frequencies. The decision on respective weights

given to the ID and NCD incidence should consider the different timelines at which these

diseases occur: typically, short-term for IDs and mid- to long-term for NCDs. IDs represent a

substantial socioeconomic burden (42), which can justify the implementation of teleworking

as an intervention to reduce disease spread. Part of this burden is related to sick leave (43),

which can lead companies to act in order to minimise disease incidence among their

employees. However, when designing a teleworking policy, deciders need to account for

feasibility, legal and ethical criteria. For instance, teleworkers need to have access to

telework equipment, which was not always straightforward during the COVID-19 pandemic

(44).

Future studies should better characterize the distribution of telework among population and

identify its association with NCDs and mental health outcomes based on individual

characteristics (age, gender, job…). It is also important to explore how telework dynamics

can influence the transmission of infectious diseases among workers.

Current lack of evidence, especially data to assess the impact of exposure to telework more

finely, prevented us from relying on robust exposure-response functions linking NCDs and

telework. However, pending such data, our work provides a framework to balance the

infectious and NCD risks in an epidemiologic context that could easily be re-used with

updated data.

6. Conclusions

In our rapid review, we identified three studies, two being longitudinal. This very low number underlines the need for more data to monitor the health impacts of teleworking. In particular, further studies should characterise the relationship between telework frequency and NCDs. To this end, it is crucial to collect data outside the context of the COVID-19 pandemic, since additional stressors during this period may have modulated the relationship between telework and health. In addition, the mechanisms by which teleworking impacts mental and physical health should be better characterised (e.g. unadapted workstation for MSDs). Lastly, further efforts are needed to identify the individual factors affecting the exposure-response relationships both during and outside of epidemic contexts.

Our innovative approach, which attempts to combine short- and longer-term consequences of teleworking in a unique framework, could serve as a basis to develop tools for employers and policymakers. Such tools could be used to quantify the impact of telework on employee health and identify optimal telework strategies to limit health adverse events and improve employee well-being.

7. References

1. International Labour Organization. COVID-19: Guidance for labour statistics data
collection: Defining and measuring remote work, telework, work at home and home-
based work [Internet]. 2020 [cited 2023 Aug 24]. Available from:
https://www.ilo.org/global/statistics-and-databases/publications/WCMS_747075/lang--
en/index.htm

2. Eurostat. Employed persons working from home as a percentage of the total
employment, by sex, age and professional status [Internet]. 2023 [cited 2023 Aug 29].
Available from:
https://ec.europa.eu/eurostat/databrowser/view/lfsa_ehomp/default/table?lang=en

3. Chocron V. Télétravail «total» chez Boursorama: les salariés commencent à quitter
Paris pour la province. Le Monde [Internet]. 2021 Sep 28. Available from:
https://www.lemonde.fr/economie/article/2021/09/28/teletravail-total-chez-boursorama-
les-salaries-commencent-a-quitter-paris-pour-la-province_6096226_3234.html

4. Hyder B. Return & Remote: How Salesforce Brings Employees Together for Modern
Work [Internet]. 2023 [cited 2023 Aug 29]. Available from:
https://www.salesforce.com/news/stories/return-remote-how-salesforce-brings-
employees-together-for-modern-work/

5. OECD. Teleworking in the COVID-19 pandemic: Trends and prospects [Internet]. OECD;
2021 [cited 2023 Oct 31]. Available from: https://www.oecd.org/coronavirus/policy-
responses/teleworking-in-the-covid-19-pandemic-trends-and-prospects-
72a416b6/#biblio-d1e1079

6. Weber C, Golding SE, Yarker J, Lewis R, Ratcliffe E, Munir F, et al. Future Teleworking
Inclinations Post-COVID-19: Examining the Role of Teleworking Conditions and
Perceived Productivity. Front Psychol [Internet]. 2022 [cited 2023 Oct 31];13. Available
from: https://www.frontiersin.org/articles/10.3389/fpsyg.2022.863197

7. Bouziri H, Smith DRM, Descatha A, Dab W, Jean K. Working from home in the time of
COVID-19: how to best preserve occupational health? Occup Environ Med. 2020
Jul;77(7):509–10. doi:10.1136/oemed-2020-106599 PubMed PMID: 32354748; PubMed
Central PMCID: PMC7231547.

8. Fisher KA. Telework Before Illness Onset Among Symptomatic Adults Aged ≥18 Years
With and Without COVID-19 in 11 Outpatient Health Care Facilities — United States,
July 2020. MMWR Morb Mortal Wkly Rep. 2020;69. doi:10.15585/mmwr.mm6944a4

9. Galmiche S, Charmet T, Schaeffer L, Paireau J, Grant R, Chény O, et al. Exposures
associated with SARS-CoV-2 infection in France: A nationwide online case-control study.
Lancet Reg Health - Eur. 2021 Jun 7;7:100148. doi:10.1016/j.lanepe.2021.100148
PubMed PMID: 34124709; PubMed Central PMCID: PMC8183123.

10. Ahmed F, Zviedrite N, Uzicanin A. Effectiveness of workplace social distancing measures
in reducing influenza transmission: a systematic review. BMC Public Health. 2018 Apr
18;18:518. doi:10.1186/s12889-018-5446-1 PubMed PMID: 29669545; PubMed Central
PMCID: PMC5907354.

11. Hill EM, Atkins BD, Keeling MJ, Dyson L, Tildesley MJ. A network modelling approach to
assess non-pharmaceutical disease controls in a worker population: An application to

SARS-CoV-2. PLoS Comput Biol. 2021 Jun 16;17(6):e1009058.
doi:10.1371/journal.pcbi.1009058 PubMed PMID: 34133427; PubMed Central PMCID:
PMC8208574.

12. Jarvis CI, Gimma A, van Zandvoort K, Wong KLM, Edmunds WJ. The impact of local and
national restrictions in response to COVID-19 on social contacts in England: a
longitudinal natural experiment. BMC Med. 2021 Feb 19;19:52. doi:10.1186/s12916-021-
01924-7 PubMed PMID: 33602244; PubMed Central PMCID: PMC7892289.

13. Mauras S, Cohen-Addad V, Duboc G, Tour MD la, Frasca P, Mathieu C, et al. Mitigating
COVID-19 outbreaks in workplaces and schools by hybrid telecommuting. PLOS Comput
Biol. 2021 Aug 26;17(8):e1009264. doi:10.1371/journal.pcbi.1009264

14. Mousa A, Winskill P, Watson OJ, Ratmann O, Monod M, Ajelli M, et al. Social contact
patterns and implications for infectious disease transmission – a systematic review and
meta-analysis of contact surveys. Rodriguez-Barraquer I, Serwadda DM, editors. eLife.
2021 Nov 25;10:e70294. doi:10.7554/eLife.70294

15. Bosetti P, Huynh BT, Abdou AY, Sanchez M, Eisenhauer C, Courtejoie N, et al.
Lockdown impact on age-specific contact patterns and behaviours, France, April 2020.
Eurosurveillance. 2021 Dec 2;26(48):2001636. doi:10.2807/1560-
7917.ES.2021.26.48.2001636 PubMed PMID: 34857064; PubMed Central PMCID:
PMC8641071.

16. Nowrouzi-Kia B, Haritos AM, Long BZS, Atikian C, Fiorini LA, Gohar B, et al. Remote
work transition amidst COVID-19: Impacts on presenteeism, absenteeism, and worker
well-being—A scoping review. PLOS ONE. 2024 Jul 18;19(7):e0307087.
doi:10.1371/journal.pone.0307087

17. Perelman J, Serranheira F, Castanheira F, Raposo JF, Aguiar P, Neves P, et al.
Teleworking: does it make workers healthier and productive? A cross-sectional study on
a Southern European population. BMC Public Health. 2024 Jul 20;24(1):1946.
doi:10.1186/s12889-024-19481-y

18. OECD. Assessing teleworking strategies for local development: a framework proposal
[Project Note] [Internet]. Paris; 2022 [cited 2023 Aug 31]. Available from:
https://www.oecd.org/cfe/leed/Assessing-teleworking-strategies-for-local-development-
PAT.pdf

19. World Health Organization and the International Labour Organization. Healthy and safe
telework: technical brief [Internet]. Geneva; 2021. Available from:
https://www.who.int/publications/i/item/9789240040977

20. Tezuka M, Nagata T, Saeki K, Tsuboi Y, Fukutani N. Association Between Abrupt
Change to Teleworking and Physical Symptoms During the Coronavirus Disease 2019
(COVID-19) Emergency Declaration in Japan. J Occup Environ Med. 2022 Jan 1;64(1):1.
doi:10.1097/JOM.0000000000002367 PubMed PMID: 34419985; PubMed Central
PMCID: PMC8715931.

21. De Macêdo TAM, Cabral ELDS, Silva Castro WR, De Souza Junior CC, Da Costa Junior
JF, Pedrosa FM, et al. Ergonomics and telework: A systematic review. Work. 2020 Sep
17;66(4):777–88. doi:10.3233/WOR-203224

22. Henke RM, Benevent R, Schulte P, Rinehart C, Crighton KA, Corcoran M. The Effects of
Telecommuting Intensity on Employee Health. Am J Health Promot. 2016 Nov
1;30(8):604–12. doi:10.4278/ajhp.141027-QUAN-544

23. Buitrago-Garcia D, Egli-Gany D, Counotte MJ, Hossmann S, Imeri H, Ipekci AM, et al.
Occurrence and transmission potential of asymptomatic and presymptomatic SARS-
CoV-2 infections: A living systematic review and meta-analysis. PLOS Med. 2020 Sep
22;17(9):e1003346. doi:10.1371/journal.pmed.1003346

24. Diekmann O, Heesterbeek JAP, Metz JAJ. On the definition and the computation of the
basic reproduction ratio R0 in models for infectious diseases in heterogeneous
populations. J Math Biol. 1990 Jun 1;28(4):365–82. doi:10.1007/BF00178324

25. Leclerc Q. Supplementary Information: Optimizing telework in an epidemic context:
contrasting the infectious and non-communicable diseases perspectives [Internet].
Zenodo;   2026   [cited   2026   Apr   27].   Available   from:
https://zenodo.org/doi/10.5281/zenodo.19738148 doi:10.5281/ZENODO.19738148

26. Chen Y, Weziak-Bialowolska D, Lee MT, Bialowolski P, Cowden RG, McNeely E, et al.
Working from home and subsequent work outcomes: Pre-pandemic evidence. PloS One.
2023;18(4):e0283788. doi:10.1371/journal.pone.0283788 PubMed PMID: 37014892;
PubMed Central PMCID: PMC10072379.

27. Matsugaki R, Muramatsu K, Tateishi S, Nagata T, Tsuji M, Hino A, et al. Association
Between Telecommuting Environment and Low Back Pain Among Japanese
Telecommuting Workers: A Cross-Sectional Study. J Occup Environ Med. 2021
Dec;63(12):e944–8. doi:10.1097/JOM.0000000000002412

28. Institute for Health Metrics and Evaluation (IHME). Global Burden of Disease
Collaborative   Network   (GBD   2021)   [Internet].   2024.   Available   from:
https://vizhub.healthdata.org/gbd-results/

29. Dhungel B, Rahman MS, Rahman MM, Bhandari AKC, Le PM, Biva NA, et al. Reliability
of Early Estimates of the Basic Reproduction Number of COVID-19: A Systematic
Review and Meta-Analysis. Int J Environ Res Public Health. 2022 Jan;19(18):18.
doi:10.3390/ijerph191811613

30. Wu Y, Kang L, Guo Z, Liu J, Liu M, Liang W. Incubation Period of COVID-19 Caused by
Unique SARS-CoV-2 Strains: A Systematic Review and Meta-analysis. JAMA Netw
Open. 2022 Aug 22;5(8):e2228008. doi:10.1001/jamanetworkopen.2022.28008

31. Kissler SM, Fauver JR, Mack C, Olesen SW, Tai C, Shiue KY, et al. Viral dynamics of
acute SARS-CoV-2 infection and applications to diagnostic and public health strategies.
Riley   S,   editor.   PLOS   Biol.   2021   Jul   12;19(7):e3001333.
doi:10.1371/journal.pbio.3001333

32. Quentin Leclerc. MESuRS-Lab/telework_health: Release - code for publication [Internet].
Zenodo;   2026   [cited   2026   Apr   27].   Available   from:
https://zenodo.org/doi/10.5281/zenodo.19738028 doi:10.5281/ZENODO.19738028

33. Variation in the COVID-19 infection–fatality ratio by age, time, and geography during the
pre-vaccine era: a systematic analysis. The Lancet. 2022 Apr 16;399(10334):1469–88.
doi:10.1016/S0140-6736(21)02867-1

34. Grant MJ, Booth A. A typology of reviews: an analysis of 14 review types and associated
methodologies. Health Inf Libr J. 2009 Jun;26(2):91–108. doi:10.1111/j.1471-
1842.2009.00848.x

35. Beckel JLO, Fisher GG. Telework and Worker Health and Well-Being: A Review and
Recommendations for Research and Practice. Int J Environ Res Public Health. 2022 Mar
24;19(7):7. doi:10.3390/ijerph19073879 PubMed PMID: 35409563; PubMed Central
PMCID: PMC8998114.

36. Furuya Y, Nakazawa S, Fukai K, Tatemichi M. Health impacts with telework on workers:
A scoping review before the COVID-19 pandemic. Front Public Health. 2022 Oct
31;10:981270. doi:10.3389/fpubh.2022.981270

37. Liu W, Xu Y, Ma D. Work-Related Mental Health Under COVID-19 Restrictions: A Mini
Literature Review. Front Public Health. 2021;9:788370. doi:10.3389/fpubh.2021.788370
PubMed PMID: 34900925; PubMed Central PMCID: PMC8651716.

38. Wels J, Wielgoszewska B, Moltrecht B, Booth C, Green MJ, Hamilton OK, et al. Home
working and social and mental wellbeing at different stages of the COVID-19 pandemic in
the UK: Evidence from 7 longitudinal population surveys. Thorley J, editor. PLOS Med.
2023 Apr 27;20(4):e1004214. doi:10.1371/journal.pmed.1004214

39. Fadel M, Bodin J, Cros F, Descatha A, Roquelaure Y. Teleworking and Musculoskeletal
Disorders: A Systematic Review. Int J Environ Res Public Health. 2023 Mar
11;20(6):4973. doi:10.3390/ijerph20064973 PubMed PMID: 36981881; PubMed Central
PMCID: PMC10049341.

40. Lunde LK, Fløvik L, Christensen JO, Johannessen HA, Finne LB, Jørgensen IL, et al.
The relationship between telework from home and employee health: a systematic review.
BMC Public Health. 2022 Jan 7;22(1):47. doi:10.1186/s12889-021-12481-2 PubMed
PMID: 34996400; PubMed Central PMCID: PMC8741267.

41. Kornadt AE, Bowen CE, Lepinteur A, D'Ambrosio C, Ratti L, Vögele C. Working from
home and well-being during the pandemic and beyond: a longitudinal analysis in five
countries. BMC Public Health. 2025 Mar 28;25(1):1183. doi:10.1186/s12889-025-22349-
4

42. Samsudin EZ, Yasin SM, Ruslan NH, Abdullah NN, Noor AFA, Hair AFA. Socioeconomic
impacts of airborne and droplet-borne infectious diseases on industries: a systematic
review. BMC Infect Dis. 2024 Jan 16;24(1):93. doi:10.1186/s12879-024-08993-y

43. Smith DRM, Jijón S, Oodally A, Shirreff G, Bouziad KA, Ante-Testard PA, et al. Sick
leave due to COVID-19 during the first pandemic wave in France, 2020. Occup Environ
Med. 2023 May 1;80(5):268–72. doi:10.1136/oemed-2022-108451 PubMed PMID:
36914254.

44. Belzunegui-Eraso A, Erro-Garcés A. Teleworking in the Context of the Covid-19 Crisis.
Sustainability. 2020 Jan;12(9):9. doi:10.3390/su12093662

8. List of abbreviations

ID: Infectious disease

IU: Inverted U-shaped

LD: Linear decrease

LI: Linear increase

LS: L-shaped

MSD: Musculoskeletal disorder

NCD: Non-communicable disease

US: U-shaped

9. Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Data, script and code availability

All code used in the analysis are available in the following GitHub repository: https://github.com/MESuRS-Lab/telework_health. The code at the time of publication is archived online at: https://doi.org/10.5281/zenodo.19738028.

Supplementary Information

The Supplementary Information for this article can be accessed online (https://doi.org/10.5281/zenodo.19738147). This file includes "Supplementary Text 1: Calculation of the basic reproduction number R0", and "Supplementary Text 2: Calculation of the exposure-response functions between teleworking and non-communicable disease risk" (including Supplementary Figures 1-4).

Competing interests

The authors declare that they have no competing interests.

Funding

This project did not receive any specific funding.

Authors' contributions

All authors contributed to the design of this study. LM, FT, EH, NL, KAB, and MBH performed the study selection for the rapid review based on our inclusion and exclusion criteria. QJL, ML, PH, AM and LT designed and implemented the mathematical model. WB, SZ and KJ contributed to data analysis. All authors contributed to the redaction of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors would like to thank Hanifa Bouziri for helpful discussions on musculoskeletal disorders, and Pascal Crépey for access to the COVID-19 incidence data.