Title: Micronutrient-deficient diets and possible environmental enteric dysfunction in Buruli ulcer endemic communities in Ghana: lower dietary diversity and reduced serum zinc and vitamin C implicate micronutrient status a possible susceptibility factor

Aloysius Loglo[1,2], Philemon Antwi[2], Abass Kabiru Mohammed[3], Samuel Osei-Mireku[3], George Amofa[4], Elizabeth Ofori[5], Jonathan Adjei[2], Michael Oppong[2] Richard O. Phillips[2, 6], Reginald Annan[7&], Barbara Engel[8&], Rachel E. Simmonds[1&*].

Short title: Role of diet in the pathogenesis of Buruli ulcer

[1] Department of Microbial Sciences, School of Biosciences, Faculty of Health and Medical Sciences, University of Surrey, Guildford, GU2 7XH United Kingdom.

[2] Kumasi Centre for Collaborative Research in Tropical Medicine (KCCR), Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.

[3] Agogo Presbyterian Hospital, Agogo, Ghana

[4] Dunkwa Government Hospital, Dunkwa, Ghana

[5] Tepa Government Hospital, Tepa, Ghana

[6] Department of Medicine, School of Medical Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.

[7] Department of Biochemistry and Biotechnology, College of Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.

[8] Department of Nutritional Sciences, School of Biosciences, University of Surrey, Guildford, GU2 7XH United Kingdom

Authors' email:

AL: aloysiusloglo@gmail.com, **PA**: antwiphilemon@gmail.com, **AKM**: abass@agogopresbyhospital.org, **SM**: mireku16@gmail.com, **GA**: gkamofa@gmail.com, **EO**: elizabethofori100@gmail.com, **JA**: jonathanadjei@gmail.com, **MO**: moppong567@gmail.com **RP:** phillips@kccr.de, **RA:** raannan.cos@knust.edu.gh, **BE**: b.engel@surrey.ac.uk, **RES**: rachel.simmonds@surrey.ac.uk

[*]**Corresponding author:** rachel.simmonds@surrey.ac.uk

Abstract

Background. The nutritional status of communities susceptible to Buruli ulcer (BU, a skin NTD caused by infection with *Mycobacterium ulcerans*) remains almost completely obscure. We have assessed the diets of BU patients vs. controls from the same BU-endemic communities, and compared their circulating biomarkers of nutrients and inflammation.

Methods/Principal Findings. We investigated two cohorts of BU patients and controls. The first were administered food frequency and multi-pass 24-hour recall questionnaires to determine patterns of foods consumed, nutrient intake and nutrient adequacy. The second used archived serum samples collected as baseline to measure the circulating concentration of zinc, vitamin C, CRP, IL-1β, IFN-γ, TNF-α and IL-6. Stunted growth was more prevalent than expected (31%), while 18% of participants were underweight and most had inadequate intake of all micronutrients except for carbohydrate. BU patients had a lower intake of zinc, selenium and vitamin B_{12}, and for selenium and vitamin B_{12} a higher proportion had dietary insufficiency (40% vs. 15% and 45% vs. 20%, respectively). In line with this, BU patients had significantly lower levels of zinc in their serum, and more had levels below the normal range (72% vs. 43%). Despite many participants having a good intake of vitamin C, serum levels were low, and lower amongst the BU patients. As expected, there was little evidence of systemic inflammation (CRP <0.6 mg/L). Elevated IL-6 levels were present in several participants suggesting that environmental enteric dysfunction may be prevalent in these communities, however this was similar in cases vs. controls.

Conclusions/Significance. Diet and nutritional status may be a contributing factor to BU pathogenesis. Protein and the micronutrients zinc, selenium, vitamin B_{12} and vitamin C may be of particular importance. Nutritional interventions may have potential for both prophylaxis and treatment of BU, which may be a cost-effective approach to achieving the NTD Roadmap goals.

Introduction

Buruli ulcer (BU) is a necrotizing skin disease caused by the non-tuberculous *Mycobacterium ulcerans* [1]. It is one of the neglected tropical diseases (NTDs) with skin manifestations [2], occurring as a painless nodule, plaque, oedema or ulcer [3]. A pathogen-derived exotoxin, mycolactone, is known to play a central role in disease pathogenesis [4, 5] due to its blockade the Sec61 translocon [6] underpinning its cytotoxic and immunosuppressive activities [7] as well as its induction of vascular dysfunction [8]. Hence, patients with BU do not have a systemic inflammatory response and generally present without fever or malaise even when ulceration is extensive [9, 10]. However, it is known that the immune system does control susceptibility to BU, with evidence from patients and animal models showing that macrophage function can influence infection outcomes [11-13].

Globally, BU is the third most prevalent mycobacterial disease and is commonly found in tropical or semitropical climates and most frequently in West Africa and Australia [1, 10, 14]. In Africa, the disease is widespread in poor rural communities with most cases found in children below 15 years. Even in high-resource settings, delays in treatment can result in permanent skin deformation or lifelong disabilities. Early case detection is currently the most effective approach for harm-minimisation as no vaccine exists.

To seek novel and cost-effective prevention approaches, and knowing that nutritional inadequacy is a risk factor for other mycobacterial infections (for example tuberculosis [15, 16] and leprosy [17, 18]) because it compromises immunity [19, 20], we postulated that this may also be the case for BU. However, there is a paucity of data for both BU patients and marginalised rural subsistence communities in West African countries. We, therefore, aimed to fill this knowledge gap by investigating the potential role of nutrition in the pathogenesis of BU.

Methods

Ethics

This study received favourable ethical opinion from the School of Medical Sciences Committee on Human Research, Publication and Ethics of KNUST (CHRPE/AP/18, CHRPE/AP/310/21) and the Faculty of Health and Medical Sciences Ethics Committee of the University of Surrey (UEC 2017 98 FHMS). All study participants gave informed consent.

Study area

BU patients were recruited from Agogo Presbyterian Hospital, Dunkwa Government Hospital, and Tepa Government Hospital in Ghana, which are located in areas of known BU endemicity. The archived serum samples were from previously published cohorts [21, 22].

Study design, population, and sampling

BU patients <5 years old, pregnant and lactating mothers were excluded from the study. Demographic and clinical data of cases was collected on a standardised skin NTDs [23] or BU01 form [21, 22]. The type and category of the lesion were assigned based on the WHO case definition [24]. Ulcerated lesions were dressed by nurses according to the WHO-recommended wound management protocol [24]. BU case confirmation was by a validated combined 16S rRNA reverse transcriptase/IS2404 qPCR assay [22] or IS2404 qPCR [21, 22].

Controls were recruited from the same study sites to ensure an unbiased reflection of the socioeconomic status, and food availability of the study population. Controls were age-sex-matched (as closely as possible) and residing in the same community of a recruited BU patient but from a different household.

Anthropometry

Anthropometric measures were obtained from as many participants as possible. Stratification of BMI was as advised by the WHO [25] including BMI-for-age estimation for participants 5-19 years [26, 27]. Anthropometry data had not been collected for the controls in Cohort 2. The height-for-age z-score (HAZ) was used to determine the stunting status of young people using AnthroPlus (WHO; Version 1.0.4.0) [27, 28].

Dietary assessment (Cohort 1)

A Ghana-specific food frequency questionnaire (FFQ) made up of 105 foods (**S1 Table**) was used to assess the frequency with which food items were consumed over the previous month at the time of recruitment (daily, weekly, monthly, and rarely/never) [29, 30]. These foods were divided into 25 food groups, using an approach adapted from a previous study of Ghanaian diets [31].

A Ghana-specific multi-pass 24-hour dietary recall questionnaire was also administered to obtain information on foods consumed for two week-days and one weekend day [32]. Household measures and food models were used to aid easy quantification of foods

consumed, which were then converted into grams using a standardized unit of measure provided on a nutrient analysis template (NAT) for each participant [33, 34]. Each participant's nutrient intake was compared to the relevant cut-off for their age (and gender if relevant), using information from the estimated average requirement (EAR) of FAO/WHO [35] or the RDA [36] if no EAR was available. The dietary diversity score (DDS) was generated from the 24-hour recall data using the embedded NAT parameters. This had 15 food groups (**S2 Table**) adapted from FAO guidelines for measuring household and individual dietary diversity [37], but with different numbers of food groups.

We assessed potential socioeconomic factors influencing the diets of participants using a questionnaire to obtain data on the gender, household income, education level and occupation of study participants. The household income was estimated based on the reported household monthly income per individual. Some of the participants depended on foodstuffs from their farms or livestock rearing. Those participants made an estimate on the current cost of the foods consumed from their farms or livestock to help ensure accurate determination of the monthly household income. Parents were allowed to make the estimation of household income for their children.

The highest educational status of the participants was categorized into elementary (none, primary or junior-high), secondary (senior-high and post-secondary) and higher (tertiary education only) [38]. The occupation was grouped into formal or informal. Formal included professionals, managers, clerical support staff, higher grade routine non-manual employees service and sales-related occupations while informal consisted of craft and related trades workers, elementary occupations (work that requires unskilled and semi-skilled labour) and farmers[38].

Quantification of nutritional and inflammatory biomarkers (Cohort 2)

Vitamin C, zinc, and inflammatory markers were assessed in archived serum samples had been stored at stable -80°C for up to 8 years. Vitamin C and CRP were assessed by competitive and sandwich ELISA, respectively (MBS2700398, MBS8807458 MyBioSource, San Diego, USA). Zinc quantification used a colorimetric assay (MBS841453 MyBioSource).

A bespoke Cytometric Bead Array (CBA) LEGENDplex™ (Biolegend, USA) was used for the simultaneous detection of four cytokines [IL-1β (B326518), IL-6 (B326531), IFN-γ (B326521) and TNF-α (B326522)] in 92 serum samples according to the manufacturer's instructions.

Vitamin C and zinc concentrations were compared to normal ranges that have been determined in and for European populations [39-42]. We have no reason to believe that the nutritional requirements for Africans would vary widely from these values.

Statistical analysis

Data were analysed using GraphPad Prism version 9.4.1 (GraphPad Software Inc., USA). All continuous variables were tested for normal distribution. Non-Gaussian data compared medians between two groups using a one- or two-tailed Mann-Whitney-U test. Multiple comparisons used the Kruskal-Wallis test together with the Dunn`s post-hoc correction. Multiple t-tests were used to compare nutrient intake between patients and controls, as many nutrients correlated with each other (determined in Pearson correlation matrix). To compare the frequency of categorical variables we used Fisher's exact test or Chi-squared where there were more than two variables.

Results

Characteristics of the study populations

We recruited 40 confirmed BU cases and 40 controls in Cohort 1, and identified archived samples from 80 confirmed BU cases and 80 controls in Cohort 2 (**Table 1**). The median age of the cohorts was 19.5 (IQR 11-34) and 14.5 (IQR 10-27) years, respectively (range 5-63 years). In both cohorts the majority of cases presented with ulcers, and lesions were predominately of Category I or II (80% and 73% respectively). Notably the rate of category III lesions was higher in the most recently recruited patients (Cohort 1), probably due to the disruption in case detection due to the COVID-19 pandemic. In line with the age profile of Cohort 1, the approximately half had (or were having) primary school education, and the socioeconomic profile was similar between cases and controls (**S3 Table**).

While the majority of the participants had normal weight categorisation, 18.1% were underweight including 22.5% and 19% of BU cases and 12.5% of controls. In Cohort 1 the difference in proportion of underweight participants was not statistically significant (**Table 1**). Among the 86 young participants for whom we had anthropometry data, 27 (31.4%) had stunted growth (**Table 1**). The relatively low numbers in the subgroups make it hard to compare BU cases and control, but in Cohort 2, 3 out of 46 BU cases (7%) were categorised as being severely stunted.

Frequency of food consumption

The FFQ gave qualitative insights into the frequency with which Cohort 1 study participants ate foods within a range of different food groups (**Figure S1**). Overall, this analysis shows that the participants in this study ate a fairly typical rural Ghanaian diet consisting of carbohydrate-rich foods. For example, whole grain cereals, refined cereals, fermented maize products, rice and pasta, roots/tubers/plantain were eaten frequently by more than half the participants (**Figure S1A-E**). This was alongside protein sources (predominantly chicken, fish and legumes **Figure S1F, M&L**), fruit (**Figure S1J**) and vegetables (**Figure S1G-I**). Some participants also partook in "Western" processed foods and drinks such as sodas and sweets/cakes (**Figure S1O&Q**), although more than three-quarters of the participants reported they never ate potatoes or processed meat (**Figure S1R&Q**). However, since the FFQ did not include a measure of quantity, it was hard to know how fully this meets the dietary requirements of the participants.

Nutrient intake and dietary diversity

A multi-pass 24-hour recall questionnaire was used to determine the overall energy and dietary macro and micro-nutrient intake of Cohort 1 study participants. This showed that, on average, the proportion of macronutrients making up the daily intake was close to expected norms (**S4 Table**). In line with the FFQ, carbohydrate was the predominant nutrient in the meals of both BU cases and controls (64% of daily energy intake), which is at the higher end of expected norms (45-65%). On the other hand, protein made up the smallest portion of energy intake (9.6%), at the lower end of the normal requirement of 10-35%.

While nutrient distribution was relatively normal, the energy intake (1683±837 Kcal, **Table 2**) for the vast majority of the participants was inadequate, with only 15% meeting this requirement (**Table 3**). For macronutrients, while almost all Cohort 1 participants had adequate dietary carbohydrate, only 49% of the study population had inadequate intake of

protein (**Table 3**). There were no differences in total daily macronutrient intake between BU cases and controls (**Table 2**)

In terms of the micronutrient intake of Cohort 1, while the majority of the cohort had adequate intakes of dietary iron, selenium, and vitamin C (74%, 72%, and 82%, respectively, **Table 3**), >60% had inadequate intakes of dietary vitamin A (60%), vitamin B_{12} (68%), vitamin E (86%), vitamin K (78%), and zinc (68%). For most micronutrients, there was no statistically significant difference in nutritional adequacy between BU cases and controls. However, a higher proportion of BU patients were below the EAR for dietary selenium and vitamin B_{12} compared with the controls (**Table 3**). Moreover, when comparing the mean intake of micronutrients calculated from the 24-hour recall interviews, selenium, vitamin B_{12}, and zinc were significantly lower in BU cases than controls (**Table 2**), and vitamin C was different with a P-value of 0.086.

There were no statistical differences in intake amongst those with different categories of BU lesions, gender, or household income (data not shown), although in some cases, this may have been driven by the power of our study.

A diverse diet is strongly associated with diet quality; therefore, we generated a dietary diversity score (DDS) for the participants. This showed the diversity of the diets among 15 food groups. Overall, the DDS of this cohort were within acceptable ranges for both groups, with only 6 BU cases and 4 controls having a DDS of <5 (**Figure 1**). However, the DSS for BU cases was significantly lower than in the controls (6.2 vs. 7.2, p = 0.0320).

Nutritional biomarkers

Based on these dietary data, we used archived serum samples to carry out a proof of principle that these dietary differences would translate to changes in serum concentrations, and selected zinc and vitamin C for analysis. The serum concentration of zinc and vitamin C was significantly lower in BU cases compared to controls (**Figure 2A**). Notably many BU patients and controls had levels below the expected normal ranges in serum, and this was significantly more frequent amongst BU patients than controls (P< 0.001, **Table 4**).

Severity of BU disease can be determined both by the different clinical presentations, and by the lesion category which is based on the longest diameter of the lesion (**Figure 2B&C**). There were no differences in the serum concentration of zinc or vitamin C between BU cases with less severe disease as defined by WHO category I, II or III lesion (**Figure 2B**), although a trend for lower vitamin C with increasing seriousness may have been limited by the few cases of Category III lesions in this cohort (n=6; 8%). However, BU patients with plaque forms had significantly lower levels of vitamin C compared to those who presented with nodules (**Figure 2C**), which are normally the first sign of infection.

Inflammatory biomarkers

The serum levels of CRP, IL-1β, IFN-γ, TNF-α and IL-6 were also assessed in a subset of Cohort 2. However, with the exception of IL-6, many samples had circulating levels below the detection limit of the assays (data not shown). Therefore, we took an approach that assigned each participant within a threshold (**Table 5**). No participants had levels of CRP that indicated the presence of even "high-normal" [43] levels of CRP. For IL-1β, we chose a cut-off of 57pg/ml, and only 7-13% of Cohort 2, respectively, were above this level, (**Table 5**). For TNF-α and IFN-γ we choose two cut-offs, with 2-5% having levels above the higher cut-

off. There was no statistically significant difference between BU cases and controls for these parameters.

For IL-6, such cut-offs are less well-defined than for CRP, but better defined than for other cytokines, including by systematic review that showed the range of IL-6 measured in cohorts of healthy patients was from 0-43.5pg/ml [44], although other reports from the global North report at the lower end of this (2-3pg/mL [43, 45, 46]). Hence, for our analysis, we decided to stratify our data based on whether they were normal (<7pg/ml), slightly (7-50pg/ml), moderately (50-500pg/ml) or highly (>500pg/ml) elevated (**Table 5**). The proportion of BU cases and controls in these different criteria were not statistically different, and neither was there a difference in serum IL-6 levels (**Figure 3A**). There was no significant increase in IL-6 levels within the presentation of BU (**Figure 3B**), nor the category of lesion (**Figure 3C**). However, like vitamin C, this may be due to the small number of category III lesions.

However, this analysis showed that more than half of both groups had evidence of elevated IL-6 compared to literature reports in European populations (**Table 5**). In order to understand if these elevated levels of IL-6 might explain the reduced zinc or vitamin C with IL-6, these were correlated (**Figure S2**). No significant associations were identified for BU patients (**Figure S2A**) or controls (**Figure S2B**).

Discussion

The intimate links between diet, the immune system, and infectious disease have been known for some time [47]. Indeed, malnutrition has been shown to be a predominant underlying cause of death from a range of infectious diseases [48, 49]. Inadequate dietary intake results in decreased immune function which leads to increased incidence of infection/disease and consequently increased severity and duration of disease as the systemic and local responses to infection are impaired. In addition, many infectious diseases lead to decreased intake or malabsorption of, or increased requirements for, proteins (amino acids), specific fatty acids, vitamins and trace elements. The increased metabolic demand for specific nutrients can also lead to their deficiency leading to a vicious cycle of the emergence of new infections or resurgence of old infectious diseases.

Perhaps because of the perceived dominant effect of the cytotoxin mycolactone in disease progression, the role of diet in BU has received little attention to date. In the current work, we show that many of those living in BU endemic regions of Ghana have diets inadequate in all macronutrients except carbohydrates, as well as vitamins A, B_{12}, E and K, and zinc. Moreover, BU patients' diets were worse than their control counterparts who had similar age and socioeconomic backgrounds and who were also from the same villages as the BU patients (and therefore had access to similar foods). Our approach to nutritional intake used two independent tools, both of which have been used previously in rural and urban Ghanaian settings [32, 50]. These were appropriate for low-literacy populations which was the case here, and for those who reside in rural and at times remote communities.

Insights into the diet of BU endemic areas

The WHO defines malnutrition as the deficiency or excess of nutrient intake, an imbalance of nutrients or impaired nutrient utilisation in children and adults. Undernutrition can manifest in different ways including wasting, stunting, underweight and micronutrient deficiencies. On the other hand, excessive nutrients can cause overweight or obesity. Women, infants, children, and adolescents are at particular risk of malnutrition, and poverty exacerbates the risk of malnutrition [51].

This study showed that those living in BU endemic regions in the Ashanti region of Ghana had a typical rural West African diet [31, 52], which is mainly carbohydrate-based. The diets of Ghanaians have been undergoing an evolution over recent years, particularly in urban areas [53], where diets are shifting towards a 'Western-like' pattern [52, 54]. However, the present study confirms that in rural areas, the dietary pattern has remained predominantly carbohydrate-based. Apart from socio-cultural beliefs [54], the accessibility and relatively cheap cost of starchy staples at the study sites may be the reason. Cereal-based meals like akple, kenkey, and rice and root tubers like yam, cassava, and cocoyam (used in preparation of staples like fufu and ampesi), were the most consumed by the study participants. Fufu is a well-known staple food in Ghana and is eaten by most ethnic groups and is prepared by pounding cassava and plantain or yam/cocoyam, and is consumed with soup. Additionally, phytate content of plant-based diets will reduce absorption of micronutrients including zinc and iron.

Despite the high proportion of calories from carbohydrate, the estimated requirement was not met by most of the Cohort 1 participants. Other studies have also reported nutritional inadequacies in rural (but not urban) Ghana [31, 50, 55]. Moreover, the energy intake was lower than reported for people in Eastern and Southern Africa, which were 2245 and 2618 Kcal/capita/day, respectively [56]. This suggests that diets in BU endemic areas are poorer than African averages, highlighting the need for further research to understand nutrition in such marginalised communities. Indeed, this is the first study to report in the diets of people in these areas of Ghana.

Protein in the diets of the study participants was inadequate in about 50% of the participants. This was in line with a cross-sectional study that found low protein consumption more often in rural Ghana compared to people residing in urban areas and migrants in Europe [31]. Moreover, in a wildlife-depleted farm-forest landscape in South-Western Ghana, between 14% and 60% of households ate less than the recommended daily allowance of protein [57]. Although livestock rearing is common in the study sites, Cohort 1 participants frequently ate fish and poultry, and rarely ate red or processed meat. The cost of meat could be a factor affecting the low protein intake, with the majority of the protein coming from starchy staples, which may then reduce the intake of some essential amino acids.

In terms of micronutrients, Cohort 1 also had a high prevalence of dietary inadequacy of vitamins A, B_{12}, E, K and zinc. In line with this, overall micronutrient intake has been shown to be inadequate in other studies of diet in West Africa and Ghana. This is thought to be because they are mostly found in the vegetables, fruits, and nuts that are found in the soups, stews, and sauces that accompany the main starchy component of a meal [58-61]. While, some micronutrients can be found in carbohydrate-rich foods, including roots and tubers that were frequently consumed by Cohort 1 participants, their low DDS could be the source of this inadequacy. This is because it suggests that many of them ate from a relatively small palette of foods on a regular basis.

In the context of a population at risk of BU, the widespread inadequacy of micronutrient intake is interesting because they are essential for immune function [20] and are also seen in patients with chronic wounds. For example, inadequacies of serum vitamins A, E, zinc, and carotenes have been reported in patients with chronic leg ulcers [62]. During the inflammatory stage of wound healing, vitamin E helps reduce free radical damage caused by neutrophils [63]. On the other hand, vitamin K is necessary for the functionally essential post-translational modification of proteins involved in blood coagulation, which is known to be linked to the histopathology of the disease [8, 64, 65]. Inadequacy in vitamin K is known to cause haemorrhage and purpura in newborns and has recently been shown to affect bone

and vascular health [66]. Zinc status has previously been reported to be low in Ghana [67] (prevalence of zinc deficiency ~22% in 2005 [68, 69]), and the prevalence of zinc deficiency in Sub-Saharan Africa varies between 15-50%. The current work found higher prevalence of zinc deficiency at 58%, further emphasising the poor quality of BU endemic diets compared to the country at large.

Potential role for environmental enteric dysfunction (EED) in malnutrition in BU endemic areas

Despite finding no overt systemic inflammation in BU cases, we did find evidence for circulating cytokines above normal levels (low pg/ml concentrations) in a proportion (approximately 10-50%) of both cases and controls of Cohort 2. Indeed, this is in line with studies of healthy controls in other low- and middle-income countries (LMICs), including Ghana [70-72]. Of particular note, elevated IL-6 levels were measured in more than half the participants. This likely reflects participants who were currently or recently co-infected with other pathogens. For instance, malaria infection can modulate circulating cytokines [73]. However, it could also reflect a prevalence of environmental enteric dysfunction (EED), a syndrome linked to areas of poor sanitation and hygiene that is characterised by weight loss, and histopathological evidence of chronic infection without a clearly diagnosed infection being present [74]. Notably it can cause stunting in children and here approximately one third of the young participants had stunted growth, adding further weight to this hypothesis. This is higher than national and regional prevalence, which in 2022 was reported to be 17% (20% in rural areas) and 17% in the Ashanti region in under 5 year olds, respectively [75]. One of the consequences of EED is malabsorption of nutrients (with the best evidence for zinc) [76], which could potentially affect the interpretation of our findings. On the other hand, IL-6 levels did not correlate inversely with those of zinc, suggesting that the degree of systemic inflammation in patients was not sufficient to cause the physiological and molecular changes that resulted in hypozincaemia [77]. Moreover, they were observed in the absence of a high level of CRP, which would be an indication of systemic inflammation [78]. Additional specific studies aimed at identifying EED, co-infections and systemic inflammation would be required to find out if this is a general feature of the population and whether this influences BU susceptibility and/or pathogenesis.

One interesting and unexpected finding was that, despite vitamin C having similar prevalence of dietary sufficiency (82%) and generally a good intake in their diets, many participants had serum vitamin C levels below the assumed normal range (55-88% depending on the cut-off used). This could potentially be attributed to these measurements being taken from two independent cohorts. However, both cohorts came from the same distribution of BU-endemic communities surrounding Agogo, Tepa and Dunkwa, ruling out local availability of food. Moreover, since BU disease has become less common, both cohorts were recruited over a long period (Cohort 1; July 2018 to Mary 2022, Cohort 2; October 2008 to June 2016), and therefore experienced similar seasonal variation in food availability. Therefore, it seems that even in the setting of adequate vitamin C intake, this did not translate to normal circulating vitamin C. This could be due to several factors, including that the main source of vitamin C is likely to be stews that may have been boiled for extensive periods, meaning that the intake was an over-estimate as the nutrient was destroyed [79]. It is also possible that participants over-estimated their intake during the 24-hour recall for psychological reasons or that EED could contribute since intestinal inflammation may be a confounding factor in vitamin C status [67, 74] as it is absorbed in the jejunum and ileum.

Insights into the role of diet in the pathogenesis of BU

While we could see no difference in the patterns of foods consumed between the BU cases and controls in the FFQ, the 24-hour recall showed that the cases consumed diets with a significantly lower DDS than the controls. This is in line with our findings that macronutrient intake (predominately carbohydrate) was similar between BU cases and controls, but the intake of micronutrients was different. In other words, it suggests that the lower dietary diversity of the BU patients could be one cause of the lower micronutrient intake due to the narrower range of foods eaten. In the context of a predominantly carbohydrate diet, a recent mouse study is of interest when considering underlying risk factors for BU in this population. Here, a ketogenic diet was modelled using b-hydroxybutyrate (the main ketone body produced in such diets), and this suppressed skin ulceration by stopping *M. ulcerans* growth and mycolactone synthesis with contrasting results observed in mice with a carbohydrate-rich diet [69]. Therefore, the bacterium seems to require a diet rich in carbohydrates for growth and mycolactone production, which is more than adequately provided by the diet of the participants.

When considering micronutrient differences in detail, cases' diets contained significantly less selenium and vitamin B_{12} and they were more likely to have dietary insufficiencies of these micronutrients. Vitamin B_{12} deficiency is prevalent in regions with a high burden of NTDs including helminth infection, Chagas disease and leprosy [80]. However, vitamin B_{12} was recently shown to be an important host-requirement for Mtb infection, as it has lost the biosynthetic machinery for this vitamin [81]. In contrast, selenium, a trace element, is required to clear mycobacteria [15]. In *M. tuberculosis* infection, selenium was shown to have a restricting function against MTB by regulating autophagy in macrophages contributing to low levels in patients compared to controls [82].

Zinc

Another of the largest differences in micronutrient intake between BU cases and controls was in zinc, which was 20% lower in cases. Here we were able to confirm, in an independent cohort, that serum levels were similarly 25% lower in cases than control. Moreover, BU patients were much more likely to have serum zinc levels below that of the assumed normal range than control counterparts. We are confident this represents a genuine micronutrient deficiency, since CRP levels were extremely low in both groups (below 0.05 mg/dL) so we did not need to correct this [83]. Moreover, our IL-6 data supports that both cases and controls had equal exposure to other infections and/or EED, and hence BU is associated with an additional reduction in zinc. Our findings relating to circulating cytokines are in line with previous reports of BU, which were found to be either unchanged or reduced compared to control groups [70, 84]. This is thought to be due to the anti-inflammatory activity of mycolactone, which blocks the Sec61 translocon, causing a sharp reduction in the production of proinflammatory cytokines that are reliant on Sec61 for secretion [6]. Therefore, while it is difficult to separate cause vs. effect for micronutrient measurement, our finding that BU patients' nutritional intake of zinc was lower than controls supports a nutritional component to the susceptibility to disease.

For BU patients, inadequate zinc may be implicated in disease susceptibility and progress due to the role that this mineral plays particularly in the immune response. Indeed, zinc supplementation has been reported to be an effective solution for immune recovery and wound healing [85]. Zinc deficiency has been shown to decrease the production of TNF-α, IFN-γ and IL-2 by peripheral blood mononuclear cells [86]. Zinc deficiency is reported to

increase susceptibility to leishmaniasis and TB [87], and in TB and leprosy hypozincaemia has been observed at the time of diagnosis [17]. In leprosy, there is a gradual reduction in serum zinc concentration as severity moves from Tuberculoid Leprosy to Lepromatous Leprosy, affecting Th1/ Th2 cytokine responses [88]. Furthermore, in humans, low serum zinc concentrations in cases with cutaneous leishmaniasis increase during antimonial therapy. This may be due to pre-existing zinc deficiencies or related to redistribution associated with the APR of visceral leishmaniasis [87, 89].

Vitamin C

The findings for vitamin C are more difficult to interpret. When comparing BU cases with controls, intake of vitamin C was >30% higher in BU cases of Cohort 1 although this did not reach statistical significance. On the other hand, serum levels were 20% lower in Cohort 2, and a higher proportion of BU cases had abnormal serum levels that was highly significant. Furthermore, within the BU cases, there was also an association between BU lesion category and serum levels of vitamin C (in Cohort 2). Indeed, those with the more serious plaque form of the disease had significantly lower vitamin C serum levels than those with its earliest presentation, the nodule. Since all nodules are Category I lesions, it is notable that the few Category III lesions in Cohort 2 were all associated with vitamin C levels below the lower limit of the normal range. While these were approximately half those of the Category I group, this difference did not reach statistical significance, perhaps due to the low numbers in this sub-group.

Therefore, while acknowledging that dietary and biomarker data were collected in different cohorts, our data suggests that the presence of *M. ulcerans* infection itself can affect vitamin C status. Notably, others have found this to be the case in patients with sepsis [90] and a cross-sectional analysis of the NHANES database found a negative correlation between inflammatory markers and vitamin C [39] possibly due to increased demands from the inflammatory response. Although vitamin C levels did not correlate with IL-6 in the current work, it may be that the presence of chronic, localised inflammation in BU patients is influencing the vitamin C levels in these patients.

Vitamin C is well-known to have important functions in skin health [91] as it is required for the production and stability of collagen [91, 92]. Furthermore, a deficiency in vitamin C dysregulates both phagocytosis and the respiratory burst [93]. The stimulation of proinflammatory and Th1 cytokines is lower during vitamin C deficiency as it alters redox signalling, essential during differentiation of Th cells [94]. We speculate that mycolactone's known effect on extracellular collagen [8, 95] could synergise with these lower vitamin C levels to have serious consequences for skin integrity.

Limitations and conclusion

There are some limitations that should be noted in this study. Here, dietary patterns and dietary intake of the study group were assessed using methodological tools that rely on the memory of the subjects, which can be biased. It is known that participants tend to underestimate or overestimate dietary recalls, especially in food frequency questionnaires, even though efforts were made to reduce possible biases to the minimum. The study also includes two independent cohorts of BU patients and controls that were matched as closely as possible. In Cohort 2, anthropometry data was not collected when these were recruited, and so could not be reported It was not possible to determine the inflammation markers in all patients, and our attempt to measure vitamin B_{12} in this same cohort was subject to assay failure.

Despite these limitations, these findings align with two recent independent reports of BU patients in Côte d'Ivoire [96, 97]. While there was no information on controls, in one study, 51 BU patients also ate predominantly starchy foods and minimal proteins; indeed, dairy products, fruits, vegetables, and vegetable oils were consumed by only 8% of the cases [97]. Their food choices were found to be influenced by societal norms where food types were prohibited in their communities, including certain animal and plant proteins, vegetables, and fruits. Being underweight tended to be higher in Côte d'Ivoire vs Ghanaian BU cases (33-53% vs 19-23%), and acute malnutrition was also reported [97]. Interestingly, CRP levels >20 mg/L were reported in 7-20% of the cases, suggesting that EED may be more prevalent in Côte d'Ivoire. However, in a subset of these patients and similar to here, zinc was found to be below normal levels in all BU patients [96]. Importantly, nutritional assistance provided at one health centre significantly improved wound healing alongside marked improvement in anthropometrics and anaemia compared to another where no nutritional support was included in BU patients' management package [97]. Taken together, the findings strongly support large scale randomised control studies for nutritional intervention in at-risk communities for BU and nutritional support for all BU patients, alongside a detailed assessment of the prevalence of EED and how this may influence susceptibility to BU and other NTDs.

Acknowledgement

The authors would like to thank all study participants and community leaders. We would like to express our utmost gratitude to all nurses, diseases control officers, and physiotherapists from Agogo Presbyterian Hospital, Tepa Government Hospital and Dunkwa Government Hospital for their support in patient management. We are grateful to the District Directors and the National Buruli ulcer Control programme. We thank Dr Peter Williams of University of Surrey, FHMS for statistical advice. This work was funded by a Wellcome Trust Investigator Award to Prof Rachel Simmonds (WT202843_Z_16),the Nestle Foundation (Lausanne, Switzerland) and the University of Surrey.

Author contributions

Conceptualization: RES, RP, BE, AL, RA

Formal Analysis: AL, RES

Funding Acquisition: AL, RA, RP, RES

Investigation: AL, PA, AKM, GA, EO, SM, JA, MO

Methodology: AL, RES

Project Administration: AL, RES, RP, BE, RA

Resources: AL, RES, RP

Supervision: RES, RA, BE, RP

Validation: RES, BE, RA, AL

Visualization: AL, RES

Writing—original draft preparation: AL, RES

Writing—review and editing, RA, RES, BE, AL, RP

Figure legends

Figure 1. Comparison of dietary diversity between BU cases and controls.

The dietary diversity score (DDS) was generated from 24-hour recall data obtained from all 80 Cohort 1 participants (40 BU cases, 40 controls), which assessed whether each participant ate any food from a particular group, out of 15, over the three days of the multi-pass (2 weekdays and one weekend day). The food groups for the were cereals, roots & tubers, green leafy vegetables, other vegetables, fruits, meat, fish & seafood, eggs, pulses & nuts, milk & milk products, oils & fats, sugar, condiments, drinks & beverages, and 'other'. Dietary diversity between BU cases and controls was compared using a two-tailed Mann-Whitney test. A p-value ≤ 0.05 was considered statistically significant.

Figure 2. BU patients have significantly lower zinc and vitamin C serum levels than controls.

The serum concentrations of zinc and vitamin C were assessed in BU cases and controls from Cohort 2 (40 BU cases, 40 controls). Individual data representing the average of duplicate assays is displayed along with the median and interquartile range. The expected normal ranges for each analyte are indicated with the dotted line (zinc 65-130µg/dL; vitamin C 0.4/0.6-2 mg/dL). P-values are displayed, with P < 0.05 considered significant. **A.** Data were compared using a one-tailed Mann-Whitney test. **B&C.** Subgroup analysis for BU cases with different clinical forms of the disease (N = 19, 18, and 42 for nodule, plaque and ulcer respectively, **B**) and different WHO lesion categories at presentation (N = 42, 32, and 6 for category I, category II, and category III lesions respectively, **C**). Data were compared between groups using the Kruskal-Wallis test and Dunn post-hoc test for non-parametric data.

Figure 3. IL-6 levels are not significantly different between BU patients vs controls, between BU lesion forms or lesion category. Serum IL-6 was quantified in 45 BU patients and 47 controls and presented on a $\log_{10}$ scale. **A.** Data for patients and controls were compared using a two-tailed Mann-Whitney test. **B.** Levels were compared between BU patients with different clinical presentations of the disease (Nodules, N=6; Plaques; N=11 and Ulcer; N=27), using the Kruskal-Wallis test with Dunn's correction for multiple comparisons. **C.** Levels were compared between BU patients with different WHO lesion category at presentation (Category I; N=21, Category II; N=21 and Category III; N=3), using the Kruskal-Wallis test with Dunn's correction for multiple comparisons. GraphPad Prism version 9.4.1 (GraphPad Software Inc., USA) was used for all analyses. P values are displayed, with P < 0.05 considered significant.

Figure S1. The most and least frequently consumed food groups by the study participants.

A food frequency questionnaire was put to all 80 Cohort 1 participants (40 BU cases, 40 controls), covering a total of 105 foods from 25 food groups. There are 4 possible answers for frequency of consumption for each food, namely daily, weekly, monthly, and rarely/never. This figure visualises the distribution of the most frequent consumption of one or more foods in that food group in the past month amongst the participants. The red line represents the frequency of consumption, and its length is proportional to the number of participants with that maximal frequency. Data from the food groups showed that at least 60% of the participants ate foods from these food groups at least weekly, where similar numbers consumed foods monthly, rarely or never vs. weekly or daily (N-Q), and those where at least 60% of the participants ate them monthly, rarely or never (R-Y).

Figure S2. No correlation between nutrition and inflammation markers in BU cases and controls.

Serum concentrations of vitamin C, zinc and IL-6 were analysed using Spearman's correlation, and are presented as a heatmap. Spearman's correlation coefficients are given along with P-value (in brackets).

Table 1. Characteristics of study participants.

Variables	Cohort 1			Cohort 2		
	BU cases N=40	Controls N=40	P-value	BU cases N=80	Controls N=80	P-value
Age range (years)	5-63	5-58	0.530[a]	5-56	5-62	0.9599[a]
Median (IQR)	17 (10-38)	21.5 (13-29)		14.5 (10-25)	14.5 (10-28)	
Gender, N (%)						
Male	19 (48%)	22 (55%)	0.654[b]	39 (49)	40 (50)	>0.9999[b]
Female	21 (52%)	18 (45%)		41 (51)	40 (50)	
Lesion type, N (%)						
Nodule	3 (8%)	-		19 (24%)	-	
Plaque	11 (28%)	-		18 (23%)	-	
Oedema	0 (0%)	-		1 (1%)	-	
Ulcer	25 (63%)	-		42 (53%)	-	
Osteomyelitis	1 (3%)	-		0 (0%)	-	
Lesion Category, N (%)						
I	14 (35%)	-		42 (53%)	-	
II	18 (45%)	-		32 (40%)	-	
III	8 (20%)	-		6 (8%)	-	
Waist-hip ratio[d]						
Median	0.88	0.90	0.643[a]			
IQR	0.73-1.02	0.77-0.96				
Weight categorisation[d,e], N (%)						
All	40	40		69		
Underweight	9 (22.5%)	5 (12.5%)	0.582[c]	13 (19%)	N/A	
Normal	27 (67.5%)	31 (77.5%)		51 (74%)	N/A	
Overweight	3 (7.5%)	2 (5%)		5 (7%)	N/A	
Obese	1 (2.5%)	2 (5%)				
Adult (≥20yr)	18	20		23		
Underweight	3 (16.7%)	3 (15%)		0 (0%)	N/A	
Normal	13 (72.2%)	15 (75%)		19 (83%)	N/A	
Overweight	1 (5.6%)	1 (5%)		4 (17%)	N/A	
Obese	1 (5.6%)	1 (5%)				
Young people (5-19yr)	22	18		46		
Underweight	6 (27.3%)	1 (5.6%)		13 (28%)	N/A	
Normal	14 (77.8%)	15 (83.3%)		32 (70%)	N/A	
Overweight	2 (11.1%)	1 (5.6%)		1 (2%)	N/A	
Obese	0 (0%)	1 (5.6%)				
Height-for-age status (5-19yr)[d,f], N (%)						
Not stunted	15 (68.2%)	14 (77.8%)	0.734[b]	30 (65%)	N/A	
Moderately stunted	7 (31.8%)	4 (22.2%)		13 (28%)	N/A	
Severely stunted	0 (0%)	0 (0%)		3 (7%)		

Counts of demographic, clinical, and socioeconomic characteristics of study participants. [a]; Two-tailed Mann-Whitney tests, [b]; Fisher's exact test, [c]; χ^2 test (p-value < 0.05 was considered statistically significant), [d]; Anthropometric data was available for 69 BU patients and no controls in Cohort 2, [e]; Weight categorisation used BMI-for-age for study participants aged 5-19 [31], and BMI for adults, [f]; stunting according to WHO AnthroPlus. Abbreviation: SD; Standard Deviation, IQR; Interquartile range, N/A: not available.

Table 2. Comparison of nutrient intake amounts between Cohort 1 participants.

Variables	All N=80 Mean± SD	BU Cases N=40 Mean± SD	Controls N=40 Mean± SD	P-value
Energy (Kcal)	1683±837	1702±994.4	1664±655.9	0.840
Carbohydrate (g)	291±259	258.8±118.4	322.7±345.7	0.274
Protein (g)	42.1±19.6	38.58±16.35	45.63±22.05	0.109
Fat (g)	54.1±68.6	60.76±95.15	47.52±19.88	0.394
Fibre (g)	23.3±10.7	23.44±10.11	23.23±11.29	0.930
Folate (µg)	296±176	302.2±181.5	289.8±172.6	0.754
Iron (mg)	9.90±4.68	9.50±4.59	10.29±4.80	0.450
Selenium (µg)	65.00±40.5	52.14±31.85	77.35±44.53	**0.005**
Vitamin A (µg)	610±719	599.8±655.1	620.0±785.8	0.901
Vitamin B_{12} (µg)	1.61±2.11	0.89±0.92	2.332±2.68	**0.002**
Vitamin C (mg)	110±77.0	124.7±91.33	95.01±56.54	0.086
Vitamin E (mg)	6.21±3.43	6.079±3.964	6.340±2.86	0.736
Vitamin K (µg)	50.00±45.3	48.97±39.29	50.94±51.14	0.850
Zinc (mg)	6.20±2.50	5.57±1.70	6.90±2.96	**0.011**

Data were compared between cases and controls using an unpaired t-test. P-value <0.05 are indicated in bold text.

Table 3. Percentage adequacy of nutrient intake in Cohort 1 participants.

Variables	Frequency	All N=80	BU cases N=40	Controls N=40	P- value[a]
Energy (Kcal)	Adequate	12 (15%)	6 (15%)	6 (15%)	1.000
	Inadequate	68 (85%)	34 (85%)	34 (85%)	
Carbohydrate (g)	Adequate	79 (99%)	40 (100%)	39 (98%)	1.000
	Inadequate	1 (1%)	0	1 (2%)	
Protein (g)	Adequate	41 (51%)	18 (45%)	23 (57%)	0.371
	Inadequate	39 (49%)	22 (55%)	17 (43%)	
Fibre (g)	Adequate	28 (35%)	17 (43%)	11 (28%)	0.241
	Inadequate	52 (65%)	23 (58%)	29 (73%)	
Folate (µg)	Adequate	40 (50%)	21 (52%)	19 (47%)	0.823
	Inadequate	40 (50%)	19 (48%)	21 (53%)	
Iron (mg)	Adequate	59 (74%)	29 (72%)	30 (75%)	1.000
	Inadequate	21 (26%)	11(14%)	10 (25%)	
Selenium (µg)	Adequate	58 (72%)	24 (60%)	34 (85%)	**0.023**
	Inadequate	22 (28%)	16 (40%)	6 (15%)	
Vitamin A (µg)	Adequate	32 (40%)	18 (45%)	14 (35%)	0.494
	Inadequate	48 (60%)	22 (55%)	26 (65%)	
Vitamin B_{12} (µg)	Adequate	26 (32%)	8 (20%)	18 (45%)	**0.031**
	Inadequate	54 (68%)	32 (80%)	22 (55%)	
Vitamin C (mg)	Adequate	66 (82%)	36 (90%)	30 (70%)	0.139
	Inadequate	14 (18%)	4 (10%)	10 (25%)	
Vitamin E (mg)	Adequate	11 (14%)	8 (20%)	3 (7%)	0.193
	Inadequate	69 (86%)	32 (80%)	37 (93%)	
Vitamin K (µg)	Adequate	18 (22%)	11 (28%)	7 (18%)	0.423
	Inadequate	62 (78%)	29 (73%)	33 (83%)	
Zinc (mg)	Adequate	26 (32%)	12 (30%)	14 (35%)	0.812
	Inadequate	54 (68%)	28 (70%)	26 (65%)	

Nutritional adequacy was determined using the Estimated Average Requirement (EAR) for all nutrients except for energy, fibre and vitamin K that do not have an EAR, and here Recommended Dietary Allowance (RDA) was used instead. Both vary depending on age, gender, height, pregnancy, and lactation (although pregnant and lactating mothers were excluded from the study). [a] two-tailed Fisher's exact test. P-value <0.05 are indicated in bold text.

Table 4. Serum zinc and vitamin C below the normal range are more common in BU patients than controls in Cohort 2.

Markers	Description	Threshold	All, N (%)	BU Cases, N (%) N=80	Controls, N (%) N=80	P-value[a]
Zinc	Normal	>60 µg/dL	68 (42%)	22 (28%)	46 (57%)	0.0002
	Reduced	≤60 µg/dL	92 (58%)	58 (72%)	34 (43%)	
Vitamin C	Normal	>0.6 mg/dL	19 (12%)	2 (3%)	17 (21%)	0.0003
	Reduced	≤0.6 mg/dL	141 (88%)	78 (97%)	63 (79%)	
	Normal	>0.4 mg/dL	72 (45%)	16 (20%)	56 (70%)	<0.0001
	Reduced	≤0.4 mg/dL	88 (55%)	64 (80%)	24 (30%)	

[a] Fisher's exact test with P < 0.05 considered statistically significant.

Table 5. Comparable serum concentration of inflammation markers in patients and controls in Cohort 2.

Markers Total	Description	Threshold	BU Cases, N (%) 45	Controls, N (%) 47	P-value
hs-CRP	Normal	<0.6 mg/L[a]	80 (100)	80 (100)	ND
	High normal	0.6-3 mg/L	0 (0)	0 (0)	
	Minor	3-10 mg/L	0 (0)	0 (0)	
	Moderate	10-80 mg/L	0 (0)	0 (0)	
	Major	>80 mg/L	0 (0)	0 (0)	
IL-1β	Normal/slightly elevated	≤57 pg/mL[b]	42 (93)	41 (87)	0.4861
	Elevated	>57 pg/mL	3 (7)	6 (13)	
TNF-α	Normal	<7 pg/mL[c]	37 (82)	36 (77)	0.6094[e]
	Slightly elevated	7-50 pg/mL	6 (13)	9 (19)	
	Elevated	>50 pg/mL	2 (5)	2 (4)	
IFN-γ	Normal	<5 pg/mL[d]	32 (71)	34 (72)	>0.9999[e]
	Slightly elevated	5-50 pg/mL	12 (27)	12 (26)	
	Elevated	>50 pg/mL	1 (2)	1 (2)	
IL-6	Normal	<7 pg/mL	21 (47)	22 (49)	0.8645[f]
	Slightly elevated	7-50 pg/mL	15 (33)	16 (36)	
	Elevated	50-500 pg/mL	6 (13)	6 (13)	
	Highly elevated	>500 pg/mL	3 (7)	1 (2)	

Inflammation markers serum level concentration of study participants grouped according to thresholds that indicate the degree to which the marker was elevated. Numbers in each category were compared using contingency tables, using Fisher's exact or chi-square tests with P < 0.05 considered statistically significant. [a] Includes 67 patients and 72 controls whose CRP levels were lower than the detection limit of the assay. [b] Includes 32 patients and 36 controls whose IL-1β levels were lower than the detection limit of the assay. [c] Includes 28 patients and 30 controls whose TNFα levels were lower than the detection limit of the assay. [d] Includes 18 patients and 24 controls whose IFNγ levels were lower than the detection limit of the assay. [e] Participants in slightly elevated and elevated categories had to be combined for this analysis. [f] Elevated and highly elevated categories had to be combined for this analysis.

S1 Table. List of foods used in the FFQ administration.

Food group	Food items
Whole grain cereals	Oats porridge, maize porridge, millet porridge, wheat porridge, whole meal bread/roll, Tuo zafi (T.Z.)
Refined cereals	Sugar coated cereals, cornflakes, white bread/roll, pancake, flour chips/achumo, kaafa
Fermented maize products	Kenkey, Banku, Akple
Rice and pasta	White rice cooked, local rice cooked, noodles, macaroni
Roots, tubers and plantain	Yam, cassava or cocoyam, plantain, fufu, kokonte, gari
Potatoes	Boiled instant potatoes/oven baked potatoes, fried potatoes chips, roast potatoes
Nuts and seeds	Cashew nuts, groundnut/pea nuts, almond/hazelnut, pistachio, coconut, sultana/raisins/currants
Legumes	Kidney beans, soya beans, black eyed beans, baked beans, chicks peas, runner/green beans, ground nut soup
Dairy products	Milk, full fat yoghurt, full fat cheese, low fat cheese, butter/margarine, cocoa, horlicks, milkshake/fula
Eggs	Egg boiled/poached, egg fried/omelette
Red meat	Beef/lamb/pork, bush meat, rabbit
Poultry/other white meat	Chicken/other poultry, snail
Processed meat	Burger, bacon, ham, sausage/luncheon meat/corned beef
Fish *and shellfish*	Fish, fish fingers, fish fried in butter, crabs, prawn, octopus
Vegetarian mixed dishes	Jollof rice, fried rice, waakye, soya bean curd/tofu, beans stew
Vegetable soups, stews, sauces	Tomato stew, melon seed and spinach stew, palmnut soup, okro soup/stew, light soup, kontomire (cocoyam leaves)
Vegetables	Sweet corn, okro/nkruma, garden egg/aubergine, avocado, garden peas/peas, mushroom, broccoli/cauliflower, carrot, cabbage, brussels sprouts, ayoyo, spring onions, onions, tomatoes, tin-tomato, sweet peppers, lettuce, cucumber, coleslaw, beetroot, green pepper, Ghanaian green pepper (Kpakposhitor),
Fruits	Banana, pears, pomegranates, mangoes, pineapple, grapefruits, orange/satsuma, grapes, melon, apple, peach/plum/nectarine, strawberries/cherries, pawpaw
Sweet spreads	Jam, marmalade, honey, peanut butter
Cakes and sweets, and sweet puddings	Sugar added to cereals, tea, coffee, cakes/scones/doughnuts, ice cream/frozen desserts, sweets biscuits, chocolate coated sweet biscuits, chocolate/chocolate bar, custard, rice pudding, sweets, toffees, mints
Savoury snacks	Crisp/pocket snacks, fried plantain chips, salty biscuits, regular popcorn, buttered popcorn
Condiments	Hot pepper sauce, tomato ketchup, salad pepper (powdered)
Alcoholic beverages	Beer/larger/cider, port/sherry/liqueur, spirits, pito/solom, palmwine (alcohol)
Sodas and juices	Real fruit juice, fruit squash, fizzy soft drink, low calorie fizzy drink, bissap (sobolo), liha (sprout maize), palmwine (fresh)
Coffee and tea	Tea, Coffee

S2 Table. Food groups used in the Dietary Diversity Score, generated from 24-hour recall data.

cereals	roots & tubers	green leafy vegetables
other vegetables	fruits	meat
fish & seafood	eggs	pulses & nuts
milk & milk products	oils & fats	sugar
condiments	drinks & beverages	'other'

S3 Table. Socioeconomic status of Cohort 1 study participants.

	Cohort 1		
Variables	**BU cases** n=40	**Controls** n=40	**P-value**
Education			0.236[a]
Primary	24 (58.5%)	18 (45%)	
Secondary	8 (19.5%)	16 (40%)	
Tertiary	0	2 (5%)	
None	9 (22%)	4 (10%)	
Marital Status			0.769[a]
Schooling	20 (50%)	22 (55%)	
Single	6 (15%)	6 (15%)	
Married	13 (33%)	12 (30%)	
Divorced	1 (3%)	0	
Household monthly income			0.478[a]
<GH₵ 50	6 (15%)	4 (10%)	
GH₵ 50-200	10 (25%)	15 (38%)	
GH₵201-800	23 (58%)	21(53%)	
>GH₵800	0	0	
Occupation			
Formal	0	0	0.236[a]
Informal	15 (38%)	17 (43%)	
Attending School	20 (50%)	22 (55%)	
Unemployed	5 (13%)	1 (3%)	

[a]; T χ² test. p-value < 0.05 was statistically significant.

S4 Table. Mean percentage of macronutrients in the diet of Cohort 1 participants.

| Nutrients | Mean nutrient intake (%) | | |
	All N=80	BU Cases N=40	Controls N=40
Energy (Kcal)	1683 (100%)	1705 (100%)	1664 (100%)
Carbohydrate (g)	291 (64.0%)	258.8 (60.0%)	322.7 (67.9%)
Protein (g)	42.1 (9.0%)	38.58 (9.0%)	45.63 (9.6%)
Fats (g)	54.1 (27.0%)	60.76 (31.0%)	47.52(22.5%)

Unit conversions: 1g of carbohydrate is equivalent to 4 Kcal, 1g of protein is equivalent to 4 Kcal and 1g of fats is equivalent to 9 Kcal, which were then used to calculate percentages.

References

1.   Yotsu RR, Suzuki K, Simmonds RE, Bedimo R, Ablordey A, Yeboah-Manu D, et al. Buruli Ulcer: a Review of the Current Knowledge. Curr Trop Med Rep. 2018;5(4):247-56. doi: 10.1007/s40475-018-0166-2.

2.   Yotsu RR, Fuller LC, Murdoch ME, van Brakel WH, Revankar C, Barogui MYT, et al. A global call for action to tackle skin-related neglected tropical diseases (skin NTDs) through integration: An ambitious step change. PLoS Negl Trop Dis. 2023;17(6):e0011357. doi: 10.1371/journal.pntd.0011357.

3.   Portaels F, Silva MT, Meyers WM. Buruli ulcer. Clin Dermatol. 2009;27(3):291-305. doi: 10.1016/j.clindermatol.2008.09.021.

4.   George KM, Chatterjee D, Gunawardana G, Welty D, Hayman J, Lee R, et al. Mycolactone: a polyketide toxin from Mycobacterium ulcerans required for virulence. Science. 1999;283(5403):854-7. doi: 10.1126/science.283.5403.854.

5.   Sarfo FS, Phillips R, Wansbrough-Jones M, Simmonds RE. Recent advances: role of mycolactone in the pathogenesis and monitoring of Mycobacterium ulcerans infection/Buruli ulcer disease. Cell Microbiol. 2016;18(1):17-29. doi: 10.1111/cmi.12547.

6.   Hall BS, Hill K, McKenna M, Ogbechi J, High S, Willis AE, et al. The pathogenic mechanism of the Mycobacterium ulcerans virulence factor, mycolactone, depends on blockade of protein translocation into the ER. PLoS Pathog. 2014;10(4):e1004061. doi: 10.1371/journal.ppat.1004061.

7.   Demangel C, High S. Sec61 blockade by mycolactone: A central mechanism in Buruli ulcer disease. Biol Cell. 2018;110(11):237-48. doi: 10.1111/boc.201800030.

8.   Hsieh LT, Hall BS, Newcombe J, Mendum TA, Umrania Y, Deery MJ, et al. Mycolactone causes catastrophic Sec61-dependent loss of the endothelial glycocalyx and basement membrane: a new indirect mechanism driving tissue necrosis in Mycobacterium ulcerans infection. bioRxiv. 2023. doi: 10.1101/2023.02.21.529382.

9.   Fevereiro J, Fraga AG, Pedrosa J. The Immunology of Buruli Ulcer. In: Pluschke G, Röltgen K, editors. Buruli ulcer: Mycobacterium ulcerans disease. Cham (CH): Springer; 2019. p. 135-58. doi: 10.1007/978-3-030-11114-4_8.

10.   Wansbrough-Jones M, Phillips R. Buruli ulcer: emerging from obscurity. Lancet. 2006;367(9525):1849-58. doi: 10.1016/s0140-6736(06)68807-7.

11.   Foulon M, Robbe-Saule M, Manry J, Esnault L, Boucaud Y, Alcaïs A, et al. Mycolactone toxin induces an inflammatory response by targeting the IL-1β pathway: Mechanistic insight into Buruli ulcer pathophysiology. PLoS Pathog. 2020;16(12):e1009107. doi: 10.1371/journal.ppat.1009107.

12.   Capela C, Dossou AD, Silva-Gomes R, Sopoh GE, Makoutode M, Menino JF, et al. Genetic Variation in Autophagy-Related Genes Influences the Risk and Phenotype of Buruli Ulcer. PLoS Negl Trop Dis. 2016;10(4):e0004671. doi: 10.1371/journal.pntd.0004671.

13.   Bieri R, Bolz M, Ruf MT, Pluschke G. Interferon-γ Is a Crucial Activator of Early Host Immune Defense against Mycobacterium ulcerans Infection in Mice. PLoS Negl Trop Dis. 2016;10(2):e0004450. doi: 10.1371/journal.pntd.0004450.

14.   Zingue D, Bouam A, Tian RBD, Drancourt M. Buruli Ulcer, a Prototype for Ecosystem-Related Infection, Caused by Mycobacterium ulcerans. Clin Microbiol Rev. 2018;31(1). doi: 10.1128/cmr.00045-17.

15.   Gupta KB, Gupta R, Atreja A, Verma M, Vishvkarma S. Tuberculosis and nutrition. Lung India. 2009;26(1):9-16. doi: 10.4103/0970-2113.45198.

16.   Schwenk A, Macallan DC. Tuberculosis, malnutrition and wasting. Curr Opin Clin Nutr Metab Care. 2000;3(4):285-91. doi: 10.1097/00075197-200007000-00008.

17.   Dwivedi VP, Banerjee A, Das I, Saha A, Dutta M, Bhardwaj B, et al. Diet and nutrition: An important risk factor in leprosy. Microb Pathog. 2019;137:103714. doi: 10.1016/j.micpath.2019.103714.

18.   Passos Vázquez CM, Mendes Netto RS, Ferreira Barbosa KB, Rodrigues de Moura T, de Almeida RP, Duthie MS, et al. Micronutrients influencing the immune response in leprosy. Nutr Hosp. 2014;29(1):26-36. doi: 10.3305/nh.2014.29.1.6988.

19.   Raiten DJ, Sakr Ashour FA, Ross AC, Meydani SN, Dawson HD, Stephensen CB, et al. Inflammation and Nutritional Science for Programs/Policies and Interpretation of Research Evidence (INSPIRE). J Nutr. 2015;145(5):1039s-108s. doi: 10.3945/jn.114.194571.

20.   Gombart AF, Pierre A, Maggini S. A review of micronutrients and the immune system–working in harmony to reduce the risk of infection. Nutrients. 2020;12(1):236. doi: 10.3390/nu12010236.

21.   Loglo AD, Frimpong M, Duah MS, Sarfo F, Sarpong FN, Agbavor B, et al. IFN-γ and IL-5 whole blood response directed against mycolactone polyketide synthase domains in patients with Mycobacterium ulcerans infection. PeerJ. 2018;6:e5294. doi: 10.7717/peerj.5294.

22.   Sarpong-Duah M, Frimpong M, Beissner M, Saar M, Laing K, Sarpong F, et al. Clearance of viable Mycobacterium ulcerans from Buruli ulcer lesions during antibiotic treatment as determined by combined 16S rRNA reverse transcriptase/IS 2404 qPCR assay. PLoS Negl Trop Dis. 2017;11(7):e0005695. doi: 10.1371/journal.pntd.0005695.

23.   WHO. Skin NTDs clinical and treatment form [Document]. Genevea: World Health Organization 2020 [cited 2024 14th June]. Available from: https://www.who.int/docs/default-source/ntds/buruli-ulcer/recording-and-reporting-forms/1a-skin-ntd-bu01-2020-09-02.pdf.

24.   WHO. Treatment of Mycobacterium ulcerans disease (Buruli ulcer); guidance for health workers [Guidance document]. Geneva: Wold Health Organisation; 2012 [cited 2024 14th Jun]. Available from: https://www.who.int/publications/i/item/9789241503402.

25.   Nishida C, Uauy R, Kumanyika S, Shetty P. The joint WHO/FAO expert consultation on diet, nutrition and the prevention of chronic diseases: process, product and policy implications. Public Health Nutr. 2004;7(1a):245-50. doi: 10.1079/phn2003592.

26.   WHO. Growth reference data for 5–19 years: BMI-for-age (5–19 years) [Internet]. Geneva: World Health Organization 2024 [cited 2024 14th June]. Available from: https://www.who.int/tools/growth-reference-data-for-5to19-years/indicators/bmi-for-age.

27.   de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO growth reference for school-aged children and adolescents. Bull World Health Organ. 2007;85(9):660-7. doi: 10.2471/blt.07.043497.

28.   WHO. WHO AnthroPlus for personal computers Manual: Software for assessing growth of the world's children and adolescents Geneva: WHO; 2009 [cited 2024 14th June]. Available from: http://www.who.int/growthref/tools/en/

29.   Mantey AA, Annan RA, Lutterodt HE, Twumasi P. Iron status predicts cognitive test performance of primary school children from Kumasi, Ghana. PLoS One. 2021;16(5):e0251335.

30.   Azupogo F, Abizari A-R, Aurino E, Gelli A, Osendarp SJ, Bras H, et al. Malnutrition, hypertension risk, and correlates: An analysis of the 2014 ghana demographic and health survey data for 15–19 years adolescent boys and girls. Nutrients. 2020;12(9):2737.

31.   Galbete C, Nicolaou M, Meeks KA, Aikins Ad-G, Addo J, Amoah SK, et al. Food consumption, nutrient intake, and dietary patterns in Ghanaian migrants in Europe and their compatriots in Ghana. Food Nutr Res. 2017.

32.   Annan RA, Apprey C, Asamoah-Boakye O, Okonogi S, Yamauchi T, Sakurai T. The relationship between dietary micronutrients intake and cognition test performance among school-aged children in government-owned primary schools in Kumasi metropolis, Ghana. Food Sci Nutr. 2019;7(9):3042-51.

33.   Science F, Nutrition Department UoG. The nutrient analysis template software excel spreadsheet for Ghanaian foods. University of Ghana Accra; 2010.

34.   Opoku K. Estimation of food portion sizes: A comparison between the use of household measures and a photographic food atlas in a rural population in Ghana [PhD??? Thesis]: University of Ghana; 2017. Available from: Is this available??

35.   WHO, FAO. Vitamin and mineral requirements in human nutrition: report of a joint FAO/WHO expert consultation, Bangkok, Thailand, 21–30 September 1998. 2nd Edition [Report]. Geneva: World Health Organization 1998 [cited 2024 14th June]. Available from: https://www.who.int/publications/i/item/9241546123.

36.	Dietary reference values for food energy and nutrients for the United Kingdom. Report of the Panel on Dietary Reference Values of the Committee on Medical Aspects of Food Policy. Rep Health Soc Subj (Lond). 1991;41:1-210. doi: PMID: 1961974.

37.	FAO. Guidelines for measuring household and individual dietary diversity [Report]. Food and Agriculture Organization of the United Nations; 2010 [cited 2024 14th June]. Available from: https://www.fao.org/4/i1983e/i1983e00.pdf.

38.	Addo J, Agyemang C, de-Graft Aikins A, Beune E, Schulze MB, Danquah I, et al. Association between socioeconomic position and the prevalence of type 2 diabetes in Ghanaians in different geographic locations: the RODAM study. J Epidemiol Community Health. 2017;71(7):633-9. doi: 10.1136/jech-2016-208322. eng.

39.	Crook J, Horgas A, Yoon S-J, Grundmann O, Johnson-Mallard V. Insufficient vitamin C levels among adults in the United States: results from the NHANES surveys, 2003–2006. Nutrients. 2021;13(11):3910. doi: 10.3390/nu13113910.

40.	Gordon BL, Galati JS, Yang S, Longman RS, Lukin D, Scherl EJ, et al. Prevalence and factors associated with vitamin C deficiency in inflammatory bowel disease. World J Gastroenterol. 2022;28(33):4834-45. doi: 10.3748/wjg.v28.i33.4834.

41.	Rutkowski M, Grzegorczyk K. [Vitamin C in medicine: "normal concentration" in serum]. Pol Merkur Lekarski. 1999;6(31):57-60. Polish.

42.	King JC, Brown KH, Gibson RS, Krebs NF, Lowe NM, Siekmann JH, et al. Biomarkers of Nutrition for Development (BOND)-Zinc Review. J Nutr. 2015;146(4):858s-85s. doi: 10.3945/jn.115.220079.

43.	Held C, White HD, Stewart RAH, Budaj A, Cannon CP, Hochman JS, et al. Inflammatory Biomarkers Interleukin-6 and C-Reactive Protein and Outcomes in Stable Coronary Heart Disease: Experiences From the STABILITY (Stabilization of Atherosclerotic Plaque by Initiation of Darapladib Therapy) Trial. J Am Heart Assoc. 2017;6(10):e005077. doi: 10.1161/jaha.116.005077.

44.	Said EA, Al-Reesi I, Al-Shizawi N, Jaju S, Al-Balushi MS, Koh CY, et al. Defining IL-6 levels in healthy individuals: A meta-analysis. J Med Virol. 2021;93(6):3915-24. doi: 10.1002/jmv.26654.

45.	Kim HO, Kim H-S, Youn J-C, Shin E-C, Park S. Serum cytokine profiles in healthy young and elderly population assessed using multiplexed bead-based immunoassays. J Transl Med. 2011;9:1-7. doi: 10.1186/1479-5876-9-113.

46.	Spranger J, Kroke A, Mohlig M, Hoffmann K, Bergmann MM, Ristow M, et al. Inflammatory cytokines and the risk to develop type 2 diabetes: results of the prospective population-based European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam Study. Diabetes. 2003;52(3):812-7. doi: 10.2337/diabetes.52.3.812.

47.	Scrimshaw NS, San Giovanni JP. Synergism of nutrition, infection, and immunity: an overview. Am J Clin Nutr. 1997;66(2):464s-77s. doi: 10.1093/ajcn/66.2.464S.

48.	Caulfield LE, de Onis M, Blössner M, Black RE. Undernutrition as an underlying cause of child deaths associated with diarrhea, pneumonia, malaria, and measles. Am J Clin Nutr. 2004;80(1):193-8. doi: 10.1093/ajcn/80.1.193.

49.	WHO. Communicable Diseases and Severe Food Shortage: WHO Technical Note [Technical Note]. Geneva: World Health Organization; 2010 [cited 2024 14th June]. Available from: https://www.ncbi.nlm.nih.gov/books/NBK304204/.

50.	Ayensu J, Annan R, Lutterodt H, Edusei A, Peng LS. Prevalence of anaemia and low intake of dietary nutrients in pregnant women living in rural and urban areas in the Ashanti region of Ghana. PLoS One. 2020;15(1):e0226026. doi: 10.1371/journal.pone.0226026.

51.	WHO. Malnutrition [Internet]. Geneva: World Health Organization 2024 [cited 2024 14th June]. Available from: https://www.who.int/news-room/fact-sheets/detail/malnutrition/.

52.	Bosu WK. An overview of the nutrition transition in West Africa: implications for non-communicable diseases. Proc Nutr Soc. 2015;74(4):466-77. doi: 10.1017/S0029665114001669.

53.	Aberman N-L, Billings L, Hidrobo M, Verstraeten R, Salm L, van den Bold M. Building evidence for policy and program action for nutrition in Ghana [PHND Project Note].

International Food Policy Research Institute; 2018 [cited 2024 14th June]. Available from: http://ebrary.ifpri.org/cdm/singleitem/collection/p15738coll2/id/132558.

54.   Oniang'o RK, Mutuku JM, Malaba SJ. Contemporary African food habits and their nutritional and health implications. Asia Pac J Clin Nutr. 2003;12(3):331-6. doi: PMID: 14505997.

55.   Amugsi DA, Mittelmark MB, Lartey A. Dietary Diversity is a Predictor of Acute Malnutrition in Rural but Not in Urban Settings: Evidence from Ghana. Journal of Advances in Medicine and Medical Research. 2014;4(25):4310-24. doi: 10.9734/BJMMR/2014/10014.

56.   van Wesenbeeck CF, Keyzer MA, Nubé M. Estimation of undernutrition and mean calorie intake in Africa: methodology, findings and implications. Int J Health Geogr. 2009;8:37. doi: 10.1186/1476-072x-8-37.

57.   Heinbuch U. Animal protein sources for rural and urban populations in Ghana [Report]. Cotonou: Programme for the Integrated Development of Artisanal Fisheries in West Africa of the Food and Agriculture Organization of the United Nations; 1994 [cited 2024 14th June]. Available from: https://www.fao.org/4/ad341e/ad341e00.pdf.

58.   Chadare FJ, Affonfere M, Aidé ES, Fassinou FK, Salako KV, Pereko K, et al. Current state of nutrition in West Africa and projections to 2030. Global Food Security. 2022;32:100602. doi: 10.1016/j.gfs.2021.100602.

59.   Chagomoka T, Drescher A, Glaser R, Marschner B, Schlesinger J, Nyandoro G. Vegetable production, consumption and its contribution to diets along the urban–rural continuum in Northern Ghana. Afr J Food Agric Nutr Dev. 2015;15(4):10352-67. doi: 10.18697/ajfand.71.15530.

60.   FAO, IFAD, UNICEF, WFP, WHO. The state of food security and nutrition in the world 2018: building climate resilience for food security and nutrition [Report]. Rome: Food and Agriculture Organization of the United Nations; 2018 [cited 2024 14th June]. Available from: http://www.fao.org/3/I9553EN/i9553en.pdf.

61.   2020 global nutrition report: action on equity to end malnutrition [Report]. Bristol: Development Initiatives; 2020. Available from: https://globalnutritionreport.org/reports/2020-global-nutrition-report/.

62.   Rojas AI, Phillips TJ. Patients with chronic leg ulcers show diminished levels of vitamins A and E, carotenes, and zinc. Dermatol Surg. 1999;25(8):601-4. doi: 10.1046/j.1524-4725.1999.99074.x.

63.   Sinno S, Lee DS, Khachemoune A. Vitamins and cutaneous wound healing. J Wound Care. 2011;20(6):287-93. doi: 10.12968/jowc.2011.20.6.287.

64.   Hsieh LT, Dos Santos SJ, Hall BS, Ogbechi J, Loglo AD, Salguero FJ, et al. Aberrant stromal tissue factor localisation and mycolactone-driven vascular dysfunction, exacerbated by IL-1β, are linked to fibrin formation in Buruli ulcer lesions. PLoS Pathog. 2022;18(1):e1010280. doi: 10.1371/journal.ppat.1010280.

65.   Ogbechi J, Ruf MT, Hall BS, Bodman-Smith K, Vogel M, Wu HL, et al. Mycolactone-Dependent Depletion of Endothelial Cell Thrombomodulin Is Strongly Associated with Fibrin Deposition in Buruli Ulcer Lesions. PLoS Pathog. 2015;11(7):e1005011. doi: 10.1371/journal.ppat.1005011.

66.   Vermeer C, Shearer MJ, Zittermann A, Bolton-Smith C, Szulc P, Hodges S, et al. Beyond deficiency: potential benefits of increased intakes of vitamin K for bone and vascular health. Eur J Nutr. 2004;43(6):325-35. doi: 10.1007/s00394-004-0480-4.

67.   Gibson RS. Zinc: the missing link in combating micronutrient malnutrition in developing countries. Proc Nutr Soc. 2006;65(1):51-60. doi: 10.1079/pns2005474.

68.   Ritchie H, Roser M. Micronutrient deficiency [Internet]. UK: OurWorldInData.org; 2017 [cited 2024 14th June]. Available from: https://ourworldindata.org/micronutrient-deficiency.

69.   Wessells KR, Brown KH. Estimating the global prevalence of zinc deficiency: results based on zinc availability in national food supplies and the prevalence of stunting. PLoS One. 2012;7(11):e50568. doi: 10.1371/journal.pone.0050568.

70.   Phillips RO, Sarfo FS, Landier J, Oldenburg R, Frimpong M, Wansbrough-Jones M, et al. Combined inflammatory and metabolic defects reflected by reduced serum protein levels

in patients with Buruli ulcer disease. PLoS Negl Trop Dis. 2014;8(4):e2786. doi: 10.1371/journal.pntd.0002786.

71. Kumar NP, Moideen K, Banurekha VV, Nair D, Babu S. Plasma Proinflammatory Cytokines Are Markers of Disease Severity and Bacterial Burden in Pulmonary Tuberculosis. Open Forum Infect Dis. 2019;6(7):ofz257. doi: 10.1093/ofid/ofz257.

72. Joshi L, Ponnana M, Sivangala R, Chelluri LK, Nallari P, Penmetsa S, et al. Evaluation of TNF-α, IL-10 and IL-6 cytokine production and their correlation with genotype variants amongst tuberculosis patients and their household contacts. PLoS One. 2015;10(9):e0137727. doi: 10.1371/journal.pone.0137727.

73. Ozarslan N, Robinson JF, Gaw SL. Circulating Monocytes, Tissue Macrophages, and Malaria. J Trop Med. 2019;2019:3720838. doi: 10.1155/2019/3720838.

74. Tickell KD, Atlas HE, Walson JL. Environmental enteric dysfunction: a review of potential mechanisms, consequences and management strategies. BMC Med. 2019;17(1):181. doi: 10.1186/s12916-019-1417-3.

75. GSS, ICF. Ghana Demographic and Health Survey 2022 [Report]. Accra Ghana and Rockville Maryland USA2024 [cited 2024 14th June]. Available from: https://dhsprogram.com/pubs/pdf/PR149/PR149.pdf.

76. Kelly P. The contribution of environmental enteropathy to the global problem of micronutrient deficiency. Proc Nutr Soc. 2021;80(3):303-10. doi: 10.1017/S0029665121000549.

77. Tanaka T, Narazaki M, Kishimoto T. IL-6 in inflammation, immunity, and disease. Cold Spring Harb Perspect Biol. 2014;6(10):a016295. doi: 10.1101/cshperspect.a016295.

78. Del Giudice M, Gangestad SW. Rethinking IL-6 and CRP: Why they are more than inflammatory biomarkers, and why it matters. Brain Behav Immun. 2018;70:61-75. doi: 10.1016/j.bbi.2018.02.013.

79. Lee S, Choi Y, Jeong HS, Lee J, Sung J. Effect of different cooking methods on the content of vitamins and true retention in selected vegetables. Food Sci Biotechnol. 2018;27(2):333-42. doi: 10.1007/s10068-017-0281-1.

80. Gibson RS, Charrondiere UR, Bell W. Measurement errors in dietary assessment using self-reported 24-hour recalls in low-income countries and strategies for their prevention. Adv Nutr. 2017;8(6):980-91. doi: 10.3945/an.117.016980.

81. Campos-Pardos E, Uranga S, Picó A, Gómez AB, Gonzalo-Asensio J. Dependency on host vitamin B12 has shaped Mycobacterium tuberculosis complex evolution. Nat Commun. 2024;15(1):2161. doi: 10.1038/s41467-024-46449-8.

82. Chen W, Liu Z, Zheng Y, Wei B, Shi J, Shao B, et al. Selenium donor restricts the intracellular growth of Mycobacterium tuberculosis through the induction of c-Jun-mediated both canonical autophagy and LC3-associated phagocytosis of alveolar macrophages. Microb Pathog. 2021;161:105269. doi: 10.1016/j.micpath.2021.105269.

83. McMillan DC, Maguire D, Talwar D. Relationship between nutritional status and the systemic inflammatory response: micronutrients. Proc Nutr Soc. 2019;78(1):56-67. doi: 10.1017/S0029665118002501.

84. Phillips R, Sarfo FS, Guenin-Macé L, Decalf J, Wansbrough-Jones M, Albert ML, et al. Immunosuppressive signature of cutaneous Mycobacterium ulcerans infection in the peripheral blood of patients with buruli ulcer disease. J Infect Dis. 2009;200(11):1675-84. doi: 10.1086/646615.

85. Blass SC, Goost H, Tolba RH, Stoffel-Wagner B, Kabir K, Burger C, et al. Time to wound closure in trauma patients with disorders in wound healing is shortened by supplements containing antioxidant micronutrients and glutamine: a PRCT. Clin Nutr. 2012;31(4):469-75. doi: 10.1016/j.clnu.2012.01.002.

86. Prasad AS. Effects of zinc deficiency on Th1 and Th2 cytokine shifts. J Infect Dis. 2000;182(Supplement_1):S62-S8. doi: 10.1086/315916.

87. Cuevas LE, Koyanagi A. Zinc and infection: a review. Ann Trop Paediatr. 2005;25(3):149-60. doi: 10.1179/146532805X58076.

88. Shankar AH, Prasad AS. Zinc and immune function: the biological basis of altered resistance to infection. Am J Clin Nutr. 1998;68(2):447S-63S. doi: 10.1093/ajcn/68.2.447S.

89.   Dehghan M, Mente A, Zhang X, Swaminathan S, Li W, Mohan V, et al. Associations of fats and carbohydrate intake with cardiovascular disease and mortality in 18 countries from five continents (PURE): a prospective cohort study. Lancet. 2017;390(10107):2050-62. doi: 10.1016/S0140-6736(17)32252-3.

90.   Carr AC, Rosengrave PC, Bayer S, Chambers S, Mehrtens J, Shaw GM. Hypovitaminosis C and vitamin C deficiency in critically ill patients despite recommended enteral and parenteral intakes. Crit Care. 2017;21:1-10. doi: 10.1186/s13054-017-1891-y.

91.   Pullar JM, Carr AC, Vissers M. The roles of vitamin C in skin health. Nutrients. 2017;9(8):866. doi: 10.3390/nu9080866.

92.   Peterkofsky B. Ascorbate requirement for hydroxylation and secretion of procollagen: relationship to inhibition of collagen synthesis in scurvy. Am J Clin Nutr. 1991;54(6):1135S-40S. doi: 10.1093/ajcn/54.6.1135s.

93.   Carr AC, Maggini S. Vitamin C and Immune Function. Nutrients. 2017;9(11). doi: 10.3390/nu9111211. eng.

94.   Elmadfa I, Meyer AL. The Role of the Status of Selected Micronutrients in Shaping the Immune Function. Endocr Metab Immune Disord Drug Targets. 2019;19(8):1100-15. doi: 10.2174/1871530319666190529101816. eng.

95.   Guenin-Macé L, Veyron-Churlet R, Thoulouze MI, Romet-Lemonne G, Hong H, Leadlay PF, et al. Mycolactone activation of Wiskott-Aldrich syndrome proteins underpins Buruli ulcer formation. J Clin Invest. 2013;123(4):1501-12. doi: 10.1172/jci66576.

96.   Ugai K, Koffi DY, Kouadio K, Yao A, Yotsu RR. Nutritional status and wound healing in patients with Mycobacterium ulcerans disease (Buruli ulcer): a pilot study from rural Côte d'Ivoire. Eur J Dermatol. 2022;32(2):227-36. doi: 10.1684/ejd.2022.4225.

97.   Koffi Y, Konan A, Delmont J, Adjet A, Rey J. [Nutritional approach to management of Buruli ulcer in Côte d'Ivoire]. Med Sante Trop. 2019;29(4):409-14. doi: 10.1684/mst.2019.0935. French.