Co-development of gut microbial metabolism and visual neural circuitry over human infancy

[1]Kevin S. Bonham*, [2]Emma T. Margolis*, [1]Guilherme Fahur Bottino, [3]Fadheela Patel, [1]Shelley McCann, [3]Michal R. Zieff, [3]Kirsten A. Donald, [2]Laurel J. Gabard-Durnam†, [1]Vanja Klepac-Ceraj†

*These authors contributed equally

†co-corresponding authors

[1]Wellesley College, Wellesley, MA, USA
[2]Northeastern University, Boston, MA, USA
[3]University of Cape Town, Cape Town, Western Cape, ZAF

ORCID numbers:

Kevin S. Bonham:	0000-0003-3200-7533
Emma T. Margolis:	0000-0002-2036-8078
Guilherme Fahur Bottino:	0000-0003-1953-1576
Fadheela Patel:	0000-0001-5177-7416
Michal Zieff:	0000-0001-9352-9947
Shelley McCann:	0000-0002-9753-7968
Kirsten A. Donald:	0000-0002-0276-9660
Laurel J. Gabard-Durnam	0000-0002-4564-8068
Vanja Klepac-Ceraj:	0000-0001-5387-5706

Running title: Infant gut microbiome associates with visual neurodevelopment

Keywords: visual cortex development, metagenome, infant gut microbiome, feature set enrichment analysis, visual-evoked potential

Abstract

Infancy is a time of elevated neuroplasticity supporting rapid brain and sensory development. The gut microbiome, also undergoing extensive developmental changes in early life, may influence brain development through metabolism of neuroactive compounds. Here, we leverage longitudinal data from 194 infants across the first 18 months of life to show that microbial genes encoding enzymes key to modulating early neuroplasticity are associated with visual cortical neurodevelopment, measured by the visual-evoked potential (VEP). Neuroactive compounds included neurotransmitters GABA and glutamate, the amino acid tryptophan, and short-chain fatty acids involved in myelination, including acetate and butyrate. Microbial gene sets around 4 months of age were strongly associated with the VEP from around 9 to 14 months of age and showed more associations than concurrently measured gene sets, suggesting microbial metabolism in early life may affect subsequent neural plasticity and development.

Introduction

The gut microbiome in early life has potential long-term implications for brain and body health. One important way this influence can occur is through interactions with the central nervous system as a "microbial-gut-brain axis" [1,2]. The metabolic potential of the microorganisms that inhabit the gut vastly exceeds that of human cells alone, with microbial genes outnumbering host genes by a hundredfold [3]. In particular, gut microbes have the ability to metabolize and synthesize many neuroactive compounds [4]. Extensive work in preclinical models suggests that these neuroactive compounds can influence the brain through both direct and indirect routes. For example, major neurotransmitters (e.g., glutamate, γ-aminobutyric acid (GABA), serotonin, and dopamine) are readily synthesized and degraded by intestinal microbes and can enter circulation and pass the blood-brain barrier to influence central nervous function [5–8]. Glutamatergic/GABA-ergic signaling is critical for balancing the brain's excitatory and inhibitory neurotransmission levels, and bi-directional glutamatergic/GABA-ergic signaling between the gut microbiome and brain is implicated in several physical and mental health conditions [9,10]. Similarly, the gut and the microbiome are critical to the regulation of metabolism for the neurotransmitters serotonin and dopamine, particularly through the metabolism of dietary tryptophan [11]. Moreover, short-chain fatty acids (SCFAs) produced by the gut microbiome may impact the brain directly by modulating neurotrophic factors, glial and microglial maturation and myelination, and neuroinflammation [12,13]. Other indirect pathways for gut microbial influence on the brain include vagus nerve stimulation, neuroendocrine modulation and immune system regulation [1].

Rapidly growing literature connects the metabolic potential of the gut microbiome and brain function in humans (reviewed in [6,14,15]), but the overwhelming majority of this research is performed in adults. Importantly, both the gut microbiome and the brain undergo dramatic and rapid development over the first postnatal years [16–19]. However, very little is currently known about how gut-brain influences emerge or change during this critical window for both systems [20–22]. Interrogating this early co-development in humans is, therefore, key to both understanding adaptive gut-brain function and behavior and informing strategies to support it. Specifically, the visual cortex has been shown to be sensitive to gut microbiome modulations in adults [23] and in rodents [24], yet the visual cortex undergoes its most rapid period of plasticity and maturation over infancy when the microbiome changes most significantly [25–27]. Visual cortical functional maturation can be robustly indexed via electroencephalography (EEG) with the visual-evoked potential (VEP) response to visual stimuli from birth. The VEP is especially useful for indexing visual neurodevelopment as its morphology includes amplitude deflections and latencies to those deflections that reflect maturation of function and structure, respectively.

Here, we investigated the longitudinal co-development of microbial metabolic potential - specifically genes encoding enzymes that metabolize neuroactive compounds - and visual neurodevelopment as indexed by the VEP in a longitudinal community sample of 194 infants from Gugulethu in Cape Town, South Africa, recruited as part of a prospective longitudinal study, called "Khula" [28]. Stool samples and EEG were collected at up to 3 visits in the first 18 months of life. Shotgun metagenomic sequencing was used to obtain microbial gene sequences from infant stool samples. To index visual cortical functional development, latencies and peak

amplitudes were extracted from each component of the VEP (i.e., first negative-going deflection,
N1; first positive-going deflection, P1; and second negative-going deflection, N2), producing six
VEP features of interest. We evaluated the concurrent association between microbial genes and
VEP amplitudes and latencies and tested prospective influences of microbial genes from early
visits on VEP changes at later visits to reveal the temporal dynamics of gut-brain co-
development within individuals during this most critical window of plasticity in both systems.

Results

The brain and microbiome develop rapidly in the first months of life

To investigate the co-development of the gut microbiome and visual neurodevelopment, we collected stool and measured the VEP in a longitudinal cohort of 194 children in South Africa in the first 18 months of life (Figure 1A, B, Table 1; visit 1, N = 119, age 3.6 ± 0.7 months, visit 2, N = 144, age 8.7 ± 1.4 months, visit 3, N = 130, age 14.2 ± 1.0 months). As expected for children at this age [16,29,30], the microbial composition was developmentally dependent, with the first principle coordinate axis for both taxonomic profiles (Figure 1C; variance explained = 15.1%; R = -0.50) and functional profiles (Figure 1D; variance explained = 12.9%; R = -0.57) driven strongly by the age of the participant at the time of collection. Similarly, both amplitude and latency VEP features were also strongly correlated with age as expected [16,18]. That is, N1 amplitude became more negative (N1: b=-0.07, p<.05), corrected P1 amplitude became smaller (P1: b=-0.50, p<.05) and corrected N2 amplitude became smaller (N2: b=0.52, p<.05) as infants got older, and all latencies became shorter (N1: b=-0.79, p<.05; P1: b=-1.21, p<.05; N2: b=-4.10, p<.05) (Figure 1E).

Fig. 1: The gut microbiome and VEP both develop over the first 18 months of life.
(A) Study design; participants (N=194) were seen up to 3 times over the first 18 months of life. Stool samples and EEG data were collected, generating microbial functional profiles (stool) and VEP waveforms (EEG) used in subsequent analyses. (B) Longitudinal sampling of study participants; Density plots (top) for stool and EEG collection show the ages represented in each visit. The scatter plot (bottom) shows individual participant visits. Dotted lines connect separate visits for the same participant. When stool and EEG data were collected for the same visit (purple) but not on the same day, dot represents the median age of collection, and vertical bars in blue and red represent stool and EEG collections respectively. (C) Principal coordinate analysis (PCoA) by multidimensional scaling (MDS) on Bray-Curtis dissimilarity of taxonomic profiles; percent variance explained (fraction of positive eigenvalues) by each of the first two axes are indicated on the x and y axes respectively. (D) PCoA of microbial functional profiles (UniRef90s). (E) Average (solid lines) ± standard deviation (dotted lines) for VEP waveforms at each visit.

Table 1: Participant demographics

	Overall (N=194)
Mean (SD) Age at EEG Data Collection (months)	
Visit 1 (N=97)	3.7 (0.85)
Visit 2 (N=129)	8.6 (1.46)
Visit 3 (N=130)	14.1 (1.03)
Mean (SD) Age at Stool Data Collection (months)	
Visit 1 (N=119)	3.6 (0.76)
Visit 2 (N=105)	8.8 (1.43)
Visit 3 (N=91)	14.0 (1.24)
Maternal Place of Birth	
South Africa	191 (98.5%)
In the African Continent (not South Africa)	3 (1.5%)
Primary Spoken Language	
Xhosa Language	187 (96.4%)
Sotho Language	2 (1.0%)
Zulu Language	1 (0.5%)
English Language	2 (1.0%)
Ndebele Language	1 (0.5%)
Afrikaans Language	1 (0.5%)
Maternal Age at Infant Birth (years)	
Mean (SD)	29.2 (5.63)
Median [Min, Max]	29.0 [18.0, 41.0]
Missing	1 (0.5%)
Maternal Educational Attainment[a]	
Completed Grade 6 (Standard 4) to Grade 7 (Standard 5)	4 (2.1%)
Completed Grade 8 (Standard 6) to Grade 11 (Standard 9) i.e., high school without matriculating	78 (40.2%)
Completed Grade 12 (Standard 10) i.e., high school	88 (45.4%)
Part of university/ college/ post-matric education	13 (6.7%)
Completed university/ college/ post-matric education	11 (5.7%)
Maternal Monthly Income[b] (South African Rand/ZAR)	
Less than R1000 per month	97 (50.0%)
R1000 - R5000 per month	76 (39.2%)

Table 1: Participant demographics

	Overall (N=194)
R5000 - R10,000 per month	16 (8.2%)
More than R10,000 per month	0 (0%)
Unknown	5 (2.6%)
Infant Biological Sex	
Female	91 (46.9%)
Male	103 (53.1%)

[a] The South African Educational System was formerly divided into years called standards, similarly to the way the United States Educational System is divided into grades. The equivalent in terms of standards is provided in parentheses next to each mentioned grade. "University/College/Post-Matric Education" refers to tertiary or post-secondary education as defined by the World Bank.
[b] At the time of writing (1/16/24), 1 US Dollar = 18.87 South African Rand (ZAR).

Microbial genes with neuroactive potential are associated with concurrently measured visual development

To test whether microbial metabolic potential was related to early life brain activity, we performed feature set enrichment analysis (FSEA) using groups of potentially neuroactive microbial genes and the concurrently measured VEP amplitude and latency features [4,17]. For each gene set that had at least 5 genes represented in a given comparison group, logistic regression was performed using VEP features as predictors and the presence or absence of each microbial gene in the metagenome as the response to determine concurrent associations (see Methods). Z statistics for in-set genes were compared to all genes using a permutation test to determine significance of the associations as described in [31]. Of the 35 genesets assessed, 19 had sufficient representation to test, and of those, 18 were significantly associated with at least one EEG feature during at least one visit within the 18-month window after correcting for false discovery rate (Benjamini-Hochberg, q < 0.2; Figure 2, Table S2). Microbial genes involved in synthesis or degradation of molecules with neuroactive potential across all categories considered (i.e., neurotransmitters, amino acid metabolism, SCFAs, other) were associated with both VEP amplitudes and latencies at each visit, demonstrating widespread associations between early life gut microbiome and visual cortex neurodevelopment. The number of these associations increased over time (visit 1 had 6 associations, visit 2 had 24, and visit 3 had 37). To differentiate whether this cross-sectional finding indicated the early microbiome (~4 months) was sparsely related to visual cortical development at 4 months only or more broadly at all subsequent time points, longitudinal analyses were performed (results below).

Fig. 2: Concurrent feature set enrichment analysis of microbial neuroactive genes and VEP for three visits.
FSEA results for all genesets where at least one visit had a significant hit (q < 0.2) with at least one VEP latency (A) or amplitude (B). Dots indicate the Z-statistic from logistic regression for each gene in a gene set. Vertical bars indicate the median Z-statistic for the gene set as a whole. Y-axis position for each gene

149 set indicates visit number. Visit 1 for inositol degradation and DOPAC synthesis were not tested, since
150 there were fewer than 5 genes from those genesets present in the sample (See Methods).
151
Specifically, across the gene sets involved in neurotransmitter synthesis and degradation,
glutamate synthesis/degradation and GABA synthesis showed associations with all VEP
features, primarily at the 2nd and 3rd visit (mean ages 8.6 and 14.1 months, respectively; Table
S2). The regulation of this balance between glutamate and GABA (i.e., excitatory/inhibitory
balance) is important for modulating neuroplasticity in the visual cortex during infancy [32,33], and
suggests that microbial metabolism for these compounds is related to the VEP as an index of
this changing neuroplasticity. Gene sets involved in tryptophan metabolism and associated
pathways (i.e., quinolinic acid) were also strongly concurrently related to VEP development.
Tryptophan metabolism is linked to the metabolism of the neurotransmitter serotonin as well as
other neuroactive molecules such as kynurenic acid (an SMDAR antagonist) [11], both of which
are also implicated in early neuroplasticity and neurotransmitter regulation. While quinolinic acid
is part of the kynurenine pathway and is a neurotoxin that can cause neuronal dysfunction, it
may also play a role in glutamate uptake in the brain [34,35]. Specifically, tryptophan metabolism
genes were associated with VEP latencies just after each VEP component showed its greatest
window of developmental change (components emerge sequentially as follows: P1, N1, N2),
that is, of P1 at visit 1, N1 and P1 at visit 2, and N2 at visit 3. This pathway may thus relate to
processes stabilizing the neural circuitry (i.e. downregulating neuroplasticity) underlying each
VEP component, an account consistent with recently observed effects of serotonin within the
visual cortex in rodents [36].
171
Short-chain fatty acids (SCFAs) are important in downregulating neuroinflammation and
promoting myelination within the brain, and several SCFA-metabolizing gene sets were found to
have multiple associations with VEP features. Specifically, acetate synthesis was strongly
associated with almost all VEP features (Table S2). Acetate is a critical component required for
the increased lipid synthesis that happens during postnatal myelination in the brain [37].
Circulating butyrate also increases myelination [38], and butyrate synthesis was associated with
P1 and N2 amplitudes and latencies around the end of the first year of life (visits 2 and 3) when
visual cortex is most actively undergoing myelination. Lastly, propionate synthesis/degradation
was significantly associated with VEP latencies at every visit over the 18-month window (N1 at
visits 1 and 2, and both P1 and N2 at visit 3). Though its relation to myelinating
oligodendrocytes remains unclear, it is known to protect myelinating Schwann cells outside of
the brain from oxidative stress [39]. Acetate, butyrate, and propionate are all also widely regarded
as neuroprotective by promoting healthy microglial development and downregulating
neuroinflammation that interferes with myelination [40]. VEP latency features specifically reflect
myelination (Margolis et al, 2024), and accordingly, these SCFA metabolizing genes showed
almost double the concurrent associations with VEP latencies than amplitudes (11 associations
with latencies, 6 associations with amplitudes).
189
190 Finally, within the remaining gene sets tested, we observed robust associations in particular
191 between menaquinone (Vitamin K2) gene sets and the VEP features over this infancy window.
192 This is reassuring as vitamin K2 specifically is posited to promote healthy vision both outside of

193   the brain through effects on the retina and within the brain where it can protect neural circuits
194   from oxidative stress [41].
195
196   Notably, across significant gene set associations with VEP features, the P1 and N2 component
197   amplitudes and latencies were consistently the most sensitive to these microbial gene sets.
198   Both P1 and N2 components are known to show the most dramatic changes with development
199   during the first year of life [18] and may best reflect underlying visual learning and plasticity at this
200   stage (Table S1).

Microbial metabolic potential predicts future brain development in infancy

Given that differences in metabolic potential of the early life microbiome may exert their effects
over time, we sought to determine whether microbial genes at early time points were associated
with subsequent VEP development. To investigate this, we performed FSEA on stool samples
collected at visit 1 with visit 2 VEP (age at stool collection = 3.6 ± 0.8 months, age at VEP = 8.6
± 1.5 months) or visit 3 VEP (age at stool collection = 3.7 ± 0.7 months, age at VEP = 14.1 ± 1.1
months), as well as visit 2 stool samples with visit 3 VEP (age at stool collection = 8.9 ± 1.5
months, age at VEP = 14.3 ± 1.0 months; Table S3) (Figure 3A). All gene sets except DOPAC
synthesis that had a significant hit with concurrently measured VEP were also significantly
associated with at least one future VEP feature (Figure 3B, Tables 2-5). Notably, the quantity of
those associations increased substantially for all longitudinal comparisons compared to
concurrent comparisons. For example, only 6 visit 1 microbial gene sets were associated with
visit 1 VEP, and each of those was only associated with a single concurrently measured VEP
feature. By contrast, these longitudinal analyses revealed that visit 1 microbial gene sets show a
much richer pattern of associations with future VEP feature development. Specifically, 13 visit 1
gene sets were associated with visit 2 VEP features, and 11 were associated with visit 3 VEP
features, the majority (9/13 for visit 2, 8/11 for visit 3) were associated with at least 2 VEP
features, and nearly half (6/13 for visit 2, 5/11 for visit 3) were associated with more than 2
future VEP features.
220
Fig. 3: Gut microbial genes predict future VEP latencies and amplitudes
(A) Age distributions for stool samples (left) and VEP (right) for each longitudinal comparison (same
individual) tested, V1 stool →V2 VEP, V1 stool →V3 VEP, and V2 stool →V3 VEP. As in Figure 2, (B) and
(C) show FSEA results for all genesets where at least one visit had a significant hit (q < 0.2) with at least
one VEP latency or amplitude respectively. Dots indicate the Z-statistic from logistic regression for each
gene in a gene set. Vertical bars indicate the median Z-statistic for the gene set as a whole. The Y-axis
position for each gene set indicates longitudinal comparison. V1 → V2 and V1 → 3 for inositol
degradation and DOPAC synthesis were not tested, since there were fewer than 5 genes from those
genesets present in the sample (See Methods).
230
231

Table 2: Longitudinal FSEA, visit 1 stool -> visit 2 VEP

Gene set	Feature	Component	Enrichment	Q value
GABA synthesis	amplitude	P1	-0.398729884	0.080789189
		N2	0.366103605	0.112944828
Glutamate synthesis	latency	N1	-0.282915658	0.04452
		P1	0.262506611	0.080789189
	amplitude	P1	0.231653804	0.102313043
Glutamate degradation	latency	P1	0.336966501	0.159815385
		N2	0.325268237	0.180080899
	amplitude	P1	0.3256632	0.176009302
Tryptophan synthesis	latency	N1	-0.221130346	0.027257143
		P1	0.167147007	0.080789189
		N2	0.19001832	0.048457143
	amplitude	N2	-0.163072333	0.102313043
Quinolinic acid synthesis	amplitude	N1	-0.273526842	0.160915663
Quinolinic acid degradation	latency	P1	0.197973559	0.102313043
		N2	0.195068231	0.111888889
Acetate synthesis	latency	N1	-0.219089592	0.029353846
		P1	0.192652389	0.061153846
		N2	0.215744459	0.0318
	amplitude	P1	0.186013239	0.0742
		N2	-0.196033217	0.054927273
Butyrate synthesis	amplitude	N1	-0.31934762	0.169849412
	latency	N1	0.271100718	0.122307692
Isovaleric acid synthesis	amplitude	N1	0.241218831	0.180080899
		P1	-0.240952289	0.160915663
Menaquinone synthesis	latency	N1	-0.139349027	0.159815385
Inositol synthesis	amplitude	N1	-0.429526827	0.180430435
	latency	N1	-0.421912686	0.183435789
p-Cresol synthesis	amplitude	N1	-0.366578927	0.182593548
		P1	0.452086746	0.101457143
		N2	-0.516450705	0.04452
17-beta-Estradiol degradation	latency	N2	0.355044972	0.180080899

Table 3: Longitudinal FSEA, visit 1 stool -> visit 3 VEP

Gene set	Feature	Component	Enrichment	Q value
GABA synthesis	latency	P1	0.434338693	0.04452
	amplitude	P1	0.409900164	0.0742
		N2	-0.506575916	0.02544
Glutamate synthesis	latency	N1	-0.303501108	0.029353846
	amplitude	N1	-0.234405478	0.111888889
Tryptophan synthesis	latency	N1	-0.187530614	0.04452
	amplitude	P1	-0.217612428	0.0159
Quinolinic acid degradation	latency	P1	-0.186280973	0.109392
		N2	-0.262447851	0.183435789
	amplitude	P1	-0.168943012	0.160915663
Acetate synthesis	latency	N1	-0.172876999	0.080789189
Propionate synthesis	amplitude	P1	0.358659876	0.080789189
Propionate degradation	latency	N1	-0.509754961	0.109392
		P1	0.534551664	0.088061538
	amplitude	P1	0.738806805	0
		N2	-0.666912201	0.027257143
Butyrate synthesis	latency	N1	0.337326194	0.122307692
		N2	-0.323276487	0.149852055
	amplitude	P1	0.368868764	0.101457143
Menaquinone synthesis	latency	P1	0.15826763	0.087031579
		N1	-0.141748648	0.139740845
	amplitude	P1	0.148803868	0.113186441
		N2	-0.156702017	0.102313043
Inositol synthesis	amplitude	N2	-0.453803843	0.121045161
	latency	N1	-0.283505043	0.180430435
ClpB	amplitude	P1	-0.328420633	0.112944828
		N2	0.35858587	0.080012903

234
235

Table 4: Longitudinal FSEA, visit 2 stool -> visit 3 VEP

Gene set	Feature	Component	Enrichment	Q value
GABA synthesis	latency	N1	-0.376215897	0.107325
Glutamate synthesis	latency	N1	-0.16762197	0.160915663
		P1	0.297968912	0.0159
	amplitude	P1	0.23424529	0.03816
Glutamate degradation	latency	P1	0.317418013	0.14045
	amplitude	N2	0.282946442	0.187555102
Tryptophan synthesis	amplitude	N1	-0.155489037	0.060834783
		P1	0.186539112	0.029353846
Quinolinic acid synthesis	latency	N1	-0.31863828	0.061056
		P1	0.333137823	0.043725
	amplitude	P1	0.257761121	0.139182609
Acetate synthesis	latency	N1	-0.16732426	0.061056
		P1	0.257469673	0
		N2	-0.137377692	0.152983784
	amplitude	P1	0.146955492	0.121045161
Propionate synthesis	amplitude	P1	0.299415199	0.126236364
Propionate degradation	latency	N1	-0.550397358	0.0742
		N2	0.449253743	0.160915663
	amplitude	P1	0.50881464	0.111888889
		N2	-0.422998998	0.187555102
Butyrate synthesis	latency	N1	-0.331642845	0.080789189
		P1	0.333561852	0.0742
		N2	-0.290812478	0.122307692
	amplitude	N2	0.306668162	0.111888889
Isovaleric acid synthesis	latency	P1	0.231685269	0.155652632
	amplitude	P1	0.246936789	0.112944828
Menaquinone synthesis	latency	N1	0.152879564	0.121045161
	amplitude	N1	-0.164433164	0.107325
		P1	0.201456836	0.029353846
		N2	-0.163403497	0.112167273
Inositol degradation	latency	N1	-0.638125991	0.029353846
		P1	0.532338179	0.101457143
	amplitude	P1	0.641392999	0.029353846
p-Cresol synthesis	latency	P1	0.311879942	0.139740845
		N2	-0.31528647	0.139182609
	amplitude	N1	-0.303976467	0.165814286
S-Adenosylmethionine synthesis	amplitude	N1	-0.222889447	0.155652632
		P1	0.229572468	0.13479403
17-beta-Estradiol degradation	amplitude	P1	0.281861389	0.187555102
ClpB	latency	P1	0.268785756	0.180430435

Longitudinally, the early microbiome (visit 1) was related to VEP features at visit 2 and visit 3

fairly evenly (12/28 visit 1 microbiome associations to visit 3 VEP latencies, 15/30 associations

to visit 3 amplitudes), suggesting early microbiome metabolism in the first 6 months of life is

associated with visual neurodevelopment over the next year. Microbiome metabolism from visit

2 was associated with similar numbers of visit 3 VEP features as visit 1 microbiome (17 visit 3

latency features, 18 visit 3 amplitude features), suggesting continued co-development of these

systems as the microbiome changes over the first postnatal year. Neurotransmitters GABA and

glutamate, tryptophan metabolism (tryptophan and quinolinic acid), SCFAs including acetate,
butyrate, and propionate, as well as menaquinone (Vitamin K2) were again all significantly
associated with multiple VEP features across multiple longitudinal comparisons.
247
Importantly, the nature and identity of the longitudinal associations varied over development for
many gene sets, indicating temporal specificity to these associations. With respect to the
neurotransmitter-related pathways, amongst the associations between GABA synthesis genes
and future VEP features, GABA genes specifically from visit 1 showed the greatest number of
associations with future VEP features (5/6 GABA associations) at visits 2 and 3 equally, and the
majority of these associations were VEP amplitudes (reflecting development of
neurotransmission including excitatory/inhibitory balance). To a lesser extent, glutamate
metabolism genes followed a similar temporal pattern (8/13 glutamate associations involved
visit 1 genes) but did not relate to amplitudes or latencies differentially like GABA. This pattern
of results suggests early (within the first 6 postnatal months) microbiome GABA/glutamate
dynamics and especially GABA are most relevant for changes to visual cortex function over the
following year. This pattern is consistent with GABA's role as a driver of visual neuroplasticity
over early critical periods for this circuitry with long-term consequences for brain function
[32,33,42,43].
262
Tryptophan related pathway genes (responsible for generating serotonin amongst other
products) from visit 1 were also responsible for the majority of associations with future VEP
features (tryptophan: 6/8 associations; quinolinic acid: 6/9 associations). In contrast to GABA
but similar to glutamate, tryptophan related gene associations were largely shorter-term
associations with VEP features at the visit immediately following gene set measurement
(approximately 5 months later; 12/17 associations), indicating dynamic co-development over the
first 18 months of life. Across neurotransmitter-related gene set associations (GABA, glutamate,
tryptophan/serotonin), there was thus a clear pattern whereby early (~4 months old) microbiome
gene sets showed the largest number of associations with subsequent VEP feature
development.
273
SCFAs showed a different developmental pattern of associations with future VEP features.
Specifically, propionate and butyrate metabolism genes from both visit 1 and visit 2 showed
associations with future VEP features, but here the effects were almost entirely observed for
VEP features at visit 3 (10/10 propionate and 7/8 butyrate associations). Indeed, acetate and
butyrate metabolism genes were doubly associated with future VEP latencies compared to
amplitude features. This pattern of results is consistent with these SCFA roles in myelination
occurring over the second half of the developmental window studied.
281
282 Finally, menaquinone (Vitamin K2) metabolism genes followed a similar pattern to the SCFAs in
283 that genes from visit 1 and visit 2 were largely associated with future VEP features at visit 3 (8/9
284 menaquinone associations), indicating persistent associations of this early microbiome gene set
285 with individual differences in VEP features in the second year of life.
286

It is possible that extrinsic factors related to development mutually influence both the gut
microbiome and neural development, though we additionally tested whether VEP features were
associated with microbial metabolism at a future visit, and found substantially fewer
associations (29 total associations, compared with 95 when analyzing early stool samples with
future VEP). While this does not prove a causal relationship, it is consistent with the hypothesis
that microbial metabolism influences brain development.

Discussion

The past decade has seen a remarkable growth in our understanding of the relationships
between the gut microbiome and the brain. However, a great deal of that investigation has
focused on neuropsychiatric disorders or adult populations, limiting their potential to explain how
these associations emerge through development. Here we address this key open question by
leveraging a rich longitudinal dataset over the first year of life, the time of greatest
developmental change for both microbiome and brain as foundational sensory
neurodevelopment unfolds.
301
Specifically, we have shown a robust, prospective relationship between microbial genes
involved in the metabolism of neuroactive compounds and the development of visual cortical
function as measured by the Visual-Evoked Potential (VEP) electrophysiological response. We
found that the microbial metabolism is more strongly associated with future measures of VEP
than those collected concurrently. While not dispositive, this would be the predicted outcome if
microbial genes are causally influencing brain development. Nor did we see the same rich set of
associations in the converse analyses examining whether VEP related to future microbiome
properties. Microbial metabolism within the first 6 months shows the most associations with
subsequent visual cortical function, suggesting the early postnatal microbiome may play a
particularly important role in the codevelopment of these systems. This interpretation is also
supported by prior research showing microbiome associations with behavioral readouts of
neurocognition are stronger prospectively than concurrently [44]. Moreover, specific associations
between gene sets and VEP features showed temporal specificity within the 18-month
developmental window assessed, suggesting that the impact of early microbial metabolism on
the brain is developmentally dependent.
317
Notably, the gene sets most associated with visual functional neurodevelopment over infancy
are for the metabolism of molecules with known links to developmental neuroplasticity [32,33,45].
Specifically, we observed gene sets related to glutamate and GABA that are central to E/I
balance in the cortex. Developmental changes in E/I balance modulate the degree of
neuroplasticity in the mammalian cortex, including regulating the start and progression of critical
period neuroplasticity mechanisms in visual cortex [33]. Gut production of GABA may influence
cortical GABA levels via active transport from bloodstream to brain [46–48]. Recent evidence
suggests gut-derived glutamate may also influence brain levels and function [7–9] and can operate
via indirect mechanisms (either transformation into GABA or via regulating glutamate levels in
the bloodstream that impact glutamate transfer from brain to bloodstream). Moreover,
tryptophan related pathway genes were identified here that are responsible for generating

serotonin, a key neuroplasticity regulator with potent effects noted for visual cortex plasticity [49,50].
Nearly all of the body's serotonin is produced in the gut by enterochromaffin cells, and this
biosynthesis is regulated by microbes [51,52], making this pathway an especially promising
candidate intervention target for further research in development. Finally, the short-chain fatty
acids we observed (i.e., propionate, acetate, and butyrate) can pass the blood-brain barrier to
directly influence myelination-related processes. Myelination is important for down-regulating
plasticity in neural circuitry over development, stabilizing and protecting circuits that have been
shaped by early experience (Takesian and Hensch 2013). Taken together, the pattern of results
suggests early postnatal microbiome-derived metabolites relate to key neuroplasticity regulation
processes within the cortex.
339
Our study is a substantial advance over prior work on the microbial-gut-brain axis in early life
due to the sequencing method, the large number of participants, and the longitudinal study
design. The use of shotgun metagenomic sequencing enables direct interrogation of microbial
metabolic potential. Prior research primarily used amplicon (16S rRNA gene) sequencing, which
enables lower-resolution taxonomic identification and is restricted to inferring metabolic potential
based on taxonomy. Moreover, while several studies in infancy have inferred gut-brain
associations by linking microbiome measures to subsequent neurodevelopmental measures
using behavioral assessments (e.g., Bayley Scales of Infant Development), noting associations
with visually-mediated cognition [44], this study assessed the development of the visual cortex
directly by measuring the Visual-Evoked Potential. The VEP is advantageous because features
reflect largely neurotransmission (amplitudes) or structural (e.g., myelination, latencies) changes
over this developmental window, facilitating some specificity in interpretation of the associations
observed. Moreover, the VEP can be indexed with fidelity from birth, providing a continuous
measure of visual cortical function across the study age-range. Additionally, this study involved
a large number of participants (194) contributing dense longitudinal data, with three time points,
all taken in the first 18 months of an infant's life. While prior work focused on single time point
measures of microbiome and neurodevelopment [53], longitudinal associations allowed us to
investigate the changing relation between gut microbial metabolism and the development of
visual neurocircuitry over time.
359
One limitation of this study is the fact that we are only able to observe the genomic composition
of the microbiome, rather than the concentration of metabolites themselves. This prevents us
from determining the concentration of these molecules in the gastrointestinal tract, blood, and
brain, as the abundance of these genes does not provide information about their activity, their
interactions with other metabolic pathways (including those of the host), or absorption by colonic
epithelial cells. Moreover, the relationship between gene abundance and molecule
concentration may be counterintuitive, since the relationship between degradation and synthesis
of metabolites occurs both at the individual organism level and at the community level. For
example, genes for breaking down a molecule may be prevalent if that molecule is at high
concentrations, or the molecules may be rapidly degraded by other members of the community
the moment they are produced. Furthermore, it may be that the relationship between metabolite
and brain development remains the same over time, while the relationship between molecules
and microbial selection changes at different stages of life. Addressing these limitations in

humans is challenging, even if looking at stool metabolites, since overall exposure throughout
the gastrointestinal tract is not necessarily reflected in the final concentration of those molecules
in stool. Therefore, metabolites from blood plasma could provide more accurate systemic
concentrations of molecules, but challenges remain on how to interpret them in humans [54,55].
377
Given that the VEP is evolutionarily conserved in mammals and can be accurately measured
during development, the hypotheses generated in humans in this study are readily testable
mechanistically using *in vivo* models in future research. Germ-free or defined-microbiome
animals [56,57] may be supplemented with specific molecules such as SCFAs, or colonized with
microbial species lacking or providing specific metabolic pathways and molecule concentrations
in tissues from the gut to the brain can be directly assessed. Uncovering relationships between
microbial metabolism and specific molecules may also generate hypotheses that can be
confirmed in human data. This study, therefore, provides a foundation for deep investigation of
the relationships between the human gut microbiome and brain development.
387

Materials and methods

Cohort, ethics, etc

Participants and Study Design
Infants were recruited from local community clinics in Gugulethu, an informal settlement in Cape
Town, South Africa, as part of a prospective longitudinal study (most enrollments happened
prenatally with 16% of infants enrolled shortly after birth [28]). The first language for the majority of
residents in this area is Xhosa. Study procedures were offered in English or Xhosa depending
on the language preference of the mother. This study was approved by the relevant university
Health Research Ethics Committees (University of Cape Town study number: 666/2021).
Informed consent was collected from mothers on behalf of themself and their infants.
Demographic information, including maternal place of birth, primary spoken language, maternal
age at enrollment, maternal educational attainment, and maternal income, were collected at
enrollment (see Table 1).
401
402 Families were invited to participate in three in-lab study visits over their infant's first year and a
403 half of life. At the first in-lab study visit (hereafter visit 1), occurring when infants were between
404 approximately 2 months and 6 months of age, the following data were collected: the infants' age
405 (in months), sex, infant electroencephalography (EEG), and infant stool samples.
406
At the second study visit (hereafter visit 2), occurring when infants were between approximately
6 months and 12 months of age (age in months: M=8.60, SD=1.48, range=5.41-12.00) and at
the third study visit (hereafter visit 3), occurring when infants were between approximately 12
months and 17 months of age (age in months: M=14.10, SD=1.04, range=12.10-17.00), infant
EEG and stool samples were collected again. At visits in which infants were unable to complete
both EEG and stool samples on the same day, EEG and stool samples were collected on
different days. For concurrent time point analyses, infants with EEG and stool collected more
than two months apart were excluded. Not all infants had EEG and microbiome data collected at
all three time points or contributed usable data at all three-time points.
416

417   All enrolled infants received a comprehensive medical exam at each visit, which included
418   assessments of eye-related conditions. Several infants (n=3) were identified as having eye-
419   related anomalies during the medical exam, and they were excluded from any further analyses.

EEG Processing

EEG Data Acquisition

Electroencephalography (EEG) data were acquired from infants while they were seated in their caregiver's lap in a dimly-lit, quiet room using a 128-channel high density HydroCel Geodesic Sensor Net (EGI, Eugene, OR), amplified with a NetAmps 400 high-input amplifier, and recorded via an Electrical Geodesics, Inc. (EGI, Eugene, OR) system with a 1000 Hz sampling rate. EEG data were online referenced to the vertex (channel Cz) through the EGI Netstation software. Impedances were kept below 100KΩ in accordance with the impedance capabilities of the high-impedance amplifiers. Geodesic Sensor Nets with modified tall pedestals designed for improving inclusion of infants with thick/curly/tall hair were used as needed across participants (Mlandu et al., 2024). Shea moisture leave-in castor oil conditioner was applied to hair across the scalp prior to net placement to improve both impedances and participant comfort (Mlandu et al., 2024). This leave-in conditioner contains insulating ingredients so there is no risk of electrical bridging and has not been found to disrupt the EEG signal during testing (unpublished data). Conditioning hair in this way allows for nets to lay closer to the scalp for curly/coily hair types and makes for far more comfortable net removal at the end of testing.

The Visual-Evoked Potential (VEP) task was presented using Eprime 3.0 software (Psychology Software Tools, Pittsburgh, PA) on a Lenovo desktop computer with an external monitor 19.5 inches on the diagonal facing the infant (with monitor approximately 65 cm away from the infant). A standard phase-reversal VEP was induced with a black and white checkerboard (1cm x 1 cm squares within the board) stimulus that alternated presentation (black squares became white, white squares became black) every 500 milliseconds for a total of 100 trials. Participant looking was monitored by video and by an assistant throughout data collection. If the participant looked away during the VEP task, the task was rerun.

EEG Data Pre-Processing

VEP data were exported from native Netstation .mff format to .raw format and then pre-processed using the HAPPE+ER pipeline within the HAPPE v3.3 software, an automated open-source EEG processing software validated for infant data [58]. A subset of the 128 channels were selected for pre-processing that excluded the rim electrodes as these are typically artifact-laden (channels excluded from pre-processing included in Table S4).The HAPPE pre-processing pipeline was run with user-selected specifications outlined in Table S4.

Pre-processed VEP data were considered usable and moved forward to VEP extraction if HAPPE pre-processing ran successfully, at least 15 trials were retained following bad trial rejection, and at least one good channel was kept within the visual ROI. Note that channels marked bad during pre-processing had their data interpolated as part of standard preprocessing pipelines for ERPs [58]. Interpolated channels were included in analyses here as is typically done in developmental samples, and given the low overall rates of interpolation present (e.g., an average of between 4 to 5 of 5 possible good channels in the region of interest were retained at each visit time point).

Visual-Evoked Potentials (VEPs)

VEP waveforms were extracted and quantified using the HAPPE+ER v3.3 GenerateERPs script [58]. Electrodes in the occipital region were selected as a region of interest (i.e., E70, E71, E75,

E76, E83). The VEP waveform has three main components to be quantified: a negative N1
peak, a positive P1 peak, and a negative N2 peak. Due to normative maturation of the
waveforms as infants age, one set of user-specified windows for calculating component features
was used for visit 1 and 2 and another was used for visit 3. For visits 1 and 2, the window for
calculating features for the N1 component was 40-100 ms, 75-175 ms for the P1 component,
and 100-325 ms for the N2 component. For visit 3, the window for calculating features for the
N1 component was 35-80 ms, 75-130 ms for the P1 component, and 100-275 ms for the N2
component. HAPPE+ER parameters used in extracting the ERPs are summarized in Table S5.
474
To correct for the potential influence of earlier components on later components, corrected
amplitudes and latencies were calculated and used in all analyses. Specifically, the P1
amplitude was corrected for the N1 amplitude (corrected P1 amplitude = P1 - N1 amplitude); the
P1 latency was corrected for the N1 latency (corrected P1 latency = P1 - N1 latency); the N2
amplitude was corrected for the P1 amplitude (corrected N2 amplitude = N2 - P1 amplitude),
and the N2 latency was corrected for the P1 latency (corrected N2 latency = N2 - P1 latency).
481
All VEPs were visually inspected to ensure that the automatically extracted values were correct
and were adjusted if observable peaks occurred outside the automated window bounds.
Participants were considered to have failed this visual inspection and were subsequently
removed from the data set if their VEP did not produce three discernible peaks. VEP waveforms
of included participants by time point are included in Figure 1E. 97 infants provided usable VEP
data at visit 1, 130 infants provided usable VEP data at visit 2, and 131 infants provided usable
VEP data at visit 3. For included participants, EEG data quality metrics are summarized in Table
S6. T-tests for data quality metrics (i.e., number of trials collected, number of trials retained,
number of channels retained in the ROI, Pearson's r for data pre- vs. post-wavelet thresholding
at 5, 8, 12, and 20 Hz) were run between each visit combination (i.e., visit 1 vs. visit 2, visit 1 vs.
visit 3, visit 2 vs. visit 3). For visits that differed in data quality, follow-up post hoc correlations
were run for the data quality measure with each VEP feature at each visit in the T-test. In no
case did the data quality metric relate to VEP features at multiple visits, making it highly
unlikely the data quality difference contributed to results.

Biospecimens and sequencing

Sample Collection
497
498 Stool samples (n=315) were collected in the clinic by the research assistant directly from the
499 diaper, transferred to Zymo DNA/RNA Shield™ Fecal collection Tubes (#R1101, Zymo
500 Research Corp., Irvine, USA) and immediately frozen at -80 ˚C. Stool samples were not
501 collected if the participant had taken antibiotics within the two weeks prior to sampling.
502

DNA Extraction
503
DNA extraction was performed at Medical Microbiology, University of Cape Town, South Africa,
from stool samples collected in DNA/RNA Shield™ Fecal collection tube using the Zymo
Research Fecal DNA MiniPrep kit (# D4300, Zymo Research Corp., Irvine, USA) following
manufacturer's protocol. To assess the extraction process's quality, ZymoBIOMICS® Microbial
Community Standards (#D6300 and #D6310, Zymo Research Corp., Irvine, USA) were
incorporated and subjected to the identical process as the stool samples. The DNA yield and
purity were determined using the NanoDrop® ND -1000 (Nanodrop Technologies Inc.
Wilmington, USA).

Sequencing

Shotgun metagenomic sequencing was performed on all samples at the Integrated Microbiome
Research Resource (IMR, Dalhousie University, NS, Canada). A pooled library (max 96
samples per run) was prepared using the Illumina Nextera Flex Kit for MiSeq and NextSeq from
1 ng of each sample. Samples were then pooled onto a plate and sequenced on the Illumina
NextSeq 2000 platform using 150+150 bp paired-end P3 cells, generating 24M million raw
reads and 3.6 Gb of sequence per sample [59].

Statistics / computational analysis

Age-Related Changes in VEP Features

To determine age-related changes in VEP features, six linear mixed models with each VEP
feature as the outcome (i.e., N1 amplitude/latency, P1 amplitude/latency, N2 amplitude/latency)
were run using the *lme4* package [60] in R with age in months as the predictor of interest and
number of retained trials as a covariate.

Metagenome processing

Raw metagenomic sequence reads ($2.5 \times 10^7 \pm 1.4 \times 10^7$ reads/sample) were processed using
tools from the bioBakery as previously described [17,61]. Briefly, KneadData v0.10.0 was used with
default parameters to trim low-quality reads and remove human sequences (using reference
database hg37). Next, MetaPhlAn v3.1.0 (using database mpa_v31_CHOCOPhlAn_201901)
was used with default parameters to map microbial marker genes to generate taxonomic
profiles. Taxonomic profiles and raw reads were passed to HUMAnN v3.7 to generate stratified
functional profiles.

Microbial community analysis

Principal coordinates analysis was performed in the julia programming language [62] using the
Microbiome.jl package [63]. Bray-Curtis dissimilarity (Distances.jl) was calculated across all pairs
of samples, filtering for species-level classification. Classical multidimensional scaling was
performed on the dissimilarity matrix (MultivariateStats.jl), and axes with negative eigenvalues
were discarded.

Feature Set Enrichment Analysis (FSEA)

Potentially neuroactive genesets were extracted from Supplementary Dataset 1 from [4]. Gut-
brain modules provide Kegg Orthologue IDs (KOs) [64,65], which were mapped to UniRef90 IDs
using the utility mapping file provided with HUMAnN v3.1 [61]. For each stool / VEP pair, logistic
regression (LR) was performed, linking the presence or absence of that UniRef in a sample with
each VEP feature (N1, P1, and N2 latencies and amplitudes), controlling for the age at which
the stool sample was collected, the number of retained VEP trials, and the difference in age
between the stool collection and VEP measurement. For concurrently collected stool and VEP
comparisons (Figure 2), participants whose stool collection and VEP measurements were more
than two months apart were excluded.

(1) UniRef ~ vep + age_months + n_trials + age_diff

FSEA was performed on each geneset that had at least five members in each comparison
group according to the procedure set out in [31]. Briefly, enrichment scores (ES) are calculated
based on the rank order of z-statistics from the LR for each UniRef. A permutation test was then
performed where the ES for 5000 random samples of ranks of the same length as the gene set
are calculated, and the pseudo-p value is the fraction of permutations where the permutation ES
has a greater absolute value than the true ES.

Benjamini-Hochberg FDR correction was performed separately on all concurrently tested
geneset / VEP feature combinations, and all longitudinal geneset / VEP feature combinations.
Corrected p-values (q-values) less than 0.2 were considered statistically significant.

For longitudinal comparisons, all participants that had a stool sample collected at one visit and
a VEP assessment at a subsequent visit were included (visit 1 stool → visit 2 VEP, N = 84; v1 →
v2, N = 76; v2 → v3, N = 69). A total of 95 geneset / VEP features were significant when using an
FDR-corrected p-value cutoff of q < 0.2. To ensure the robustness of these findings, we
randomly permuted participant IDs between the stool and VEP assessments and repeated the
analysis. Over 10 random permutations, a mean of 19.5 significant associations were identified,
suggesting that FDR correction is correctly calibrating the false-positive rate.

Data and code availability

Code for initial processing of data and for analyses performed in this manuscript are available
on github and archived on Zenodo [66]. Input data will be archived on Dryad and downloadable via
included scripts in analysis code upon publication.

References

1. Rhee, S. H., Pothoulakis, C. & Mayer, E. A. Principles and clinical implications of the brain-
gut-enteric microbiota axis. *Nat. Rev. Gastroenterol. Hepatol.* **6**, 306–314 (2009).
2. Collins, S. M., Surette, M. & Bercik, P. The interplay between the intestinal microbiota and
the brain. *Nat. Rev. Microbiol.* **10**, 735–742 (2012).
3. Gilbert, J. A. *et al.* Current understanding of the human microbiome. *Nat. Med.* **24**, 392–400
(2018).
4. Valles-Colomer, M. *et al.* The neuroactive potential of the human gut microbiota in quality of
life and depression. *Nat. Microbiol.* **4**, 623–632 (2019).
5. Janik, R. *et al.* Magnetic resonance spectroscopy reveals oral Lactobacillus promotion of
increases in brain GABA, N-acetyl aspartate and glutamate. *NeuroImage* **125**, 988–995
(2016).
6. Ahmed, H. *et al.* Microbiota-derived metabolites as drivers of gut–brain communication. *Gut
Microbes* **14**, 2102878 (2022).
7. Matsumoto, M. *et al.* Cerebral low-molecular metabolites influenced by intestinal microbiota:

a pilot study. *Front. Syst. Neurosci.* **7**, 9 (2013).

8. Kawase, T. *et al.* Gut microbiota of mice putatively modifies amino acid metabolism in the
host brain. *Br. J. Nutr.* **117**, 775–783 (2017).

9. Filpa, V. *et al.* Role of glutamatergic neurotransmission in the enteric nervous system and
brain-gut axis in health and disease. *Neuropharmacology* **111**, 14–33 (2016).

10. Baj, A. *et al.* Glutamatergic Signaling Along The Microbiota-Gut-Brain Axis. *Int. J. Mol. Sci.*
20, 1482 (2019).

11. Agus, A., Planchais, J. & Sokol, H. Gut Microbiota Regulation of Tryptophan Metabolism in
Health and Disease. *Cell Host Microbe* **23**, 716–724 (2018).

12. Dalile, B., Van Oudenhove, L., Vervliet, B. & Verbeke, K. The role of short-chain fatty acids
in microbiota-gut-brain communication. *Nat. Rev. Gastroenterol. Hepatol.* **16**, 461–478
(2019).

13. Erny, D. *et al.* Microbiota-derived acetate enables the metabolic fitness of the brain innate
immune system during health and disease. *Cell Metab.* **33**, 2260-2276.e7 (2021).

14. Parker, A., Fonseca, S. & Carding, S. R. Gut microbes and metabolites as modulators of
blood-brain barrier integrity and brain health. *Gut Microbes* **11**, 135–157 (2019).

15. Aburto, M. R. & Cryan, J. F. Gastrointestinal and brain barriers: unlocking gates of
communication across the microbiota–gut–brain axis. *Nat. Rev. Gastroenterol. Hepatol.* **21**,
222–247 (2024).

16. Yassour, M. *et al.* Natural history of the infant gut microbiome and impact of antibiotic
treatment on bacterial strain diversity and stability. *Sci. Transl. Med.* **8**, 343ra81 (2016).

17. Bonham, K. *et al.* Gut-resident microorganisms and their genes are associated with
cognition and neuroanatomy in children. *Sci. Adv.* **9**, eadi0497 (2023).

18. Lippé, S., Roy, M.-S., Perchet, C. & Lassonde, M. Electrophysiological Markers of
Visuocortical Development. *Cereb. Cortex* **17**, 100–107 (2007).

19. Dean, D. C. *et al.* Modeling healthy male white matter and myelin development: 3 through
60 months of age. *NeuroImage* **84**, 742–752 (2014).

20. Ahrens, A. P. *et al.* Infant microbes and metabolites point to childhood neurodevelopmental
disorders. *Cell* **187**, 1853-1873.e15 (2024).

21. Meyer, K. *et al.* Association of the Gut Microbiota With Cognitive Function in Midlife. *JAMA*
Netw. Open **5**, e2143941 (2022).

22. Callaghan, B. L. *et al.* Mind and gut: Associations between mood and gastrointestinal
distress in children exposed to adversity. *Dev. Psychopathol.* **32**, 309–328 (2020).

23. Canipe, L. G., Sioda, M. & Cheatham, C. L. Diversity of the gut-microbiome related to
cognitive behavioral outcomes in healthy older adults. *Arch. Gerontol. Geriatr.* **96**, 104464
(2021).

24. Lupori, L. *et al.* The gut microbiota of environmentally enriched mice regulates visual cortical
plasticity. *Cell Rep.* **38**, 110212 (2022).

25. Deen, B. *et al.* Organization of high-level visual cortex in human infants. *Nat. Commun.* **8**,
13995 (2017).

26. Ellis, C. T. *et al.* Retinotopic organization of visual cortex in human infants. *Neuron* **109**,
2616-2626.e6 (2021).

27. Kiorpes, L. Visual development in primates: Neural mechanisms and critical periods. *Dev.*
Neurobiol. **75**, 1080–1090 (2015).

28. Zieff, M. R. *et al.* Characterizing developing executive functions in the first 1000 days in
South Africa and Malawi: The Khula Study. *Wellcome Open Res.* **9**, 157 (2024).
29. Koenig, J. E. *et al.* Succession of microbial consortia in the developing infant gut
microbiome. *Proc. Natl. Acad. Sci. U. S. A.* **108 Suppl 1**, 4578–4585 (2011).
30. Bäckhed, F. *et al.* Dynamics and stabilization of the human gut microbiome during the first
year of life. *Cell Host Microbe* **17**, 690–703 (2015).
31. Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for
interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci.* **102**, 15545–15550
(2005).
32. Fagiolini, M. & Hensch, T. K. Inhibitory threshold for critical-period activation in primary
visual cortex. *Nature* **404**, 183–186 (2000).
33. Hensch, T. K. *et al.* Local GABA Circuit Control of Experience-Dependent Plasticity in
Developing Visual Cortex. *Science* **282**, 1504–1508 (1998).
34. Lugo-Huitrón, R. *et al.* Quinolinic Acid: An Endogenous Neurotoxin with Multiple Targets.
Oxid. Med. Cell. Longev. **2013**, 104024 (2013).
35. Tavares, R. G. *et al.* Quinolinic acid inhibits glutamate uptake into synaptic vesicles from rat
brain. *NeuroReport* **11**, 249 (2000).
36. Hong, S. Z. *et al.* Norepinephrine potentiates and serotonin depresses visual cortical
responses by transforming eligibility traces. *Nat. Commun.* **13**, 3202 (2022).
37. Madhavarao, C. N. *et al.* Defective N-acetylaspartate catabolism reduces brain acetate
levels and myelin lipid synthesis in Canavan's disease. *Proc. Natl. Acad. Sci.* **102**, 5221–
5226 (2005).
38. Chen, T., Noto, D., Hoshino, Y., Mizuno, M. & Miyake, S. Butyrate suppresses
demyelination and enhances remyelination. *J. Neuroinflammation* **16**, 165 (2019).
39. Grüter, T. *et al.* Propionate exerts neuroprotective and neuroregenerative effects in the
peripheral nervous system. *Proc. Natl. Acad. Sci. U. S. A.* **120**, (2023).
40. Caetano-Silva, M. E. *et al.* Inhibition of inflammatory microglia by dietary fiber and short-
chain fatty acids. *Sci. Rep.* **13**, 2819 (2023).
41. Li, J. *et al.* Novel Role of Vitamin K in Preventing Oxidative Injury to Developing
Oligodendrocytes and Neurons. *J. Neurosci.* **23**, 5816–5826 (2003).
42. Gabard-Durnam, L. & McLaughlin, K. A. Sensitive periods in human development: charting
a course for the future. *Curr. Opin. Behav. Sci.* **36**, 120–128 (2020).
43. Margolis, E. T. & Gabard-Durnam, L. J. Prenatal influences on postnatal neuroplasticity:
Integrating DOHaD and sensitive/critical period frameworks to understand biological
embedding in early development. *Infancy* **n/a**,.
44. Carlson, A. L. *et al.* Infant Gut Microbiome Associated With Cognitive Development. *Biol.
Psychiatry* **83**, 148–159 (2018).
45. Takesian, A. E., Bogart, L. J., Lichtman, J. W. & Hensch, T. K. Inhibitory circuit gating of
auditory critical-period plasticity. *Nat. Neurosci.* **21**, 218–227 (2018).
46. Takanaga, H., Ohtsuki, S., Hosoya Ki, null & Terasaki, T. GAT2/BGT-1 as a system
responsible for the transport of gamma-aminobutyric acid at the mouse blood-brain barrier.
J. Cereb. Blood Flow Metab. Off. J. Int. Soc. Cereb. Blood Flow Metab. **21**, 1232–1239
(2001).
47. Al-Sarraf, H. Transport of 14C-γ-aminobutyric acid into brain, cerebrospinal fluid and choroid

plexus in neonatal and adult rats. *Dev. Brain Res.* **139**, 121–129 (2002).

48. Shyamaladevi, N., Jayakumar, A. R., Sujatha, R., Paul, V. & Subramanian, E. H. Evidence

that nitric oxide production increases γ-amino butyric acid permeability of blood-brain

barrier. *Brain Res. Bull.* **57**, 231–236 (2002).

49. Vetencourt, J. F. M., Tiraboschi, E., Spolidoro, M., Castrén, E. & Maffei, L. Serotonin

triggers a transient epigenetic mechanism that reinstates adult visual cortex plasticity in rats.

Eur. J. Neurosci. **33**, 49–57 (2011).

50. Gu, Q. & Singer, W. Involvement of Serotonin in Developmental Plasticity of Kitten Visual

Cortex. *Eur. J. Neurosci.* **7**, 1146–1153 (1995).

51. Yano, J. M. *et al.* Indigenous bacteria from the gut microbiota regulate host serotonin

biosynthesis. *Cell* **161**, 264–276 (2015).

52. Jameson, K. G. & Hsiao, E. Y. Linking the Gut Microbiota to a Brain Neurotransmitter.

Trends Neurosci. **41**, 413–414 (2018).

53. Gao, W. *et al.* Gut microbiome and brain functional connectivity in infants-a preliminary

study focusing on the amygdala. *Psychopharmacology (Berl.)* **236**, 1641–1651 (2019).

54. Deng, K. *et al.* Comparison of fecal and blood metabolome reveals inconsistent associations

of the gut microbiota with cardiometabolic diseases. *Nat. Commun.* **14**, 571 (2023).

55. Dekkers, K. F. *et al.* An online atlas of human plasma metabolite signatures of gut

microbiome composition. *Nat. Commun.* **13**, 5370 (2022).

56. Wymore Brand, M. *et al.* The Altered Schaedler Flora: Continued Applications of a Defined

Murine Microbial Community. *ILAR J.* **56**, 169–178 (2015).

57. Kennedy, E. A., King, K. Y. & Baldridge, M. T. Mouse Microbiota Models: Comparing Germ-

Free Mice and Antibiotics Treatment as Tools for Modifying Gut Bacteria. *Front. Physiol.* **9**,

(2018).

58. Monachino, A. D., Lopez, K. L., Pierce, L. J. & Gabard-Durnam, L. J. The HAPPE plus

Event-Related (HAPPE+ER) software: A standardized preprocessing pipeline for event-

related potential analyses. *Model. Neural Dev.* **57**, 101140 (2022).

59. Comeau, A. M. & Filloramo, G. V. Preparing multiplexed WGS/MetaG libraries with the

Illumina DNA Prep kit for the Illumina NextSeq or MiSeq. (2023).

60. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using

lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

61. Beghini, F. *et al.* Integrating taxonomic, functional, and strain-level profiling of diverse

microbial communities with bioBakery 3. *Elife* **10**, (2021).

62. Bezanson, J., Karpinski, S., Shah, V. B. & Edelman, A. Julia: A fast dynamic language for

technical computing. (2012).

63. Bonham, K., Kayisire, A., Luo, A. & Klepac-Ceraj, V. Microbiome.jl and BiobakeryUtils.jl -

Julia packages for working with microbial community data. *J. Open Source Softw.* **6**, 3876

(2021).

64. Suzek, B. E., Huang, H., McGarvey, P., Mazumder, R. & Wu, C. H. UniRef: comprehensive

and non-redundant UniProt reference clusters. *Bioinformatics* **23**, 1282–1288 (2007).

65. Kanehisa, M., Goto, S., Kawashima, S., Okuno, Y. & Hattori, M. The KEGG resource for

deciphering the genome. *Nucleic Acids Res.* **32**, D277–D280 (2004).

66. Bonham, K. Klepac-Ceraj-Lab/EEGMicrobiome: Initial submission. Zenodo

https://doi.org/10.5281/zenodo.13625448 (2024).