Equity, cost and disability adjusted life years (DALYs) of tuberculosis treatment supported by digital adherence technologies and differentiated care in Ethiopia: a trial-based distributional cost-effectiveness analysis

Nicola Foster[1,2], Amare W Tadesse[1], Mahilet Belachew[3], Mamush Sahlie[3], Christopher Finn McQuaid[1], Lara Goscé[1], Ahmed Bedru[3], Tofik Abdurhman[3], Demekech. G Umeta[3], Amanuel Shiferaw[3], Gedion T Weldemichael[3], Taye Letta Janfa[4], Norma Madden[5], Salome Charalambous[6], Job van Rest[5], Kristian van Kalmthout[5], Degu Jerene[5], Katherine L Fielding[1,7]

[1]Tuberculosis Centre, Department of Infectious Disease Epidemiology, London School of Hygiene & Tropical Medicine, London, United Kingdom
[2]UCL Respiratory, University College London, London, United Kingdom
[3]KNCV Tuberculosis Plus, Addis Ababa, Ethiopia
[4]National Tuberculosis Control Program, Ethiopian Ministry of Health, Addis Ababa, Ethiopia
[5]KNCV Tuberculosis Plus, The Hague, the Netherlands
[6]Aurum Institute, Johannesburg, South Africa
[7]School of Public Health, University of Witwatersrand, Johannesburg, South Africa

Corresponding author:
Nicola Foster,
London School of Hygiene & Tropical Medicine,
Keppel St WC1E 7HT
London, UK
nicola.foster@lshtm.ac.uk
ORCID: 0000-0003-4630-6243

Abstract

Summary

Background

Evidence of the cost-effectiveness of digital adherence technologies (DATs) for supporting

tuberculosis treatment has been inconclusive and primarily omitted patient-incurred costs. We

aimed to assess the societal costs, equity impact and cost-effectiveness of DATs and differentiated

care compared to routine care in Ethiopia.

Methods

We conducted a distributional cost-effectiveness analysis using data from the cluster randomised

trial that evaluated the implementation of labels and pillbox followed by differentiated care to

support tuberculosis treatment adherence in 78 health facilities in Ethiopia. We estimated the costs,

cost per disability-adjusted life year (DALYs) averted and equity impact of the implementation of the

DATs interventions. Costs and DALYs were estimated at a participant level based on patient events

collected during the trial and the trial endpoints for intention-to-treat population. Uncertainty in

cost-effectiveness estimates were assessed by plotting cost-effectiveness acceptability frontiers. The

trial is registered with Pan African Clinical Trial Registry (PACTR) PACTR202008776694999, registered

on 11 August 2020 at https://pactr.samrc.ac.za/TrialDisplay.aspx?TrialID=12241 and has been

completed.

Findings

The mean total societal treatment cost per trial participant was US$507 (95%CI: 458; 555) in the SOC,

US$196 (95%CI: 190; 218) in the labels and US$206 (95%CI: 167; 213) in the pillbox study arms. We

estimated that there was a 49-56% probability that the implementation of the DAT interventions,

would improve the cost-effectiveness of tuberculosis treatment at a cost-effectiveness threshold of

US$100. There was no difference in DALYs between socio-economic position groups (p=0.920),

however, patient costs were less concentrated among those relatively poor in the intervention arms

– labels (illness concentration index [ICI]=0.03 (95%CI: 0.01; 0.05)) and pillbox (ICI=0.01 (95%CI:-0.01;

0.02)); compared to the SOC (ICI=-0.05 (95%CI: -0.07; -0.02). Between group comparison (p<0.001).

Interpretation

DAT interventions were cost-saving and reduced the inequitable distribution of patient costs

compared to the SOC. This highlights the potential value of interventions that reduce health service

visits in improving the equitable distribution of health services.

Funding

Unitaid (Grant Agreement Number: 2019-33-ASCENT).

Research in context

Evidence before this study

In November 2022, we searched PubMed and MedRxiv for English-language studies published
between January 2000 and current, using the terms "tuberculosis" AND "cost" AND ("Digital
Adherence Technologies" OR "DATS" OR "99DOTS" OR "Pillbox"). This search was repeated as part of
a systematic review in April 2023 followed by an update in May 2024. Twenty-nine relevant studies
have been identified, estimating the costs of DATS, though many did not assess the full economic
costs of implementation. Only two studies included an assessment of patient-incurred costs, and
none considering the equity distribution of costs or outcomes using an asset-based index.

Added value of this study

The ASCENT study provides robust evidence using a comprehensive economic evaluation framework,
that DATs decreased the cost of tuberculosis treatment in Ethiopia for a cohort of adults with
pulmonary tuberculosis. There was a 49-56% probability of DATs improving the cost-effectiveness of
tuberculosis treatment and there was no significant difference in disability adjusted life years (DALYs)
between study arms. The implementation of DATs did not change the distribution of costs or DALYs
between people with tuberculosis (PWTB) of different household socio-economic position, however
it did reduce the magnitude of patient costs among PWTB in the lower socio-economic position (SEP)
quintile.

Implications of all the available evidence

While there is limited evidence of the effectiveness of Digital Adherence Technologies (DATs), this
study is the first to show what impact the DATs may have on the costs of treatment, by reducing the
number of healthcare visits leading to cost savings. There is further evidence that DATs may reduce
the burden of patient costs on those who are least wealthy. We recommend that future investments
in DATs for tuberculosis treatment support consider how healthcare providers integrate DATs for
tuberculosis treatment support in the health facility workflow and how this translates to cost savings.

Key words

Digital adherence technology; Tuberculosis; Treatment adherence; Equity; Ethiopia

Word count 4497

Introduction

The 2018 World Health Assembly on Digital Health highlighted the potential of digital technologies to

advance universal health coverage, while cautioning that contributions should be carefully assessed

to ensure investments do not divert resources from alternative, more cost-effective non-digital

approaches [1]. Tuberculosis, a notifiable disease of public health importance, requiring adherence

to treatment schedules of 6 months or more, may benefit from innovation to support treatment

adherence. In 2022, 7.5 million people globally started treatment for tuberculosis, and with an

estimated 1.3 million deaths, it remains one of the deadliest infectious diseases [2]. In Ethiopia the

tuberculosis incidence rate is 126 (85-176) cases per 100,000 population compared to the global

average of 133. Disease risk factors include undernourishment, HIV, alcohol use disorders, diabetes,

and smoking. Of individuals newly started on tuberculosis treatment in 2021, 87% successfully

completed. Current treatment options for drug-susceptible tuberculosis (DS-TB) include a six-month

course of treatment or a 6-20-month course for rifampicin-resistant tuberculosis (RR-TB).

Intermittent dosing of tuberculosis medication as is observed with suboptimal dosing is possible due

to the post-antibiotic effect of drugs included in first-line treatment. However, the treatment

regimen will be less bacteriostatic when dosing do not coincide with the metabolic activity of semi-

dormant persistent tubercle bacilli [3]. Consistent adherence is therefore recommended to reduce

the risk of treatment failure, relapse and acquired drug resistance, especially among persons with

initial pulmonary cavitation or HIV disease .

Historically, directly observed therapy short course (DOTS) was used to achieve high rates of

treatment success, necessitating persons with TB (PWTB) attending the health facility daily to be

observed when taking their medication. Digital solutions for adherence monitoring has made

community-based, patient-centered treatment adherence support possible. A range of digital

adherence technologies (DATs) have been assessed with limited evidence of improvements in end of

treatment outcomes [4–6] and cost-effectiveness [7–10]. Interventions range from video-supported

therapy and digital pillboxes to SMS-based interventions. The use of these interventions may be

dependent on PWTB's access to digital devices, and concerns have been raised about equity

implications [11]. Since lower household socio-economic position (SEP) increases individuals' risk of

contracting tuberculosis disease [12], simultaneously increasing the risk for suboptimal treatment

adherence and poor health outcomes, it is important to assess the equity implications of digital

interventions. As part of the adherence support coalition to end tuberculosis (ASCENT) project, a

pragmatic cluster randomised trial was conducted to assess the effectiveness and cost-effectiveness
of digital pillboxes (pillbox) and medication labels (labels) with daily monitoring using a web-based
platform to inform differentiated care, in reducing a composite unfavourable outcome, including
tuberculosis recurrence in Ethiopia. The trial did not find evidence of a difference in the composite
unfavourable outcomes between the labels (adjusted odd ratio [aOR]=1.14; 95%CI:0.83-1.61; p=0.62)
or pillbox study arms (aOR=1.04; 95%CI:0.74-1.45; p=0.95) when compared against the standard of
care (SOC) [13]. We conducted a trial-based distributional cost-effectiveness analysis to assess the
equity impact, costs and cost-effectiveness of digital adherence technologies with differentiated care
compared to the SOC in Ethiopia.

136
137

Methods

Study design and population.

We conducted a trial-based distributional cost-effectiveness analysis using participant-level data,
following a pre-specified health economics analysis plan [14]. Costs and outcomes were assessed
from a societal perspective. ASCENT was a three-arm pragmatic cluster randomised trial of digital
adherence technologies (DATs) for tuberculosis treatment support, followed by differentiated care
compared to the standard of care (SOC) in Ethiopia. The unit of randomisation was health facilities,
with 78 facilities (26 per study arm) in the Addis Ababa and Oromia regions of Ethiopia enrolled.
Clusters were randomised (1:1:1) using stratified restricted randomisation, based on province and TB
notifications to provide either SOC, medication labels (labels), or digital pillboxes (pillbox;
EvriMed1000). 3,858 individuals ≥ 18 years of age, with pulmonary DS-TB were enrolled and
followed up 12 months after treatment initiation to determine outcomes. Two trial endpoint
measures were collected, (i) end-of-treatment outcome recorded in the facility TB treatment register,
and (ii) for all participants with bacteriologically confirmed tuberculosis who had either a cured or
completed end of treatment outcome, self-reported TB retreatment. For participants able to provide
a sputum sample, culture was done six months after the end of treatment to measure disease
recurrence. Data on patient events and trial endpoints were collected for all trial participants.
Provider – and patient costs were collected from the same sample of 15 health facilities (5 per study
arm), and 10 participants per facility (total of 150 observations).

159
160

Ethics

The study received ethics approval from LSHTM Ethics Committee, UK (19120–1); WHO Ethical
Review Committee, Switzerland (0003297); the Addis Ababa City administration health bureau
public emergency and health research directorate institutional review board (A/A/H/B/947/227);
and the Oromia Regional Health Bureau public emergency and health research directorate
institutional review board (BEFO/HBTFH/1–16/10322), Ethiopia. Written informed consent were
sought from individuals enrolled in the study.

Interventions

Two DAT interventions, pillbox and labels, were introduced to health facilities randomised to the
intervention arms (one DAT per intervention arm). Health facilities assigned to the SOC study arm
continued routine practice. TB focal staff providing tuberculosis care in each of the health facilities,
received training from trial staff in tuberculosis treatment adherence monitoring and the use of the
DATs. Each dose taken by participants was logged on the platform either automatically when the
participant opened the pillbox to take their medication, or when the participant texted a code on
the dose label to the dedicated toll-free number. Participants receiving care in facilities randomised
to labels, who did not have a phone were offered a pillbox. DATs were linked to an online platform
for daily monitoring of participant engagement with the DAT. Automated SMS reminders were sent
to all participants who did not open their pillbox or had not sent a text by 11:00 each day. Doses not
recorded were considered missed and the TB focal person started a differentiated response based
on data on the platform. The response included a phone call to the patient if 1-2 doses are missed;
home visits by a community healthcare provider if there is evidence of persistent missed doses (>5
consecutive doses); and a switch to DOT was considered if >14 doses were missed.

Assessment of household socio-economic position, cost and outcomes

Between May 2021 and August 2022, adults receiving tuberculosis treatment at participating health
facilities were offered enrolment into the trial. Participants were followed up for 12 months until
August 2023. Patient events were collected at enrolment, from treatment registers and at the 12-
month follow-up [14]. Household socio-economic position (SEP) as a relative measure of poverty was
estimated by calculating an asset index through principal component analysis of 27 multi-
dimensional poverty indicators collected using a survey at enrolment (table S2). Using this index,

participants were divided into quintiles of relative wealth. Costs and outcomes were compared
between quintiles using concentration index-based indices [15]. Illness concentration curves were
constructed by plotting the cumulative percentage of illhealth against the proportion of the
population ranked by household SEP. The standard illness concentration index (ICI) derived from the
curve was defined as twice the area between the concentration curve and the line of equality on the
graph [16].

Cost categories included technology, network – and support costs, health service costs of treatment,
and patient-incurred costs. Technology costs included the cost of the devices, accounting for cluster-
level re-use rates for pillboxes. Network costs for labels was the costs of two-way SMS' and monthly
hosting costs of the web-based platform for both technologies. Support costs include the cost of
training, support visits to health facilities, and 0.5 full-time equivalent of a technical officer to
support the implementation and integration of the intervention. Provider costs were estimated at
facility level for 13 health facilities. The cost of staff time was estimated using timesheets completed
by tuberculosis focal staff (table S3). A separate log kept by trial staff was used to record the number
of support visits, telephone calls and travel to health facilities. Time spent was valued using the
average wage rate for health care workers in Ethiopia. The number of visits to health facilities were
recorded prospectively for each participant, on a facility log and multiplied by the provider unit costs
to calculate total costs. Costs valued before 2023 were inflated using the average annual consumer
inflation rate in Ethiopia over the previous five years (22.2% per annum). For the cost of
hospitalisation, diagnostic tests and RR-TB treatment, we used secondary data on unit costs
multiplied by the number of events collected for each participant [17]. Data on patient-incurred costs
such as transport and other out-of-pocket costs were collected by surveying 10 trial participants in
each of the 15 health facilities (5 per study arm). Costs reported in Ethiopian Birr were converted to
2023 United States Dollars using the average annual exchange rate (1 EB = 0.018 USD).

We estimated participant-level disability-adjusted life years (DALYs) based on participants' gender,
age, and trial endpoint. DALYs are the product of years of life lost, and years lived disabled due to
tuberculosis. Years of life lost due to premature mortality were based on age- and gender-stratified
expected life expectancy for Ethiopia, if the participant died during treatment or prior to 12-month
follow-up. Years lived disabled were calculated based on time spent on treatment (including
recurrence or drug-resistant tuberculosis) multiplied by the global burden of disease disability weight
for tuberculosis disease (Eq S6) [18].

Statistical analyses

Cluster-level mean provider costs from 15 health facilities were singly imputed to the 78 trial health

facilities by province (Addis/Oromia) and study arm. Patient costs, socio-economic position and

DALYs at 12 months were jointly imputed for participants with missing values using multi-level

multiple imputation by chained equations, generating 50 imputed datasets. For each of the imputed

datasets, costs and DALYs were jointly modelled using bivariate Bayesian hierarchical models and log-

normal distributions with random effects to allow for cluster-level variances. Data were fitted using

Markov Chain Monte Carlo sampling to obtain posterior simulations. The results of the imputed

models were pooled by combining the posterior draws for all 50 imputed models. Total costs and

DALYs in each study arm were estimated by summing participant- level costs and DALYs. The mean

differences in costs and DALYs were calculated group-wise by subtracting the arithmetic mean cost in

each of the intervention groups from those in the SOC arm. The incremental cost-effectiveness ratio

(ICER) was calculated by the differences in costs divided by the difference in DALYs averted between

the intervention arms and the SOC study arm (Eq S7). The distributional incremental cost-

effectiveness ratio (dICER) was calculated as the ratio of net health benefit divided by the difference

in the illness concentration indices (ICI) between the intervention and SOC study arms (Eq S11). ICERs

were compared against a cost-effectiveness threshold for Ethiopia of $100 per DALY averted [19].

Prespecified sensitivity analyses were used to assess how changes in the analysis may affect the

results of our study; including (i) changing the number of healthcare visits in the SOC to the number

of visits observed under guidelines circumstances; (ii) increasing the cost of the intervention; (iii) only

including the costs and outcomes incurred during the treatment of trial participants until the end of

the trial; (iv) valuing participant time using the average wage rate of government employees in

Ethiopia; and (v) varying the discount rate used for costs and effects from 0% to 10%. Variance in the

posterior draws of the model compared against a range of thresholds were used to quantify the

uncertainty in our estimates in a cost-effectiveness acceptability frontier for the probabilistic

sensitivity analysis. Data were analysed using Stata 18 and R statistical software. The study is

reported using the Consolidated Health Economic Evaluation Reporting Standards (CHEERS) 2022

checklist.

Governance

The study was funded by UNITAID (2019-33-ASCENT) through the Adherence Support Coalition to

End TB (ASCENT) project. The funder had no role in study design, data collection, data analysis or

interpretation or the writing of the manuscript. The trial had independent oversight from a Technical

Advisory Group (TAG) with representatives from all five ASCENT countries, as well as a Community

Advisory Group (CAB) including representatives from people affected by tuberculosis in Ethiopia. The

Ethiopian National TB Programme was involved in the design of the economic evaluation and

received regular updates. The trial is registered with Pan African Clinical Trial Registry (PACTR)

PACTR202008776694999, registered on 11 August 2020 at

https://pactr.samrc.ac.za/TrialDisplay.aspx?TrialID=12241.

Results

Between May 2021 and August 2022, 3885 adults (≥18 years) starting DS-TB treatment were

enrolled, with follow-up visits completed in August 2023 (figure S1). Participants were excluded if

their diagnosis changed to "not TB" or if they were diagnosed with RR-TB treatment less than 28

days before treatment start or were re-enrolled leaving 1295 participants in the standard of care

(SOC), 1305 in the Labels -, and 1258 in the Pillbox arms of the study in the intention-to-treat

population.   Participants were analysed in the based on DAT received, leaving 1295 in SOC, 1152 in

labels and 1411 in pillbox (table S1). Baseline characteristics of the clusters and trial participants are

shown in Table 1.

Table 1. Baseline characteristics of the enrolled population by study arm participants received.

	Standard of care	Labels	Pillbox
Characteristics of the clusters/ health facilities			
Number of clusters	26	26	26
Region			
Addis Ababa	12 (46%)	12 (46%)	12 (46%)
Oromia	14 (54%)	14 (54%)	14 (54%)
Area			
Urban	24 (92%)	24 (92%)	25 (96%)
Rural	2 (8%)	2 (8%)	1 (4%)
Median number of individuals per cluster (IQR)	48 (41, 59)	50 (36, 65)	45 (39, 50)
Baseline characteristics of the enrolled population			
Number of trial participants	1 295	1 152	1 411
Mean age in years (SD)	35.5 (14.7)	32.4 (13.5)	34.3 (14.1)
Female	496 (38%)	487 (42%)	584 (41%)
Bacteriologically confirmed tuberculosis	806 (62%)	756 (66%)	936 (66%)
Previous tuberculosis treatment	112 (9%)	63 (5%)	81 (6%)
HIV status			
Negative	1 134 (88%)	1 012 (88%)	1 218 (86%)
Positive, not on ART	2 (0%)	4 (0%)	9 (1%)
Positive, on ART	157 (12%)	133 (12%)	182 (13%)
Unknown	2 (0%)	3 (0%)	2 (0%)
Educational level			
None (Iliterate)	198 (15%)	114 (10%)	220 (16%)
Less than primary	73 (6%)	65 (6%)	119 (8%)
Primary	635 (49%)	571 (50%)	647 (46%)
Secondary or Higher	376 (29%)	394 (34%)	419 (30%)
Unknown	13 (1%)	8 (1%)	6 (0%)
Mean number of people per room of the house (SD)	2.6 (1.7)	2.6 (1.6)	2.5 (1.5)
Relationship status and living arrangements as a proxy for social capital			
Single	309 (24%)	313 (27%)	218 (15%)
Single (independent)	193 (15%)	211 (18%)	321 (23%)
Married or cohabitating	621 (48%)	528 (46%)	146 (10%)
Separated or widowed	158 (12%)	93 (8%)	52 (4%)
Unknown	14 (1%)	7 (1%)	6 (0%)
Frequency of income received			
No income	277 (21%)	266 (23%)	324 (23%)
Irregularly	198 (15%)	127 (11%)	286 (20%)
Seasonally	243 (19%)	245 (21%)	291 (21%)
Monthly	563 (43%)	507 (44%)	504 (36%)
Unknown	14 (1%)	7 (1%)	6 (0%)
Household socio-economic position			

Poorest	253 (20%)	211 (18%)	263 (19%)
Less poor	258 (20%)	187 (16%)	280 (20%)
Middle	227 (18%)	209 (18%)	289 (20%)
More wealthy	239 (18%)	238 (21%)	249 (18%)
Wealthiest	211 (16%)	260 (23%)	257 (18%)
Unknown	107 (8%)	47 (4%)	73 (5%)

284 All data shown are n(%) or mean (SD) or n/N(%). IQR=interquartile range. ART=antiretroviral therapy. SD=standard
285 deviation. Each health facility is one cluster.
286

Baseline characteristics were similar between study arms. 62% (806/1295) of participants in the SOC,
compared to 66% (756/1152) in the labels and 66% (936/1411) in the pillbox arms had
bacteriologically confirmed tuberculosis. Slightly more participants in the SOC arm had a history of
previous tuberculosis disease at 9% (112/1295) compared to 5% (63/1152) in the labels and 6%
(81/1411) in the pillbox arms. Considering the variables describing household socio-economic
position (SEP), similar proportions of participants in the SOC 21% (277/1295) compared to the labels
23% (266/1152) and pillbox 23% (324/1411) arms of the study reported no individual income.
Participants in the SOC arm were less wealthy (20% classified as lowest wealth quintile) compared to
the labels and pillbox arms (18% and 19% respectively). Participants in the labels study arm were
wealthier (43% (498/1152) in the two wealthiest quintiles) than those in the SOC (35% (450/1295))
and pillbox (36% (506/1411)) study arms. The number of health facility visits per treatment episode
compared between study arms are shown in table S4. Compared to the SOC, there was a reduction
in the mean number of visits to health facilities in the pillbox and labels study arms throughout the
treatment period; labels compared to SOC by -20.2 (95%CI: -26.3; -14.1; $p<0.001$) and pillbox
compared to SOC -20.4 (95%CI: -26.5;-14.3; $p<0.001$) (table S4). Cohort level patient events during
treatment compared between study arms are summarised in table S5. There were more home visits
in the labels (77), and pillbox (58) compared to the SOC (5) study arms. Similarly nights in hospital
was greater in the labels (648) and pillbox arms (310) compared to the SOC (135). This was partly
explained by one healthfacility in the labels arm that admitted more people to hospital, and may also
be as a result of more intensive follow-up because of monitoring using the DATs. A total of 22
participants required a second course of tuberculosis treatment in the SOC, compared to 24 in the
labels and 31 in the pillbox study arms. With 4 participants in the SOC, 1 in labels and 2 in the pillbox
study arms starting rifampicin resistant tuberculosis treatment. The economic costs of treatment are
summarised in Table 2.

Table 2. Total and per-patient costs (2023 USD) incurred for trial participants, by cost categories and study arm.

	Standard of Care (n=1295)		Labels (n=1152)		Pillbox (n=1411)	
	Total costs [95%CI]	Cost/patient [95%CI]	Total costs [95%CI]	Cost/patient [95%CI]	Total costs [95%CI]	Cost/patient [95%CI]
Technology	N/A	N/A	$806 [fixed]	$0.70 [fixed]	$11,006 [10,752; 11,161]	$7.80 [7.62; 7.91]
Network costs	N/A	N/A	$12,534 [fixed]	$10.88 [fixed]	$9,665 [fixed]	$6.85 [fixed]
Support and training	$1,334 [1,308; 1,373]	$1.03 [1.01; 1.06]	$1,140 [1,117; 1,175]	$0.99 [0.97; 1.02]	$1,270 [1,242; 1,298]	$0.90 [0.88; 0.92]
Sub-total intervention costs	**$1,334 [1,308; 1,373]**	**$1 [1; 1]**	**$14,481 [14,135; 14,826]**	**$13 [12; 13]**	**$21,941 [21,419; 22,364]**	**$16 [15; 16]**
Health facility visits	$472,675 [448,070; 498,575]	$365 [346; 385]	$78,336 [76,032; 79,488]	$68 [66; 69]	$88,893 [87,482; 91,715]	$63 [62; 65]
Drug sensitive tuberculosis treatment	$45,325 [44,030; 45,325]	$35 [34; 35]	$40,320 [39,168; 40,320]	$35 [34; 35]	$45,325 [47,974; 49,385]	$35 [34; 35]
Smear microscopy tests	$75,110 [75,110; 76,405]	$58 [58; 59]	$66,816 [66,816; 67,968]	$58 [58; 59]	$81,838 [81,838; 83,249]	$58 [58; 59]
Hospitalisation	$1,036 [376; 1,645]	$0.80 [0.29; 1.27]	$4,712 [2,684; 6,751]	$4.09 [2.33; 5.86]	$2,300 [1,242; 3,344]	$1.63 [0.88; 2.37]
Treatment for recurrence (drug-sensitive) or multi-drug resistance tuberculosis	$41,440 [6,475; 76,405]	$32 [5; 59]	$19,584 [1,152; 40,320]	$17 [1; 35]	$28,220 [1,411; 55,029]	$20 [1; 39]
Sub-total provider costs.	**$635,586 [574,061; 698,355]**	**$491 [443; 539]**	**$209,768 [185,852; 234,847]**	**$182 [177; 204]**	**$246,576 [219,947; 282,722]**	**$178 [156; 200]**
Patient incurred costs[1]	$10,360 [10,360; 10,360]	$8 [8; 8]	$10,368 [9,216; 10,368]	$9 [8; 9]	$9,877 [8,466; 9,877]	$7 [6; 7}
Productivity costs[2]	$10,360 [10,360; 10,360]	$8 [8; 8]	$6,475 [6,475; 6,475]	$5 [5; 5]	$7,055 [7,055; 8,466]	$5 [5; 6]
Sub-total patient-incurred costs	**$20,720 [20,720; 20,720]**	**$16 [15; 16]**	**$16,843 [15,691; 16,843]**	**$14 [13; 14]**	**$16,932 [15,521; 18,343]**	**$12 [11; 13]**
Total societal costs	**$657,640 [596,089; 720,447]**	**$507 [458; 555]**	**$241,091 [215,678; 266,516]**	**$196 [190; 218]**	**$285,449 [256,887; 323,429)**	**$206 [167; 213]**

314
315
316
All costs are shown as the mean costs in 2023 USD. [1]Based on single imputation. [2]Productivity costs are based on the value of individuals' time spent travelling to health facilities or waiting at the facility. 95%CI=95% confidence interval. Technology and network costs are fixed costs per patient as primarily purchased from third party providers at a fixed rate of cost per patient, except in the technology costs for pillbox where there is variability because of differences in cluster-level device re-use rates.

Intervention costs were $13 (95%CI: 12; 13) per patient in the Labels compared to $16 (95%CI: 15; 16) in the Pillbox arm, with an empirically derived pillbox reuse rate of 2.16 (95%CI: 1.58; 2.85) included in the calculation. Per patient provider costs were much higher in the SOC $491 (95%CI: 443; 539) compared to the Labels $182 (95%CI: 177; 204) and pillbox study arms $178 (95%CI: 156; 200), driven by the larger number of health care visits in the SOC. The cost of medication, diagnostic tests, hospitalisation, and further treatment were similar between study arms. Patient-incurred costs are similar between study arms at $16 (95%CI 15; 16) per person in the SOC, $14 (95%CI: 13; 14) and $12 (95%CI: 11; 13) in the labels and pillbox arms respectively despite the large difference in the number of health facility visits in the intervention arms. The distributions of DALYs, health facility visits, patient – and provider costs by household socio-economic position (SEP) are shown in Figure 1.

[FIGURE 1]

Figure 1. illness concentration indices for (A) Disability Adjusted Life Years (DALYs); (B) Number of health facility viists; (C) Patient costs in 2023 USD; and (D) Provider costs in 2023 USD.
The concentration index (ICI) assesses if there is a within study arm difference in the outcome by ranked by household socio-economic position (SEP). Clustering was taken into account in estimating 95% confidence intervals (95%CI). A negative concentration index means that the outcome, here a negative consequence, is concentrated among less wealthy households. Conversely, a positive index value means that the outcome is concentrated among relatively wealthier households. We compared the difference in the distribution in outcome between SEP groups, where the *p*-value represents the likelihood of any observed difference between the groups being by chance.

The equity impact of the interventions in four domains, namely DALYs, health facility visits, patient-

incurred costs and provider costs are shown in figure 1, figure s2 and table S6. The illness

concentration indices (ICIs) for DALYs were -0.06 (95%CI: -0.24; 0.11) in SOC; -0.02 (95%CI: -0.17;

0.14) in labels and -0.02 (95%CI: -0.16; 0.12) in pillbox study arms, with no evidence of differences in

the distribution of DALYs within or between study arms (p=0.920). The number of health facility visits

were concentrated among the comparatively poor in the SOC -0.07 (95%CI: -0.10; -0.05) and pillbox -

0.08 (95%CI: -0.09; -0.06) arms but not in the labels arm 0.01 (95%CI: -0.00; 0.02). Patient costs were

concentrated among those relatively poor in the SOC arm -0.05 (95%CI: -0.07; -0.02), while

concentrated among the wealthy in the labels arm 0.03 (95%CI: 0.01; 0.05) and pillbox 0.01 (95%CI: -

0.01; 0.02) with evidence of differences between study arms (p<0.001). Provider costs were

concentrated among the relatively wealthy, suggesting that the wealthy most benefits from public

health expenditure, in the SOC 0.13 (95%CI: 0.08; 0.18); labels 0.07 (95%CI: 0.05; 0.08) and pillbox

0.07 (95%CI: 0.04; 0.10) arms, though there was insufficient evidence (p=0.225) of a difference

between study arms. The cost-effectiveness and distributional cost-effectiveness of the interventions

compared   to   the   SOC   are   summarised   in   Table   3.

Table 3. Incremental cost-effectiveness from the provider – and societal perspectives.

	Simple imputation of the means		Multilevel multiple imputation		Group-level summary metrics	
	Incremental costs per patient [95% CI]	DALYs averted [95% CI]	Incremental costs per patient [95% CI]	DALYs averted [95% CI]	Incremental cost-effectiveness ratio [95% CI]	Incremental distributional cost-effectiveness ratio [95% CI]
	Mean difference [95% CI]	Mean difference [95% CI]	Mean difference [95% CI]	Mean difference [95% CI]		
Provider perspective						
Labels versus SOC	-309 [-266; -335]	-0.46 [-2.07; 1.17]	7 [-191; 189]	0.04 [-1.93; 1.93]	672 [14; 1420]	18.53 [4.16; 31.85]
Pillbox versus SOC	-313 [-287; -339]	-0.32 [-1.61; 0.97]	-1 [-190; 183]	-0.01 [-1.93; 1.93]	978 [-54; 1626]	19.80 [8.88; 30.73]
Societal perspective						
Labels versus SOC	-311 [-268; -337]	-0.46 [-2.07; 1.17]	7 [-191; 189]	0.04 [-1.93; 1.93]	676 [17; 1435]	18.68 [4.30; 31.99]
Pillbox versus SOC	-301 [-291; -342]	-0.32 [-1.61; 0.97]	-1 [-190; 183]	-0.01 [-1.93; 1.93]	941 [-53; 1652]	18.96 [9.16; 30.94]

*Costs reported based on multiple multi-level imputation. In a separate analysis, we estimated costs using single imputation where the means of patient costs were used from the subsample –
the findings of this analysis are summarised in the appendix along with a detailed report of the costs estimated. The current estimates suggest that the cost-effectiveness threshold (CET) for
Ethiopia is between $10 and $255 (we used an average of $93) per DALY averted. For the comparison, we use $100 as an estimate within the range. LTFU: lost to follow-up. [f]Mean age at death
was 48.6 (SD: 18.6) years in the Standard of Care arm; 43.9 (SD: 17.8) years in the Labels - and 41.2 (SD: 14.7) in the pillbox arm. DALYs: Disability adjusted life years. SD: standard deviation.
95%CI: 95% Confidence Interval. The distributional cost-effectiveness ratio was estimated, with the efficiency impact being the net monetary benefit calculated as the change in effectiveness
multiplied by the CET minus the change in costs (EqS8); and the equity impact being the difference in health inequality measured as the difference in the illness concentration indices between
study arms. The distributional cost-effectiveness ratio is then a ratio of incremental efficiency/ equity impact (EqS12).
25

26   The difference in societal costs, using multiple imputation, was a reduction by $299 per patient in the

27   Labels and $298 in the Pillbox arms when compared against the SOC. The societal cost per DALY

28   averted was $676 (95%CI 17; 1435) in the Labels and $941 (95%CI -53; 1652) in the Pillbox study

29   arms.

31   [FIGURE 2]

Figure 2. Cost-effectiveness planes
Scatter plot of the estimated joint density of incremental costs and incremental DALYs averted, showing uncertainty in the
ICER estimate obtained from the analysis models. The dots represent the findings of a probabilistic sensitivity analysis.

When plotted on cost-effectiveness planes, we find that the ICERs are crossing the axes but slightly

concentrated in the bottom left hand quadrant suggesting that DATs are less costly and less effective

than the standard of care. ICERs in this quadrant are interpreted as cost-effective if they fall below

the cost-effectiveness threshold. Using the outcomes of the probabilistic sensitivity analysis, plotted

on the cost-effectiveness acceptability frontier, the probability that each of the interventions would

be cost-effective compared against a range of cost-effectiveness thresholds (CETs) are shown in

figure S3. Using a range of US$5 to US$900 per DALY averted for the CETs, we find that the

interventions have a 56% probability (labels) and 49% probability (pillbox) of being cost-effective

compared to SOC at a CET of US$100. The results of the distributional cost-effectiveness analysis

estimates the distributional cost-effectiveness ratio at 18.68 for labels and 18.96 for pillbox. We

compare this finding against a relative health inequality aversion parameter of 5.66. In the absence

of empirical research on the level of inequality aversion by policymakers in Ethiopia, we used an

estimate previously derived for decisions in Ethiopia [20]. As the value of the inequality aversion

parameter increases, it suggests that there is a greater willingness to achieve fewer health

improvements to reduce inequality in health or other outcomes. Our distributional cost-effectiveness

ratio estimated (18.68 and 18.96) exceeds the parameter used for previous decisions in Ethiopia for

vaccines (5.66) suggesting that policy makers would need to be more averse to inequality than in

previous decisions, to recommend the implementation of DATs based on health equity.

In the sensitivity analyses (table S7), we found that more visits in the SOC increased potential cost-

savings from implementing the interventions. If we assumed an increased cost of the pillbox, for

example in setting where reuse rate is low, the cost savings compared to the SOC reduced by 16%.

Removing the costs and DALYs associated with recurrence reduces cost savings and has minimal

effect on the DALYs averted. Valuing participants using an average wage rate increases the cost

savings to patients. Main study findings, however, remained similar under sensitivity analysis.

Discussion

In health facilities implementing DATs, trial participants made 20 fewer visits per treatment episode
compared to the standard of care (SOC), substantially reducing the cost to providers. We didn't find
evidence of differences in poor health outcomes (DALYs) between study arms or their distribution
between socio-economic position (SEP) groups. However, there was a reduction in patient-incurred
costs among the least wealthy households in the intervention compared to SOC study arms. When
assessing uncertainty, the 95% confidence intervals of the incremental cost-effectiveness ratios
(ICERs) were wide, suggesting substantial uncertainty that DATs would be cost-effective when
compared to the SOC. We estimated that the labels and pillbox arms have a 49-56% probability of
improving the cost-effectiveness of tuberculosis treatment compared to SOC at a cost-effectiveness
threshold of $100 per DALY averted. Our study follows the work of other cluster randomised trials
evaluating the effectiveness and costs of Digital Adherence Technologies (DATs) in China [5,10], South
Africa [21], Uganda [7] and Ethiopia [22], by including a detailed analysis of human resources, patient
costs and equity in addition to costs and cost-effectiveness.

Differences in provider costs between the SOC and intervention arms of the study were primarily
driven by the number of healthfacility visits per treatment episode. In a separate analysis, we
showed that health care workers spent more time with PWTB per visit in the intervention arms than
when they were attending more frequently in the SOC arm [23]. This increase in time spent
remained lower than the cumulative time of multiple visits in the SOC. Our evaluation of cost of
treatment included above service-level costs. These costs were highly variable, particularly between
urban and rural health facilities where fewer people with tuberculosis (PWTB) were accessing
services. Given that we had overhead costs from only 12 clusters, single imputation of provider costs
has likely resulted in larger variability than if we had collected data for all 78 clusters. Provider costs
reported in this study were broadly similar to the costs estimated by Terefe et al's comprehensive
assessment of the cost of tuberculosis care in Ethiopia, though we used a micro-costing approach
[17]. Our total cost estimate, includes costs incurred by all participants who received the intervention
irrespective of whether they completed treatment (ITT population). Provider costs incurred through
diagnostic tests, hospitalisation and subsequent treatment episodes were similar between study
arms, and differences in treatment costs were predominantly driven by the reduction in the number
of health facility visits.

Patient costs were similar in the intervention - compared to the SOC arms of the study, despite large

differences in facility visits. This is explained by lower transport costs per visit reported in the SOC

where there are more frequent visits compared to the intervention study arms. Furthermore, we

used the human capital approach for valuing time spent seeking care, whereby individual self-

reported income is used to value people's time. This likely does not fully captured the full cost of

time spent. Approximately 35% of participants reported no- or irregular income with consequently

no cost allocated to their time spent. Furthermore, we collected self-reported income data at

enrolment into the trial when people were on treatment and already had suffered disease-

associated income loss. A recent meta-analysis of patient costs associated with tuberculosis in

Ethiopia, found that pre-diagnosis costs were more than double post-diagnosis costs US$97.6 (56.4-

184.3) versus US$45.1 (-119.1-209.37)) [28]. Differences in time spent between study arms due to a

reduction in number of visits, would therefore only have a marginal effect on productivity costs. We

quantified this potential bias by conducting a sensitivity analysis estimating the productivity costs for

each of the study arms by valuing all participants' time equally.

Our study is the first to empirically demonstrate that DATs could be cost saving by reducing the

number of health facility visits. This observed difference is dependent on the frequency of health

facility visits in the SOC. At the time of our study, PWTB were attending health facilities on average

30 times per treatment episode, our findings may therefore not hold in settings where less frequent

facility visits are already routine practice. Furthermore, the observed difference between study arms

would have been more pronounced prior to COVID-19, when visits to health facilities were more

frequent [25]. Other studies evaluating the economic costs of DATs found them to be more costly and

effect-neutral [7,8,10]. In Uganda, the provider cost of 99DOTS was estimated to be $59 (range: $50-

$70) per participant successfully treated [7]. Similarly, in Morocco, the cost-effectiveness of a digital

pillbox from a provider perspective was found to be $434 per DALY averted among MDR-TB patients.

This was a model-based analysis where the number (and cost) of health facility visits were assumed

to be the same between study arms [8]. A study in China, considering the societal perspective, found

digital pillbox to be more costly and effect neutral with an ICER of $3,668.59 per DALY averted. Data

for assessing number of health facility visits was limited (n=120) and did not fully assess differences

in practice between studyarms [10].

Strengths of our study include it being a trial-based cost-effectiveness analysis with detailed data

collection of patient events, healthfacility visits and rigourous assessment of trial endpoints

embedded within a randomised study where the SOC reflects routine practice in Ethiopia. We

furthermore went beyond the traditional cost-effectiveness framework to analyse the equity impact

of the interventions. Study limitations include, the trial follow-up period was not long enough to

capture all health and economic consequences of the interventions – particularly tuberculosis

transmission. This question was addressed in a transmission modelling study following this study

[26]. The number of health facility visits were recorded prospectively, and patient event data

(including hospitalisations) collected at the 12 month follow-up. Self-reported patient events would

not be available for patients who died or became LTFU during treatment episode and we

underestimate the cost associated with care-seeking prior to death or LTFU. Given the

representativeness of the sample, imputed costs would undervalue true costs. Our DALY estimates

are likely to be conservative, because we only include DALYs for the current and identified future

treatment episodes and not for any longer-term sequalae of tuberculosis disease [27]. Our estimates

are similarly likely to underestimate the full costs and economic losses incurred because our

analytical time horizon, restricted to the current tuberculosis event, includes costs incurred once

participants started treatment and not during the diagnostic phase. We could only assess the equity

distribution impact of the intervention among those already accessing care, and expecting to live in

the health facility's catchment area for the duration of treatment. We may therefore be

underestimating the value of the intervention among more marginalised groups. Nevertheless, we

expect the findings to be representative of adults who are accessing care at public health clinics in

Ethiopia.

We used relative SEP based on multi-dimensional poverty assessment, to incporate equity in the

decision framework [29]. Alternative approaches are based on estimates of individual income to

assess inequality [30]. Measures of income-based inequality would correctly ascertain immediate

ability to respond to a health shock, but would fail to account for resource-sharing within families,

saving ability, and non-monetary assets that are important in assessing poverty in rural areas. Multi-

dimensional poverty metrics include locally-relevant assets such as land, livestock, social capital and

education that individuals may draw on in the event of a health shock. Our evaluation as part of the

trial, was not powered to compare outcomes between SEP groups within study arms, but sufficient

to evaluate how the concentration of the outcome between those relatively wealthy and relatively

poor varies between study arms.

In conclusion, we found the implementation of DATs to be effect-neutral but cost-saving due to a

marked reduction in healthfacility visits. These savings would only be fully realised if healthcare

providers have other productive tasks to fill their time previously spent on DOTS, or in contexts

where routine care is not daily facility-based DOTS. We found limited evidence of differences in
DALYs averted between study arms and estimate a 49-56% probability that DATs will improve the
cost-effectiveness of tuberculosis treatment at a CET of \$100 per DALY averted. The implementation
of DATs reduced the inequitable burden of patient-incurred costs.

Authors' contributions

NF, KLF, AWT and CFM designed this study. The trial was designed by KLF, KvK, JvR and DJ. All authors
have contributed to the study design. KLF provided statistical expertise in clinical trial design and
analysis. NF conducted the analysis and provided health economics expertise in design and analysis.
AWT, MB, MS, AB, TA, NM, KvK and JvR implemented the study. All authors contributed to the
interpretation of the findings and revision of the final manuscript. The authors read and approved
the final manuscript.

Acknowledgements

The authors acknowledge the contributions of the participants who enrolled in the study, the
Ethiopian tuberculosis focal persons implementing the digital adherence technologies and the
National Tuberculosis Program at the Ministry of Health Ethiopia without whom this study would not
have been possible. We acknowledge the oversight and inputs of the ASCENT Technical Advisory
Group.

Data availability

Data analysed in this study are part of the ASCENT Consortium Trials data. A minimal dataset
required   to   reproduce   the   findings   and   the   survey   materials   is   available   from
https://doi.org/10.17037/DATA.00003721 and analysis code from https://github.com/nicolacfoster.
The minimal dataset is available for non-commercial use upon request, after signing a data sharing
agreement, to studies approved by an ethics committee. Any publications arising from the shared
data must acknowledge the investigators who collected the data, the institutions involved, and the
funding sources by citing the data record and including a standard acknowledgement statement
provided.

References

1.   World Health Organisation. WHO guideline: recommendations on digital interventions for health
system strengthening. Geneva, Switzerland: WHO; 2019.

2.   World Health Organisation. Global Tuberculosis Report 2023. World Health Organisation; 2023.
Available: https://iris.who.int/bitstream/handle/10665/373828/9789240083851-
eng.pdf?sequence=1

3.   Chang KC, Leung CC, Grosset J, Yew WW. Treatment of tuberculosis and optimal dosing
schedules. Thorax. 2011;66: 997–1007. doi:10.1136/thx.2010.148585

4.   Liu X, Thompson J, Dong H. Digital adherence technologies to improve tuberculosis treatment
outcomes in China: a cluster-randomised superiority trial. Lancet Glob Health. 2023;11: 693–
703.

5.   Wei X, Hicks JP, Zhang Z, Haldane V, Pasang P, Li L, et al. Effectiveness of a comprehensive
package based on electronic medication monitors at improving treatment outcomes among
tuberculosis patients in Tibet: a multicentre randomised controlled trial. The Lancet. 2024;
S0140673623022705. doi:10.1016/S0140-6736(23)02270-5

6.   Cattamanchi A. Digital adherence technology for tuberculosis treatment supervision: A stepped-
wedge cluster-randomized trial in Uganda'. PLoS Medicine. 2021;18: 1–15.
doi:10.1371/journal.pmed.1003628.

7.   Thompson RR, Kityamuwesi A, Kuan A, Oyuku D, Tucker A, Ferguson O, et al. Cost and Cost-
Effectiveness of a Digital Adherence Technology for Tuberculosis Treatment Support in Uganda.
Value Health. 2022;25: 924–930. doi:10.1016/j.jval.2021.12.002

8.   Yang J, Kim H-Y, Park S, Sentissi I, Green N, Oh BK, et al. Cost-effectiveness of a medication event
monitoring system for tuberculosis management in Morocco. PLoS One. 2022;17: e0267292.
doi:10.1371/journal.pone.0267292

9.   Nsengiyumva NP. Evaluating the potential costs and impact of digital health technologies for
tuberculosis treatment support. Eur Respir J. 2018;52.

10.   Sweeney S, Fielding K, Liu X, Thompson JA, Dong H, Jiang S, et al. Unit costs and cost-
effectiveness of a device to improve TB treatment adherence in China. IJTLD OPEN. 2024;1: 299–
305. doi:10.5588/ijtldopen.23.0451

11.   Saunders MJ, Wingfield T, Tovar MA, Herlihy N, Rocha C, Zevallos K, et al. Mobile phone
interventions for tuberculosis should ensure access to mobile phones to enhance equity - a
prospective, observational cohort study in Peruvian shantytowns. Trop Med Int Health. 2018;23:
850–859. doi:10.1111/tmi.13087

12.   Oxlade O, Murray M. Tuberculosis and poverty: why are the poor at greater risk in India? PLoS
One. 2012;7: e47533. doi:10.1371/journal.pone.0047533

13.   Tadesse AW, Sahile M, Foster N, McQuaid CF, Weldemichael GT, Abdurhman T, et al. Cluster-
randomized trial of digital adherence technologies and differentiated care to reduce poor end-
of-treatment outcomes and recurrence among adults with drug-sensitive pulmonary TB in
Ethiopia. 2024. doi:10.1101/2024.05.09.24307117

14. Foster N, Tadesse AW, McQuaid CF, Gosce L, Abdurhman T, Assefa D, et al. Evaluating the equity
impact and cost-effectiveness of digital adherence technologies with differentiated care to
support tuberculosis treatment adherence in Ethiopia: protocol and analysis plan for the health
economics component of a cluster randomised trial. Trials. 2023;24: 292. doi:10.1186/s13063-
023-07289-x

15. Contoyannis P, Hurley J, Walli-Attaei M. When the technical is also normative: a critical
assessment of measuring health inequalities using the concentration index-based indices. Popul
Health Metrics. 2022;20: 21. doi:10.1186/s12963-022-00299-y

16. Erreygers G, Van Ourti T. Measuring socioeconomic inequality in health, health care and health
financing by means of rank-dependent indices: A recipe for good practice. Journal of Health
Economics. 2011;30: 685–694. doi:10.1016/j.jhealeco.2011.04.004

17. Minwyelet Terefe M, Eyob H, Letta T, Nadew E, Assefa T, Hailu B, et al. Efficiency of TB service
provision in the public and private health sectors in Ethiopia. int j tuberc lung dis. 2022;26:
1128–1136. doi:10.5588/ijtld.21.0481

18. Vos T, Lim SS, Abbafati C, Abbas KM, Abbasi M, Abbasifard M. Global burden of 369 Diseases and
Injuries in 204 Countries and Territories, 1990-2019: a Systematic Analysis for the Global Burden
of Disease Study 2019. Lancet. 2020;396: 1204–1222. doi:10.1016/s0140-6736(20)

19. Ochalek J, Lomas J, Claxton K. Cost per DALY averted thresholds for loaw- and middle-income
countries: evidence from cross country data.

20. Dawkins BR, Mirelman AJ, Asaria M, Johansson KA, Cookson RA. Distributional cost-effectiveness
analysis in low- and middle-income countries: illustrative example of rotavirus vaccination in
Ethiopia. Health Policy and Planning. 2018;33: 456–463. doi:10.1093/heapol/czx175

21. Charalambous S, Maraba N, Jennings L, Rabothata I, Cogill D, Mukora R, et al. Treatment
Adherence and Clinical Outcomes Amongst Drug-Susceptible TB Persons Using Medication
Monitor and Differentiated Care Approach in South Africa: Results from a Cluster Randomized
Trial. SSRN; 2024. doi:10.2139/ssrn.4756469

22. Manyazewal T, Woldeamanuel Y, Fekadu A, Holland DP, Marconi VC. Effect of Digital Medication
Event Reminder and Monitor-Observed Therapy vs Standard Directly Observed Therapy on
Health-Related Quality of Life and Catastrophic Costs in Patients With Tuberculosis: A Secondary
Analysis of a Randomized Clinical Trial. JAMA Netw Open. 2022;5: e2230509.
doi:10.1001/jamanetworkopen.2022.30509

23. Belachew M, Sahile M, Gadissa D, Abdurhman T, Bedru A, McQuaid CF, et al. Effect of
implementing digital adherence technologies on the use of health care providers' time and the
human resource cost of tuberculosis treatment adherence support in Ethiopia. Infectious
Diseases (except HIV/AIDS); 2024 Mar. doi:10.1101/2024.03.14.24304331

24. Whitehead M. The concepts and principles of equity and health. International journal of health
services. 1992.

25. Khan MS, Rego S, Rajal JB, Bond V, Fatima RK, Isani AK, et al. Mitigating the impact of COVID-19
on tuberculosis and HIV services: A cross-sectional survey of 669 health professionals in 64 low
and middle-income countries. Di Gennaro F, editor. PLoS ONE. 2021;16: e0244936.
doi:10.1371/journal.pone.0244936

26. Goscé L, Tadesse AW, Foster N, Van Kalmthout K, Van Rest J, van der Wal J, et al. Modelling the
278   epidemiological and economic impact of digital adherence technologies with differentiated care
279   for tuberculosis treatment in Ethiopia. TBC. 2024.

280   27. Menzies NA, Quaife M, Allwood BW, Byrne AL, Coussens AK, Harries AD, et al. Lifetime burden of
281   disease due to incident tuberculosis: a global reappraisal including post-tuberculosis sequelae.
282   The Lancet Global Health. 2021;9: e1679–e1687. doi:10.1016/S2214-109X(21)00367-3

283   28. Assefa DG, Dememew ZG, Zeleke ED, Manyazewal T, Bedru A. Financial burden of tuberculosis
284   diagnosis and treatment for patients in Ethiopia: a systematic review and meta-analysis. BMC
285   Public Health. 2024;24: 260. doi:10.1186/s12889-024-17713-9

286   29. Cookson R, Griffin S, Norheim OF, Culyer AJ. Distributional cost-effectiveness analysis:
287   quantifying health equity impacts and trade-offs. Oxford; 2021.

288   30. Verguet S, Laxminarayan R, Jamison DT. Universal Public Finance of Tuberculosis Treatment in
289   India: An Extended Cost-Effectiveness Analysis. Health Econ. 2015;24: 318–332.
290   doi:10.1002/hec.3019

Illness concentration index of DALYs ranked by SEP

Illness concentration index

(95% CI)

SOC   -0.06 (-0.24, 0.11)

Labels   -0.02 (-0.17, 0.14)

Pillbox   -0.02 (-0.16, 0.12)

Illness concentration index of visits ranked by SEP

Illness concentration index

(95% CI)

SOC   -0.07 (-0.10, -0.05)

Labels   0.01 (-0.00, 0.02)

Pillbox   -0.08 (-0.09, -0.06)

Between group comparison p-value<0.001

Illness concentration index of patient costs ranked by SEP

Illness concentration index

(95% CI)

SOC   -0.05 (-0.07, -0.02)

Labels   0.03 (0.01, 0.05)

Pillbox   0.01 (-0.01, 0.02)

Between group comparison p-value<0.001

Illness concentration index of provider costs ranked by SEP

Illness concentration index

(95% CI)

SOC   0.13 (0.08, 0.18)

Labels   0.07 (0.05, 0.08)

Pillbox   0.07 (0.04, 0.10)

Between group comparison p-value=0.225

)ALYs); (B) Nu   )sts in

2025 USD.

Figure 2. Cost-effectiveness planes.