Multivariable Mendelian randomization to disentangle the alcohol harm paradox.

Gemma Sawyer[1*], Hannah Sallis[1,2], Marcus Munafò[1,3], Liam Mahedy[1], Jasmine Khouja[1,3]

[1] MRC Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School,

University of Bristol, Bristol, United Kingdom

[2] Centre for Academic Mental Health, Population Health Sciences, Bristol Medical School,

University of Bristol, Bristol, United Kingdom

[3] School of Psychological Sciences, University of Bristol, Bristol, United Kingdom

*Corresponding author

Email: gemma.sawyer@bristol.ac.uk

Abstract

The alcohol harm paradox, whereby low socioeconomic position (SEP) groups experience greater

alcohol-related harms despite reporting lower alcohol consumption, is yet to be fully understood

through observational studies because key drivers are correlated and share similar confounding

structures. Multivariable Mendelian randomization (MVMR) were conducted to estimate the direct

causal effect of number of drinks per week (DPW) and years of schooling (YOS) on multiple health

outcomes. Previously published genome-wide association summary (GWAS) statistics for DPW and

YOS were utilised, and summary statistics were generated from individual-level data from UK

Biobank (N = 462,818) for all health outcomes. Inverse variance weighted analyses demonstrated

evidence for direct effects of DPW and YOS on liver diseases, mental and behavioural disorders due

to alcohol, and stroke, indicating that increasing alcohol consumption increased the likelihood of

outcomes whereas increasing years of education decreased their likelihood. There was also evidence

for a direct effect of DPW on depression, anxiety, influenza/pneumonia, and heart disease. In

contrast, there was evidence of a total, but not direct, effect of DPW on depression,

influenza/pneumonia, epilepsy, and injuries when accounting for YOS. Although caution is required

when interpreting these results due to weak instruments for alcohol, these results provide some

evidence that the alcohol harm paradox is partially due to the protective effect of additional years of

education, resulting in a reduced likelihood of higher SEP groups developing many alcohol-related

outcomes. Replication with strong instruments would be necessary to draw causal inferences.

Author Summary

Individuals from lower socioeconomic backgrounds tend to disproportionately experience

alcohol-related physical and mental ill health, despite reporting lower alcohol consumption

than those from higher socioeconomic backgrounds. This is known as the alcohol harm

paradox. One key difficulty in understanding this paradox is due to methodological

limitations establishing the relative contribution of multiple complex social behaviours. In

this study, we used genetic variants associated with alcohol consumption and years of

education to explore their direct effects on multiple health outcomes. The findings indicate

that higher alcohol consumption and fewer years of education may potentially increase the

likelihood of developing various health conditions, including liver diseases, mental and

behavioural disorders due to alcohol, and stroke. This may suggest that the alcohol harm

paradox is due to the protective effect of additional years of education amongst those from

higher socioeconomic backgrounds, reducing their likelihood of developing the health

conditions. However, these findings are exploratory and limited by various methodological

issues, suggesting these findings should be interpreted with caution. Replication and further

studies are needed.

Introduction

Alcohol-related harms are common and represent a large burden of morbidity and mortality

worldwide, contributing to 5.3% of all deaths and 5.1% of global burden of disease [1].

Moreover, such alcohol-related harms tend to be disproportionately experienced by

individuals of low socioeconomic position (SEP) despite reporting lower alcohol consumption

compared to high SEP individuals, known as the alcohol harm paradox [2–5]. Despite evidence

for this paradox, there is limited understanding of its origins.

There have been multiple explanations proposed for the alcohol harm paradox. One

explanation is that, whilst low SEP individuals report consuming less alcohol, other patterns

of drinking behaviour are more common. For example, low SEP individuals are more likely to

be in the extreme ends of drinking consumption, with a greater proportion abstaining or

drinking very lightly compared to high SEP groups, but also being more likely to drink

excessively or binge drink [6]. The possibility that drinking patterns may account for the

alcohol harm paradox has been supported by research demonstrating a greater

socioeconomic gradient for acute health outcomes which are wholly attributable to alcohol,

compared to chronic conditions which are associated with a wide range of behaviours [7]. An

alternative explanation is that individuals of low SEP may be more likely to engage in other

unhealthy behaviours, such as tobacco use, which are associated with multiple adverse health

outcomes [8]. Observational studies have supported this explanation, providing evidence that

individuals of lower SEP are increasingly likely to smoke [3]. Furthermore, adjusting for

tobacco smoking reduces the association between low SEP and hospital admissions [9], while

causal mediation analysis demonstrated that 18% of the total effect of income on alcohol-

related mortality was due to indirect effects via smoking and BMI [10]. Whilst these proposed

explanations have received some support from observational studies, they are limited in their

ability to draw causal inferences and are yet to fully account for the alcohol harm paradox.

The primary complication with disentangling the alcohol harm paradox is that a combination

of highly correlated causes, including alcohol consumption, SEP, and smoking, could be

important drivers. Therefore, it is necessary to establish the relative causal contribution of

alcohol consumption alongside these other factors on the likelihood of developing multiple

adverse health outcomes with novel causal inference methods, such as Mendelian

randomization (MR). MR relies on Mendel's law of random assortment, using genetic variants

associated with the exposure as instrumental variables to assess the total causal effect of the

exposure on the outcome [11]. Multivariable MR (MVMR) builds upon this method, using

genetic variants associated with multiple exposures to estimate the direct causal effect of

each exposure on the outcome, accounting for the effect of the other exposures [12]. The

validity of causal inferences drawn using MVMR relies on three key assumptions [13]: (1) each

exposure is robustly predicted by the genetic instruments; (2) there are no confounders of

the outcome and any of the genetic instruments; and (3) the genetic instruments only impact

the outcome through at least one of the exposures.

The current study will apply MVMR to the alcohol harm paradox to examine the direct causal

effect of alcohol consumption on adverse health outcomes whilst accounting for education

(as a proxy for SEP) and vice versa. We will first conduct MR to examine the total causal effect

of alcohol and education separately on adverse health, followed by MVMR to examine their

direct causal effects. Comparison of the MR and MVMR results will provide insights into the

relative contribution of alcohol and education. For example, if there appears to be a causal

effect of alcohol in both the MR and MVMR, this suggests that alcohol consumption impacts

the health outcome even when the impact of education is considered. However, if a causal

effect of alcohol in the MR is attenuated in the MVMR, this may indicate that the impact of

education was driving the evidence of a causal effect of alcohol in the MR.

Materials and Methods

This method was outlined in our pre-registered analysis plan

(https://doi.org/10.17605/OSF.IO/5G8W4) and the methods presented here are consistent with

this unless otherwise specified.

Exposures

Summary-level data reported by the two Genome-Wide Association Studies (GWAS) were

utilised to identify suitable genetic instruments associated with the relevant exposure

phenotypes: drinks per week (DPW) and years of schooling (YOS). In contrast with our pre-

registration, the main analysis reported here focuses solely on these two exposures and

additional supplementary analysis included cigarettes per day (CPD) as a third exposure (see

Supplementary Methods for details). We are focusing on the two exposures of DPW and YOS

because alcohol and smoking are highly correlated and both suffer from weak instrument bias

in the three-exposure MVMR, making it difficult to draw conclusions from such findings. To

aid interpretation, we simplified the analysis to a two-exposure model to explore whether

SEP plays a causal role in the alcohol harm paradox. Consequently, the results of this study

should be considered exploratory.

GWAS & Sequencing Consortium of Alcohol and Nicotine Use (GSCAN)

Liu and colleagues [14] report summary-level statistics from a GWAS meta-analysis of alcohol

use based on data from participants of European ancestry from 29 cohorts (N=941,280).

Alcohol use was defined as the reported number of drinks (combined across all alcohol types)

consumed per week (DPW) amongst active drinkers. Where individual studies recorded

categorical bins of drinks per week (e.g., 1-4, 5-9), the midpoint value was taken. Analyses

adjusted for age, sex, age and sex interaction, and 10 genetic principal components.

Independent variants were reported to be associated with DPW if they met the standard

significance threshold (p < 5x10^8). The associated SNPs explained 2.5% of the variance in DPW.

We utilised the publicly available summary statistics from this GWAS, excluding UK Biobank

(UKB), to minimise sample overlap, and 23andMe, due to data sharing restrictions

(N=226,223). Smoking (cigarettes per day) was also measured in this GWAS, as described in

the Supplementary Material (p. 1).

Social Science Genetic Association Consortium (SSGAC)

Okbay and colleagues [15] report summary-level statistics from a GWAS meta-analysis of

education, defined as years of schooling (YOS), based on data from participants of European

ancestry from 65 cohorts (N=293,723). Reported number of years of schooling was utilised

where this was available. Where individual studies reported qualifications, the International

Standard Classification of Education (ISCED; 1997) [16] was used to map each major

qualification that could be attained in each country into one of seven categories: pre-primary

education, primary education, lower secondary education, upper secondary education, post-

secondary non-tertiary education, first-stage tertiary education, and second-stage tertiary

education. This was subsequently utilised to impute a YOS value for each category. Analyses

adjusted for age, sex, age and sex interaction, ten genetic principal components, and cohort-

specific controls. Variants were reported to be associated with YOS if they met the genome-

wide significance threshold (p <5x10^8) and were independent.

Health outcomes

Individual-level data, including genetic and outcome data for all individual participants, was

obtained from UKB to generate summary statistics for multiple health outcomes.

UK Biobank

UKB is a population-based cohort, which recruited 500,000 people aged between 40-69 years

in 2006-2010. With their consent, they provided detailed information about their lifestyle,

physical measures and had blood, urine and saliva samples collected and stored for future

analysis. Using this individual-level data, we generated summary statistics by regressing each

available SNP on each health outcome through the MRC-IEU UKB GWAS pipeline [17] using a

BOLT-LMM (linear mixed model) to account for relatedness and population stratification.

Furthermore, we additionally adjusted for age, sex, and genotyping chip array. For the main

two-exposure MVMR analysis, this was conducted for the overall UKB sample; however, it

was also conducted in four smaller samples stratified by smoking status for comparison with

the supplementary three-exposure MVMR: ever smokers (defined as those reporting smoking

2 or more cigarettes), never smokers (defined as those reporting smoking 1 or less cigarettes),

current smokers, and former smokers (Figure 1).

Figure 1. Flowchart demonstrating the source of exposure and outcome data and the inclusion and exclusion of genetic variants (single

nucleotide polymorphisms; SNPs) in this analysis. Details of the three-exposure MVMR (grey dashed lines) are in the Supplementary Materials.

Our pre-registered analysis specified 15 outcomes wholly or partially attributable to alcohol,

based upon ICD-10 summary diagnoses classification codes (UKB field ID 41270; see

Supplementary Table 1). However, once all the necessary exclusions had been made, four

outcomes (alcoholic cardiomyopathy, alcoholic myopathy, alcoholic polyneuropathy, and

alcoholic poisoning) had a case number below 50 or less than 0.1% in at least one of the

stratified datasets and were therefore removed from the analysis. The remaining outcomes

included alcoholic liver disease, other liver diseases, stroke, ischaemic heart disease (IHD),

influenza/pneumonia, viral hepatitis, epilepsy, injuries, mental and behavioural disorders due

to alcohol, depression, and anxiety.

Analysis

Analyses were conducted in R version 4.1.2 [18], using the *Mendelian Randomization*[19],

TwoSampleMR [20]*, MVMR* [21] and *MR-PRESSO* [22] R packages.

Selection of genetic variants

Single nucleotide polymorphisms (SNPs) were selected for inclusion in the analysis if they

were reported to be associated with the exposure phenotype of interest (alcohol use or

education) and (conditionally) independent of any other SNP associations at the genome-

wide significance level ($p < 5\times10^{-8}$) in the relevant GWAS. These SNPs also needed to be

available in UKB, resulting in 96 DPW and 74 YOS SNPs in the univariable MR. For the MVMR

analyses, where a SNP from GSCAN (DPW) was not available in SSGAC (YOS), we selected

proxy SNPs with a minimum linkage disequilibrium (LD) R^2 of 0.8. Moreover, an additional

clumping stage was conducted in the MVMR to ensure all exposure SNPs were independent

(LD R^2 < 0.1, clumping window > 500 kb), resulting in 71 DPW and 62 YOS SNPs in the MVMR

(Figure 1; see Supplementary Table 2 for details).

We assessed instrument strength using a conditional F-statistic and instrument validity using

Cochran's Q-statistic. In line with conventional guidelines, the F-statistic should be greater

than 10 and the Q-statistic should be less than the number of SNPs [23].

Univariable MR

We estimated the total causal effects of alcohol use and education on all 11 health outcomes

using two-sample, summary level MR. The main analysis was conducted using inverse

variance weighted (IVW) regression [24], however; we also conducted sensitivity analyses

using three complementary MR methods MR-Egger [25], weighted median estimation [26],

and weighted mode estimation [27]. These methods hold different assumptions regarding

pleiotropy and therefore consistent results across the methods provide stronger evidence

that the observed effects are causal.

Multivariable MR

In MVMR analyses, we estimated the direct causal effect of alcohol use, education, and

tobacco use on all health outcomes using two sample, summary level MVMR with two

complimentary methods – MVMR-IVW and MVMR-Egger. Whilst we intended to conduct

MVMR-PRESSO in our pre-registered analysis plan, we had to remove this method due to

issues with the R package.

Results

The MR-IVW, MVMR-IVW, MR-Egger and MVMR-Egger results exploring the total and direct effects of alcohol and education are displayed in Figure 2. The complete results of the MVMR analyses exploring the direct effects of (1) alcohol and education (n = 133 SNPs) and (2) alcohol, education, and smoking (n = 171 SNPs) on health outcomes are displayed in Supplementary Tables 3-13. The results are presented in odds ratios (ORs) per standard deviation (SD) increase in either drinks per week or years of schooling.

Instrument strength and heterogeneity

Instrument strength was calculated using the conditional F-statistic. In the MR, the F-statistics suggest that the SNPs included in the analysis are strong instruments for alcohol (F = 14.87) and education (F = 35.32). The conditional F-statistics in the two exposure MVMR indicate that the included SNPs are strongly associated with education whilst accounting for the effects of alcohol (F = 17.42), but not strongly associated with alcohol while accounting for education (F = 6.32).

The Cochran's Q statistics were greater than the number of SNPs included in most of the models, indicating heterogeneity. Therefore, the results also include MVMR-Egger, which provides estimates that are robust to directional pleiotropy under the assumption that pleiotropy is uncorrelated with the strength of the association between the SNP and exposure. However, MVMR-Egger analyses have reduced statistical power compared with IVW analyses and therefore we must consider the results from both analysis when drawing conclusions. Where Egger results indicate a similar effect size but with larger confidence intervals which cross the null, we interpret this as evidence in support of the IVW analysis.

Figure 2. Forest plots displaying the effect of alcohol (drinks per week) and education (years

of schooling) on eight health outcomes using Mendelian randomization (MR) and

multivariable MR (MVMR) analysis: inverse variance weighted (IVW) and Egger regression.

Alcoholic liver disease

There was evidence for a total (OR = 36.57, 95% CI 16.46 to 81.24) and direct (OR = 50.19,

95% CI 19.35 to 130.21) effect of alcohol on alcoholic liver disease when accounting for

education. There was also evidence that education decreased the odds of alcoholic liver

diseases in the univariable (OR = 0.44, 95% CI 0.23 to 0.86) and multivariable (OR = 0.27, 95%

CI 0.14 to 0.53) analysis when accounting for alcohol (Figure 2; Supplementary Table 3). Whilst

the MR-Egger and MVMR-Egger results were consistent with the IVW results for alcohol, for

education, the effect estimates were similar but the wider confidence intervals overlapped

the null.

Mental and behavioural disorders due to alcohol

There was evidence for a total (OR = 15.88, 95% CI 10.08 to 25.01) and direct (OR = 12.89,

95% CI 7.46 to 22.27) effect of alcohol on mental and behavioural disorders due to alcohol,

as well as a total (OR = 0.60, 95% CI 0.44 to 0.83) and direct (OR = 0.51, 95% CI 0.35 to 0.75)

effect of education on mental and behavioural disorders due to alcohol (Figure 2;

Supplementary Table 6). The MR- and MVMR-Egger findings were consistent with the IVW

findings for alcohol and showed similar effect estimates for education, although the

confidence intervals crossed the null.

Other liver diseases

There was evidence for a total (OR = 1.80, 95% CI 1.21 to 2.70) and direct (OR = 1.82, 95% CI

1.12 to 2.94) effect of alcohol on other liver diseases, as well as evidence for a total (OR =

0.47, 95% CI 0.35 to 0.62) and direct (Or = 0.42, 95% CI 0.30 to 0.58) effect of education on

other liver diseases (Figure 2; Supplementary Table 4). The MR- and MVMR-Egger results

calculated similar effect estimates, although the wider confidence intervals crossed the null

for alcohol and the total effect of education.

Stroke

There was evidence for a total (OR = 1.94, 95% CI 1.46 to 2.58) and direct (OR = 1.94, 95% CI

1.30 to 2.89) effect of alcohol on stroke, when accounting for education. Also, whilst there

was no clear evidence for a total effect of education on stroke, there was some evidence of a

direct effect in the MVMR when accounting for alcohol (OR = 0.73, 95% CI 0.55 to 0.97) (Figure

2; Supplementary Table 5). The MR- and MVMR-Egger findings were similar to the IVW

findings in terms of the effect estimates. However, the confidence intervals were wider and

therefore crossed the null for education and the direct effect of alcohol.

Injuries

There was evidence for a total (OR = 1.47, 95% CI 1.24 to 1.73) and direct (1.29, 95% CI 1.03

to 1.60) effect of alcohol on injuries. However, there was no evidence for a total (OR = 0.98,

95% CI 0.85 to 1.14) or direct (OR = 0.87, 95% CI 0.75 to 1.02) effect of education (Figure 2;

Supplementary Table 7). The MR- and MVMR-Egger results supported the IVW results for

education and the total effect of alcohol; however, the confidence intervals crossed the null

for the direct effect of alcohol.

Depression

In the univariable MR, there was evidence for a total effect of alcohol on depression (OR =

1.33, 95% CI 1.09 to 1.64); however, there was no evidence for a direct effect when

accounting for education (OR = 1.11, 95% CI 0.84 to 1.48). There was evidence that education

decreased the odds of depression in the univariable (OR = 0.59, 95% CI 0.48 to 0.73) and

multivariable analyses when accounting for alcohol (OR = 0.56, 95% CI 0.45 to 0.68) (Figure 2;

Supplementary Table 8). The MR- and MVMR-Egger results supported the IVW findings of a

total, but not direct, effect of alcohol (although the wider confidence intervals crossed the

null), as well as a direct effect of education. The MR-Egger did not however support IVW

evidence for a total effect of education.

Epilepsy

Whilst there was evidence for a total effect of alcohol on epilepsy (OR = 1.91, 95% CI 1.33 to

2.75), there was no evidence for a direct effect when accounting for education (OR = 1.17,

95% CI 0.72 to 1.89). Moreover, there was no evidence for a total (OR = 0.83, 95% CI 0.60 to

1.15) or direct (OR = 0.82, 95% CI 0.58 to 1.14) effect of education on epilepsy (Figure 2;

Supplementary Table 9). The MR-Egger results supported a total effect of alcohol and also

found some evidence for a direct effect, which was not indicated in the IVW findings. The MR-

and MVMR-Egger effect estimates for education were in the opposite direction to the IVW

results, although all confidence intervals crossed the null.

Influenza and pneumonia

There was evidence for a total (OR = 1.32, 95% CI 1.03 to 1.68) but not a direct effect of alcohol

on influenza/pneumonia when accounting for education (OR = 1.28, 95% CI 0.94 to 1.73).

There was evidence that education decreased the odds of influenza/pneumonia in the

univariable (OR = 0.63, 95% CI 0.53 to 0.76) and multivariable (OR = 0.60, 95% CI 0.48 to 0.74)

analysis (Figure 2; Supplementary Table 10). The MR- and MVMR-Egger results were primarily

consistent with the IVW results, supporting a total and direct effect of education and no direct

effect of alcohol. Whilst the effect estimate for the total effect of alcohol was similar, the MR-

Egger confidence interval crossed the null.

Other outcomes

There was no evidence for a total or direct effect of alcohol on IHD (Supplementary Table 11),

viral hepatitis (Supplementary Table 12), or anxiety (Supplementary Table 13). There was

evidence for a total (OR = 0.61, 95% CI 0.53 to 0.70) and direct (OR = 0.62, 95% CI 0.52 to 0.74)

effect of education on IHD when accounting for alcohol. There was also evidence for a total

(OR = 0.67, 95% CI 0.52 to 0.86) and direct (OR = 0.62, 95% CI 0.48 to 0.80) effect of education

on anxiety. There was no evidence for a total or direct effect of education on viral hepatitis.

Overall, the MR- and MVMR-Egger results were mostly similar to the IVW results for IHD,

anxiety, and viral hepatitis. The only differences were that the MR-Egger did not support a

total effect of education on IHD or anxiety.

Discussion

The current study has found evidence for direct effects of alcohol and education on alcoholic and other liver diseases, mental and behavioural disorders due to alcohol, and stroke, indicating that alcohol use increases the likelihood of such outcomes whereas years of education decreases their likelihood. There was also evidence of a direct effect of education on depression, anxiety, influenza/pneumonia, and IHD. In contrast, there was evidence of a total, but not direct effect of alcohol on depression, influenza/pneumonia, epilepsy, and injuries when accounting for education. These results are consistent with previous findings from observational studies suggesting that both alcohol consumption and SEP may impact the likelihood of developing numerous health conditions and that the alcohol harm paradox is due to these multiplicative risk factors [3,4]. However, when interpreting these results, it is important to be cautious and consider the strength of instruments. The conditional F-statistics indicate that the instruments were not strongly associated with alcohol in the MVMR, despite strong associations in the MR. Therefore, we cannot be as confident or draw causal inferences from our estimates of direct effects; instead, our findings may be indicative of possible causal effects which would need to be replicated with strong instruments to be confident. In addition, we can only consider these results as exploratory as they were not all specified in our pre-registered analysis plan.

For multiple outcomes, there was little difference between the total and direct effects of both alcohol and education, including alcoholic liver disease, other liver diseases, and mental and behavioural disorders due to alcohol. These results suggest that alcohol use increases the likelihood of such outcomes when accounting for education, whereas years of education decreases their likelihood when accounting for alcohol use. Therefore, for these outcomes, it

is possible that greater alcohol consumption and less education both causally contribute to

their likelihood and, as such, the alcohol harm paradox may be due to a combination of these

risk factors in low SEP groups, whereas high SEP groups are somewhat protected by additional

years of education. Moreover, there was evidence for a total and direct effect of alcohol on

stroke but only evidence for a direct, but not a total, effect of education. Again, these results

indicate that alcohol use increased the likelihood of stroke, even when accounting for

education, and education decreases the likelihood of stroke. However, it may also suggest

that failing to account for alcohol in the MR masked the effect of education on stroke.

For depression and influenza/pneumonia, there was little difference between the total and

direct effects of education; however, the evidence for a total effect of alcohol was attenuated

when accounting for education. This suggests that years of education decreases the likelihood

of developing depression and influenza/pneumonia, whereas alcohol use does not directly

influence the likelihood of such outcomes when education is considered alongside. Similarly,

for injuries and epilepsy, there was evidence for a total effect of alcohol which was attenuated

when education was considered, perhaps suggesting that alcohol does not directly influence

these outcomes. However, there was no evidence of a total or direct effect of education on

injuries and epilepsy these outcomes, indicating that education also does not influence these

outcomes. It is plausible that the attenuation of the total effect of alcohol is the result of

reduced instrument strength in the MVMR compared with the MR as opposed to education

driving the total effect of alcohol identified in the MR. Replication of our results with stronger

instruments is necessary to clarify such findings. The current results also support a total and

direct effect of education only on anxiety and IHD, whereby years of education decreased the

likelihood of developing these outcomes when accounting for alcohol. There was no evidence

for a total or direct effect of alcohol on either anxiety or IHD, nor a total or direct effect of

alcohol or education on viral hepatitis.

Strengths and Limitations

This is the first study, to our knowledge, to apply MVMR, a novel causal inference method, to

understand the direct causal effects of alcohol and education on multiple alcohol-related

conditions to enhance understanding of the alcohol harm paradox. However, this study also

has multiple limitations. As previously mentioned, the conditional F-statistics indicate that

alcohol suffers from weak instruments in the MVMR, which may possibly bias our results.

Heterogeneity is indicated by the Cochran's Q statistics and, even though we applied MR-

Egger to address this, heterogeneity may remain. This indicates that the analyses may also be

biased by invalid instruments. Directional pleiotropy is similarly likely to persist in our analyses

despite applying pleiotropy-robust methods as indicated by the MR-Egger intercepts, which

are not null in most of the models. This means that the genetic instruments may influence the

adverse health outcomes within this study via pathways that do not include alcohol or

education and therefore may not tell us about the causal effect of alcohol or education on

such outcomes. Moreover, DPW is a somewhat limited phenotype to reflect alcohol. This is

because many factors influence the effects of alcohol beyond number of drinks, including

binge patterns and alcohol types. Similarly, education is one factor that contributes to an

individual's SEP; factors such as income, neighbourhood, and occupation are also important

and may show different relationships. Therefore, focusing purely on DPW and YOS may be

restricting our understanding. Combined with the exploratory nature of these analyses, these

limitations suggest that caution is highly necessary in interpreting the current findings.

Future Research and Implications

Due to limitations within the study regarding weak instruments and possible pleiotropy, the

current study should be replicated with newly available GWAS data [28] and thus potentially

valid, stronger instruments can be utilised in these analyses. Replication will help confirm

whether the results presented in this study reflect the true causal effects of alcohol and

education or are due to methodological limitations. If the current findings were supported,

this would suggest that the alcohol harm paradox may be due to the protective effect of years

of schooling on many alcohol-related conditions resulting in a reduced likelihood of higher

SEP groups developing such outcomes. Future research may therefore benefit from exploring

the mechanisms through which education impacts such health outcomes to inform

preventative policy. Beyond this, it may be useful for future work to explore alternative

exposures, such as type of alcohol and income, and outcomes, including cancers to broaden

understanding.

Conclusion

This study found evidence for a direct effect of alcohol on alcoholic and other liver diseases,

mental and behavioural disorders due to alcohol, and stroke. There was also evidence for a

total effect of alcohol on depression, epilepsy, influenza/pneumonia, and injuries, which were

attenuated when education was accounted for. Despite weak instrument bias, such results

cautiously indicate that many partially alcohol-attributable conditions are causally impacted

by years of education, which we use here as a proxy for SEP. Replication with strong

instruments is necessary however to confirm these findings and future research may

consequently endeavour to explore the causal effect of education on additional partially

alcohol-attributable conditions and identify the mechanism through which further years of

education protects against these conditions.

Acknowledgements

This research has been conducted using the UK Biobank Resource under application number

9142. This work uses data provided by patients and collected by the NHS as part of their care

and support.

Funding

GS, HS, MM, LM, and JK are all members of the Medical Research Council (MRC) Integrative

Epidemiology Unit at the University of Bristol. This publication is the work of the authors and

they will serve as guarantors for the contents of this paper. GS is supported by a Wellcome

Trust PhD studentship in Molecular, Genetic and Lifecourse Epidemiology (ref:

218495/Z/19/Z). For the purpose of Open Access, the author has applied a CC BY public

copyright licence to any Author Accepted Manuscript version arising from this submission.

References

References

1. World Health Organization. Alcohol. 2018 [cited 9 Mar 2022]. Available:
https://www.who.int/news-room/fact-sheets/detail/alcohol
2. Smith K, Foster J. Alcohol, Health Inequalities and the Harm Paradox: Why some
groups face greater problems despite consuming less alcohol. 2014. Available:
https://www.alcoholpolicy.net/2014/11/alcohol-health-inequalities-and-the-harm-
paradox-ias-report-reviews-higher-harms-in-deprived-groups.html
3. Bellis MA, Hughes K, Nicholls J, Sheron N, Gilmore I, Jones L. The alcohol harm
paradox: Using a national survey to explore how alcohol may disproportionately
impact health in deprived individuals. BMC Public Health. 2016;16.
doi:10.1186/s12889-016-2766-x
4. Beard E, Brown J, West R, Angus C, Brennan A, Holmes J, et al. Deconstructing the
Alcohol Harm Paradox: A population based survey of adults in England. PLoS One.
2016;11. doi:10.1371/journal.pone.0160666
5. Møller SP, Pisinger VSC, Christensen AI, Tolstrup JS. Socioeconomic position and
alcohol-related harm in Danish adolescents. J Epidemiol Community Health (1978).
2019;73: 839–845. doi:10.1136/jech-2018-211634
6. Lewer D, Meier P, Beard E, Boniface S, Kaner E. Unravelling the alcohol harm paradox:
A population-based study of social gradients across very heavy drinking thresholds.
BMC Public Health. 2016;16. doi:10.1186/s12889-016-3265-9
7. Sadler S, Angus C, Gavens L, Gillespie D, Holmes J, Hamilton J, et al. Understanding
the alcohol harm paradox: an analysis of sex- and condition-specific hospital
admissions by socio-economic group for alcohol-associated conditions in England.
Addiction. 2017;112: 808–817. doi:10.1111/add.13726
8. Reitsma MB, Fullman N, Ng M, Salama JS, Abajobir A, Abate KH, et al. Smoking
prevalence and attributable disease burden in 195 countries and territories, 1990-
2015: A systematic analysis from the global burden of disease study 2015. The Lancet.
2017;389: 1885–1906. doi:10.1016/S0140-6736(17)30819-X
9. Gartner A, Trefan L, Moore S, Akbari A, Paranjothy S, Farewell D. Drinking beer, wine
or spirits - Does it matter for inequalities in alcohol-related hospital admission? A
record-linked longitudinal study in Wales. BMC Public Health. 2019;19.
doi:10.1186/s12889-019-8015-3
10. Peña S, Mäkelä P, Laatikainen T, Härkänen T, Männistö S, Heliövaara M, et al. Joint
effects of alcohol use, smoking and body mass index as an explanation for the alcohol
harm paradox: causal mediation analysis of eight cohort studies. Addiction. 2021;116:
2220–2230. doi:10.1111/add.15395
11. Davey Smith G, Ebrahim S. "Mendelian randomization": Can genetic epidemiology
contribute to understanding environmental determinants of disease? Int J Epidemiol.
2003;32: 1–22. doi:10.1093/ije/dyg070
12. Sanderson E, Davey Smith G, Windmeijer F, Bowden J. An examination of
multivariable Mendelian randomization in the single-sample and two-sample
summary data settings. Int J Epidemiol. 2019;48: 713–727. doi:10.1093/ije/dyy262
13. Sanderson E, Glymour MM, Holmes M v., Kang H, Morrison J, Munafò MR, et al.
Mendelian randomization. Nature Reviews Methods Primers. 2022;2.
doi:10.1038/s43586-021-00092-5

14. Liu M, Jiang Y, Wedow R, Li Y, Brazel DM, Chen F, et al. Association studies of up to
1.2 million individuals yield new insights into the genetic etiology of tobacco and
alcohol use. Nature Genetics. Nature Publishing Group; 2019. pp. 237–244.
doi:10.1038/s41588-018-0307-5

15. Okbay A, Beauchamp JP, Fontana MA, Lee JJ, Pers TH, Rietveld CA, et al. Genome-
wide association study identifies 74 loci associated with educational attainment.
Nature. 2016;533: 539–542. doi:10.1038/nature17671

16. UNESCO. International Standard Classification of Education ISCED 1997. 2006.
Available: https://isced.uis.unesco.org/wp-
content/uploads/sites/15/2022/12/international-standard-classification-of-
education-1997-en_0.pdf

17. Mitchell R, Elsworth B, Raistrick C, Paternoster L, Hemani G, Gaunt T. MRC IEU UK
Biobank GWAS pipeline version 2. 2019.
doi:10.5523/bris.pnoat8cxo0u52p6ynfaekeigi

18. R Core Team. R: A language and environment for statistical computing. R Foundation
for Statistical Computing. 2021 [cited 18 Mar 2022]. Available: https://www.R-
project.org/

19. Yavorska OO, Burgess S. MendelianRandomization: An R package for performing
Mendelian randomization analyses using summarized data. Int J Epidemiol. 2017;46:
1734–1739. doi:10.1093/ije/dyx034

20. Hemani G, Zheng J, Elsworth B, Wade KH, Haberland V, Baird D, et al. The MR-Base
platform supports systematic causal inference across the human phenome. Elife.
2018;7. doi:10.7554/eLife.34408

21. Spiller W, Bowden J, Sanderson E. MVMR: MVMR. R package version 0.3. 2022.
Available: https://github.com/WSpiller/MVMR

22. Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in
causal relationships inferred from Mendelian randomization between complex traits
and diseases. Nat Genet. 2018;50: 693–698. doi:10.1038/s41588-018-0099-7

23. Sanderson E, Spiller W, Bowden J. Testing and correcting for weak and pleiotropic
instruments in two-sample multivariable Mendelian randomization. Stat Med.
2021;40: 5434–5452. doi:10.1002/sim.9133

24. Burgess S, Butterworth A, Thompson SG. Mendelian randomization analysis with
multiple genetic variants using summarized data. Genet Epidemiol. 2013;37: 658–
665. doi:10.1002/gepi.21758

25. Bowden J, Smith GD, Burgess S. Mendelian randomization with invalid instruments:
Effect estimation and bias detection through Egger regression. Int J Epidemiol.
2015;44: 512–525. doi:10.1093/ije/dyv080

26. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent Estimation in Mendelian
Randomization with Some Invalid Instruments Using a Weighted Median Estimator.
Genet Epidemiol. 2016;40: 304–314. doi:10.1002/gepi.21965

27. Hartwig FP, Smith GD, Bowden J. Robust inference in summary data Mendelian
randomization via the zero modal pleiotropy assumption. Int J Epidemiol. 2017;46:
1985–1998. doi:10.1093/ije/dyx102

28. Saunders GRB, Wang X, Chen F, Jang S-K, Liu M, Wang C, et al. Genetic diversity fuels
gene discovery for tobacco and alcohol use. Nature. 2022;612: 720–724.
doi:10.1038/s41586-022-05477-4